

Can novel methods be useful for pain assessment of castrated piglets?

C. Lonardi¹, A. Scollo¹, S. Normando², M. Brcic¹ and F. Gottardo^{1†}

¹Department of Animal Medicine, Production and Health, University of Padova, Viale dell'Università 16, 35020 Agripolis Legnaro (PD), Italy; ²Department of Comparative Biomedicine and Food Science, University of Padova, Viale dell'Università 16, 35020 Agripolis Legnaro (PD), Italy

(Received 10 January 2014; Accepted 18 November 2014; First published online 5 January 2015)

Given that surgical castration is a painful practice performed on millions of pigs every year, a need to identify novel reliable pain assessment tools exists in order to test anaesthetic and analgesic protocols that may reduce related pain. Two treatments were considered: handling (H) and surgical castration (C). Physiological (cortisol, lactate, glycaemia, rectal and eye temperature) and behavioural variables (latency to move after treatment and alterations in posture and walking) were analysed. Cortisol showed the greatest level in C piglets within 20 min after the surgical procedure and a positive correlation with glucose concentration. Eye temperature was higher in C piglets, and the same difference was detected for rectal temperature 3 h after castration. Behavioural parameters revealed that C piglets had longer latency to move and a higher percentage of them showed alterations in posture and walking. Results of this study showed that, in castrated piglets behavioural and physiological alterations occur mainly in the first 3 h from treatment. Latency to move, alterations in posture and walking, and eye temperature appear to give additional and useful information in piglet pain assessment. However, differently from the behavioural parameters considered, eye temperature involves several manipulations of the animals and a long process to acquire the data.

Keywords: animal welfare, piglets, castration, pain assessment, thermography

Implications

Surgical castration is a painful practice performed on millions of pigs every year and finding novel easily applicable pain assessment tools is necessary for the assessment of pain reduction protocols. Results of this study showed that in castrated piglets behavioural and physiological alterations occur mainly in the first 3 h from treatment. Latency to move, alterations in posture and walking, and IR thermography for the detection of eye temperature are promising tools for pain assessment in piglets. However, IR thermography needs additional handling of the animals during the process for temperature acquiring.

Introduction

Surgical castration in piglets is the most commonly performed management practice in order to avoid boar taint in swine meat. It is estimated that 100 million male pigs are castrated every year in the 25 EU member states (European Food Safety Authority, 2004). This practice is widely executed without pain

relief, even though it appears to cause distress regardless of piglet age (Carrol *et al.*, 2006; von Borell *et al.*, 2009). Moreover, it represents an animal welfare issue with an increasing negative public perception owing to animal suffering (Guatteo *et al.* 2012; Sutherland *et al.*, 2012); hence, the European Declaration on alternatives to surgical castration of pigs (EU Commission, 2010) exhorts to abandon the practice by 2018. However, the same document recognises the unavoidability of castration in case of pig meat produced for traditional products, in order to meet their current quality standards. For this reason pigs reared under protocols for 'traditional guaranteed specialties' or for 'geographical indications' (Protected Geographical Indication or Protected Designation of Origin (PDO)) should be therefore submitted to surgical castration using analgesia and/or anaesthesia. The waiver include the Italian heavy pig production for the typical cured ham in which castration is unavoidable considering that pigs are slaughtered at around 160 kg and 9 months of age, when the sexual maturity has been reached and might be responsible for pronounced boar taint, but an appropriate protocol for pain reduction during castration is required.

Pain is a complex and individual sensation, which means that it is difficult to measure and compare among animals

† E-mail: flaviana.gottardo@unipd.it

(Sneddon and Gentle, 2000) and handling could also interfere with the behavioural and physiological responses of young pigs (Prunier *et al.*, 2005). During the last years, methods to reduce pain with anaesthesia and/or analgesia were investigated (Sutherland *et al.*, 2010; Hansson *et al.*, 2011; Kluivers-Poodt *et al.*, 2012). In order to monitor suffering after mutilation, several pain indicators are commonly considered such as physiological (including a clinical and a neuroendocrine approach) and behavioural parameters (Weary *et al.*, 2006; Sutherland *et al.*, 2012). Within physiological parameters, clinical indicators such as rectal temperature, heart rate, respiratory rate and blood pressure are feasible to measure in restrained awake piglets (White *et al.*, 1995), while capillary perfusion and pupillary dilatation are usually assessed during general anaesthesia (Hodgson, 2007) or sedation (Axiak *et al.*, 2007). Regarding neuroendocrine indicators, surgical castration has shown to cause the activation of the hypothalamus–pituitary–adrenal and the sympathetic axes (Prunier *et al.*, 2005). This leads to several consequences such as the increase of ACTH and plasma cortisol up to 60 min after castration (Marchant-Forde *et al.*, 2009; Keita *et al.*, 2010; Sutherland *et al.*, 2012) with a possible second peak 3 h later (Moya *et al.*, 2008), the increase of lactate owing to glycogen mobilisation (Prunier *et al.*, 2005), and the rapid and transient increase of epinephrine followed by a longer lasting increase in plasma noradrenaline (Prunier *et al.*, 2002; Mühlbauer *et al.*, 2010). Moreover, variations of sympathetic and parasympathetic systems seem to alter eye temperature in animal subjected to painful practice, as recently described in cattle by Stewart *et al.* (2010a and 2010b). Surgical castration also causes behavioural changes (Taylor *et al.*, 2001; Hay *et al.*, 2003). Changes in behaviour are considered relevant parameters to assess the overall pain and discomfort following this practice and are studied with methodologies such as scan or focal sampling (Hay *et al.*, 2003; Moya *et al.*, 2008; Sutherland *et al.*, 2012). Alteration of latency to move is considered to indicate distress in pigs (Chaloupková *et al.*, 2007). Increased restlessness in animals suffering from pain could be interpreted as an adaptation attempt to stop other animals from inflicting more pain (Hay *et al.*, 2003). Moreover, stiffness can be considered as a protective mechanism, allowing the animals to avoid or reduce the stimulation of the painful tissues after castration (Molony and Kent, 1997; Mellor *et al.*, 2000). Moreover, Keita *et al.* (2010) observed that an animal in pain assumes a different standing position and changes walking behaviour in order to relieve pain after castration.

The present study aimed to test if some innovative parameters such as latency to move, differences in postures and walking, and eye temperature are useful for detection of pain in piglets in response to surgical castration. These parameters have been compared with those generally used as indicators of pain. Considering the pressure of the European Declaration (EU Commission, 2010), which is requiring research on alternatives to surgical castration in piglets used for production of traditional high quality raw ham, the study

was conducted in the specific rearing context of the heavy pig, with the perspective to subsequently test analgesic and anaesthetic protocols.

Material and methods

Animal, housing and surgical procedure

All procedures carried out on animals were approved by the Ethical Committee for Animal Experiments of the University of Padova with the permission number 56 BIS-2012.

The study was conducted in a 2500 sow commercial pig farm located in the north-east of Italy. Data were collected over a 16-week period. All piglets in the study belonged to a commercial hybrid (75% Large White and 25% Belgian Landrace) selected for the production of the PDO cured ham.

Farrowing accommodations consisted of eight identical rooms. Each room held 12 (1.50 × 2.00 m) farrowing crates, with two rows of six crates separated by a corridor. Crates had fully slatted floors made of wire mesh covered with a plastic carpet. Ventilation and temperature were automatically controlled by fans and air heating, and an electric radiant lamp was used to heat the creep area in each pen. Immediately after farrowing, fostering was performed in litters for size and BW of piglets. According to the regular managerial practices of the hosting farm, at day 4 after birth, all piglets were subjected to intramuscular injection of iron dextran and males were castrated immediately after. Piglets were not submitted to tail docking nor tooth resection. All male piglets within litter were randomly allocated either to handling (H) or to surgical castration (C) experimental treatments excluding underweight or clinically sick piglets. The veterinarian entered the farrowing crate, took each piglet from under the sow or from the nest and individually identified piglets with a different coloured number on their back depending on treatment group. Surgical castration of the C group was performed following the common castration technique without using anaesthesia and analgesia, as allowed by the European Directive in force (EU Council Directive 2008/120/EC, 2008) if castration is carried out within the 7th day of life. The piglets were restrained between the handler's legs in a head down position, then an incision on each side of the scrotum was carried out with a scalpel and the testes removed by cutting the spermatic cord. Finally, a chlorhexidine-based antiseptic was applied on both the open wounds. For handled piglets (H) castration was simulated only by manipulation without making an incision, whereas iron injection and antiseptic application were performed as for the C group. After the procedure, piglets were returned to the farrowing crate.

To avoid confounding effects and excessive stress owing to manipulations during data collection, different sets of piglets were used for different experimental measurements, except for rectal and eye temperature. For each parameter considered, H and C piglets were randomised within each litter, choosing to have a higher number of castrated piglets (one or two H v. two or three C piglets). Within litter, H and C piglets were numbered consecutively and they were submitted to experimental measurements at different time points from treatment with the same order.

Experimental measurements

Plasma cortisol, lactate and glucose levels were measured on samples collected from 32 male piglets (12 H and 20 C) belonging to eight randomly chosen litters. Blood samples were collected from the anterior vena cava using 2.5 ml syringe and then stored in vacuum tubes without anticoagulant (Vacutest Kima srl, Arzergrande, PD, Italy). Blood was sampled 1 h before treatment (−1 h) to obtain individual basal levels and then repeated within 20 min after castration (20 min). To decrease the stress owing to repeated blood sampling, half the piglets were sampled 3 h after treatment (3 h), whereas the other half were sampled 5 h after treatment (5 h); 24 h after treatment all the piglets were sampled again. Glucose levels were measured immediately after sampling at the farm using a human self-monitoring system for assessing glycemic status (Accu-Chek®), never used in swine but already validated in cattle as a convenient field monitor (Rumsey *et al.*, 1999). The system allowed the measurement of glucose levels instantaneously using a drop of blood that remained in the syringe after the collection. For cortisol and lactate, blood samples were stored at 4°C and transferred to the laboratory within 8 h where they were centrifuged at 2500 × g for 10 min at 20°C. Serum cortisol concentration was determined with chemiluminescent assay (LKC01; Medical System, Genova, Italy) performed with automated analyzer Immulite One (Siemens Healthcare Diagnostics Products Ltd, Gwynedd, UK). The assay sensitivity was 5.5 nmol/l. The intra- and inter-assay CVs were 5.8% and 6.3%, respectively. The method was linear for up to 1380 nmol/l. Lactate was determined with a commercial colorimetric assay kit (L-Lactate; Randox Laboratories Ltd, Crumlin, UK) performed with an automated analyzer Cobas 501 (Roche Diagnostics, Mannheim, Germany). The assay sensitivity was 0.165 mmol/l. The intra- and inter-assay CVs were 0.9% and 5.7%, respectively. The method was linear for 0.165 to 19.7 mmol/l.

Rectal temperature was measured using a digital paediatric thermometer that was disinfected after each measurement and lubricated with gel for rectal examination in order to decrease the stress to the animals. In the first part of the experiment, rectal temperature was measured on 10 H and 14 C piglets belonging to six litters. The measurement was performed 1 h before treatment in order to obtain the basal temperature (−1 h), then it was repeated within 20 min, 4 and 24 h after treatment. On the basis of the results of the first set of piglets, showing an increment of temperature around 4 h after treatment, a second set of piglets (19 H and 43 C piglets belonging to 16 litters) was submitted to temperature detection 1 h before and 1, 3, 5 and 24 h after treatment.

Eye temperature was measured on 12 H and on 16 C piglets belonging to seven litters. The measurements were carried out 1 h before and repeated within 20 min, 2 and 4 h after treatment. The same piglets were submitted also to measurement of rectal temperature in order to correlate eye to rectal temperature. Piglets eye images were scanned using a portable IR camera (ThermaCam P25; Flir Systems, Boston,

MA, USA), which was calibrated to room temperature and absorptive conditions on each sampling day. While piglets were restrained by an operator, a second operator took two pictures of piglets' left half of the face. To reduce the effects of environmental factors on thermographic readings, all images were scanned within the barn at a right angle and at the same distance from the subject for all the piglets and all the time of measurement. The settings of the camera were as follows: range of temperature 7 to 34°C; emissivity of skin 0.93; reflected air temperature 20°C; distance between camera and skin surface 0.8 m; and field of view 23°.

The detector consisted of a focal plane array uncoiled microbolometer with 320 × 240 pixels resolution, thermal sensitivity of 0.08°C at 30°C, spatial resolution (instantaneous field of view) of 1.3 mrad, spectral range between 7.5 and 13 µm, accuracy ±2°C. Automatic corrections based on user input were conducted for reflected ambient temperature, distance, relative humidity and atmospheric transmission. Maximum and mean eye temperatures (°C) of the medial posterior palpebral margin of the lower eyelid and the lacrimal caruncle (Stewart *et al.*, 2008) were then extracted from images. Image analysis was performed using the ThermaCam Researcher Basic software (Flir Systems) only on images depicting completely open eyes.

Latency to move was measured on 287 piglets (130 H and 157 C) from 48 litters. Latency to move was defined as the time required by piglets to start walking when placed back in the farrowing crate after H or C treatment. This measurement was performed once by two veterinarians. The first one took all the male piglets of the litter away from each sow (one at a time), applied the treatments (H or C) and right after the treatment placed back each piglet in its farrowing crate. The other veterinarian (always the same one) started the stopwatch at the moment of the placement of the piglet in the farrowing crate. Every piglet was placed in the same point on the back of the farrowing crate at a constant distance from the sow which was standing and from the heated nest.

Occurrence of alterations in posture and gait such as weaker and protracted forward hind limbs (hind limbs positioned more forward than normal under the belly), hind limb non-weight bearing (walking with one hind limb off the ground), hind tip-toe walking pattern (contact with the floor only by the distal part of the hoof) and kyphosis (hunchback) were observed on a total of 97 (20 H and 77 C) male piglets belonging to 25 litters. Direct observations were carried out always by the same veterinarian 1 h before, within 20 min and 1, 2, 3, 5 and 24 h after treatment. As for latency, all piglets of the litter were taken from the sow and placed back one at a time with the same order by one veterinarian while the other carried out the observations. Each piglet was observed for a maximum of 1 min. When a piglet showed at least one alteration it was classified as 'altered', otherwise as 'normal'.

Statistical analysis

All statistical analyses were performed using SAS (SAS 9.3 Institute Inc., Cary, NC, USA). Data on cortisol, lactate, glucose,

rectal temperature and eye temperature were analysed using an unequally spaced repeated measures model (PROC MIXED), assuming as covariance structure among repeated measures a spatial power law (Littell *et al.*, 1998 and 2006) at different times that included the fixed effects of treatment (H v. C), time and their interaction, the litter was the random effect and the animal the repeated factor. For statistical differences ($P < 0.05$) showed by time and time \times treatment effects, contrast were also calculated. The variable latency to move was analysed using a mixed model procedure considering the treatment as fixed effect and the litter as random. Normality of residuals was evaluated by Shapiro–Wilk test (PROC UNIVARIATE) and values ≥ 0.9 were considered as normal, whereas not normally distributed variables (glucose and latency to move) were log transformed before analysis. Independence of residuals was graphically checked.

Pearson correlations within treatment were also calculated among blood parameters (cortisol, lactate and glucose) and between rectal and eye temperature. Percentage of animals with locomotory alterations were analysed by a repeated GLM (PROC GENMOD) with the Poisson distribution and the log link function. The model considered the effects of treatment (H v. C), time and their interaction and piglet within litter was the repeated effect.

Results

Cortisol levels were significantly affected by the interaction between time and treatment ($P < 0.001$) and its concentration showed the highest value within 20 min after surgery in C piglets (Figure 1a). Time of sampling affected significantly lactate concentrations ($P < 0.001$) and the lowest values were observed 24 h after treatments (Figure 1b). Glucose was also affected only by time ($P < 0.001$; Figure 1c). In particular, the concentration was higher before and within 20 min compared with 3, 5 and 24 h after treatment. In C piglets glucose was positively correlated to lactate ($r_p = 0.30$; $P < 0.05$) and cortisol concentrations ($r_p = 0.38$; $P < 0.01$).

In this first set of piglets, rectal temperature was significantly affected by time ($P < 0.05$), but not by treatment or interaction between time and treatment (Figure 2a). In particular, rectal temperature at 4 h after treatment was higher compared with the baseline and with those at 20 min and 24 h after treatment. In the second set of piglets, rectal temperature was affected by interaction between time and treatment (Figure 2b), in particular at 3 h after treatment it was higher in castrated compared with handled piglets ($P < 0.01$).

Mean eye temperature was affected by time of measurement ($P < 0.001$) and by the interaction between time of measurement and treatment ($P < 0.05$). Baseline mean eye temperature was lower and similar between H and C piglets, the highest value was observed for castrated piglets 4 h after castration (Figure 2c). Maximum eye temperatures were significantly affected by treatment ($P < 0.05$) and by time




Figure 1 Effect of interaction treatment \times time on (a) cortisol (LS means \pm s.e.m.) in handled (H) and castrated (C) piglets at different time of sampling, and effect of time of sampling on (b) lactate (LS means \pm s.e.m.) and (c) glucose (back transformed LS means \pm lower and upper 95% confidence interval). LS means with different letters differ significantly per $P < 0.05$. LS = least square.

of measurement ($P < 0.001$) but not by their interaction. Maximum eye temperature was higher in castrated piglets compared with handled (35.6 ± 0.08 v. 35.4 ± 0.09 ; $P < 0.05$) and it was higher 4 h after treatments compared with the previous measurements (Figure 2d). Rectal temperature was correlated with maximum eye temperature in both H ($r_p = 0.45$; $P < 0.01$) and C piglets ($r_p = 0.31$; $P < 0.01$) and with mean eye temperature in both H ($r_p = 0.38$; $P < 0.05$) and C piglets ($r_p = 0.28$; $P < 0.05$).

Latency to move was significantly affected by treatment ($P < 0.01$), and H piglets hesitated less before walking after being handled, whereas C piglets showed a latency of almost 4 s (Figure 3). Prevalence of alterations in locomotory behaviours (postures and gait) were the highest within 20 min from castration (Figure 4). Moreover, surgically castrated




Figure 2 Effect of time of measurement on (a) rectal temperature in piglets of the first trial (LS means \pm s.e.m.), and effect of the interaction treatment \times time on (b) rectal temperature in handled (H) and castrated (C) piglets of the second trial (LS means \pm s.e.m.). Effect of the interaction treatment \times time on (c) mean eye temperature in handled (H) and castrated (C) piglets (LS means \pm s.e.m.), and effect of time of measurement on (d) maximum (max) eye temperature in piglets (LS means \pm s.e.m.). LS means with different letters differ significantly per $P < 0.05$. LS = least square.




Figure 3 Latency to move in piglets (LS means \pm s.e.m.) immediately after handling (H) or surgical castration (C). LS means with different letters differ significantly ($P < 0.01$). LS = least square.

piglets had a higher risk to show alterations than handled piglets within 20 min, 1 and 3 h after treatment (Table 1). At the other two observations carried out 5- and 24-h post-treatment the risk was not statistically different between handled and castrated piglets.

Discussion

The present study aimed to test if some innovative parameters such as latency to move, differences in postures and walking, and eye temperature are useful for detection of pain in piglets in response to surgical castration, besides the physiological indicators commonly used for pain assessment. Regarding blood parameters, the highest cortisol concentration observed for castrated piglets at the first sampling




Figure 4 Prevalence (%) of locomotory behavior alterations in handled (H) and castrated (C) piglets at different time of observation.

after surgical castration (within 20 min) is in line with previous studies that reported increased cortisol levels for up to 120 min after castration (Prunier *et al.*, 2005; Waldmann *et al.*, 2010; Sutherland *et al.*, 2012). This is indicative of acute stress and could be related to an increased adrenal output owing to pain and tissue damage (Prunier *et al.*, 2005). After a painful event, epinephrine stimulates the mobilisation of muscular glycogen and its consequent hepatic metabolism, resulting in an increment of plasma lactate and glucose concentrations (Mayes, 1995). Lactate and glucose concentrations were therefore expected to vary between treatments and to be higher in castrated piglets, likely experiencing greater visceral pain. The lack of a significant treatment effect on glucose could be because of insufficient hepatic glycogen stores in 4-day-old piglets as hypothesised by Prunier *et al.* (2005). The high lactate levels in

Table 1 Relative risk (RR) of locomotory behavior alterations in surgically castrated piglets (C) compared with handled piglets (H) ($P = 0.037$) at different time of observation

Time of observation ¹	RR	Confidence interval (95%)		P-value
		Lower limit	Upper limit	
Within 20 min	12.30	2.45	61.64	0.002
1 h	8.22	1.63	41.54	0.011
3 h	9.11	1.76	47.21	0.008
5 h	3.79	0.74	19.22	0.108
24 h	1.23	0.55	2.73	0.617

¹Time of observation: right after treatment (within 20 min), 1, 3, 5 and 24 h after treatment.

piglets of both treatments in the current study, that were in the range of peak values observed 5 min after castration (Prunier *et al.*, 2005) or other acute stressors (Merlot *et al.*, 2011), could be related to a high level of stress owing to manipulation.

In the present study, castrated piglets had higher rectal temperature than handled ones only 3 h after treatment. Body temperature was reported to increase in response to anxiogenic or stress-inducing stimuli and injury (surgery and trauma) for the activation of the endogenous mechanisms related to inflammation (Takakazu *et al.*, 2001; Olivier *et al.*, 2003; Roth *et al.*, 2009). Inflammatory mediators such as TNF- α and IL-1 β are considered the main endogenous pyrogens (Roth *et al.*, 2009) and they were observed to increase in piglets 3 h after castration (Moya *et al.*, 2008). This could explain the tardive hyperthermia observed in the present study in castrated piglets, even if other external factors interfering with body temperature (exposure to heat lamps or time from milk intake) cannot be completely excluded. Moreover, the stress due to extra manipulation for temperature measurement could be associated to the late (5 and 24-h post-treatment) increase in rectal temperatures in both, handled and castrated piglets.

Maximum eye temperature was also higher in castrated compared with handled piglets and it increased in both treatments over time. Stewart *et al.* (2010a and 2010b) and Dockweiler *et al.* (2013) described an increase of eye temperature in calves right after castration (up to 10 min) due to an increase in parasympathetic nervous system activity associated with deep visceral pain. Results from these studies suggest that it would be beneficial to investigate eye temperature in a closer time interval to castration increasing the number of samplings but at the same time avoiding repeated handling that would not allow discrimination between pain and handling stress. Considering eye thermography, presence of different amounts of tears might have also influenced eye temperature as reported by Kamao *et al.* (2011) studying the dry eye syndrome in humans. In this case, it could be speculated that the difference between castrated and handled piglets detected in our study would be even underestimated.

Among behavioural parameters, latency to move differed between treatments. Castrated piglets needed more time to

start walking and this gives evidence that castration causes pain in the region of the hind legs. Similar hypothesis were formulated by McGlone *et al.* (1993) and Keita *et al.* (2010) who reported that walking is easily involved in alterations probably due to pain at the hind region subsequent to surgical castration of the piglet. Further support to this assumption comes from results of this study pointing out that none of the piglets showed locomotory alterations before treatments and that castrated piglets had a higher risk of occurrence of weaker and protracted forward hind limbs, hind limb non-weight bearing, hind tip-toe walking or kyphosis within 20 min, 1 and 3 h after treatment. Similar results were found by Lonardi *et al.* (2011) and Kluivers-Poodt *et al.* (2013) although they are in contrast with those of Hay *et al.* (2003) who reported abnormal behaviour for a few days after treatment.

Results of the study pointed out that detectable alterations of behaviour and of physiological parameters occur in the first 3 h from castration. Latency to move, alterations in posture and walking, and eye temperature acquired by IR thermography appear to be novel applicable pain assessment measures. The first two measurements are particularly promising since they require minimal handling of the piglets although they should be validated with inter- and intra-observer reliability tests. On the other hand, acquisition of data on eye temperature requires animals to be constrained in standardised conditions and needs further refinement of the methodology on pigs.

Acknowledgement

The authors would like to thank Animal Ethics Pty Ltd (Australia) for funding this project. The authors are grateful to Dr Meredith Sheil for advice and comments.

References

- Axiak SM, Jaggin N, Wenger S, Doherr MG and Schatzmann U 2007. Anaesthesia for castration of piglets: comparison between intranasal and intramuscular application of Ketamine, clomazepam and azaperone. *Schweizerisches Archiv für Tierheilkunde und Tierheilkunde* 149, 395–402.
- Carroll JA, Berg EL, Strauch TA, Roberts MP and Kattesh HG 2006. Hormonal profiles, behavioral responses, and short-term growth performance after castration of pigs at three, six, nine, or twelve days of age. *Journal of Animal Science* 84, 1271–1278.
- Chaloupková H, Illmann G, Neuhauserová K, Tománek M and Valis L 2007. Prewaning housing effects on behavior and physiological measures in pigs during the suckling and fattening periods. *Journal of Animal Science* 85, 1741–1749.
- Dockweiler JC, Coetzee JF, Edwards-Callaway LN, Bello NM, Glynn HD, Allen KA, Theurer ME, Jones ML, Miller KA and Bergamasco L 2013. Effect of castration method on neurohormonal and electroencephalographic stress indicators in Holstein calves of different ages. *Journal of Dairy Science* 96, 4340–4354.
- European Food Safety Authority 2004. Welfare aspects of the castration of piglets. *The EFSA Journal* 91, 1–18.
- EU Commission 2010. European Declaration on alternatives to surgical castration of pigs. Retrieved August 26, 2013, from http://ec.europa.eu/food/animal/welfare/farm/docs/castration_pigs_declaration_en.pdf
- EU Council Directive 2008/120/EC 2008. Laying down minimum standards for the protection of pigs (Codified version). *Official Journal of the European Union L* 47, 10.

- Guatteo R, Levionnois O, Fournier D, Guémené D, Latouche K, Letierrier C, Mormède P, Prunier A, Servière J, Terlouw C and Le Neindre P 2012. Minimising pain in farm animals: the 3S approach – 'Suppress, Substitute, Soothe'. *Animal* 6, 1261–1274.
- Hansson M, Lundeheim N, Nyman G and Johansson G 2011. Effect of local anaesthesia and/or analgesia on pain responses induced by piglet castration. *Acta Veterinaria Scandinavica* 53, 1–9.
- Hay M, Vulin A, Genin S, Sales P and Prunier A 2003. Assessment of pain induced by castration in piglets: behavioral and physiological responses over the subsequent 5 days. *Applied Animal Behaviour Science* 82, 201–218.
- Hodgson DS 2007. Comparison of isoflurane and sevoflurane for short-term anesthesia in piglets. *Veterinary Anaesthesia and Analgesia* 34, 117–124.
- Kamao TK, Yamaguchi M, Kawasaki S, Mizoue S, Shiraiishi A and Ohashi Y 2011. Screening for dry eye with newly developed ocular surface thermographer. *American Journal of Ophthalmology* 151, 782–791.
- Keita A, Pagot E, Prunier A and Guidarini C 2010. Pre-emptive meloxicam for postoperative analgesia in piglets undergoing surgical castration. *Veterinary Anaesthesia and Analgesia* 37, 367–374.
- Kluivers-Poodt M, Zonderland JJ, Verbraak J, Lambooi E and Hellebrekers LJ 2012. Pain behaviour after castration of piglets; effect of pain relief with lidocaine and/or meloxicam. *Animal* 7, 1158–1162.
- Kluivers-Poodt M, Houx BB, Robben SRM, Koop G, Lambooi E and Hellebrekers LJ 2012. Effects of a local anaesthetic and NSAID in castration of piglets, on the acute pain responses, growth and mortality. *Animal* 6, 1469–1475.
- Lonardi C, Brscic M, Scollo A and Gottardo F 2011. A preliminary study on the assessment of pain induced by mutilation in piglets. Proceedings of the 5th International Conference on the Assessment of Animal Welfare at Farm and Group Level, 8–11 August 2011, Guelph, Canada, 97pp.
- Littell RC, Henry PR and Ammerman CB 1998. Statistical analysis of repeated measures data using SAS procedures. *Journal of Animal Science* 76, 1216–1231.
- Littell RC, Milliken GA, Stroup WW and Wolfinger RD 2006. SAS[®] System for Mixed Models, 2nd edition. SAS Institute Inc., Cary, NC.
- Marchant-Forde JN, Lay DC, McMunn KA, Cheng HW, Pajor EA and Marchant-Forde RM 2009. Postnatal piglet husbandry practices and well-being: the effects of alternative techniques delivered separately. *Journal of Animal Science* 87, 1479–1492.
- Mayes PA 1995. Metabolisme du glycogène. In *Précis de Biochimie de Harper* (ed. Les Presses de l'Université Laval), pp. 207–216. De Boeck Université, St-Foy, Canada.
- McGlone JJ, Nicholson RI, Hellman JM and Herzog DN 1993. The development of pain in young pigs associated with castration and attempts to prevent castration-induced behavioral changes. *Journal of Animal Science* 71, 1441–1446.
- Mellor DJ, Cook CJ and Stafford KJ 2000. Quantifying some responses to pain as a stressor. In *The biology of animal stress: basic principles and implications for welfare* (ed. GP Moberg and JA Mench), pp. 173–198. CAB International, Wallingford, UK.
- Merlot E, Mounier AM and Prunier A 2011. Endocrine response of gilts to various common stressor: a comparison of indicators and methods of analysis. *Physiology & Behavior* 102, 259–265.
- Molony V and Kent JE 1997. Assessment of acute pain in farm animals using behavioral and physiological measurements. *Journal of Animal Science* 75, 266–272.
- Moya SL, Boyle LA, Lynch PB and Arkins S 2008. Effect of surgical castration on the behavioural and acute phase responses of 5-day-old piglets. *Applied Animal Behaviour Science* 111, 133–145.
- Mühlbauer I, Zöls S, Otten W, Palzer A, Ritzmann M and Heinritzi K 2010. Examination of CO₂ gas anesthesia during piglet castration. Proceedings of the 21st International Pig Veterinary Society Congress, 18–21 July 2010, Vancouver, Canada, 247pp.
- Olivier B, Zethof T, Pattija T, van Boogaerta M, van Oorschota R, Leahyc C, Oostinga R, Bouwknecht A, Veeninfg J, van der Gugtena J and Groeninka L 2003. Stress-induced hyperthermia and anxiety: pharmacological validation. *European Journal of Pharmacology* 463, 117–132.
- Prunier A, Hay M and Servière J 2002. Evaluation et prévention de la douleur induite par les interventions de convenance chez le porcelet. *Journées de la Recherche Porcine* 34, 257–268.
- Prunier A, Mounier AM and Hay M 2005. Effects of castration, tooth resection, or tail docking on plasma metabolites and stress hormones in young pigs. *Journal of Animal Science* 83, 216–222.
- Roth J, Rummel C, Barth SW, Gerstberger R and Hubschle T 2009. Molecular aspects of fever and hyperthermia. *Immunology and Allergy Clinics of North America* 29, 229–245.
- Rumsey TS, Kahl S and Elsasser TH 1999. Field method for monitoring blood glucose in beef cattle. *Journal of Animal Science* 77, 2194–2200.
- Sneddon LU and Gentle MJ 2000. Pain in farm animals, Proceedings of Workshop 5 on Sustainable Animal Production, 4–5 September 2000, Mariensee, Germany, pp. 9–18.
- Stewart M, Stafford KJ, Dowling SK, Schaefer AL and Webstera JR 2008. Eye temperature and heart rate variability of calves disbudded with or without local anaesthetic. *Physiology & Behavior* 93, 789–797.
- Stewart M, Verkerk GA, Stafford KJ, Schaefer AL and Webster JR 2010a. Non-invasive assessment of autonomic activity for evaluation of pain in calves, using surgical castration as a model. *Journal of Dairy Science* 93, 3602–3609.
- Stewart M, Webster JR, Stafford KJ, Schaefer AL and Verkerk GA 2010b. Technical note: Effects of an epinephrine infusion on eye temperature and heart rate variability in bull calves. *Journal of Dairy Science* 93, 5252–5257.
- Sutherland MA, Davis BL, Brooks TA and McGlone JJ 2010. Physiology and behavior of pigs before and after castration: effects of two topical anesthetics. *Animal* 4, 2071–2079.
- Sutherland MA, Davis BL, Brooks TA and Coetzee JF 2012. The physiological and behavioral response of pigs castrated with and without anesthesia or analgesia. *Journal of Animal Science* 90, 2211–2221.
- Takakazu O, Kae O and Tetsuro H 2001. Mechanisms and mediators of psychological stress-induced rise in core temperature. *Psychosomatic Medicine* 63, 476–486.
- Taylor AA, Weary DM, Lessard M and Braithwaite LA 2001. Behavioural responses of piglets to castration: the effect of pig age. *Applied Animal Behaviour Science* 73, 35–45.
- von Borell E, Baumgartner J, Giersing M, Jäggin N, Prunier A, Tuytens FAM and Edwards SA 2009. Animal welfare implications of surgical castration and its alternatives in pigs. *Animal* 3, 1488–1496.
- Waldmann KH, Rittershaus D, Kietzmann M and Puppe B 2010. Evaluation of different topical anaesthetics for pain reduction during surgical castration of male piglets, Proceedings of the 21st International Pig Veterinary Society Congress, 18–21 July 2010, Vancouver, Canada, 249pp.
- Weary DM, Niel L, Flower FC and Fraser D 2006. Identifying and preventing pain in animals. *Applied Animal Behaviour Science* 100, 64–76.
- White RG, DeShazer JA, Tressler CJ, Borchert GM, Davey S, Waninge A, Parkhurst AM, Milanuk MJ and Clemens ET 1995. Vocalization and physiological response of pigs during castration with or without a local anesthetic. *Journal of Animal Science* 73, 381–386.