

Demographics of extra-articular calcaneal fractures: including a review of the literature on treatment and outcome

Tim Schepers · Abida Z. Ginai ·
Esther M. M. Van Lieshout · Peter Patka

Received: 25 June 2007 / Published online: 19 December 2007
© The Author(s) 2007

Abstract

Introduction Extra-articular calcaneal fractures represent 25–40% of all calcaneal fractures and an even higher percentage of up to 60% is seen in children. A disproportionately small part of the literature on calcaneal fractures involves the extra-articular type. The aim of this study was to investigate the incidence of extra-articular calcaneal fractures in a Level 1 trauma centre, define the distribution of the various types of fractures and compare patient demographics between extra- and intra-articular calcaneal fractures. In addition the literature was reviewed for the most common types of extra-articular calcaneal fractures with regard to incidence, treatment and clinical outcome.

Methods The radiological records between 2003 and 2005 were reviewed for intra- and extra-articular calcaneal fractures. Patient gender-distribution and age were compared. A literature search was conducted for the treatment of extra-articular calcaneal fractures.

Results In this 3-year study period a total of 49 patients with 50 extra-articular calcaneal fractures and 91 patients with 101 intra-articular fractures were identified. The median age for the first group was 32.7 years, and for the second group 40.3 years; $P = 0.04$. Male predominance was significantly less pronounced for extra-articular (63%) compared with intra-articular fractures (79%; $P = 0.04$).

Conclusion One-third of all calcaneal fractures are extra-articular. Significant differences exist between the intra- and extra-articular groups, in terms of lower age and male–female ratio. The literature study shows inconsistencies in treatment options, but most extra-articular fractures are well manageable conservatively.

Keywords Calcaneus · Fracture · Extra-articular · Treatment · Outcome

Introduction

The proportion of calcaneal fractures without involvement of the posterior subtalar joint in adults is reported to be approximately 25% [4, 12, 26]. In children this number is higher and ranges from 25 to 63%, with increased percentages at lower age [14, 41, 67]. A disproportionately small part of the literature on calcaneal fractures involves the extra-articular type [17]. Controversy exists in the treatment of calcaneal fractures in general, but to a lesser extent in the extra-articular type compared with intra-articular calcaneal fractures [47].

Extra-articular fractures of the calcaneus are compression or avulsion type fractures. According to several classifications, i.e. by Böhler [8], Essex-Lopresti [26] and others [66, 67, 73], they are historically divided into three different groups: anterior portion, tuberosity (posterior calcaneal) and sustentaculum tali fractures (Table 1, Fig. 1). Several decades later additional calcaneal avulsion fractures of the calcaneus were identified [13, 55, 58].

The aim of the current study was to assess the incidence of different types of extra-articular calcaneal fractures in a Level-1 trauma centre, and comparing patients with an extra- or intra-articular fracture. In addition, a review of the

T. Schepers (✉) · E. M. M. Van Lieshout · P. Patka
Department of Surgery-Traumatology, Erasmus MC,
University Medical Centre, Room H-974,
P.O. Box 2040, 3000 CA Rotterdam, The Netherlands
e-mail: t.schepers@erasmusmc.nl

A. Z. Ginai
Department of Radiology, Erasmus MC,
University Medical Centre, Rotterdam, The Netherlands

Table 1 Classification of extra-articular calcaneal fractures according to different authors

Anatomical site	Fracture type	Böhler	Essex-Lopresti	Warrick-Bremner	Rowe	Schmidt-Weiner
Tuberosity	Avulsion “beak type”	1a/c ^a	+	+	2a	2a
	Achilles insertion	1b	–	–	2b	2b
	Vertical-shear	4	–	+	3	3
	Medial process	2	+	–	1a	1a
	Horizontal	–	+	–	–	–
	Apophysis	–	–	–	–	1a
	Bone loss	–	–	–	–	6
Sustentaculum tali		3	+	+	1b	1b
Anterior	Calcaneocuboid	–	+	+	–	–
	Anterior process	–	+	–	1c	1c
	Inferolateral	–	–	–	–	1d
Various	Avulsion fractures	–	–	–	–	1e

Different fracture types are numbered as stated in the original classification. If the original classification did not use numbers, the presence of a certain fracture type is presented as – (no) or + (yes)

^a Undisplaced (1a) or displaced (1c) “Beak” fracture according to Böhler




Fig. 1 Combined classification of extra-articular calcaneal fractures according to several authors (Table 1). 1 Undisplaced (1a) beak fracture; undisplaced (1b) avulsion fracture at Achilles-tendon insertion site; displaced (1c) “beak” avulsion fracture postero-superior aspect tuberosity; 2 medial (2a) or lateral (2b) process fracture; 3 isolated sustentaculum tali fracture; 4a vertical fracture; 4b, apophysis avulsion fracture; 5a, anterior superior avulsion or compression fracture; 5b, calcaneocuboid joint fracture; 5c, distal inferolateral portion fracture; 6a, extensor digitorum avulsion fracture; 6b, plantar fascia avulsion fracture; 6c, posterior capsular avulsion fracture

literature was conducted concerning the incidence, treatment and outcome of common extra-articular calcaneal fractures.

Patients and methods

Patients

All radiological records between January 1, 2003 and December 31, 2005 were electronically reviewed for the combination “calcaneus” AND “fracture” (in Dutch) at a

Level 1 Trauma centre, with ~24,000 patients presenting at the ED annually. The starting point of this search coincided with the introduction of digital radiographs [Picture Archiving and Communication System PACS]. These records were reviewed manually and when a fracture of the calcaneus was suspected, the original radiographs were re-examined by two reviewers (TS, AG). Patient who had radiographs taken on more than one occasion, were included only once.

Fractures of the calcaneus were classified as extra- or intra-articular, with involvement of the posterior talocalcaneal joint in the latter. The extra-articular fractures were further divided into anterior calcaneal portion fractures (anterior process, calcaneocuboid joint fractures), sustentaculum tali fractures (mid calcaneal portion) posterior calcaneal portion fractures (tuberosity avulsion, vertical, lateral and medial process fractures) and a group consisting of various avulsion fractures.

Statistical analysis

Statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) version 12.0 (SPSS, Chicago, IL, USA). The median age between groups was compared with the non-parametric Mann–Whitney *U* test. The difference in gender distribution and number of bilateral fractures between groups was analysed using the Chi-square test. Correlations were considered statistically significant with a *P* value of <0.05.

Literature search

The literature was reviewed for the various types of extra-articular calcaneal fractures; their incidence, treatment and

outcome. This search was conducted in the electronic databases of Embase, Cochrane Library and Pubmed using the following search-terms and Boolean operators: (“calcaneus” or “os calcis” or “calcaneum” OR “calcaneal”) and “fracture(s)” and/or “avulsion” and/or “extra-articular” up to July 2007. In addition a hand search of reference lists of eligible manuscripts and review articles was conducted to identify additional studies.

Results

Patients

A total of 540 patients with the terms “calcaneus” AND “fracture” mentioned in their radiological records were retrieved in the three year study period. In 140 cases a fracture of the calcaneus was encountered and in an additional 97 patients a calcaneal fracture was sustained prior to the investigatory period; meaning these were control radiographs to assess healing or in cases with persisting physical complaints. These patients were excluded from further analysis.

Out of the 140 patients with a calcaneal fracture, 49 patients sustained 50 extra-articular fractures. The different types of extra-articular calcaneal fractures and their frequency of appearance are shown in Table 2. The most commonly encountered extra-articular calcaneal fractures were the anterior process ($n = 19$), the extensor digitorum brevis avulsion ($n = 8$) and tuberosity medial process fractures ($n = 8$).

Patients with an intra-articular fracture had a median age of 40.3 years, compared with those with an extra-articular

fracture 32.7 ($P = 0.04$). The proportion of male patients was 79% in the first group, and in 63% the latter. This difference was statistically significant ($P = 0.04$). A total of ten patients with an intra-articular fracture sustained a bilateral fracture (11%), in the extra-articular group one patient (2%) sustained a bilateral fracture ($P = 0.06$).

Literature review

Calcaneal tuberosity fractures

Hippocrates already described tuberosity fractures of the calcaneus, resulting in gangrene and death if treated wrongly [25]. These fractures can be divided into avulsion (“beak” and Achilles avulsion) and shear-compression (vertical, lateral and medial process) fractures, although combinations have been described [68]. Tuberosity fractures, mainly shear-compression, comprise 12–40% of all calcaneal fractures encountered [9, 26, 73].

Fractures of the tuberosity result mainly from a fall from height or striking the heel on a ledge (Fig. 2a, b) [9]. Medial process fractures are thought to be the first stage of complex fractures [68], but can occur as avulsion fractures of the medial plantar fascia insertion [26] or abductor hallucis muscle insertion at the medial process [58]. Lateral process fractures and trochlear process fractures are rare [53].

The avulsion fractures of the postero-superior portion of the tuberosity (“beak” fractures) are caused by contraction of the calf-muscles while landing on the forefoot [20, 70]. Instead of rupturing the Achilles-tendon a fragment of approximately 2–3 cm is avulsed of the calcaneus (“Boyer fracture”) [6, 16, 26, 70]. Fractures alike type 1a are probably caused by deeper fibres of the soleus muscle, tilting the fragment upwards or by a more proximal insertion of the Achilles-tendon, as the Achilles-tendon usually inserts at level 1b (Fig. 1) [48, 62]. These avulsion fractures are more common in elderly women, in which osteoporosis plays an important role [6, 24, 44, 48, 57, 62, 65, 68, 70]. Other trauma mechanisms for the avulsion fracture are isolated muscle contractions, as seen in an athlete starting a run, or by direct trauma to the heel [32].

Two additional distinct avulsion fracture types of the tuberosity exist, namely the neuropathic Charcot-like fractures in diabetics and avulsion fractures of the apophysis of the calcaneus [2, 6, 35, 43, 50, 52]. The type 1 “Iowa fracture” according to Hedlund et al. [35] is an extra-articular tuberosity avulsion fracture near the insertion of the Achilles-tendon, occurring in patients with a history of diabetes.

The epiphysiolysis, a Salter-Harris type III fracture, occurs in children [2, 7, 40, 52]. It can be mistaken for Sever's disease (apophysitis by repeated stress) if the fracture is not displaced [7].

Table 2 Extra-articular calcaneal fractures ($n = 50$) seen between 2003 and 2005 in 49 patients

Fracture type	Numeric	Fractures (n)
Tuberosity avulsion	1a	1
Medial tuberosity process	2a	8
Lateral tuberosity process	2b	1
Sustentaculum tali fracture	3	1
Sustentaculum avulsion	3	2
Vertical tuberosity	4a	2
Anterior superior process	5a	19
Calcaneocuboid joint	5b	1
Distal inferolateral	5c	3
Extensor digitorum brevis	6a	8
Posterior capsular	6b	1
Plantar fascia	6c	3

Distribution of different extra-articular fracture types (see Fig. 1) and the number of times they were encountered

Fig. 2 Examples of several subtypes of extra-articular calcaneal fractures. **a** Medial process fracture, **b** vertical tuberosity fracture, **c** anterior process (Degan type I) fracture, **d** avulsion fracture at the insertion site of the extensor digitorum brevis muscle



Tuberosity avulsion fractures treated inadequately lead to stiffness, pain and swelling at the insertion site of the Achilles-tendon. Residual complaints after conservative treatment are muscle weakness especially in plantar flexion, resulting in difficulty in climbing stairs or while walking on toes [6, 62]. Most displaced tuberosity avulsion fractures are treated by open reduction and internal fixation (ORIF) with (tension-)wiring or screw fixation and an after-treatment in a above or below-knee cast for 4 or more weeks with the foot in plantar flexion (Table 3). Patients with low physical demands or fractures with minimally displaced fractures are manageable with conservative treatment [62, 65]. Medial process and minimally displaced vertical tuberosity fractures are managed conservatively with satisfactory outcome [53, 68].

Sustentaculum tali fractures

One of the earliest descriptions of sustentacular fractures was by Abel in 1878, who described three cases by direct force [25]. Isolated fractures of the sustentaculum tali are rare. Their incidence ranges from 0.3 to 4% [9, 26, 67, 73].

According to Essex-Lopresti the sustentaculum tali fracture is an intra-articular one, whereas all other classifications group the sustentacular fractures as extra-articular

[26, 66, 67]. Treatment of isolated sustentaculum tali fractures is usually conservative with satisfactory results, as the sustentaculum tali remains in position by strong fixation to the talus by the deltoid ligament and the interosseous ligament [8, 15, 17, 31, 37, 72].

Anterior process fractures

The anterior process is a bony prominence situated at the anterior-superior front of the os calcis and goes by several names [30, 33]. It is connected to both the cuboid and the navicular bone through the bifurcate ligament [21]. Anterior process, or anterior-superior promontory, fractures (Fig. 2c) are overlooked or misdiagnosed as ankle-sprains in up to 40% of the cases in the Emergency Department [5, 21, 27, 28, 30, 42, 46, 71]. Anterior process fractures represent 8–13% of all calcaneal fractures [26, 73]. Approximately 5% of patients with the history of an ankle-sprain has an anterior processus fracture, with a predominance in women [1, 10, 18, 27, 28, 54, 56].

Almost simultaneously Drachtler and Christopher reported this type of fracture for the first time in 1931 [18, 28, 64]. Different trauma mechanisms were postulated: forcible plantar-flexion (inversion-adduction, equinovarus, supination) of the foot [5, 10, 18, 21, 28, 30] and, less

Table 3 Treatment and outcome of tuberosity avulsion fractures in the literature

Author (year)	Treatment		Outcome	
	N	Description	N	Description
Partridge (1852) [57]	1	Plaster leg cast	1	Satisfactory
Coote (1867) [20]	1	Iron splint	1	Satisfactory
Thomson (1900) [70]	1	ORIF wiring	1	Satisfactory
Eisendrath (1905) [24]	2	ORIF suturing	2	Satisfactory
Cahill (1917) [16]	1	ORIF suturing	1	Satisfactory
Mooney (1935) [52]	1 ^a	ORIF suturing	1	Satisfactory
Rothberg (1939) [65]	1	ORIF suturing	1	Satisfactory
Struppler (1939) [69]	1	Pin and plaster	1	Satisfactory
Dieterle (1940) [22]	1	ORIF suturing	1	Satisfactory
Arner (1959) [3]	3	ORIF		NS
Protheroe (1969) [62]	2	Conservative	1	Satisfactory
	3	ORIF	3	Satisfactory
Lowy (1969) [48]	1	Conservative	0	Satisfactory
	3	ORIF	3	Satisfactory
Lyngstadaas (1971) [49]	2	ORIF screws		NS
Alvarez F (1989) [2]	1 ^a	ORIF screws	1	Satisfactory
Kathol (1991) [43]	14 ^b	NS		NS
Biehl (1993) [6]	3 ^b	Conservative	3	Satisfactory
	1	ORIF	1	Complicated
Birtwistle (1995) [7]	1 ^a	ORIF screws	1	Satisfactory
Cole (1995) [19]	1	Conservative	1	Satisfactory
	3	ORIF	3	Satisfactory
Levi (1997) [44]	1	ORIF wiring	1	Satisfactory
Hedlund (1998) [35]	12 ^b	NS, 4/12 LTF	7	Satisfactory
Squires (2001) [68]	2	Conservative	0	Satisfactory ^c
	1	ORIF	1	Satisfactory
Glanzmann (2005) [32]	1	ORIF (Mitek-Anchor)	1	Satisfactory
Imai (2007) [40]	1 ^a	ORIF	1	Satisfactory
Overall	11	Conservative	7	Satisfactory
	29	ORIF	23	Satisfactory
	26	NS	31	NS

LTF lost to follow-up, N number of patients treated, NS not specified

^a Avulsion fracture of the apophysis (epiphysiolysis) tuberosity of the calcaneus

^b Neuropathic avulsion fractures of the tuberosity of the calcaneus

^c Due to comorbidity

common, dorsiflexion–eversion trauma or forceful abduction of the forefoot with the heel fixed to the ground [21, 27, 61]. The first mechanism results in an avulsion fracture, the latter in a compression fracture [21]. Concomitant occurrence of additional injuries (talar, navicular, cuboid and calcaneal fracture or ligamentous injuries) or calcaneo-navicular coalition are seen frequently [56, 59].

Clinical presentation includes: immediate pain at the dorsolateral side of the foot, losing the ability to bear weight after a few hours, swelling and later discoloration [10]. Tenderness is usually found by placing the thumb on the lateral malleolus and the third digit on the base of the fifth metatarsal, with the index finger pointing out the area of maximum pain (“Gellman grip”) [28].

Radiographic investigation involves a lateral view of the calcaneus and a 45° inverted oblique view or a 10°–15°

cephalad, 10°–15° posterior view [27, 61]. Other views are less sensitive [63]. If there is high suspicion magnetic resonance imaging can be considered [56, 64]. The anterior process fracture must be distinguished from a calcaneus secundarius, which can be found in 2–5% on autopsy [36, 38, 39, 51]. Others consider the secondary ossicle to be a non-united anterior process fracture [64]. The fractures can be divided into three types according to Degan: Type I; an undisplaced fracture of only the tip of the anterior process, type II; displaced fracture of the anterior process not involving the calcaneocuboid joint, type III; large displaced fracture involving the calcaneocuboid joint [21]. Type I and II are sprain-fractures caused by inversion trauma, and type III is a compression fracture [61].

A successful treatment of anterior process fractures relies almost completely on the early detection of the fracture [10].

If not treated well this injury can cause persistent complaints for up to 18 months [46]. Conservative treatment, usually with a short leg cast for 4 weeks and increased weight-bearing, results in satisfactory outcome in most cases (Table 4) [10, 21, 36, 64]. In Degan type III fractures ORIF can be considered [39, 71]. The two most prognostic features of outcome are the type of injury and whether the fracture unites [21]. Overall outcome is satisfactory, but non-union rates range from zero to 40%, depending on adequately made diagnosis and treatment [21, 28, 64]. “It is such injuries that very probably lead to the well-known expression that a bad sprain is worse than a fracture” [63].

Anterior avulsion fractures

The extensor digitorum brevis originates from the dorsolateral portion of the anterior portion of the calcaneus, and inserts at the lateral sides of the extensor digitorum longus

tendons of the second to fourth toe, extending them with the foot in dorsal flexion.

If the foot is subjected to forceful inversion an avulsion fracture of the insertion of the extensor muscle may occur (Fig. 2d). Clinically it is difficult to distinguish between an anterior process (avulsion) fracture and an extensor digitorum brevis insertion avulsion fracture [34]. The haematoma, the point of maximum swelling and the point of maximum tenderness are situated in the same area as in anterior process fractures [55].

Norfray et al. [55] found 10 cases in a series of 100 consecutive patients suspected for an ankle fracture. The avulsion fracture was best seen at the dorsoplantar view of the foot and on the anteroposterior view of the ankle. An additional view recommended is the reversed oblique view of the foot, with the foot in approximately 35°–40° exorotation, e.g. supination [13, 29]. Brijis identified 62 avulsions of the anterolateral aspect of the calcaneus, situated more peripherally

Table 4 Treatment and outcome of anterior process fractures in the literature

Author (year)	Treatment		Outcome	
	N	Description	N	Description
Drachtler (1931) [23]	26	4 weeks cast and NWB	26	Satisfactory
Christopher (1931) [18]	2	Cast or taping	2	Satisfactory
	1	No immobilisation	0	Satisfactory
Gellman (1951) [28]	6	Short leg cast	6	Satisfactory
Bradford (1951) [10]	26	Plaster 4–6 weeks	25	Satisfactory
			1	Non-union
Backman (1953) [5]	20	Tape or short leg cast	20	Satisfactory
Piatt (1956) [60]	10	NS	1	Satisfactory
			2	Non-union
Garvin (1957) [27]	12	8 weeks short leg cast	12	Satisfactory
Levine (1959) [45]	1	None	1	Non-union
Hellpap (1962) [36]	47	4 weeks cast and NWB	47	NS
Carey (1965) [17]	47	Short leg cast 5 weeks	16	NS
			30	Satisfactory
Hunt (1970) [39]	1	ORIF	1	Satisfactory
Degan (1982) [21]	25	Short leg cast 5–6 weeks 7 excisions (5 non-union)	23	Satisfactory
Renfrew (1985) [63]	7	Cast or taping	4	NS
			3	Satisfactory
Trnka (1998) [71]	1	Excision of non-union	1	Satisfactory
Robbins (1999) [64]	3	Immobilisation	3	Satisfactory
		2	Delayed treatment	2
Hodge (1999) [38]	1	None	1	Non-union
Pillai (2005) [61]	1	ORIF	1	Satisfactory
Ouellette (2006) [56]	14	NS	14	NS
Petrover (2007) [59]	15	NS	15	NS
Overall	224	Conservative	146	Satisfactory
<i>LTF</i> lost to follow-up, <i>N</i> number of patients treated, <i>NS</i> not specified, <i>NWB</i> non-weight-bearing	42	ORIF	2	Satisfactory
		NS	130	NS

than fractures caused by avulsion of the extensor digitorum brevis. Brijis [13] suggested that these small fractures are caused by avulsions at the insertion site of the bifurcate ligament. Usually the avulsion is an isolated fracture and avulsed fragments resolved within a few months [11, 13, 29, 55]. Treatment is usually conservatively consisting of 2 weeks non-weight bearing, bandaging and early active motion exercises, and outcome is favourable in all patients [13, 29, 55].

Discussion

Thirty-five percent of all calcaneal fractures in the current study occurred outside the posterior subtalar joint. This percentage is slightly higher than previous incidences presented in the literature [4, 12, 26]. These percentages were calculated before the description of various types of avulsion fractures and might have been underscoring the number of extra-articular calcaneal fractures [13, 55].

Patients with an extra-articular calcaneal fractures were significantly younger, male predominance was less pronounced and the occurrence of a bilateral extra-articular calcaneal fractures is lower. This is explained largely by the difference in trauma-mechanism. The intra-articular calcaneal fractures occur frequently as work-related high-energy accidents and the extra-articular mainly as distortion injuries [5, 10, 18, 21, 28, 30, 55].

A total of 66 tuberosity avulsion fractures were retrieved from the literature, including four epiphysiolyse. Patients treated conservatively reached a satisfactory result in 64% compared with 88% of patients treated operatively. Patients return to their daily activities earlier after operative treatment. No randomised trials exist on the subject and half of the patients were not available for follow-up. Nevertheless the tendency nowadays is to operate these avulsion fractures.

Some 268 cases with an anterior process fractures were identified, with 90% of the patients reaching satisfactory result after conservative treatment. As fragments are usually too small to fixate, operative treatment is not treatment of choice. Important in obtaining best outcome is an early detection of the injury and adequate immobilisation.

The 78 anterior avulsion fractures from the insertion of the extensor digitorum brevis muscle or bifurcate ligament described in the literature did well in 100% after conservative treatment, and only one large fragment was corrected operatively.

Conclusion

One-third of the calcaneal fractures are extra-articular. Statistically significant differences exist in patient

characteristics, age and gender, between the intra- and extra-articular fractures. Evidence-based treatment options of extra-articular calcaneal fractures do not exist. Due to the low incidence of this type of fractures only retrospective case series and one prospective study regarding treatment options were found in the literature. Most extra-articular calcaneal fractures can be treated conservatively with overall satisfactory results. Displaced posterior superior avulsion fractures, displaced medial process and vertical fractures of the tuberosity and large anterior process fractures are best treated by ORIF according to the available literature.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Agnholt J, Nielsen S, Christensen H (1988) Lesion of the ligamentum bifurcatum in ankle sprain. *Arch Orthop Trauma Surg* 107(5):326–328
2. Alvarez Fernandez JL, Villalba Vaquero M, Gomez Cimiano J (1989) Epiphysiolyse of the great tuberosity of the calcaneum: brief report. *J Bone Joint Surg Br* 71(2):321
3. Arner O, Lindholm A (1959) Avulsion fracture of the os calcaneus. *Acta Chir Scand* 117:258–260
4. Atkins RM, Allen PE, Livingstone JA (2001) Demographic features of intra-articular fractures of the calcaneum. *Foot Ankle Surg* 7:77–84
5. Backman S, Johnson SR (1953) Torsion of the foot causing fracture of the anterior calcaneal process. *Acta Chir Scand* 105(6):460–466
6. Biehl WC 3rd, Morgan JM, Wagner FW Jr, Gabriel R (1993) Neuropathic calcaneal tuberosity avulsion fractures. *Clin Orthop Relat Res* 296:8–13
7. Birtwistle SJ, Jacobs L (1995) An avulsion fracture of the calcaneal apophysis in a young gymnast. *Injury* 26(6):409–410
8. Böhler L (1931) Diagnosis, pathology and treatment of fractures of the os calcis. *J Bone Joint Surg* 1:75–89
9. Böhler L (1957) Die Technik der Knochenbruchbehandlung. Verlag Wilhelm Maudrich, Wien Band II, pp 2148–2217
10. Bradford CH, Larsen I (1951) Sprain-fractures of the anterior lip of the os calcis. *N Engl J Med* 244(26):970–972
11. Brandser E, Braksiek R, El-Khoury G, Saltzman C, Marsh J, Clark W, Prokuski L (1997) Missed fractures on emergency room radiographs: an analysis of 433 patients. *Emerg Radiol* 4(5):295–302
12. Bremner AE, Warrick CK (1951) Fractures of the calcaneus. *J Fac Radiol* 2(3):235–241
13. Brijis S, Brijis A (1992) Calcaneal avulsion: a frequent traumatic foot lesion. *Rofo* 156(5):495–496
14. Brunet JA (2000) Calcaneal fractures in children. Long-term results of treatment. *J Bone Joint Surg Br* 82(2):211–216
15. Burdeaux BD Jr (1993) The medial approach for calcaneal fractures. *Clin Orthop Relat Res* 290:96–107
16. Cahill GF (1917) Fractures of the os calcis. *Ann Surg* 66(6):711–717
17. Carey DJ, Lance EM, Wade PA (1965) Extra-articular fractures of the os calcis—a follow-up study. *J Trauma* 8:362–372
18. Christopher F (1931) Fracture of the anterior process of the calcaneus. *J Bone Joint Surg Am* 1:877–879

19. Cole RJ, Brown HP, Stein RE, Pearce RG (1995) Avulsion fracture of the tuberosity of the calcaneus in children. A report of four cases and review of the literature. *J Bone Joint Surg Am* 77(10):1568–1571
20. Coote H (1867) Fracture of the os calcis from muscular action. *Lancet* 89(2270):270–271
21. Degan TJ, Morrey BF, Braun DP (1982) Surgical excision for anterior-process fractures of the calcaneus. *J Bone Joint Surg Am* 64(4):519–524
22. Dieterle J (1940) A case of so-called “open-beak” fracture of the os calcis. *J Bone Joint Surg Am* 2:740
23. Drachtler H (1931) Fractures of the anterior superior portion of the os calcis due to indirect violence. *Am J Roentgenol* 25(5):629–631
24. Eisendrath DN (1905) Fractures of the tarsal bones. *Ann Surg* 41(3):363–371
25. Ely LW (1906) Old fracture of the tarsus. *Ann Surg* 45(1):69–89
26. Essex-Lopresti P (1952) Mechanism, reduction technique and results in fractures of os calcis. *Br J Surg* 3:395–419
27. Garvin EJ, Rominger CJ (1957) Fractures of the anterior process of the calcaneus. *Am J Surg* 94(3):468–471
28. Gellman M (1951) Fractures of the anterior process of the calcaneus. *J Bone Joint Surg Am* 33-A(2):382–386
29. Geusens E, Geyskens W, Brys P, Janzing H (2000) The role of the reversed oblique radiograph in trauma of the foot and ankle. *Eur Radiol* 10(3):476–479
30. Gilheany M (2002) Injuries to the anterior process of the calcaneum. *Foot* 1:142–149
31. Gilmer PW, Herzenberg J, Frank JL, Silverman P, Martinez S, Goldner JL (1986) Computerized tomographic analysis of acute calcaneal fractures. *Foot Ankle* 6(4):184–193
32. Glanzmann M, Vereb L, Habegger R (2005) [Avulsion fracture of the calcaneal tuberosity in athletes]. *Unfallchirurg* 108(4):325–326
33. Golder WA (2004) Anterior process of the calcaneus: a clinical-radiological contribution to anatomical vocabulary. *Surg Radiol Anat* 26(3):163–166
34. Hall FM (1999) Calcaneal avulsion fractures. *AJR Am J Roentgenol* 173(3):854–855
35. Hedlund LJ, Maki DD, Griffiths HJ (1998) Calcaneal fractures in diabetic patients. *J Diabetes Complications* 12(2):81–87
36. Hellpap W (1962) Bone injuries of the anterior calcaneus process. *Arch Orthop Unfallchir* 5:329–338
37. Herzenberg JE (1986) CT of calcaneal fractures. *AJR Am J Roentgenol* 146(3):644–645
38. Hodge JC (1999) Anterior process fracture or calcaneus secundarius: a case report. *J Emerg Med* 17(2):305–309
39. Hunt DD (1970) Compression fracture of the anterior articular surface of the calcaneus. *J Bone Joint Surg Am* 52(8):1637–1642
40. Imai Y, Kitano T, Nakagawa K, Takaoka K (2007) Calcaneal apophyseal avulsion fracture. *Arch Orthop Trauma Surg* 127(5):331–333
41. Inokuchi S, Usami N, Hiraishi E, Hashimoto T (1998) Calcaneal fractures in children. *J Pediatr Orthop* 18(4):469–474
42. Judd DB, Kim DH (2002) Foot fractures frequently misdiagnosed as ankle sprains. *Am Fam Physician* 66(5):785–794
43. Kathol MH, el-Khoury GY, Moore TE, Marsh JL (1991) Calcaneal insufficiency avulsion fractures in patients with diabetes mellitus. *Radiology* 180(3):725–729
44. Levi N, Garde L, Kofoed H (1997) Avulsion fracture of the calcaneus: report of a case using a new tension band technique. *J Orthop Trauma* 11(1):61–62
45. Levine J, Kenin A, Spinner M (1959) Nonunion of a fracture of the anterior superior process of the calcaneus; case report. *J Bone Joint Surg Am* 41-A(1):178–180
46. Lindsay WR, Dewar FP (1958) Fractures of the os calcis. *Am J Surg* 95(4):555–576
47. Lowery RB, Calhoun JH (1996) Fractures of the calcaneus. Part II: Treatment. *Foot Ankle Int* 17(6):360–366
48. Lowy M (1969) Avulsion fractures of the calcaneus. *J Bone Joint Surg Br* 51(3):494–497
49. Lyngstadaas S (1971) Treatment of avulsion fractures of the tuber calcanei. *Acta Chir Scand* 137(6):579–581
50. Matsumura H, Jimbo Y, Kato T, Imai S (1998) Spontaneous calcaneal fracture after deep heel burns with diabetes. *Burns* 24(7):683–686
51. Mercer J (1931) The secondary os calcis. *J Anat* 66(1):84–97
52. Mooney V (1935) Avulsion of the epiphysis of the os calcis. *J Bone Joint Surg* 17(4):1056–1057
53. Moore GE (1933) Fractures of the tuber calcanei involving the medial and lateral processes. *Surg Gynecol Obstet* 5:400–405
54. Nielsen S, Agnholt J, Christensen H (1987) Radiologic findings in lesions of the ligamentum bifurcatum of the midfoot. *Skeletal Radiol* 16(2):114–116
55. Norfray JF, Rogers LF, Adamo GP, Groves HC, Heiser WJ (1980) Common calcaneal avulsion fracture. *AJR Am J Roentgenol* 134(1):119–123
56. Ouellette H, Salamipour H, Thomas BJ, Kassarian A, Torriani M (2006) Incidence and MR imaging features of fractures of the anterior process of calcaneus in a consecutive patient population with ankle and foot symptoms. *Skeletal Radiol* 35(11):833–837
57. Partridge (1852) Fracture of the os calcis. *Lancet* 60(1512):177
58. Pelletier JP, Kanat IO (1990) Avulsion fracture of the calcaneus at the origin of the abductor hallucis muscle. *J Foot Surg* 29(3):268–271
59. Petrover D, Schweitzer ME, Laredo JD (2007) Anterior process calcaneal fractures: a systematic evaluation of associated conditions. *Skeletal Radiol* 36(7):627–632
60. Piatt AD (1956) Fracture of the promontory of the calcaneus. *Radiology* 67(3):386–390
61. Pillai A, Arora J, Williams C, Ferdinand R (2005) The sprain fracture of the calcaneus revisited. *Foot* 1:198–201
62. Protheroe K (1969) Avulsion fractures of the calcaneus. *J Bone Joint Surg Br* 51(1):118–122
63. Renfrew DL, el-Khoury GY (1985) Anterior process fractures of the calcaneus. *Skeletal Radiol* 14(2):121–125
64. Robbins MI, Wilson MG, Sella EJ (1999) MR imaging of antero-superior calcaneal process fractures. *AJR Am J Roentgenol* 172(2):475–479
65. Rothberg A (1939) Avulsion fracture of the os calcis. *J Bone Joint Surg Am* 2:218–220
66. Rowe C, Sakellarides H, Freeman P, Sorbie C (1963) Fractures of the os calcis. *JAMA* 184(12):98–101
67. Schmidt TL, Weiner DS (1982) Calcaneal fractures in children. An evaluation of the nature of the injury in 56 children. *Clin Orthop Relat Res* 171:150–155
68. Squires B, Allen PE, Livingstone J, Atkins RM (2001) Fractures of the tuberosity of the calcaneus. *J Bone Joint Surg Br* 83(1):55–61
69. Struppler V (1939) Avulsion fracture of the calcaneal tuberosity. *Arch Orthop Trauma Surg* 39(5):651–658
70. Thomson R (1900) A rare case of fracture of the os calcis by muscular force. *Lancet* 156(4025):133–134
71. Trnka HJ, Zettl R, Ritschl P (1998) Fracture of the anterior superior process of the calcaneus: an often misdiagnosed fracture. *Arch Orthop Trauma Surg* 117(4–5):300–302
72. Vollrath T, Eberle C, Grauer W (1987) Computed tomography of intra-articular calcaneal fractures. *Rofu* 146(4):400–403
73. Warrick CK, Bremner AE (1953) Fractures of the calcaneum, with an atlas illustrating the various types of fracture. *J Bone Joint Surg Br* 35-B(1):33–45