

EPJ Web of Conferences **95**,04002 (2015)

DOI: 10.1051/epjconf/20159504002

© Owned by the authors, published by EDP Sciences, 2015

# Hadronic resonance production measured by the ALICE detector at LHC energies

A. Badalá<sup>1,a</sup>, on behalf of the ALICE Collaboration<sup>1</sup>*INFN - Sezione di Catania , Via S. Sofia 64, 95123, Catania (Italy)*

**Abstract.** Hadronic resonances are a valuable tool to study the properties of the medium formed in heavy-ion collisions. In particular, they can provide information on particle-formation mechanisms and on the properties of the medium at freeze-out, and furthermore they contribute to the systematic study of energy loss and recombination. Measurements of resonances in pp and in p–Pb collisions provide a necessary baseline for heavy-ion data and help to disentangle initial-state effects from medium-induced effects. In this proceedings the latest ALICE results on hadronic resonance production in pp, p–Pb and Pb–Pb collisions at LHC energies will be presented. In particular, the production of the  $K^*(892)^0$  and  $\phi(1020)$  resonances at mid-rapidity has been studied in different collision systems at LHC energies, reconstructing the resonances via their hadronic decay in a wide momentum range. The resonance transverse momentum spectra, mean transverse momenta, ratio to stable particles and nuclear modification factor will be discussed.

## 1 Introduction

In ultrarelativistic heavy-ion collisions a hot and dense state of matter [1, 2], the quark-gluon-plasma, is expected to be produced. At a critical temperature of about 160 MeV a cross-over transition between the partonic (i.e. a system with deconfined quarks) and hadronic phases is expected to take place. Resonances, with lifetimes comparable to that of the fireball (in the range of few fm/c to some tens of fm/c) are sensitive probes for different phases of the evolution of the medium produced in ultrarelativistic heavy-ion collisions. Modifications of the yield,  $\langle p_T \rangle$  and ratio of the yields of resonances to stable particles can provide information about the regeneration and re-scattering effects in the hadronic phase. In fact the final reconstructible resonance yields depend not only on the chemical freeze-out temperature but also on the scattering cross section of the resonance decay particle and the timescale between the chemical and the kinetic freeze-out, which control the fraction of 'undetected' particles. However resonances may be regenerated by pseudo-elastic interactions in the hadronic medium, a process driven by the cross-section of the interacting hadrons.

In heavy ion collisions information on particle formation mechanisms can be derived from the comparison of resonance production with that of long lived hadrons with similar mass but different baryon number and strangeness content. Finally, a contribution to the systematic study of the in-medium parton energy loss can be obtained from the measurement of the resonance production at

---

<sup>a</sup>e-mail: [Angela.Badala@ct.infn.it](mailto:Angela.Badala@ct.infn.it)

high- $p_T$ . Particularly interesting in this respect is the study of the  $\phi$  meson with its hidden strangeness content.

Both meson and baryon resonances have been measured by the ALICE experiment [3] in different collisions systems (pp, p-Pb, Pb-Pb) at LHC energies [4–7]. Resonance measurements in pp and p-Pb systems are useful as reference and to disentangle initial-state effects from genuine in-medium effects, which may occur in Pb-Pb collisions. In this paper, focus is given to the meson resonances  $K^*(892)^0$  and  $\phi(1020)$ , reconstructed at mid-rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV [6] and in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV.

## 2 Data analysis and resonance reconstruction


The results reported here refer to analyses carried out on samples of minimum-bias pp data at  $\sqrt{s} = 2.76$  TeV and 7 TeV (about 33 and 80 million events, respectively) and of minimum bias Pb-Pb and p-Pb data at  $\sqrt{s_{NN}} = 2.76$  TeV (about 13 million events) and  $\sqrt{s_{NN}} = 5.02$  TeV (about 90 million events), collected using the ALICE detector [3]. More information about the ALICE performance can be found in [8].

The  $K^*(892)^0$  and  $\phi(1020)$  production has been measured reconstructing the resonances through their main hadronic decay:  $K^*(892)^0 \rightarrow K^+\pi^-$  ( $\bar{K}^*(892)^0 \rightarrow K^-\pi^+$ ) and  $\phi \rightarrow K^+K^-$ . All measurements of  $K^*(892)^0$  and  $\bar{K}^*(892)^0$  are averaged and these mesons are referred to as  $K^{*0}$  in the following. In pp and in Pb-Pb resonances are measured in one unit of rapidity  $|y| < 0.5$  in the centre-of-mass reference frame, while in p-Pb the rapidity range is restricted to  $-0.5 < y < 0$ , in order to ensure the best detector acceptance with the shifted centre-of-mass of the system. The position of the primary vertex along the beam axis is reconstructed using the tracks reconstructed in the Inner Tracking System (ITS) and in the Time Projection Chamber (TPC) and is required to be within 10 cm from the center of the ALICE detector. Two scintillator hodoscopes, the V0 detectors, covering the pseudo-rapidity ranges  $-3.7 < \eta < -1.7$  (V0C) and  $2.8 < \eta < 5.1$  (V0A), were used for event triggering and the definition of centrality and multiplicity classes respectively in Pb-Pb [13] and p-Pb [14] collisions.

Identification of pions and kaons is carried out using the measurement of the specific energy loss  $dE/dx$  in the TPC. In p-Pb data an improvement in the significance of the signal has been achieved using the information of the Time-Of-Flight (TOF) detector, for tracks for which it is available. Resonances are reconstructed by computing the invariant mass spectrum of all primary track pairs and then subtracting the combinatorial background, estimated by event-mixing or like-sign techniques. The position of the peak and its width, in the invariant mass distribution of  $K^{*0}(\phi)$  have been extracted fitting the signal, after the subtraction of the combinatorial background, with a relativistic Breit-Wigner (Voigtian) plus a polynomial for the residual background. In all the collisions systems mass and width of  $K^{*0}(\phi)$  are found to be close to PDG value. In particular, in Pb-Pb collisions no mass shift or broadening has been observed.

## 3 Results

The procedure used to estimate the  $K^{*0}$  and  $\phi$  yield has been extensively explained in [5, 6]. In p-Pb collisions to extract the particle yields and the  $\langle p_T \rangle$ , the spectra are fitted using a Levy-Tsallis parameterization [15]. To extract the  $dN/dy$  the measured  $p_T$  distributions are integrated, while the fits are used to estimate the resonance yield at low and high  $p_T$ , where no signal could be measured. It may be noted that the extrapolated fraction of the total yield for the  $K^{*0}$  is lower than 0.1%.



**Figure 1.** (Left panel)  $(p+\bar{p})/\phi$  ratio as a function of transverse momentum  $p_T$  measured in p–Pb 0-5% (red full circle) and 60-80% (purple hollow cross) V0A multiplicity classes, compared to pp (pink hollow circle), 0-10% (black hollow squares) and 80-90% (black hollow cross) Pb–Pb collisions. (Right panel)  $K^{*0}/K^-$  and  $\phi/K^-$  ratios as a function of the cube root of the charged particle multiplicity density  $dN_{ch}/d\eta$  for pp, p–Pb and Pb–Pb collisions, respectively, at  $\sqrt{s} = 7$  TeV and  $\sqrt{s_{NN}} = 5.02$  and 2.76 TeV. The values given by a grand-canonical thermal model with chemical freeze-out temperature of 156 MeV are also shown [21].

### 3.1 Mean transverse momentum

Information on the particle production mechanisms can be obtained from the mean transverse momentum,  $\langle p_T \rangle$ . This has been measured for  $\pi^\pm$ ,  $K^\pm$ , (anti)protons,  $K^{*0}$ ,  $\phi$  in pp, p–Pb and Pb–Pb collisions and for  $\Lambda$  in p–Pb collisions [5, 6, 14, 16, 17]. In central Pb–Pb collisions, particles with similar mass ( $K^{*0}$ , p and  $\phi$ ) have similar  $\langle p_T \rangle$  [6]. This is consistent with hydrodynamical particle production, where the  $p_T$  distribution is mainly determined by particle mass.


On the other hand, in p–Pb and pp collisions, the mass ordering is only approximate:  $\langle p_T \rangle$  of resonances is larger than the  $\langle p_T \rangle$  for p and  $\Lambda$ . A similar trend is observed in pp at 7 TeV, where  $\langle p_T \rangle_\phi > \langle p_T \rangle_{K^{*0}} > \langle p_T \rangle_p$ . The question remains open whether the mesonic resonances deviate from mass ordering or the baryons, namely p and  $\Lambda$ , do, instead. However a plot of  $\langle p_T \rangle$  as a function of the particle mass suggests the possibility of two different trends: one for the mesons (including the resonances) and another for the baryons. Furthermore the values of  $\langle p_T \rangle$  for the highest-multiplicity event class in p–Pb collisions reach (or even exceed) the values measured in central Pb–Pb collisions. All these observations suggests the possibility of a different production mechanisms in Pb–Pb and p–Pb collisions [18].

### 3.2 Particle ratios

The baryon to meson ratios are useful quantities to study the hadron mechanism production. Particularly interesting in this respect is the comparison of the yield of proton and  $\phi$ , which have a similar mass. In the left panel of fig. 1 the  $(p+\bar{p})/\phi$  ratio as a function of transverse momentum  $p_T$  for different collisions systems and centrality is shown. In central (0-10%) Pb–Pb collisions this ratio is flat below 3 GeV/c, suggesting that the low- $p_T$  spectral shapes of the p and  $\phi$  are mainly determined by the particle mass. The trend of distribution of the  $(p+\bar{p})/\phi$  ratio in p–Pb collisions for all event multiplicity

classes steeply decreases with  $p_T$ , similarly to those observed for peripheral Pb–Pb collisions and for pp collisions. In central p–Pb collisions (i.e. for 0–5% V0A multiplicity event class) the ratio shows a hint of flattening for  $p_T < 1.5$  GeV/c.

In order to check the presence of a suppression in the production of the resonances and to study whether the strength of the suppression is related to the system size, the  $K^{*0}/K^-$  and  $\phi/K^-$  ratios have been reported as a function of the cube root of the charged particle multiplicity density  $((dN_{ch}/d\eta)^{1/3})$ , which is a good approximation for the system radius [19, 20], for pp, p–Pb and Pb–Pb collisions, respectively, at  $\sqrt{s} = 7$  TeV and  $\sqrt{s_{NN}} = 5.02$  and 2.76 TeV (Fig. 1, right panel). In p–Pb collisions  $\phi/K^-$  is rather independent from event multiplicity class and  $K^{*0}/K^-$  lies on the interpolation from pp to peripheral Pb–Pb collisions. The  $\phi/K^-$  in central Pb–Pb collisions is almost flat and it is consistent with the prediction of a grand-canonical thermal model [21], which has a chemical freeze-out temperature of 156 MeV and a baryochemical potential of 0 MeV and does not include re-scattering effects. On the contrary, the  $K^{*0}/K^-$  exhibits a clear suppression with the increase of the fireball size, i.e. going from peripheral to most central Pb–Pb collisions, where the measured ratio is about 60% of the predicted thermal model value. Considering the factor of about 10 between the lifetimes of the two resonances, the origin of the differences in the  $K^{*0}$  and  $\phi$  production could be related to a large modification of the  $K^*$  yield due to the pion rescattering mechanism ( $\sigma(\pi,\pi)$ ), which destroys the pion-kaon correlation of the  $K^{*0}$  decay products.




**Figure 2.** Transverse-momentum distribution of  $K^{*0}$  and  $\phi$  resonances in central (a) and peripheral (b) Pb–Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV, compared to the blast-wave expectation (see text). The lower panels show the ratios of the measured distributions to the prediction.

### 3.3 Transverse momentum spectra and interactions in the hadronic phase

According to UrQMD calculations [22, 23] the hadronic rescattering effect is expected to be momentum dependent with greater strength at low  $p_T$  ( $p_T < 2$  GeV/c). To investigate the  $p_T$  dependence

of the observed suppression the blast-wave model [24] is used to generate an expected transverse-momentum distribution without re-scattering effects for  $K^{*0}$  and  $\phi$  at kinetic freeze-out. In fig. 2 the transverse momentum distribution of  $K^{*0}$  and  $\phi$  resonances in central (0-20%) and peripheral (60-80%) Pb–Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV are compared to the blast-wave prediction for the spectral shape. The parameters of the blast-wave curves are obtained from a simultaneous fit to the  $p_T$  distributions of charged particles (pions, kaons and protons) in Pb–Pb collisions at the same collision energy [17]. The curves are normalized to the expected resonance yields estimated by multiplying the measured yield of charged kaons in Pb–Pb collisions [17] by the  $K^{*0}/K$  and  $\phi/K$  ratios given by a thermal-model fit to ALICE data [21]. In the  $p_T$  range less than 2 GeV/c the  $\phi$  data are satisfactorily described by the prediction in both central and peripheral collisions. The same conclusions hold for  $K^{*0}$  in peripheral collisions, where the data/theory ratio does not appear to deviate significantly from unity. On the other hand, for  $p_T < 2$  GeV/c in central collisions, the  $K^{*0}$  appears suppressed by a factor 0.6. The deviation from unity is about 3 times larger than the uncertainties, suggesting that  $K^{*0}$  has undergone non-negligible re-scattering effects.

By assuming a chemical freeze-out temperature of 156 MeV, a model-dependent estimate of 2 fm/c as the limit of the time between the chemical and kinetic freeze-out has been extracted [6] using the measured  $K^{*0}/K^-$  ratio.



**Figure 3.** Nuclear modification factor of  $\phi$  in 0-5% central Pb–Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV ( $R_{AA}$ , left panel) and in minimum bias p–Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV ( $R_{pPb}$ , right panel) compared to that of identified stable hadrons.

### 3.4 Nuclear modification factor

Parton in-medium energy loss at high  $p_T$  is usually studied by the so-called nuclear modification factor  $R_{AA}$ . It is defined as  $R_{AA}(p_T) = \frac{dN_{AA}/d p_T}{\langle T_{AA} \rangle d\sigma_{pp}/d p_T}$ , where  $N_{AA}$  and  $\sigma_{pp}$  represent the charged particle yield in nucleus-nucleus collisions and the cross section in pp collisions, respectively.  $T_{AA}$  is the nuclear overlap function, computed in the framework of a Glauber model [9]. In central AA collisions a suppression of the production of high  $p_T$  particles has been observed already at RHIC energies. An increase of this effect has been reported by the ALICE collaboration [10, 11], consistent with the formation of a coloured, dense fireball at these collision energies.

The nuclear modification factors of  $K^{*0}$  and  $\phi$  have been computed for Pb–Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV ( $R_{AA}$ ) and for p–Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV ( $R_{pPb}$ , only for  $\phi$ ). In the left panel of fig.3 the nuclear modification factor of  $\phi$  in 0-5% central Pb–Pb collisions at

$\sqrt{s_{NN}} = 2.76$  TeV is shown compared to that of identified stable hadrons. In central collisions the suppression at  $p_T > 8$  GeV/c of the  $\phi$  is consistent with that measured for the stable hadrons ( $\pi$ , K and p), thus supporting once more the observation of the flavour-independence of the partonic energy loss in the medium. However, a large baryon/meson dependence is observed for low  $p_T$  in particular  $R_{AA}(p) > R_{AA}(\pi)$ . It is worth noting that the  $R_{AA}$  of the  $\phi$  meson is slightly larger than that of  $\pi$  and lower than the  $R_{AA}$  of the p. The production of  $\phi$  in minimum bias p-Pb and pp collisions are compared by computing the  $R_{pPb}$  (fig. 3, right panel). The reference pp spectrum at  $\sqrt{s} = 5.02$  TeV has been obtained from the interpolation of the spectra measured in pp at 2.76 TeV and 7 TeV, following the same procedure described in [12] for identified charged hadrons. The trend of the  $R_{pPb}$  for the  $\phi$  exhibits a moderate Cronin peak (reaching a value of about 1.2) for  $3 < p_T < 6$  GeV/c. Moreover, no Cronin peak is observed for the pions, while a stronger Cronin peak is present for the protons in the same  $p_T$  range. For the  $\phi$  and the stable hadrons no suppression is seen at high- $p_T$  ( $p_T > 8$  GeV/c) in p-Pb collisions compared to pp.

## 4 Conclusions

The latest results on  $K^*(892)^0$  and  $\phi(1020)$  resonance production, measured by the ALICE detector in p-Pb and Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV and 5.02 TeV, respectively, have been presented. In particular the spectra at high transverse momentum have been measured (up to 10 GeV/c for  $K^{*0}$  and up to 21 GeV/c for  $\phi$ ). The nuclear modification factors of these resonances have been computed for Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV ( $R_{AA}$ ) and for p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV ( $R_{pPb}$ , only for  $\phi$ ). In central Pb-Pb collisions, high- $p_T$  resonances are strongly suppressed as for other stable hadrons, while for the  $\phi$  and the stable hadrons no suppression is seen at high- $p_T$  ( $p_T > 8$  GeV/c) in p-Pb collisions compared to pp.

In p-Pb collisions the mean  $p_T$  of  $K^{*0}$  and  $\phi$  does not follow the same mass ordering as other long lived particles, while in central Pb-Pb collisions the  $\langle p_T \rangle$  is compatible with that of protons.

The ratios of resonances to stable hadrons have been measured and compared in different collision systems. The  $K^{*0}/K^-$  is suppressed in central Pb-Pb collisions, consistent with dominant re-scattering of  $K^{*0}$  decay daughters in the hadronic phase, while the  $\phi/K^-$  is not suppressed consistent with  $\phi$  longer lifetime. The  $(p+\bar{p})/\phi$  ratio is flat for  $p_T < 3-4$  GeV/c in central Pb-Pb collisions, suggesting that the low- $p_T$  spectral shapes of the p and  $\phi$  are mainly determined by the similar particle mass. In pp and in peripheral Pb-Pb collisions as well in p-Pb collisions for all event multiplicity classes a steep decrease with  $p_T$  is observed for the  $(p+\bar{p})/\phi$  ratio. Only in central p-Pb collisions (i.e. for 0-5% V0A multiplicity event class) and for low  $p_T$  ( $p_T < 1.5$  GeV/c) the ratio shows a hint of flattening, which could suggest the onset of a collective behaviour.

## References

- [1] S. Borsanyi *et al.*, J. High Energy Phys. **11** (2010), **077** (2010); S. Borsanyi *et al.*, J. High Energy Phys. **09** (2010), **073** (2010);
- [2] P. Petreczky, Proc. of Science (Confinement X) **028** (2012)
- [3] K. Aamodt *et al.* (ALICE Collaboration), J. Instrum. **3**, S08002 (2008).
- [4] B. Abelev *et al.* (ALICE Collaboration) Eur. Phys. J. C **71**, 1594 (2011)
- [5] B. Abelev *et al.* (ALICE Collaboration) Eur. Phys. J. C **72**, 2183 (2012)
- [6] B. Abelev *et al.* (ALICE Collaboration) submitted to Phys. Rev. C, arXiv:1404.0495
- [7] B. Abelev *et al.* (ALICE Collaboration), submitted to Eur. Phys. J. C, arXiv:1406.3206
- [8] B. Abelev *et al.* (ALICE Collaboration), Int. J. Mod. Phys. A **29** 1430044 (2014)

- [9] M. Miller *et al.*, Ann. Rev. Nucl. Part. Sci. **57**, 205 (2007)
- [10] K. Aamodt *et al.* (ALICE Collaboration), Phys. Lett. B **696**, 30 (2011)
- [11] B. Abelev *et al.* (ALICE Collaboration), Phys. Lett. B **720**, 52 (2013)
- [12] M. Knichel (for the ALICE Collaboration), Quark Matter 2014 proceedings, Nucl. Phys. A 2014  
In press, arXiv:1408:0216
- [13] B. Abelev *et al.* (ALICE Collaboration) Phys. Rev. Lett. **106**, 032301 (2011)
- [14] B. Abelev *et al.* (ALICE Collaboration) Phys. Lett. B **728** 25 (2014)
- [15] C. Tsallis, J. Stat. Phys. **42**, 479 (1988)
- [16] F. Bellini (for the ALICE Collaboration) Quark Matter 2014 proceedings, Nucl. Phys. A 2014  
In press, DOI: 10.1016/j.nuclphysa.2014.08.031
- [17] B. Abelev *et al.* (ALICE Collaboration), Phys. Rev. C **88**, 044910 (2013)
- [18] B. Abelev *et al.* (ALICE Collaboration), Phys. Lett. B **727**, 371(2013).
- [19] K. Aamodt *et al.* (ALICE Collaboration), Phys. Lett. B **696**, 328 (2011)
- [20] M.A. Lisa *et al.*, Annu. Rev. Nucl. Part. Sci. **55**, 357 (2005)
- [21] J. Stachel *et al.*, J. Phys. Conf. Ser. **509**, 012019 (2014)
- [22] S. Bass *et al.*, Prog. Part. Nucl. Phys. **41**, 255 (1998)
- [23] M. Bleicher *et al.*, J. Phys. G **25**, 1859 (1999)
- [24] E. Schnedermann, J. Sollfrank and U. Heinz, Phys. Rev. C **48**, 2462 (1993).