



Animal (2007), 1:7, pp 1026–1032 © The Animal Consortium 2007
doi: 10.1017/S1751731107000389

The reactivity of neonatal rabbits to the mammary pheromone as a probe for viability

G. Coureaud^{1†}, L. Fortun-Lamothe², D. Langlois³ and B. Schaal¹

¹Equipe d'Ethologie et de Psychobiologie Sensorielle, Centre Européen des Sciences du Goût, UMR 5170 CNRS/UB/INRA, 15 rue Picardet, 21000 Dijon, France;

²Equipe Tissus animaux, nutrition, digestion, écosystème et métabolisme, UMR 1289 INRA/INP-ENSAT/ENVIT, Chem. de Borde Rouge BP52627, 31326 Castanet Tolosan, France; ³Unité Flaviv, INRA, 17 rue Sully, 21065 Dijon, France

(Received 12 March 2007; Accepted 30 May 2007)

Newborn rabbits depend on a daily nursing interaction with the mother to gain milk and to survive. During this interaction, they localise and seize the nipples displaying a typical behaviour triggered by maternal odour cues. The mammary pheromone constitutes such a signal in domestic rabbits: it elicits sucking-related movements in more than 90% of the pups. However, some newborns remain unresponsive to the presentation of the pheromone, even pups apparently healthy and highly motivated to suck. The main goal of the present study was therefore to explore the link between the unresponsiveness of rabbit pups to the mammary pheromone and their growth and survival in breeding conditions. To that end, 293 newborns from 30 litters were tested for their head searching-oral grasping responses to the mammary pheromone on days 1 and 3, and their milk intake and mortality were followed up from days 1 to 21. It was hypothesised that unresponsive newborns would have subsequent difficulties in finding nipples, sucking and surviving. Early weight and success in milk intake were further considered as mediating factors in growth and viability. The results showed that pups that were unresponsive to the mammary pheromone on day 1 were less successful in gaining milk and had a higher rate of mortality than the responsive pups. However, this impact was modulated by the weight of pups: it appeared only in the lightest newborns. Moreover, this impact vanished on day 3. On the other hand, the pup weight and sucking success on days 1 to 3 strongly influenced viability and growth during the period extending from days 1 to 21. Taken together, the results show that the day-1 responsiveness of rabbit pups to the mammary pheromone can be considered as an indicator of individual viability in pups having a small weight (<48 g on day 1). The predictive validity of the pups' pheromonal reactivity seems however time-limited as it works only during the first, but crucial, postnatal days.

Keywords: growth, mother–young relationships, pheromones, rabbits, viability

Introduction

In the domestic rabbit, the first days after birth and the weeks around weaning are periods of major challenges to the viability of offspring. Under various breeding conditions, it was noted that the first wave of losses generally occurred between birth and day 10 (e.g. Lebas, 1974; Pizzi and Crimella, 1984; Gualterio *et al.*, 1988; Seitz 1997; Coureaud *et al.*, 2000a). The survival of individual newborns depends then on their ability to orient to the mother's abdomen, to locate nipples and to suck efficiently in a highly constrained time span. Indeed, rabbit females nurse their litter once a day for less than 5 min (Zarrow *et al.*, 1965; Gonzalez-Mariscal, 2007). To locate the nipples rapidly, pups rely on odour cues emitted on the surface of the females' abdomen and/or in milk (e.g. Schley, 1976; Hudson and Distel, 1983;

Keil *et al.*, 1990; Coureaud *et al.*, 2001). These odour cues may be learned before birth (e.g. Bilkó *et al.*, 1994; Coureaud *et al.*, 2002) or in the first postnatal hours or days (e.g. Hudson, 1985; Kindermann *et al.*, 1991; Coureaud *et al.*, 2006b). Some of these odour signals are however active immediately after birth, seemingly in total independence from any postnatal or prenatal experience. The mammary pheromone belongs to this latter category. It is carried in rabbit milk and is an extremely potent activator of typical orocephalic movements that pups generally display to find the nipples under the nursing female (Coureaud, 2001; Coureaud *et al.*, 2003 and 2006a; Schaal *et al.*, 2003).

Although the bulk of rabbit pups (>90%) are highly responsive to the mammary pheromone within seconds of its presentation, a small proportion of them remains unresponsive. This observation of pheromonal unresponsiveness is even maintained under circumstances when the pups are

[†] Email: coureaud@cesg.cnrs.fr

experimentally aroused and highly motivated to suck (i.e. in the minutes preceding the daily nursing). The perceptual or motor correlates of such unresponsiveness are unknown. Yet, it may be interesting to assess whether the pups that do not react in the typical way to the mammary pheromone undergo a particular developmental trajectory, in terms of adaptation to the nursing interaction. Thus, the present study aimed to evaluate whether the responsiveness displayed by rabbit newborns to the mammary pheromone on postnatal days 1 or 3 was related with milk intake, growth and viability along most of the suckling period (i.e. between days 1 and 21).

Material and methods

Animals and housing conditions

Thirty multiparous does (strain INRA A1077; inseminated with PS19 semen, Grimaud Frères Sélection, France) and their litters were used in the breeding unit of INRA (Castanet-Tolosan, France). The does were housed in individual cages (0.28 × 0.46 × 0.69 m) to which a nest box (0.35 × 0.25 × 0.15 m) containing shavings (Copeaux Gascons, Cobois Diffusion, Lafitte sur lot, France) was appended 2 days before parturition. The animals were kept under constant light/dark cycle (16 h/8 h; lights on at 0600 h) and ambient temperature (20 to 24°C). Water and pelleted food (Rablo-Maternité; Groupe Coopératif Occitan, Castelnaudary, France) were provided *ad libitum*. The access to the nest was free for the mother, except on days 1 and 3 when the responsiveness to the mammary pheromone was tested (see section Behavioural assay). In total, 293 newborns from 30 litters were involved in the study. They were individually identified by ear-rings (Boucles simplex baby, Etablissement Chevillot, Albi, France) on the day of birth (day 0). On day 1, immediately after the behavioural assay and before weighing, the litters were randomly culled or completed to 10 pups (except for two litters culled to nine) to balance between litters pup competition under the mother (five pups of medium weight born to three females excluded from the protocol were fostered to two experimental females; these pups were excluded from the analyses).

Stimulus

The odour stimulus was the female rabbit's mammary pheromone, i.e. 2-methylbut-2-enal (Sigma-Aldrich, Saint-Quentin Fallavier, France). It was diluted in distilled water at 10⁻⁶ g/ml, a concentration optimally efficient to release typical responses in 2- to 3-day-old pups (Coureaud *et al.*, 2004). The stimulus was presented to the pups' nose on a glass-stick (length × diameter: 15 × 0.3 cm), dipped in a standardised manner into the pheromonal solution.

Behavioural assay and variables

The responsiveness to the mammary pheromone was assayed on individual pups on day 1 ($n = 293$), then on day

3 ($n = 288$ pups; five pups died between days 1 and 3). The behavioural assay (e.g. Coureaud *et al.*, 2003 and 2006a; Schaal *et al.*, 2003) consisted in holding a pup in one gloved hand with its head left free to move. When the pup came to rest, the tip of the glass-stick carrying the mammary pheromone was positioned 0.5 cm in front of its muzzle, avoiding any contact. The duration of the odour presentation was set at 15 s.

Three levels of response were considered: (1) no response (0), meaning that the pup expressed merely sniffing or at most random head movements; (2) searching (S), that is extension of the head towards the stimulus, with low amplitude horizontal and vertical scanning movements leading to the eventual contact with the stick; and (3) grasping (G), that is searching movements terminated by one or several oral seizing of the stick. The pup responsiveness to the mammary pheromone was tested 10 min before the daily nursing (occurring, for a given litter, each day at regular time, between 0830 and 1130 h). Therefore, the nest box was closed in the evening of the day preceding the test.

Milk intake, growth and mortality

Immediately after behavioural testing, the pups of a litter were individually weighed (balance Balea P113 445; accuracy: 0.1 g; Saint Mathieu de Trévières, France) before being put back to the nest.

Ten minutes later, the nest was opened to allow the female's entry for suckling her litter. Almost all females readily jumped into the nest and nursed (two females on day 1 and four females on day 3 were reluctant to nurse during the time scheduled; their litters were accordingly excluded from the milk intake analyses). Five minutes after the end of the nursing episode, the pups were weighed individually to determine their milk intake. The weight was also measured after nursing on day 21. Mortality was checked daily from birth through day 21.

Regarding the does, their weight was controlled immediately after parturition, then on days 1 and 3, and their milk production was estimated on days 1 and 3 by the milk intake of their litter (measured by weighing the entire litter before/after the daily nursing; Maertens *et al.*, 2006). This allowed comparing the females' lactation in the present conditions with that of females from the same strain bred in other rabbitries.

Statistical analyses

Analyses were made with the Statistical Analysis Systems Institute (1985). The rates of responsiveness to the mammary pheromone and of pup mortality were analysed using the McNemar test (for related samples) and Fisher's exact test (for independent samples). Milk intake was examined by analysis of variance, using the general linear model. To run the analyses, pup weight was transformed in three-level variables (light, medium, heavy; referred to

Table 1 Milk intake (g) on day 1 as a function of the responsiveness to the mammary pheromone (MP) and weight on day 1

	Responsiveness to the MP on day 1 [†]			Pr > f
	0 (n)	S (n)	G (n)	
Weight on day 1				
<48 g	0.2 ^{a,A} ± 0.2 (5)	7.4 ^b ± 1.4 (14)	7.2 ^{b,A} ± 0.5 (73)	<0.01
48–57 g	7.0 ^B ± 1.5 (11)	9.6 ± 1.4 (23)	9.2 ^B ± 0.7 (59)	NS
>57 g	10.3 ^B ± 2.1 (11)	8.8 ± 1.4 (21)	10.8 ^B ± 0.7 (56)	NS
Pr > f	<0.05	NS	<0.001	

^{a,b,A,B}Means with different superscripts differ at $P < 0.05$. The interaction between weight and responsiveness to the MP was significant ($P < 0.05$).
[†](0) no response, (S) searching, (G) grasping.

three subgroups of pups in Tables 1–4). The model was as follows:

$$Y_{ijk} = \mu + Li + Rj + Wk + (RW)jk + e_{ijk},$$

where L is the litter effect (30 levels corresponding to the 30 litters); R the responsiveness to the mammary pheromone (three levels: 0, S and G); W the pup weight class (three levels: light, medium and heavy); RW the interaction between the responsiveness to the pheromone and the weight at birth; and e the residual error. When the interaction was significant, the effect of the responsiveness to the mammary pheromone was analysed for each weight class.

For the growth performance (i.e. days 1 to 21 and days 3 to 21 weight gains), the milk intake on days 1 or 3 was added to the previous model. To that goal, the milk intake was also transformed in a three-level variable (low, medium, high; referred to three subgroups in Tables 3c and 4c).

Results

General performances of females and pups

The mean weight of the females at parturition was 3508 ± 60 g, and the mean litter size and weight at birth were 12.5 ± 0.4 and 635 ± 36 g, respectively. The proportion of pups dead at birth was $4 \pm 1.6\%$ (following the trends observed in rabbit breeding; Xiccato *et al.*, 2004).

Regarding the sucking success, 14% of the pups did not get milk on day 1. This proportion decreased significantly on day 3 to reach 2% ($P < 0.001$). The milk intake per pup was, on average, 8.7 ± 0.3 g on day 1 and 15.7 ± 0.4 g on day 3. Regarding the females, the total milk production per doe was 85 ± 4.4 and 153 ± 5.5 g on days 1 and 3, with a high inter-individual variability (range: 36–130 g on day 1 and 68–203 g on day 3).

Along the day 0–21 period, the pup mortality was 6.8% ($n \pm 20/293$). Most of the pups (60%) died during the first postnatal week, a common observation in rabbitries (e.g. Venge, 1963; Lebas, 1974; Coureaud *et al.*, 2000a; Drummond *et al.*, 2000). The mean weight gain of pups across the day 1–21 period, calculated by subtracting each pup individual weight on day 1 from the weight on day 21, was

Table 2 Milk intake (g) on day 3 as a function of the responsiveness to the mammary pheromone (MP) and weight on day 3

	Responsiveness to the MP on day 3 [†]			Pr > f
	0 (n)	S (n)	G (n)	
Weight on day 3				
<64 g	13.7 ± 0.9 (6)	– (0)	12.4 ^A ± 0.8 (79)	NS
64–77 g	17.6 ± 1.9 (7)	16.0 ± 3.0 (2)	16.1 ^B ± 0.6 (76)	NS
>77 g	17.6 ± 1.3 (8)	23.3 ± 2.1 (4)	18.5 ^C ± 0.7 (68)	NS
Pr > f	NS	–	<0.001	

^{A,B,C}Means with different superscripts differ at $P < 0.05$. The interaction between weight and responsiveness to the MP was not significant ($P = 0.53$).
[†](0) no response, (S) searching, (G) grasping.

275 ± 4 g (this was also in line with data from other colonies raising the same breed; Xiccato *et al.*, 2004).

Immediate impact of pup responsiveness to the mammary pheromone

Among the 288 pups tested on day 1 and then on day 3, 8.7% and 7.2%, respectively, did not respond to the mammary pheromone. This rate of non-response was stable between days 1 and 3 ($\chi^2 < 0.5$, $P > 0.05$). The remaining pups responded strongly to the mammary pheromone by searching (S), and more often by grasping (G) (more than 70% on days 1 and 3; comparisons between S and G pups: $\chi^2 > 137$, $P < 0.001$ on day 1 as on day 3). The rate of pups that only searched decreased from days 1 to 3 (21.2% to 3.1%; $\chi^2 = 43.3$, $P < 0.001$) while the pups that grasped increased correlatively (70.1% to 89.6%; $\chi^2 = 49.1$, $P < 0.001$; Figure 1). Taken together, the pups responding to the pheromone (by S or G) were as frequent on day 1 than on day 3 (91.3% *v.* 92.8%; $\chi^2 = 0.5$, $P > 0.05$).

The milk intake on day 1 was affected by an interactive effect of the responsiveness to the mammary pheromone and the weight of the pups ($P < 0.05$). In light pups (<48 g), the sucking success was low for those who were unresponsive, as compared with the responding pups displaying either searching or grasping (comparisons between 0, S and G pups: $P < 0.01$). In contrast, the milk intake of

Table 3 Growth and mortality of pups between days 1 and 21 according to the responsiveness to the mammary pheromone (MP), weight and milk intake on day 1

	(a) Responsiveness to the MP on day 1 [†]			Pr > f
	0	S	G	
Mortality 1–7 days (%)	14.3 ^a (4/28)	0 ^b (0/61)	3.9 ^b (8/204)	<0.05
Mortality 8–21 days (%)	0 (0/24)	1.6 (1/61)	3.6 (7/196)	NS
Mortality 1–21 days (%)	14.3 (4/28)	1.6 (1/61)	7.4 ^b (15/204)	=0.06
Weight gain 1–21 days (g)	298 ± 12 (24)	286 ± 7 (60)	269 ± 4 (189)	NS
	(b) Weight on day 1 (g)			
	<48 g	48–57 g	>57 g	
Mortality 1–7 days (%)	9.2 ^a (9/98)	2.1 ^{ab} (2/97)	1.0 ^b (1/98)	<0.05
Mortality 8–21 days (%)	8.5 ^a (8/89)	0 ^b (0/95)	0 ^b (0/97)	<0.001
Mortality 1–21 days (%)	17.3 ^a (17/98)	2.1 ^b (2/97)	1.0 ^b (1/98)	<0.001
Weight gain 1–21 days (g)	228 ^a ± 6 (81)	280 ^b ± 4 (95)	310 ^c ± 6 (97)	<0.001
	(c) Milk intake on day 1 (g)			
	<7 g	7–11 g	>11 g	
Mortality 1–7 days (%)	12.0 ^a (11/92)	0 ^b (0/86)	0 ^b (0/95)	<0.001
Mortality 8–21 days (%)	6.2 ^a (5/81)	1.2 ^b (1/86)	0 ^b (0/95)	<0.01
Mortality 1–21 days (%)	17.4 ^a (16/92)	1.2 ^b (1/86)	0 ^b (0/95)	<0.001
Weight gain 1–21 days (g)	254 ± 7 (76)	273 ^b ± 7 (85)	291 ^c ± 6 (95)	<0.001

^{a,b,c}Means with different superscripts differ at $P < 0.05$. Regarding the weight gain, the interactions between the pup weight, the responsiveness to the MP and the milk intake on day 1 and on day 3 were not significant.

[†](0) no response, (S) searching, (G) grasping.

Table 4 Growth and mortality of pups between days 3 and 21 according to the responsiveness to the mammary pheromone (MP), weight and milk intake on day 3

	(a) Responsiveness to the MP on day 3 [†]			Pr > f
	0	S	G	
Mortality 3–7 days (%)	0 (0/21)	0 (0/9)	2.7 (7/258)	NS
Mortality 8–21 days (%)	0 (0/21)	0 (0/9)	3.2 (8/251)	NS
Mortality 3–21 days (%)	0 (0/21)	0 (0/9)	5.8 (15/258)	NS
Weight gain 3–21 days (g)	276 ± 13 (21)	308 ± 19 (9)	254 ± 4 (243)	NS
	(b) Weight on day 3 (g)			
	<64 g	64–77 g	>77 g	
Mortality 3–7 days (%)	6.3 ^a (6/96)	1.0 ^b (1/96)	0 ^b (0/96)	<0.05
Mortality 8–21 days (%)	6.7 ^a (6/90)	2.1 ^{ab} (2/95)	0 ^b (0/96)	<0.05
Mortality 3–21 days (%)	12.5 ^a (12/96)	3.1 ^b (3/96)	0 ^b (0/96)	<0.001
Weight gain 3–21 days (g)	222a ± 6 (84)	262 ^b ± 5 (93)	385 ^c ± 6 (96)	<0.001
	(c) Milk intake on day 3 (g)			
	<15 g	15–18 g	>18 g	
Mortality 3–7 days (%)	5.6 ^a (5/90)	0 ^b (0/76)	0 ^b (0/84)	<0.05
Mortality 8–21 days (%)	5.9 (5/85)	1.3 (1/76)	0 (0/84)	<0.05
Mortality 3–21 days (%)	11.1 ^a (10/90)	1.3 ^b (1/76)	0 ^b (0/84)	<0.001
Weight gain 3–21 days (g)	235 ± 6 (80)	259 ^a ± 5 (75)	288 ^b ± 6 (84)	<0.05

^{a,b,c}Means with different superscripts differ at $P < 0.05$. Regarding the weight gain, the interactions between the pup weight, the responsiveness to the MP and the milk intake on day 1 and on day 3 were not significant.

[†](0) no response, (S) searching, (G) grasping.




Figure 1 Frequency (%) of rabbit newborns displaying only sniffing (0), sniffing and searching (S), or sniffing, searching and grasping (G) to the mammary pheromone on days 1 and 3 ($n=288$ pups, from 30 litters, tested on both days). a, b, c and * indicate significant differences ($P<0.05$) among types of responses on a given day, or between days for a given response, respectively.

the medium and heavy pups (≥ 48 g) was similar whatever the responsiveness to the mammary pheromone (Table 1). The weight by itself was also influential, at least in unresponsive pups or in pups responding by grasping. In these subgroups, the lightest pups were less successful in gaining milk than the medium and heaviest pups (difference in milk gain between light *v.* medium, and light *v.* heavy pups: -97.2% and -98.1% for unresponsive pups, and -22% and -33% for G pups; $P<0.05$ in all cases; Table 1).

The results were clearly different on day 3, however. The interaction effect of the reactivity to the mammary pheromone and the pre-nursing weight on milk intake did no more reach significance. However, milk intake was still influenced by pup weight, at least in G pups (i.e. 89.2% of the pups). Thus, lighter pups gained less milk than medium pups, which themselves were less efficient than heavier pups ($r=0.49$; $P<0.001$; Table 2).

To sum up, pup responsiveness to the mammary pheromone was linked to milk intake on day 1, but no more on day 3. This pheromone-milk intake link on day 1 is effective in the lightest pups, leading to significantly lower milk intake in unresponsive pups.

Delayed impact of the early responsiveness to the mammary pheromone

(1) *Pup survival during the day 1–21 period.* Over the first postnatal week, pups who were unresponsive on day 1 had a higher rate of mortality (14.3%) than those who searched or grasped (0% and 3.9% for S and G pups, respectively; Table 3a). Moreover, but regardless of the response to the mammary pheromone, pup nursing weight and milk intake on day 1 were related to pup mortality: more losses were recorded for light, as opposed to medium/heavy pups (Table 3b), and for those pups who took less than 7 g of milk on day 1 (Table 3c). It may be noted that the four pups that died among the unresponsive group were lighter than the surviving pups in this group (36.0 ± 9 *v.* 59.2 ± 2 g,

$P<0.001$). The same trend was seen in the pups who responded by oral grasping to the mammary pheromone (dying *v.* surviving pups: 34.2 ± 5 *v.* 51.9 ± 1 g, respectively; $P<0.001$).

After day 7, the responsiveness to the mammary pheromone initially displayed on day 1 did no more appear to impact on pup survival. However, the mortality during the day 8–21 period, and more generally during the day 1–21 period, was influenced by the weight and milk intake of the pups on day 1: the pups that died were the lightest and the most unsuccessful in gaining milk (Table 3b and c).

On day 3, the link between pup pheromonal response and survival seemed to fade (for days 3 to 7, 8 to 21 and 3 to 21; Table 4a). However, the pup weight and milk intake on day 3 remained associated with survival on the following days, in the sense that the lightest and least successful suckers survived less than the other pups (Table 4b and c).

(2) *Pup growth during the day 1–21 period.* The mean weight (before nursing) of the pups was 52 ± 0.7 g on day 1, 70 ± 1 g on day 3 and 328 ± 4.1 g on day 21. Under our conditions, no link appeared between the day-1 response to the mammary pheromone and the weight gain of the pups until day 21 (Table 3a). However, the weight and milk intake had a strong impact, a low weight or a weak sucking success on day 1 affecting significantly the growth across the day 1–21 period ($r=0.60$, $P<0.001$ and $r=0.25$, $P<0.001$, respectively; Table 3b and c).

Similarly, pup response to the mammary pheromone on day 3 did not appear to be correlated with later growth (Table 4a). In contrast, the weight gain during the day 3–21 period was influenced both by the weight and by the milk intake of the pups on day 3 ($r=0.52$, $P<0.001$ and $r=0.45$, $P<0.001$, respectively; Table 4b and c).

Discussion

The present study aimed to evaluate whether a link could be evidenced between the response of rabbit pups to the pure mammary pheromone and the rate of survival and growth in breeding conditions. It was based on the hypothesis that the unresponsiveness to the pheromone on the first postnatal day should predict non-response to this signal in the nursing context.

In the present conditions, about 90% of the tested pups responded to the mammary pheromone on days 1 or 3. This high level of response corroborates previous observations on the strong releasing potency of this compound (Coureaud *et al.*, 2003, 2004 and 2006a; Schaal *et al.*, 2003; Montigny *et al.*, 2006). In the same time, the behavioural assay using the mammary pheromone permitted to target pups that are unresponsive to the pheromone on days 1 and 3. The follow-up of these non-responsive pups allowed gauging their differential developmental trajectory in terms of milk intake, growth and viability.

A superficial look on the longitudinal results would suggest that unresponsive pups are those who will die

between birth and day 21 (6.8%). However, it comes out that the birth weight and milk intake have buffering effects in pups who do not respond to the mammary pheromone. The pups that died were not only those who failed to respond to the mammary pheromone on day 1 (80% responded). Unresponsive pups on day 1 were indeed not all ineffective in gaining milk (only 18.5% did not get milk on day 1), nor were they all dying out (85.7% survived). In fact, the impact of a non-response to the mammary pheromone rests on the pup weight (only the lightest unresponsive pups display difficulties to suck and survive) and on the period when pheromonal reactivity is assessed (the impact of non-response on day 1 is limited to the day 1 milk intake and the days 1 to 7 survival). Factors other than pheromonal perception certainly influence early survival, among which pup maturity at birth, females' parity and behavioural treatment of the nest and litter are the most noteworthy (Coudert, 1982; Verga *et al.*, 1986; Fortun-Lamothe, 1998; Canali *et al.*, 1991; Hamilton *et al.*, 1997; Coureaud *et al.*, 2000a). Lastly, it must be stated that the present behavioural assay is rather punctual, being run once for 15 s, and that it may therefore not be ubiquitously sensitive for the varied risk factors affecting viability.

However, it appears that for the lightest pups, responding to the mammary pheromone on day 1 bears a clear advantage to gain milk. Those pups are indeed as successful in sucking as their heavier, pheromone-unresponsive or responsive littermates. Thus, for the lightest pups, the day 1 pheromonal responsiveness, weight and milk intake are additive in improving viability. The impact of the two latter factors was already known (McNitt and Moody, 1987; Coureaud *et al.*, 2000b; Poigner *et al.*, 2000; Perrier *et al.*, 2003). But here, we add the perceptual ability of day-1 pups as an additional predictor, which is especially valid for the thinnest pups during the first postnatal week.

The causes of such individual differences in the early behavioural impact of the mammary pheromone are unclear. Pheromonal unresponsiveness of pups may first relate to uneven maturation among littermates in terms of perceptual, integrative or motor performances. It may also be due to birth-related factors, such as delayed recuperation from the stress of delivery or from obstructed respiration. Anyway, defective response to the pure mammary pheromone on day 1 may reflect general difficulty to respond to odour cues of the nursing female, increasing the risk to miss the first nursing sessions and to rapidly die (failing to suck on two to three consecutive nursing bouts is generally lethal; e.g. Venge, 1963; McNitt and Moody, 1987; Coureaud *et al.*, 2000b). Here, 100% of the pups who were pheromone-unresponsive on day 1 and who died during the experimental period, died effectively before day 7. However, the majority of pups who were unresponsive to the pure pheromone administered once on day 1 became responsive on day 3 and survived ($n = 22/27$), suggesting that a functional catch up is possible. Such recuperative outcome has then most chance to turn into an adaptive end in the nursing competition (e.g. Drummond *et al.*, 2000; Bautista

et al., 2005). Moreover, after the day-1 test and during the next sucking bouts (days 2 to 3), some individuals could find the nipples and gain milk in tracing circumstantial maternal odorants that they have learned as directional cues (e.g. Hudson, 1985; Coureaud *et al.*, 2006b).

To sum up, the responsiveness of newborn rabbits to the mammary pheromone appears to be an indicator for individual viability, at least for the lightest pups and when their responsiveness is checked on postnatal day 1. The use of a pheromone to probe viability in rabbitries remains now to be ascertained. Pheromone testing on day 1 may permit selection among the lightest pups those who can support the competition for milk with heavier newborns. Conversely, pheromone-response defective pups on day 1 being at higher mortality risk, they might be reunited with light pups to be artificially fed or fostered under one female, allowing alleviation of competition and thus improved welfare and survival.

Acknowledgements

The authors acknowledge the important technical assistance and generous help of B. Lamboley-Gaüzère, C. Bannelier, J. Dedapper, A. Lapanouse and P. Aymard.

References

- Bautista A, Mendoza-Degante M, Coureaud G, Martina-Gomez M and Hudson R 2005. Scramble competition in newborn domestic rabbits for an unusually restricted milk supply. *Animal Behaviour* 70, 997–1002.
- Bilkó A, Altbäcker V and Hudson R 1994. Transmission of food preference in the rabbit: the means of information transfer. *Physiology and Behavior* 56, 907–912.
- Canali E, Ferrante V, Todeschini R, Verga M and Carezzi C 1991. Rabbit nest construction and its relationship with litter development. *Applied Animal Behaviour Science* 16, 285–293.
- Coudert P 1982. Analyse de l'origine des pertes à la maternité. *Cuniculture* 45, 136–139.
- Coureaud G 2001. [Olfactory regulation of sucking in newborn rabbit: ethological and chemical characterization of a pheromonal signal]. PhD thesis, University of Paris.
- Coureaud G, Schaal B, Coudert P, Hudson R, Rideaud P and Orgeur P 2000a. Mimicking natural nursing conditions promotes early pup survival in domestic rabbits. *Ethology* 106, 207–225.
- Coureaud G, Schaal B, Coudert P, Rideaud P, Fortun-Lamothe L, Hudson R and Orgeur P 2000b. Immediate postnatal sucking in the rabbit: its influence on pup survival and growth. *Reproduction Nutrition Development* 40, 19–32.
- Coureaud G, Schaal B, Langlois D and Perrier G 2001. Responsiveness of newborn rabbits to surface odour cues from females differing in lactational state and to milk. *Animal Behaviour* 61, 153–162.
- Coureaud G, Schaal B, Hudson R, Orgeur P and Coudert P 2002. Transnatal olfactory continuity in the rabbit: behavioral evidence and short-term consequence of its disruption. *Developmental Psychobiology* 40, 372–390.
- Coureaud G, Langlois D, Perrier G and Schaal B 2003. A single-key odorant accounts for the pheromonal effect of rabbit milk: further test of the mammary pheromone's activity against a wide sample of volatiles from milk. *Chemoecology* 13, 187–192.
- Coureaud G, Langlois D, Sicard G and Schaal B 2004. Newborn rabbit responsiveness to the mammary pheromone is concentration-dependent. *Chemical Senses* 29, 341–350.
- Coureaud G, Langlois D, Perrier G and Schaal B 2006a. Convergent changes in the maternal emission and pup reception of the rabbit mammary pheromone. *Chemoecology* 16, 169–174.

- Coureaud G, Moncomble AS, Montigny D, Dewas M, Perrier G and Schaal B 2006b. A pheromone that rapidly promotes learning in the newborn. *Current Biology* 16, 1956–1961.
- Drummond H, Vázquez E, Sanchez-Colón S, Martínez-Gómez M and Hudson R 2000. Competition for milk in the domestic rabbit: survivors benefit from littermate deaths. *Ethology* 106, 511–526.
- Fortun-Lamothe L 1998. Effects of pre-mating energy intake on reproductive performance of rabbit does. *Animal Science* 66, 236–269.
- Gonzalez-Mariscal G 2007. Mother rabbits and their offspring: timing is everything. *Developmental Psychobiology* 49, 71–76.
- Gualterio L, Valentini A and Bagliacca M 1988. Effect of season and of parturition order on mortality rate at birth and in the nest. *Proceedings of the fourth world rabbit congress, Budapest, Hungary*, pp. 182–188.
- Hamilton HH, Lukefahr SD and McNitt JI 1997. Maternal nest quality and its influence on litter survival and weaning performance in commercial rabbits. *Journal of Animal Science* 75, 926–933.
- Hudson R 1985. Do newborn rabbits learn the odor stimuli releasing nipple-search behavior? *Developmental Psychobiology* 18, 575–585.
- Hudson R and Distel H 1983. Nipple location by newborn rabbits: evidence for pheromonal guidance. *Behaviour* 82, 260–275.
- Keil W, von Stralendorff F and Hudson R 1990. A behavioral bioassay for analysis of rabbit nipple-search pheromone. *Physiology and Behavior* 47, 525–529.
- Kindermann U, Gervais R and Hudson R 1991. Rapid odor conditioning in newborn rabbits: Amnesic effect of hypothermia. *Physiology and Behavior* 50, 457–460.
- Lebas F 1974. La mortalité des lapereaux sous la mère. *Etude monographique (première partie)*. *Cuniculture* 1, 8–11.
- Maertens L, Lebas F and Szendró ZS 2006. Rabbit milk: a review of quantity, quality and non-dietary affecting factors. *World Rabbit Science* 14, 205–230.
- McNitt JI and Moody GL 1987. Nest box behavior of the domestic rabbit. *Journal of Applied Rabbit Research* 10, 159–162.
- Montigny D, Coureaud G and Schaal B 2006. Newborn rabbit response to the mammary pheromone: from automatism to prandial control. *Physiology and Behavior* 89, 742–749.
- Perrier G, Jouanno M and Drouet JP 2003. Influence de l'homogénéité et de la taille de portée sur la croissance et la viabilité des lapereaux de faible poids à la naissance. In *10èmes Journées de la Recherche Cunicole* (ed. G Bolet), pp. 119–122. ITAVI, Paris, France.
- Pizzi F and Crimella C 1984. Osservazione sull'allattamento controllato in conigliocultura intensiva. *Atti della Società Italiana di Scienze Veterinarie* 38, 504–507.
- Poigner J, Szendro ZS, Lévai A, Radnai I and Biro-Németh E 2000. Effect of birth weight and litter size on growth and mortality in rabbits. *World Rabbit Science* 8, 17–22.
- Schaal B, Coureaud G, Langlois D, Giniès C, Sémon E and Perrier G 2003. Chemical and behavioural characterization of the rabbit mammary pheromone. *Nature* 424, 68–72.
- Schley P 1976. [Investigations on the artificial rearing of domestic rabbits.]. *Habilitationschrift, Universität Giessen, Germany*.
- Seitz K 1997. [Investigations on the sucking behaviour of domestic rabbit females, and on the milk intake, growth and mortality of the young.] *Doctoral Dissertation, Justus-Liebig-Universität, Giessen*.
- Statistical Analysis Systems Institute 1985. SAS version 6.12. SAS Institute Inc., Cary, USA.
- Venge O 1963. The influence of nursing behaviour and milk production on early growth in rabbits. *Animal Behaviour* 11, 500–506.
- Verga M, Canali E, Pizzi F and Crimella C 1986. Induced reactions in young rabbits of dams of different parity and reared on two different nursing schedules. *Applied Animal Behaviour Science* 16, 285–293.
- Xiccato G, Trocino A, Sartori A and Queaque PI 2004. Effect of parity order and litter weaning age on the performance and body energy balance of rabbit does. *Livestock Production Science* 85, 239–251.
- Zarrow MX, Denenberg VH and Anderson CO 1965. Rabbit: frequency of suckling in the pup. *Science* 150, 1835–1836.