

Longitudinal muscle gene expression patterns associated with differential intramuscular fat in cattle

N. J. Hudson^{1†}, A. Reverter¹, P. L. Greenwood², B. Guo¹, L. M. Cafe² and B. P. Dalrymple¹

¹Computational and Systems Biology, CSIRO Agriculture Flagship, 306 Carmody Road, St Lucia, Brisbane, QLD 4067, Australia; ²NSW Department of Primary Industries Beef Industry Centre, University of New England, Armidale, NSW 2350, Australia

(Received 25 March 2014; Accepted 10 September 2014; First published online 13 November 2014)

Intramuscular fat (IMF) can improve meat product quality through its impact on flavour and juiciness. High marbling cuts can command premium prices in some countries and grading systems, but there is substantial cost involved in choosing to grain feed animals in an effort to deposit more IMF. There would be value in developing methods to predict predisposition to 'marble' well. Unfortunately, the biological mechanisms underpinning marbling remain a mystery: the key adipocyte cell populations have not been defined, there are no reliable DNA markers, no known (if any) causal mutations and gene expression analyses in the main have tended to characterise increases in expression of end-point fat metabolism proteins such as the fatty acid-binding proteins. To shed light on expression-based markers of marbling potential, we contrasted LD gene expression in high IMF Wagyu cross animals with a low IMF Piedmontese cross at various time points. The expected divergence in the fat metabolism genes FABP4, THRSP, CIDEA and ACACA between the breeds occurs surprisingly late in postnatal development at about 20 months. On the other hand, divergent expression of WISP2, RAI14 and CYP4F2 was discovered in animals at or before 12 months of age, suggesting these genes may have potential as earlier predictors of marbling potential. In line with other researchers, we found intriguing links between IMF development and connective tissue remodelling. WISP2 – a novel adipokine highly expressed and secreted by adipose precursor cells and an inhibitor of the pro-fibrotic connective tissue growth factor – emerges as a particularly attractive candidate. It is relatively upregulated in high marbling Wagyu before admission to feedlotting, somewhere between 7 and 12 months. This difference is subsequently maintained until 25 months, but not thereafter. RAI14, thought to play a role in porcine adipocyte differentiation and with links to retinoic acid metabolism, has an unusual expression profile. Its expression level increases monotonically with postnatal development, and is always higher in Wagyu than Piedmontese. Strong, sustained upregulation of the anti-inflammatory CYP4F2 in Piedmontese is consistent with Wagyu adiposity being a pro-inflammatory state. Application of regulatory impact factor analysis, a network method for identifying causal effector molecules, suggests marbling roles for transcription factors previously implicated in (1) the formation of liposarcoma (unconstrained fatty masses) (YEATS4, MDM2), (2) adipogenesis (CREBL2, SP1, STAT1) and (3) inflammation (ISGF3G, HOXB13, PML).

Keywords: marbling, cattle, Piedmontese, Wagyu, adipocyte

Implications

Intramuscular fat (IMF) improves meat product quality through enhanced flavour and juiciness. High marbling cuts command premium prices in some countries and grading systems, but there is substantial monetary cost involved in grain feeding animals to enhance IMF. There is need for new methods to rationally select high-potential animals for sound investment in long feeding. We also see value in developing a better understanding of fundamental marbling biology. In this article, we contrast genome-wide gene expression patterns in two breed crosses divergent for marbling in an

attempt to identify new early markers of marbling potential and help improve our biological understanding in this enigmatic area. We propose skeletal muscle expression of *WISP2* between 7 and 12 months postnatal to be an attractive new candidate for future marbling potential, presumably by quantifying adipocyte precursor cell number and/or activity.

Introduction

In the beef industry, intramuscular fat (IMF) – or 'marbling' – is highly valued as it confers desirable organoleptic properties of taste, juiciness and tenderness that command premium prices in international markets. A better understanding of the

[†] E-mail: Nick.Hudson@csiro.au

molecular drivers of marbling may ultimately help reduce variation in carcass composition and also ensure high cost grain during extended feedlotting is not wasted on low IMF potential individuals. IMF manifests as white flecks or streaks of adipose tissue between bundles of muscle fibres (Harper and Pethick, 2004). From several biological perspectives cattle marbling remains intractable. It has not yielded to genome-wide association studies – that is, there are no reliable DNA markers that explain substantial amounts of phenotypic across populations and breeds (Pannier *et al.*, 2010). However, application of the latest thinking in network theory has been exploited to reverse engineer a plausible set of contributing regulatory molecules (*PPARGC1A*, *HNF4G*, *FOXP3*) based on single nucleotide polymorphism association data across three breed types, *Bos taurus*, *Bos indicus* and tropical composites (Ramayo-Caldas *et al.*, 2014).

Similarly, there are no known causal mutations, the contributing cell populations have not been well defined (Harper and Pethick, 2004; Bonnet *et al.*, 2010), and gene expression analyses have tended to document increases in the expression of end-point fatty acid metabolism proteins such as members of the fatty acid-binding protein family (Wang *et al.*, 2009; De Jager *et al.*, 2013). However, genetics is a strong driver of marbling, such that in elite Wagyu animals IMF can exceed 50% in skeletal muscle (Gotoh *et al.*, 2009), whereas in other breeds such as Brahman the figure averages <5%. IMF is also influenced by environmental factors such as diet, particularly through vitamin A restriction. Indeed, the retinoic acid (RA) axis is involved in adipogenic phenotypes in general (Bonet *et al.*, 2012).

Here, we make use of a previously analysed gene expression experiment contrasting high marbling Wagyu × Hereford (W × H) crossbred cattle *v.* lower marbling Piedmontese × Hereford (P × H) crosses at 10 developmental time points. These include three prenatal time points coincident with the two major waves of myogenesis, and the onset and end of functional differentiation (Hudson *et al.*, 2013).

The comparison of high and low marbling cattle breeds allows an examination of early developmental processes, including the impact of precursor cell populations (as inferred via the gene expression profiles). In addition, we can also examine later environmental influences such as major nutritional transitions relating to birth, weaning and feedlotting. The multi-dimensional nature of the experimental design also allows more sophisticated analyses than simple computation of differential expression (DE).

High throughput gene expression analysis has previously been used to contrast these two breeds in the context of differential fat development (Wang *et al.*, 2009). However, the previous publication (Wang *et al.*, 2009) was based on a far less comprehensive cDNA array. Highlighted genes included *THRSP* and *FABP4*. In a separate experiment based on 48 Brahman cattle, we correlated individual gene expression measures to individual IMF% (De Jager *et al.*, 2013). This approach identified the gene set that best correlates to IMF% is triacylglyceride (TAG) deposition, and includes *FABP4*. A number of other groups have identified

similar sets of genes in various production species based on similar analyses performed on similar experimental designs (Saez *et al.*, 2009; Li *et al.*, 2013; Wang *et al.*, 2013; Ren *et al.*, 2014). However, to date the simplest measures of DE and correlations of abundance to phenotype have yielded only modest regulatory insight. Various analytical reasons can be given for this (Hudson *et al.*, 2012).

The purpose of this study is to identify early muscle gene expression markers for future marbling potential and also to shed light on molecular mechanisms of marbling development. Here, a combination of a denser array platform detailing the expression of most of the transcriptome, some targeted analytical questions, coupled with application of a set of higher-order network inference algorithms uniquely developed by our group (1) reinforces a likely link between marbling adipocyte biology and extracellular matrix (ECM) remodelling (*WISP2*), (2) allows the detection of novel candidate early diagnostic molecular markers for future marbling potential (e.g. *WISP2*, *RAI14*), (3) is consistent with a relatively pro-inflammatory state in the Wagyu compared with Piedmontese (*CYP4F2*) and (4) identifies a small set of DNA-binding transcription factors (TF) that have apparently been re-wired into a 'pro-marbling' regulatory circuit in the Wagyu breed cross (e.g. *ISGF3G*, *YEATS4*, *MDM2* and others).

Material and methods

Details relating to the animal experiment, RNA extraction, microarray hybridisation and data normalisation can be found in the studies by Wang *et al.* (2005 and 2009), Lehnert *et al.* (2007), Hudson *et al.* (2009a and 2009b). As previously described in a study by Lehnert *et al.* (2007), Hereford cows were artificially inseminated or mated to one of five different Wagyu sires (W × H) or one of six different Piedmontese sires (P × H). All Piedmontese sires were homozygous for the *MSTN* mutation in exon 3. In all, three fetuses from each sire breed were acquired by caesarean section; three newborn calves per sire breed were euthanised by lethal injection within 24 h of birth. *Longissimus* muscle was dissected immediately after death of the fetuses or calves, and tissue samples frozen in liquid nitrogen. The animals were bred and grown within pasture-based nutritional systems followed by feedlot finishing (Greenwood *et al.*, 2006). Each cohort grazed as a single group during backgrounding. Feedlotting took place at the Tullimba feedlot (Kingston, NSW). Details regarding nutrition and growth can be found in the study by Greenwood *et al.* (2006). The prenatal time points coincide with primary myogenesis, secondary myogenesis and onset of functional differentiation. The remainder comprised birth, 3 month, 7 month (weaning), 12 month, 20 month, 25 month (feedlot entry) and 30 month animals. In attempting to infer the development of marbling in cattle, we applied a number of analyses on both within- and between-breed basis. The various analytical steps are documented below.

Identification of expression-based markers for current marbling status

To prioritise genes whose expression profile might reflect current marbling status, we made an assumption that lipid content would increase its proportional representation in a given mass of postnatal bovine muscle over developmental time, via hyperplasia and hypertrophy of marbling adipocytes. Predicated on this assumption, we asked the question ‘the expression of which genes possess a positive postnatal slope in both breeds, with a steeper slope in the Wagyu?’. Within each breed and gene, we fitted a simple linear regression as follows:

$$y = a + x\beta + e$$

where y is vector of expression abundance across the seven postnatal time points; a the y-axis intercept; x the index of the seven time points (without loss of generality assumed equally spaced); β the regression slope; and e the random residual associated with the goodness of fit. We selected genes for which the P -value associated with the estimate of β was <0.01 in both populations and the magnitude of the estimate greater in Wagyu than in Piedmontese. We then ranked the output on the extent of the slope difference (the ‘differential regression’) and submitted the list to GOrilla for functional enrichment analysis. We also asked the related – but more stringent – question ‘which genes show a monotonic increase at all postnatal time points, and are always higher in the Wagyu compared to the Piedmontese at each time point?’.

Identification of expression-based markers for future marbling potential

To identify genes whose expression profiles might allow a better prediction of future marbling potential, we examined the expression profiles for genes that diverge between the breeds earlier in postnatal development than the end-point fat metabolism genes that diverge only at 20 to 25 months. Given the proposed role of ECM remodelling in preparing the tissue environment for invasion and development of marbling adipocytes (Harper and Pethick, 2004), we focused on genes whose encoded proteins have been annotated as ECM remodellers. Lists of significantly DE genes were identified at each postnatal time point as previously described and explored for functional enrichment. Gene annotation to ECM was determined by uploading the gene lists to DAVID software (Huang da *et al.*, 2009) and augmented by literature searches and manual curation.

Wagyu v. Piedmontese DE

To further understand the transcriptomic differences between Wagyu and Piedmontese muscle, we examined those genes that were both significantly differentially expressed, as well as being annotated by DAVID software as involved in fat metabolism. In a time course experiment there are a number of different legitimate ways of computing DE. For this analysis, we were interested in ‘overall’ patterns of DE so we computed a ‘surface-based’ version of DE as previously described

according to a study by Hudson *et al.* (2012) that incorporates the shape of the profile across the 10 points. The surface of the DE of the i th gene, (SDE_i), is defined as the sum of squared differences between the expression of the i th gene in the two breeds (P, Piedmontese; W, Wagyu) across the 10 time points. This sum follows a χ^2 distribution with 9 d.f.:

$$SDE_i = \sum_{t=1}^{10} (P_{i,t} - W_{i,t})^2 \sim \chi_{9d.f.}^2$$

We have previously discovered that ‘weighing’ DE for average abundance (by multiplying the two) gives a much more compelling description of the change in phenotype (Hudson *et al.*, 2009a and 2013), so we applied this correction to the SDE measure. We ranked the output on extent of this weighted SDE metric and explored it for genes annotated as being involved in fat metabolism.

The transcriptional regulation of marbling

Although molecular correlates of marbling have been previously identified, far less is known about the causal transcriptional regulation of the phenotype. To identify transcription factor regulatory circuits that have apparently been re-wired in the Wagyu breed, we applied our recently developed regulatory impact factor (RIF) analysis (Hudson *et al.*, 2009a and 2013; Reverter *et al.*, 2010). This procedure exploits global patterns of differential co-expression (or ‘differential wiring’) to infer those regulatory molecules whose behaviour is systematically different in a contrast of interest, in this case the Wagyu *v.* Piedmontese breed crosses. Herein, the experimental contrast was P *v.* W (i.e. Piedmontese *v.* Wagyu) and the RIF metrics for the r th regulator ($r = 1, 2, \dots, 920$) were computed using the following formulae:

$$RIF1_r = \frac{1}{n_{DE}} \sum_{j=1}^{j=n_{DE}} x_j \times d_j \times DC_{rj}^2$$

and

$$RIF2_r = \frac{1}{n_{DE}} \sum_{j=1}^{j=n_{DE}} \left[\left(x_j^P \times r_{rj}^P \right)^2 - \left(x_j^W \times r_{rj}^W \right)^2 \right]$$

where n_{DE} represented the number of DE genes; x_j the average expression of the j th DE gene across all time points; d_j the DE of the j th gene in the P *v.* W contrast; DC_{rj} the differential co-expression between the r th regulator and the j th DE gene, and computed from the difference between r_{rj}^P and r_{rj}^W , the correlation co-expression between the r th regulator and the j th DE gene in the P and W samples, respectively; finally, x_j^P and x_j^W the average expression of the j th DE or TS gene in the P and W samples, respectively.

Results

All normalised gene expression values are expressed on a log 2 scale. The profile of postnatal IMF content over development is schematically characterised in Figure 1a.




Figure 1 A schematic diagram illustrating the early and more rapid accumulation of marbling fat in Wagyu (red) compared with Piedmontese (blue) at the phenotype level (a). Normalised mean expression levels (expressed as log₂ values) mRNA expression profiles for *FABP4* representing lipid co-expression module 1 (b) and *S100G* representing lipid co-expression module 2 (c) show a marked breed divergence relatively late (25 months) before the profiles tend to reconverge at 30 months at a time even though the intramuscular fat phenotype itself is most divergent at this time.

This was based on the study by Pethick *et al.* (2004), which documented the relationship between carcass weight and IMF% in crossbred Angus × Hereford, Angus and Japanese Black × Holstein cross cattle. We previously identified two closely related sets of fat metabolism genes whose expression was highly correlated (co-expressed) across development (Hudson *et al.*, 2009b). One gene sets comprised *ADIPOQ*, *PLS1*, *PLIN1*, *CIDEA*, *FABP4* (Figure 1b) and *TUSC5*, whereas

the other *ADIG*, *THRSP* and *S100G* (Figure 1c). The expression of these genes reflects TAG deposition rate, which reflects current IMF% in some circumstances.

Expression-based markers for current marbling status and deposition rate

In an attempt to identify estimators of the IMF contrast between the W × H and P × H crosses, we identified those genes that showed a positive regression in the postnatal samples of both, but which was steeper in the Wagyu, then ranked the list on the extent of the breed difference. We submitted this ranked list to GOrilla (Eden *et al.*, 2007) and explored it for functional enrichment. This enriched for fatty acid metabolism with *THRSP*, *PCK1*, *ADIPOQ*, *ADIG*, *SCD*, *GOS2*, *ACSM1*, *PLIN1*, *CIDEA* and *TUSC5*, all in the top 25 (Table 1; Supplementary Material S1¹). *THRSP* was ranked in first position illustrating that it met this criterion the best. From this perspective, one could conclude that *THRSP* is the most robust of the end-point characterisers of current marbling status in contrasting these two breeds, in line with a previous publication based on a cDNA array (Wang *et al.*, 2009). However, further exploration of these profiles, including *THRSP*, showed they did not increase monotonically at each postnatal time point and also that the profiles converged during feedlotting, just before slaughter, when the difference in the marbling phenotype is at its most apparent (Figure 2a). This implies the profiles do not provide an unequivocal indication of current IMF%, but are influenced by environmental factors. As previously suggested they most likely represent TAG deposition rate.

In an attempt to overcome this possible limitation, we asked the related question 'which genes, if any, increase expression monotonically at all postnatal time points (i.e. 280 days through to 30 months) in both breeds, and are also always relatively higher in W × H than P × H at each time point'. We asked this question on the basis that we know IMF content to increase over postnatal time. Only two genes met these very stringent criteria (*LACTB2* and *RAI14*). Of these, retinoic acid-induced 14 (*RAI14*) appears particularly noteworthy as it has a potential functional link to marbling through its involvement in both the RA axis and adipocyte biology. The maintenance of the cross divergence in the expression profile until 30 months contrasts with the fatty acid metabolism genes (Figure 2d).

Expression-based prognostics for future marbling potential

In order to identify a set of genes that might better predict future marbling potential, we built an analysis on the

¹Within each breed and gene, we fitted a simple linear regression as follows: $y = a + x < \beta > + e$, where 'y' is the vector of expression abundance across the seven postnatal time points; 'a' the y-axis intercept; 'x' the index of the seven time points (without loss of generality assumed equally spaced); '< β >' the regression of the slope; and 'e' the random residual associated with the goodness of fit. We selected genes for which the P-value associated with the estimate of '< β >' was 0.01 in both populations and the magnitude of the estimate greater in Wagyu than Piedmontese. We then ranked the output on the extent of the slope difference (the differential regression) and submitted the list to GOrilla for functional enrichment analysis.

Table 1 Top 10 extreme 'differential regression' Wagyu minus Piedmontese in the postnatal samples

Gene name	Gene function
<i>THRSP</i>	Thyroid hormone responsive, highly expressed in adipocytes, regulation of triglyceride synthesis
<i>PCK1</i>	Phosphoenolpyruvate carboxykinase 1, regulation of gluconeogenesis and adipogenesis, PPAR signalling
<i>CACNA2D3</i>	Voltage-dependent calcium channel
<i>FGG</i>	Fibrinogen γ -chain, complement and coagulation
<i>ADIPOQ</i>	Exclusively expressed in adipocytes, the encoded protein circulates in the plasma
<i>H2A.1</i>	Histone cluster 1, H2AM, nucleosome structure
<i>FBXO32</i>	Atrogin, role in muscle atrophy, phosphorylation-dependent ubiquitination
<i>ADIG</i>	Adipocyte-specific factor, plays a role in adipocyte differentiation
<i>HIST1H4I</i>	Histone cluster 1, H24I, nucleosome structure
<i>MAL2</i>	Component of lipid rafts, found in cytoplasm and membrane




Figure 2 Microarray-based normalised mean expression levels (expressed as log₂ values) of genes encoding proteins annotated as being involved in fat metabolism that are divergent between the breeds based on the following criteria: differential regression in the postnatal samples illustrated by *THRSP* (a), extreme surface differential expression illustrated by *CYP4F2* (b), early postnatal divergence and role in connective tissue remodelling illustrated by *WISP2* (c) and monotonic increase in all the postnatal time points in both breed crosses, but always higher in the Wagyu illustrated by *RAI14* (d). The horizontal arrow in (a) denotes all the postnatal time points, that is, from birth to 30 months. The standard errors on or between gene within time point basis ranged from 1.13 to 1.19.

assumption that ECM remodelling most likely precedes marbling, providing a favourable tissue environmental niche for the marbling adipocytes to divide and grow. Consequently, we aimed to identify those genes which both diverge between the breeds earlier than the classic end-point fatty acid metabolism genes (i.e. before 20 months), prioritising those (if any)

annotated as being involved in matrix remodelling. This analysis prioritised *WISP2* (Figure 2c).

W × *H* v. *P* × *H* DE

In the absence of any particular expectation of expression profile in the two crosses, we also explored overall patterns

Table 2 Top 10 extreme 'surface DE' accounting for abundance when comparing Wagyu v. Piedmontese muscle

Gene name (rank)	Gene function
<i>CYP4F2</i> (39)	Fatty acid and arachidonic acid metabolism. Inactivates the pro-inflammatory leukotriene B4
<i>FABP4</i> (64)	Binds long chain fatty acids, role in the uptake, transport and metabolism of lipids
<i>ANGPTL4</i> (75)	Encodes a serum hormone directly involved in glucose homeostasis, a target of the peroxisome proliferator activators
<i>SCD</i> (79)	Catalyses the rate limiting step in the synthesis of unsaturated fatty acids
<i>FASN</i> (95)	Catalyses the synthesis of palmitate from acetyl CoA into long chain saturated fatty acids
<i>ELOVL6</i> (126)	A fatty acid elongase that uses malonyl CoA as a two carbon donor in the rate limiting step of fatty acid elongation
<i>LPL</i> (144)	Triglyceride hydrolase
<i>ADIPOQ</i> (156)	Encodes a circulating protein involved in metabolism
<i>PPARG-TSEN2</i> (161)	Regulation of adipocyte differentiation
<i>VLDLR</i> (166)	VLDL triglyceride metabolism

DE = differential expression; VLDL = very low-density lipoprotein.

The illustration of those genes annotated as encoding proteins with roles in fat metabolism. All but *CYP4F2* are relatively upregulated in high IMF Wagyu.

of DE of genes encoding proteins annotated as being involved in fatty acid metabolism. We ranked on the extent of surface DE. This prioritised the following genes encoding products with roles in fat metabolism as being most informative between breeds: *CYP4F2* (Figure 2b), *VLDLR*, *ANGPTL4*, *FABP4*, *SCD*, *FASN*, *ELOVL6*, *LPL*, *ADIPOQ* and *PPARG-TSEN2* (Table 2; Supplementary Material S2²). Except for the first three, they are all 'end-point' genes (De Jager *et al.*, 2013). *CYP4F2*, the top-ranked gene, is most divergent between the two crosses at 7 months. However, unlike the other fat metabolism genes, which were more highly expressed in W × H animals, it was actually 16-fold higher in the lower IMF P × H animals. A number of other cytochrome P450 family members (e.g. *CYP1A1*, *CYP4B1*) are also highly DE between the breeds, although the direction of DE is not always consistent.

The transcriptional regulation of marbling

We ranked the RIF output, and explored those TF among the extremes of the ranked list that have been annotated as being involved in the transcriptional regulation of fat metabolism (Table 3; Supplementary Material S3³). As previously reported (Hudson *et al.*, 2009a), *MSTN* possesses the most extreme differential network value, reflecting the site of the causal mutation in Piedmontese, and its position as pre-eminent driver of the breed phenotype difference. Further manual exploration of this list with a view to identifying transcriptional regulators involved in adipogenic phenotypes highlighted *YEATS4* (16th), *STAT1*, (18th) and *MDM2* (34th).

To provide the genome-wide context for the prioritised DE genes, we plotted MA plots at 7, 12 and 25 months

²The average DE column (column D) indicates whether the gene is up or down on average. The DE is expressed Piedmontese minus Wagyu, so negative values represent higher expression in Wagyu. The top 500 corrected surface DE ranged from 63 to 886 with an average of 123. The original data are intensity signals, so this derived measure lacks a physical unit.

³RIF was calculated following the study by Hudson *et al.* (2009a). So we could rank on absolute RIF score, we squared the values to yield positive values. Genes encoding proteins annotated as involved in fat metabolism are highlighted. GDF8 is the alias for myostatin.

Table 3 Transcriptome wiring analysis

Gene name (rank)	Gene function
<i>ISGFG3</i> (8)	Inflammation
<i>CREBL2</i> (15)	Fat cell differentiation
<i>HOXB13</i> (16)	Pathologies characterised by inflammation
<i>YEATS4</i> (17)	Liposarcoma, fatty masses in skeletal muscle
<i>STAT1</i> (20)	Adipogenesis, inflammation
<i>PML</i> (25)	Inflammation
<i>NR1I2</i> (32)	Retinoic acid signalling
<i>SP1</i> (33)	Fatty acid metabolism
<i>NFYC</i> (35)	Fatty acid metabolism, cooperates with SP1
<i>MDM2</i> (42)	Liposarcoma, fatty masses in skeletal muscle

Tabulated are those transcriptional regulators who are in the top 5% (i.e. the top 46 out of 920 regulators) in terms of change of network connectivity between Piedmontese and Wagyu and who have been previously annotated as having a role relating to fat metabolism. *MSTN*, the causal double muscling mutation in Piedmontese, was ranked first by this analysis.

(Figure 3a, b and c, respectively), highlighting those genes whose expression level was detected as relevant to fat metabolism.

Discussion

In this paper, we describe a combination of expression-based analyses aimed at defining genes related to marbling development. We were particularly interested in the identification of early markers of marbling potential. In comparing the breeds, we prioritised genes showing at least one of extreme DE; dramatic changes in regression; monotonic increase postnatally but always higher in the W × H; substantial global network re-wiring as estimated by genome-wide differential co-expression. A major strength of the experimental design under consideration is its comprehensiveness (seven postnatal time points, 10 developmental time points in total) and the extent of the breed divergence in the phenotype, which provides a favourable signal to noise.




Figure 3 Microarray-based genome-wide MA plots based on normalised mean expression levels (expressed as log₂ values) at 7 month (a), 12 month (b) and 25 month (c) postnatal highlighting the extent and timing of differential expression (DE) of fat module 1 (*ADIPOQ*, *PLS1*, *PLIN1*, *CIDEA*, *FABP4* and *TUSC5*), fat module 2 (*ADIG*, *THRSP*, *S100G*), *WISP2*, *RAI14* and *CYP4F2*. Other DE functional groups not highlighted include those encoding muscle slow twitch fibre subunits, immune proteins and mitochondrial proteins.

Caveats

A weakness is that a component of the *relative* increase in marbling observed in the W×H will be attributable to the impact of the *MSTN* mutation on leanness in the Piedmontese, and that this *MSTN* signal interferes with interpretation. Thus, it is not clear whether any given fat metabolism gene expression

change reflects *MSTN*-induced leanness in Piedmontese, or enhanced marbling in Wagyu, or some combination therein. This issue potentially complicates the generalisation of the gene expression findings reported here to other breeds. In an attempt to generate a gene list that might be attributable to impaired *MSTN* signalling, we examined the gene expression output for a *MSTN* knockout mouse model (Welle *et al.*, 2009). Surprisingly, no genes annotated in fatty acid metabolism displayed a downregulation in expression in this particular model. Unfortunately, this provides no information for us to refine our molecular understanding of the impact of *MSTN* on leanness observed at the phenotype level. It is unlikely that differences in the statistical methods used to establish significance of DE account for the fat metabolism discrepancy between our cattle study and this mouse study.

Marbling physiology and development

In postnatal ruminant muscle, adipocytes undergo hyperplasia before a period of infilling hypertrophy. The relative contributions of these two processes to marbling are not fully known. However, both processes – either in isolation or in combination – would be expected to systematically increase IMF% over postnatal development. In principle, this should be reflected by corresponding increases in the expression of a set of ‘adipocyte fat metabolism genes’. Indeed, examples of these, which tend to relate to TAG deposition, have been previously identified using DE (Wang *et al.*, 2009), co-expression (Hudson *et al.*, 2009b) and individual correlations of gene expression to phenotype (De Jager *et al.*, 2013). In the latter study, a TAG deposition gene set comprised *THRSP*, *FABP4*, *CIDEA*, *DGAT2*, *PLIN1*, *PCK1*, *AGPAT2*, *ACSM1*, *CIDEA*, *ADIPOQ*, *S100G*, *TUSC5*, *ADIG*, *PLS1* and *GPAM*. Of these, *ADIPOQ* and *THRSP* were previously found to correlate with individual IMF% according to a study by Wang *et al.* (2009), and a subset were previously found to be co-expressed in two separate network modules (*ADIPOQ*, *PLS1*, *PLIN1*, *CIDEA*, *FABP4* and *TUSC5* on the one hand and *ADIG*, *THRSP* and *S100G* on the other) in the muscle co-expression network according to a study by Hudson *et al.* (2009b).

Some authors have argued that the best predictor of future marbling potential is current marbling status (Pethick *et al.*, 2004). If this is correct, the molecular profiles of the endpoint fatty acid metabolism genes would be among the most valuable future predictors of marbling potential from a molecular perspective, but only in circumstances where animals are actively depositing fat. From a prognostic perspective, a problem is that these probe-based profiles (e.g. *FABP4*) do not diverge between the breeds until very late in the postnatal time course, that is, 20 to 25 months. This means they may have limited practical value, as substantial economic investment in the animal has already been made by this time.

Early gene expression markers for marbling development

Given these issues, we developed a number of simple analytical questions with a view to identifying novel gene

expression markers of marbling adipocytes. By exploring positive regressions over postnatal development, we hoped one could identify hitherto unrecognised adipocyte markers, in addition to the conventional fat metabolism genes. Not surprisingly, enriched among the genes with positive slopes in both breeds postnatally, but steeper in the W × H, were the expected fat metabolism genes. *THRSP* was the single best performer, topping the ranked list. This prioritisation of *THRSP* on differential regression supports a previous finding based on DE using a cDNA array (Wang *et al.*, 2009). However, the expression of *THRSP* fluctuates along the developmental time course, implying it is quite sensitive to environmental and nutritional signals.

Genes with more robust expression profiles, that are less sensitive to nutrition and that do not converge later on in the time course, may be of value. This logic provides the basis for our next steps. There are various lines of evidence linking adipogenesis and ECM biology. One is that during adipogenesis the ECM develops from a fibrillar to laminar structure as cells move from commitment to growth and triglyceride storage (Mariman and Wang, 2010). In addition, adipogenesis tends to proceed in a collagen-rich ECM environment (Sato-Kusubata *et al.*, 2011). Further, mature adipocytes invest a substantial amount of energy on ECM maintenance (Mariman and Wang, 2010). Along these lines, a recent publication using gene expression analysis on Korean Hanwoo cattle found a connective tissue remodeller *ADAMTS4* to be a prominent candidate for marbling development (Lee *et al.*, 2010). Given the proposed role of ECM remodelling in driving the marbling tissue deposition environment, we undertook a deeper exploration of our data.

We attempted to identify genes annotated as ECM remodellers that diverge earlier than the 20 months displayed by the end-point fat metabolism genes. This process prioritised *WISP2* that diverges much earlier, certainly by the 12-month time point and potentially as early as soon after 7 months. *WISP2* is known to be highly expressed in human pre-adipocytes. It has recently been discovered to encode a novel secreted adipokine protein that is a negative regulator of pre-adipocyte commitment (Hammarstedt *et al.*, 2013). It has also been shown to inhibit the pro-fibrotic connective tissue growth factor (Sabbah *et al.*, 2011). Furthermore, *WISP2* was recently identified as one of the top candidate secreted proteins in the development of obesity, as determined by fat depot DE between obese and lean humans (Dahlman *et al.*, 2012).

Therefore, it appears that the protein encoded by *WISP2* has a number of biochemical properties that may lend itself to being a reasonable diagnostic marker for future marbling potential. Its expression level in 12-month bovine muscle apparently represents the number and/or activity of precursor adipocytes and their biological connection to the surrounding ECM. Its expression level would be predicted to be maximally informative in muscle tissue samples somewhere between 7- and 12-month old animals. *WISP2* loses its descriptive ability later on in the time course in that, similar to the fat metabolism genes it also converges at the 30-month

time point when the difference in IMF phenotype is actually expressed most strongly. This may reflect the fact that the muscle systems have ceased active matrix remodelling by this point and/or that precursor adipocytes have diminished in activity/number.

In a further effort to overcome issues of environmental and nutritional noise, we asked a more stringent question of the data. Which genes, if any, show a monotonic increase at all postnatal time points, in addition to being higher in Wagyu than the Piedmontese at those time points. Notable among the two genes that meet these criteria was *RAI14*. This gene has recently been discovered to be expressed in porcine adipocytes treated with an inhibitor of the microRNA-103 implicated in adipocyte differentiation (Li *et al.*, 2010). The divergence between Wagyu and Piedmontese *RAI14* is strongly maintained through to the 30-month time point. The expression of this gene would be hypothesised to reflect IMF % increases in a manner that is relatively insensitive to nutrition.

Differentially expressed fat metabolism genes

We were able to identify numerous differentially expressed fat metabolism genes, irrespective of the exact nature of their expression profile in the two breeds. The majority of these are relatively upregulated in the high IMF Wagyu. Having said this, the single largest discriminator between the breeds was *CYP4F2*, a monooxygenase that can inactivate leukotriene B₄, a mediator of inflammation (Hardwick, 2008). This molecule is relatively upregulated in the Piedmontese. Its role in metabolising both pro- and anti-inflammatory agents indicates it may function in both the activation and resolution of inflammation (Hardwick, 2008). *CYP4F2* is also involved in long chain and very long chain fatty acid metabolism whereupon the hydroxylated fatty acids are converted to dicarboxylic acids, which are preferentially metabolised in the peroxisome to shorter chain fatty acids that in turn can be processed in the mitochondrion (Hardwick, 2008). *CYP4F2* has high affinity for arachidonic acid (Hardwick, 2008), a compound previously explored in the marbling literature, and is also responsive to RA stimulation (Kalsotra *et al.*, 2008). Given the strong relationship between inflammation and adiposity, the observation that an inhibitor of inflammation is relatively downregulated in the high fat, pro-inflammatory Wagyu is noteworthy.

Transcriptional regulatory network

One means of inferring causal regulation of biological processes is to exploit global patterns of differential co-expression of TF and other regulators through RIF analysis (Hudson *et al.*, 2009a; Reverter *et al.*, 2010). RIF has been used to unravel the causal basis of Parkinson's (Rhinn *et al.*, 2012) and Alzheimer's (Rhinn *et al.*, 2013) diseases from brain transcriptome data, showing it generalises very well across species, tissues and data structure. This approach identified key roles for a number of TF relating to inflammation, fatty acid metabolism and production of fatty masses called liposarcoma.

These exemplar TF molecules all possess substantially different patterns of network connectivity between the breeds even though they are not necessarily differentially expressed themselves. The purpose of this final aspect of our analysis is not to identify expression-based markers, but to shed light on earlier mechanisms. The biological origin of any observed differential connectivity between breeds is unknown, with possibilities including breed by developmental changes in (1) cellular localisation, (2) phosphorylation status, (3) ligand binding or (4) possible mutations in the coding sequence. *YEATS4* and *MDM2* are of particular interest as they have both been observed to be amplified and overexpressed in liposarcoma (Italiano *et al.*, 2008), where proliferating fatty masses occur in an uncontrolled manner in a skeletal muscle environment. We have used a number of methods to highlight genes behaving differently between the breed crosses at a number of different development stages.

Conclusions

We conclude that divergent *WISP2* expression implicates differential number and/or activity of precursor adipocyte cell populations and ECM remodelling, *CYP4F2* may reflect divergent levels of systemic inflammation, *RAI14* the differential involvement of the RA axis, and *YEATS4* and *MDM2* the differential TF regulatory activity in two molecules previously associated with liposarcoma.

The Future

The challenges for the future include the elucidation of a detailed molecular path from causal mutation (if any) to ultimate phenotype in high marbling breeds such as the Wagyu. From a more practical perspective, even earlier diagnostics of marbling potential would be useful, particularly tests that may aid decisions to short or long feed in a commercial feedlot. Given gene expression is an invasive procedure requiring tissue biopsies, connecting the muscle-derived expression data to a blood-borne diagnostic would be of great utility. In terms of the specific molecules whose abundance at key time points has been highlighted here for potential predictive purposes (particularly *WISP2*, but also potentially *RAI14* and *CYP4F2*), their performance in other commercially important muscles (in addition to LD) and in additional breeds would be an important validation step.

Acknowledgements

We wish to thank Dr Gregory Harper and Dr Aaron Ingham for stimulating insights and discussion relating to an earlier version of this manuscript. The animal experimentation and tissue sampling associated with this study was conducted within the Australian Cooperative Research Centre for Cattle and Beef Quality. The analysis was partially supported by the CSIRO Food Futures Flagship.

Supplementary materials

For supplementary material/s referred to in this article, please visit <http://dx.doi.org/10.1017/S1751731114002754>

References

- Bonet ML, Ribot J and Palou A 2012. Lipid metabolism in mammalian tissues and its control by retinoic acid. *Biochimica et Biophysica Acta* 1821, 177–189.
- Bonnet M, Cassar-Malek I, Chilliard Y and Picard B 2010. Ontogenesis of muscle and adipose tissues and their interactions in ruminants and other species. *Animal* 4, 1093–1109.
- Dahlman I, Elsen M, Tennagels N, Korn M, Brockmann B, Sell H, Eckel J and Arner P 2012. Functional annotation of the human fat cell secretome. *Archives of Physiology and Biochemistry* 118, 84–91.
- De Jager N, Hudson NJ, Reverter A, Barnard R, Cafe LM, Greenwood PL, Dalrymple BP 2013. Gene expression phenotypes for lipid metabolism and intramuscular fat in skeletal muscle of cattle. *Journal of Animal Science* 91, 1112–1128.
- Eden E, Lipson D, Yogev S and Yakhini Z 2007. Discovering motifs in ranked lists of DNA sequences. *PLoS Computational Biology* 3, e39.
- Gotoh T, Albrecht E, Teuscher F, Kawabata K, Sakashita K, Iwamoto H and Wegner J 2009. Differences in muscle and fat accretion in Japanese black and European cattle. *Meat Science* 82, 300–308.
- Greenwood PL, Cafe LM, Hearnshaw H, Hennessy DW, Thompson RF and Morris SG 2006. Long-term consequences of birth weight and growth to weaning on carcass, yield and beef quality characteristics of Piedmontese- and Wagyu-sired cattle. *Australian Journal of Experimental Agriculture* 46, 257–269.
- Hammarstedt A, Hedjazifar S, Jenndahl L, Gogg S, Grunberg J, Gustafson B, Klimcakova E, Stich V, Langin D, Laakso M and Smith U 2013. *WISP2* regulates preadipocyte commitment and PPARgamma activation by BMP4. *Proceedings of the National Academy of Science USA* 110, 2563–2568.
- Hardwick JP 2008. Cytochrome P450 omega hydroxylase (*CYP4*) function in fatty acid metabolism and metabolic diseases. *Biochemical Pharmacology* 75, 2263–2275.
- Harper GS and Pethick DW 2004. How might marbling begin. *Australian Journal of Experimental Agriculture* 44, 653–662.
- Huang da W, Sherman BT and Lempicki RA 2009. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 4, 44–57.
- Hudson NJ, Reverter A and Dalrymple BP 2009a. A differential wiring analysis of expression data correctly identifies the gene containing the causal mutation. *PLoS Computational Biology* 5, e1000382.
- Hudson NJ, Dalrymple BP and Reverter A 2012. Beyond differential expression: the quest for causal mutations and effector molecules. *BMC Genomics* 13, 356.
- Hudson NJ, Reverter A, Wang Y, Greenwood PL and Dalrymple BP 2009b. Inferring the transcriptional landscape of bovine skeletal muscle by integrating co-expression networks. *PLoS One* 4, e7249.
- Hudson NJ, Lyons RE, Reverter A, Greenwood PL and Dalrymple BP 2013. Inferring the in vivo cellular program of developing bovine skeletal muscle from expression data. *Gene Expression Patterns* 13, 109–125.
- Italiano A, Bianchini L, Keslair F, Bonnafous S, Cardot-Leccia N, Coindre JM, Dumollard JM, Hofman P, Leroux A, Mainguene C, Peyrottes I, Ranchere-Vince D, Terrier P, Tran A, Gual P and Peduteur F 2008. *HMG2* is the partner of *MDM2* in well-differentiated and dedifferentiated liposarcomas whereas *CDK4* belongs to a distinct inconsistent amplicon. *International Journal of Cancer* 122, 2233–2241.
- Kalsotra A, Du L, Wang Y, Ladd PA, Kikuta Y, Duvic M, Boyd AS, Keeney DS and Strobel HW 2008. Inflammation resolved by retinoid X receptor-mediated inactivation of leukotriene signaling pathways. *FASEB Journal* 22, 538–547.
- Lee SH, Gondro C, van der Werf J, Kim NK, Lim DJ, Park EW, Oh SJ, Gibson JP and Thompson JM 2010. Use of a bovine genome array to identify new biological pathways for beef marbling in Hanwoo (Korean Cattle). *BMC Genomics* 11, 623.
- Lehnert SA, Reverter A, Byrne KA, Wang Y, Natrass GS, Hudson NJ and Greenwood PL 2007. Gene expression studies of developing bovine longissimus muscle from two different beef cattle breeds. *BMC Developmental Biology* 7, 95.
- Li DL, Chen JL, Wen J, Zhao GP, Zheng MQ and Liu C 2013. Growth, carcass and meat traits and gene expression in chickens divergently selected for intramuscular fat content. *British Poultry Science* 54, 183–189.

- Li G, Wu Z, Li X, Ning X, Li Y and Yang G 2010. Biological role of microRNA-103 based on expression profile and target genes analysis in pigs. *Molecular Biology Reports* 38, 4777–4786.
- Mariman EC and Wang P 2010. Adipocyte extracellular matrix composition, dynamics and role in obesity. *Cellular and Molecular Life Sciences* 67, 1277–1292.
- Pannier L, Mullen AM, Hamill RM, Stapleton PC and Sweeney T 2010. Association analysis of single nucleotide polymorphisms in DGAT1, TG and FABP4 genes and intramuscular fat in crossbred *Bos taurus* cattle. *Meat Science* 85, 515–518.
- Pethick DW, Harper GS and Oddy VH 2004. Growth, development and nutritional manipulation of marbling in cattle: a review. *Australian Journal of Experimental Agriculture* 44, 705–715.
- Ramayo-Caldas Y, Fortes MR, Hudson NJ, Porto-Neto LR, Bolormaa S, Barendse W, Kelly M, Moore SS, Goddard ME, Lehnert SA and Reverter A 2014. A marker-derived gene network reveals the regulatory role of PPARGC1A, HNF4G and FOXP3 in intramuscular fat deposition of beef cattle. *Journal of Animal Science* 92, 2832–2845.
- Ren ZQ, Wu WJ, Liu WH, Zheng R, Li JL, Zuo B, Xu DQ, Li FE, Lei MG, Ni DB and Xiong YZ 2014. Differential expression and effect of the porcine ANGPTL4 gene on intramuscular fat. *Genetics and Molecular Research* 13, 2949–2958.
- Reverter A, Hudson NJ, Nagaraj SH, Perez-Enciso M and Dalrymple BP 2010. Regulatory impact factors: unraveling the transcriptional regulation of complex traits from expression data. *Bioinformatics* 26, 896–904.
- Rhinn H, Fujita R, Qiang L, Cheng R, Lee JH and Abeliovich A 2013. Integrative genomics identifies APOE epsilon4 effectors in Alzheimer's disease. *Nature* 500, 45–50.
- Rhinn H, Qiang L, Yamashita T, Rhee D, Zolin A, Vanti W and Abeliovich A 2012. Alternative alpha-synuclein transcript usage as a convergent mechanism in Parkinson's disease pathology. *Nature Communications* 3, 1084.
- Sabbah M, Prunier C, Ferrand N, Megalophonos V, Lambein K, De Wever O, Nazaret N, Lachuer J, Dumont S and Redeuilh G 2011. CCN5, a novel transcriptional repressor of the transforming growth factor beta signaling pathway. *Molecular and Cell Biology* 31, 1459–1469.
- Saez G, Davail S, Gentes G, Hocquette JF, Jourdan T, Degrace P and Baeza E 2009. Gene expression and protein content in relation to intramuscular fat content in Muscovy and Pekin ducks. *Poultry Science* 88, 2382–2391.
- Sato-Kusubata K, Jiang Y, Ueno Y and Chun TH 2011. Adipogenic histone mark regulation by matrix metalloproteinase 14 in collagen-rich microenvironments. *Molecular Endocrinology* 25, 745–753.
- Wang W, Xue W, Jin B, Zhang X, Ma F and Xu X 2013. Candidate gene expression affects intramuscular fat content and fatty acid composition in pigs. *Journal of Applied Genetics* 54, 113–118.
- Wang YH, Byrne KA, Reverter A, Harper GS, Taniguchi M, McWilliam SM, Mannen H, Oyama K and Lehnert SA 2005. Transcriptional profiling of skeletal muscle tissue from two breeds of cattle. *Mammalian Genome* 16, 201–210.
- Wang YH, Bower NI, Reverter A, Tan SH, De Jager N, Wang R, McWilliam SM, Cafe LM, Greenwood PL and Lehnert SA 2009. Gene expression patterns during intramuscular fat development in cattle. *Journal of Animal Science* 87, 119–130.
- Welle S, Cardillo A, Zanche M and Tawil R 2009. Skeletal muscle gene expression after myostatin knockout in mature mice. *Physiological Genomics* 38, 342–350.