

The effect of temporary deprivation of lying and feeding on the behaviour and production of lactating dairy cows

M. D. Cooper^{1,2}, D. R. Arney^{1,3} and C. J. C. Phillips^{2,4†}

¹Moulton College, Northampton, UK; ²Department of Clinical Veterinary Medicine, University of Cambridge, Cambridge, UK; ³Estonian University of Life Sciences, Tartu, Estonia; ⁴Centre for Animal Welfare and Ethics, School of Veterinary Sciences, University of Queensland, Gatton, Australia

(Received 9 February 2007; Accepted 24 October 2007)

Two experiments were conducted to examine the effects of depriving dairy cows of the ability to feed and lie down for short periods, on behaviour and production. In experiment 1, cows were deprived by confining them in pairs in a pen for 2 or 4 h, and they more frequently exhibited behaviour likely to suggest discomfort – leg stamping, repositioning themselves, shifting their weight between legs and butting. After deprivation, the cows deprived for 2 h made up their lost feeding time within 24 h, but cows deprived for 4 h did not restore their feeding time within the 41-h period of observation. Lying time was not restored in either treatment within the 41-h period. Milk yield was not affected by the treatment. However, in experiment 2, when cows were deprived of feeding and lying for 4 h, during which time their hooves were trimmed (which is likely to be a painful and stressful procedure and result in some discomfort for a period post-trimming) the evidence suggested that milk yield was reduced by approximately 2 l/day for 3 days, with corresponding increases during the subsequent 2 days. Walking speed on returning to the herd was the same as before the treatment. In summary, temporary deprivation of feeding and lying for 2 and 4 h/day induced behaviours that were indicative of discomfort and frustration but had no negative effect on milk production, except when 4 h of deprivation was accompanied by foot trimming.

Keywords: behaviour, dairy cows, feeding, lying

Introduction

Dairy cows are usually deprived of the ability to lie down and feed while they are waiting for routine veterinary inspections, such as for pregnancy diagnosis or tuberculin testing, for foot trimming, and also during milking and the holding period prior to milking. Previously we have shown that during deprivation of lying, cows increased the time spent feeding and more frequently exhibited leg stamping, weight shifting, body repositioning and butting; behaviours that suggest discomfort (a negative subjective feeling in response to an unpleasant stimuli that is sufficient in intensity for the animal to initiate a response, e.g. a change in posture, which exists until the stimuli is reduced and at the extreme would consist of pain and distress) (Cooper *et al.*, 2007). After deprivation, cows recovered some but not all of their lying time by reducing feeding and standing. However, feeding deprivation is often concomitant with lying deprivation when cows are withheld for veterinary treatment or reproductive techniques, and it is unclear as to whether this magnifies the impact on behaviour.

Dairy cows are highly motivated to feed and rest, potentially leading to a motivational conflict in cows with limited time available for both activities (Metz, 1985; Fregonesi and Leaver, 2001), exacerbated by the fact that their feed intake capacity has not increased to the same extent as their milk production (Kamphues, 1998). Depriving lactating dairy cows of opportunities for both feeding and lying will help to determine their priorities for these two behaviours. Bolinger *et al.* (1997) found that although post-deprivation feeding frequency was reduced after 4 h deprivation of lying and feeding when cows were placed in restraint yokes, feed intake and milk yield were not affected. Plasma cortisol, neutrophil-to-mononuclear cells ratio and somatic cell count (SCC) were not affected by restraint, but grooming and agonistic behaviours were increased and there was increased lying post restraint. During restraint, cows appeared to increase time spent ruminating, and spent less time ruminating afterwards. Bolinger *et al.* (1997) suspected that restrained cattle shifted their weight from one leg to another, and recommended that this be further studied.

We investigated the impact on behaviour and production of depriving cows of the ability to lie down and to feed for

† E-mail: c.phillips@uq.edu.au

2 or 4 h on a single occasion, and the impact of an additional stressful procedure – foot trimming – during 4 h of deprivation.

Material and methods

Experiment 1

An experiment was conducted over 18 consecutive days, commencing 28 June and ending 15 July, using 18 cows from the Holstein-Friesian dairy herd of Moulton College, Northampton, UK. Cows were fed a total mixed ration containing fresh weight proportions as follows: second-cut grass silage (53.5%), maize silage (30.6%), wet maize gluten (15.3%) and minerals (0.6%). Other aspects of the management of the herd were described previously (Cooper *et al.*, 2007).

The cows were in late lactation (mean \pm s.e. lactation duration 273 ± 5.1 days), with a mean \pm s.e. initial milk yield of 21.8 ± 1.12 kg/day, mean \pm s.e. body condition score of 3.00 ± 0.05 and a mean \pm s.e. time to parturition of 80.2 ± 1.9 days. Cows were allocated to nine pairs at random, and each pair was randomly assigned to one of three blocks of a replicated Latin-square design (no. 63 of Patterson and Lucas, 1962; Table 1). This design had three periods, three blocks and three treatments: control (no lying/feeding deprivation), 2 h of lying plus feeding deprivation (treatment 2-h) and 4 h of lying plus feeding deprivation (treatment 4-h).

Each pair was allocated to one of three adjacent, identical pens, as described in Cooper *et al.* (2007), at random, where the necessary treatments were applied. Pens provided *ad libitum* access to drinking water and the same total mixed ration as the rest of the herd, which was delivered at the same time. During periods of darkness, artificial lighting was provided overhead by six strip lights to facilitate video-recording. Cows spent 48 h in the experimental pens and were returned back to the herd towards the end of morning milking when the next group of six cows was collected. Each group of six cows underwent the experimental procedure three times, with each pair in the group subject to a different treatment on each occasion. The experimental cows were allowed approximately 2 h to become accustomed to the pens and each other before any behavioural observations took place. Lying deprivation was achieved by securing the cows in the deprivation area using

a non-electrified spring gate, which prevented the cows having access to the bedded area. Feeding deprivation was achieved by withdrawing feed from the barrier. The 4- and 2-h deprivation periods commenced at 1020 and 1220 h, respectively, and at 1420 h the spring gates were removed and the cows gained access to the bedded area. Following the application of treatments, the cows remained in their pairs for a further 41 h, after which they were returned to the herd.

Behavioural observations

Deprivation period. Cows were observed in the same order each time by one of three trained observers. Behaviours previously thought to be associated with discomfort, frustration or fatigue (see Cooper *et al.*, 2007) were recorded continuously as frequencies. These were leg stamping, repositioning, butting, head swinging and weight shifting. The frequency of nosing (touching with the nose) and sniffing the ground, self-licking and walking (moving all four legs slowly forward one at a time in a synchronised manner covering greater than one body length in distance) were also recorded in this way. Behaviours recorded by duration included rubbing of the head against the housing, drinking (drinking from water trough/bowl) and sleeping (lying down with neck relaxed and eyes closed). All the above behaviours not defined here have previously been defined by Cooper *et al.* (2007) – however leg stamping was previously recorded as leg stomping, and licking was termed grooming. Bouts were separated by the cow showing at least 1 s of another behaviour. Behaviours recorded by scan sampling at 5 min intervals included feeding, ruminating (chewing partially digested food that has been regurgitated) while lying or standing, and standing or lying but not ruminating, with a total of 144 measurements per cow and 2592 in total.

Post deprivation. Continuous time-lapse video-recordings (a Sanyo 3372 camera fitted with a varifocal (3.5 to 8 mm) lens (Sanyo, Watford, UK) and a HS-1024EB video-recorder (VHS Mitsubishi, Hatfield, UK) played at 2.6 mm/s) commenced immediately after the deprivation period for the subsequent 41 h when the cows remained in the pen, during which time it was expected that residual or compensatory behavioural effects of treatment may be displayed (Metz, 1985). Recordings of lying, standing and feeding behaviour were made every 5 min, except when the cows were being milked, with a total of 1476 measurements per cow and 26 568 measurements in total. Feeding was defined as the cow having food in the mouth and/or chews (Munksgaard *et al.*, 1999). Lying was defined as the body resting on the floor (Munksgaard *et al.*, 1999). For each individual cow, the total time spent in each activity was estimated by assuming it had been performing the behaviour for the remainder of the 5 min period and multiplying the number of recordings of each behaviour by the number of scans in the 41-h period. The duration of the first

Table 1 Latin-square design showing order of treatments for each pair of cows in each block for each period

Pair	Block 1			Block 2			Block 3		
	1	2	3	4	5	6	7	8	9
Period 1	C	2-h	4-h	C	2-h	4-h	4-h	C	2-h
Period 2	2-h	4-h	C	4-h	C	2-h	C	2-h	4-h
Period 3	4-h	C	2-h	2-h	4-h	C	2-h	4-h	C

C = control treatment; 2-h = 2 h deprivation treatment; 4-h = 4 h deprivation treatment.

meal post-deprivation was determined from the start and cessation of this activity.

Milk production. Milk yield was recorded on four consecutive occasions for each cow after entering the experiment – afternoon milking on day 1 (the deprivation day), morning and afternoon milking on day 2 and morning milking on day 3, prior to the cows being returned to the herd. Morning and afternoon milking took place between 0700 and 0830 h, and between 1500 and 1720 h, respectively.

Statistical analyses

The mean value of each pair of cows was used as the replicate. Most behaviours were not normally distributed across pairs as defined by the Anderson–Darling test ($P < 0.05$), so all behaviours were analysed by Friedman's non-parametric test with pair and treatment as factors. Spearman's rank order correlation coefficients were determined between behavioural variables. The number of vocalisations and the time spent sleeping were both too few to analyse by hour, and were only analysed for all 4 h together. For the post-deprivation behaviour data, graphical representation of mean time spent in each behaviour for each treatment revealed that blocks of 8-h periods were most appropriate for analysis. Standard errors for individual hours were calculated to facilitate identification of differences between treatment means. Milk production data were also not normally distributed and therefore Friedman's test was used as for behaviours, as well as the duration of the first post-derivation feeding and lying bouts. Correlation coefficients were additionally determined for production characteristics and behaviours using the same method as that used for correlations between behaviours. Because of the large number of potential correlations, a statistical probability of 0.01 was used to indicate significance, unless the correlation was considered particularly relevant to the discussion.

Experiment 2

Because it is difficult to distinguish the effects of potentially stressful events, such as foot trimming of a herd, from unrelated but concurrent events, such as changes in diet or the weather, a trimming programme that affected a small number of cows at regular intervals was selected for this purpose. Fifteen cows with overgrown front hooves were selected from the University of Cambridge dairy herd of Holstein-Friesian cows to be trimmed by veterinary students, who had no prior experience of the procedures. The cows were removed from the herd at morning milking (0700 h) in five groups of three at weekly intervals. They waited in a 4 × 4 m pen with a concrete floor and no bedding, with access to water but no food, until the students arrived at 0900 h to lead them to individual trimming crushes, where they were treated in the standing position. Here the cows were subjected to supervised trimming of the front feet for 120 min by the five-step Dutch method

(Greenough, 1997). Trimming ended 4 h after initial restraint, and the cows were then returned to the herd in a straw yard with *ad libitum* access to a total mixed ration.

To assess the effect of trimming on the cow's walking behaviour, their speed and step rate were recorded by an observer with a stopwatch over a 30 m distance as they voluntarily walked back along a passageway to their straw yard accommodation following afternoon milking 2 days before the day of trimming and on the day of trimming. Each cow's front foot was used as a reference point. Milk yield was recorded automatically in the rapid exit parlour for each cow 5 days before and 5 days after trimming.

Results

Experiment 1

Deprivation period. Cows in treatments C and 2-h both spent a median 15 min/h feeding during the 4 and 2 h, respectively, when food was available, with no differences between treatments ($P = 0.62$). There was no difference between treatments C and 2-h in the median time spent lying without ruminating (range 0 to 5 min/h for both treatments, $P > 0.25$) and lying ruminating (range 8 to 16 min/h for both treatments, $P > 0.25$) during the first 2 h (treatment 2-h cows were deprived of lying during the subsequent 2-h period). No cow from either of the deprivation groups attempted to lie down during deprivation. Time spent standing, not ruminating, was greater when the cows were undergoing deprivation (Table 2) and tended to be greater for treatment 4-h cows in hours 1 and 2, compared with those in hours 3 and 4. Standing ruminating time was also greater when the cows were undergoing deprivation, and particularly in treatment 2-h cows and treatment 4-h cows in hours 3. The total time spent standing, not ruminating, and standing and ruminating over the 4-h period both increased between treatments with length of deprivation.

Table 2 The median duration (min/h) of standing with and without ruminating recorded instantaneously every 5 min for hours 1 to 4 of the deprivation period, and the median per hour for all 4 hours, for cows in control, 2-h and 4-h treatments

Treatment	Hour				Median value for all 4 hours
	1	2	3	4	
Standing not ruminating					
Control	15.0	17.5	12.5	11.7	57.5
2 h	17.5	17.5	27.5	25.0	86.7
4 h	40.0	35.0	25.0	33.0	95.8
<i>P</i>	0.001	0.001	0.005	0.005	0.004
Standing ruminating					
Control	4.2	5.8	7.5	3.3	15.0
2 h	2.5	6.7	30.0	29.2	61.7
4 h	18.3	25.0	32.5	20.0	85.8
<i>P</i>	0.04	0.04	0.02	0.001	0.002

In the deprived cows, the frequency of leg stamping during hours 3 and 4 was approximately double that of the control group and this behaviour increased overall with duration of deprivation (Table 3). Repositioning was not observed, nor weight shifting to any major extent, for the cows in treatment C or treatment 2-h when not deprived, but both behaviours were performed more frequently during deprivation, especially repositioning. These behaviours increased overall with duration of deprivation. Head swinging, sniffing the ground and self-licking were not significantly affected by treatment. Nosing the ground was only observed in treatment C cows in hours 3, but occurred in treatment 4-h cows and, to a lesser extent, in treatment 2-h cows during deprivation. The frequency of cows rubbing their heads against the housing increased overall between treatments with length of deprivation. Butting was rare outside deprivation, and was increased more overall for cows in treatment 4-h than in 2-h. Time spent drinking and the number of times cows were observed walking were both reduced by deprivation, particularly for cows in treatment 4-h. The number of vocalisations was increased by deprivation, particularly for cows in treatment 4-h (median: 0, 0.1 and 0.5 vocalisations per 4 h for cows in treatments C, 2-h and 4-h, respectively, $P=0.004$). The time spent sleeping was reduced by deprivation (median: 3.5, 2.5 and 0 s per 4 h for cows in treatments C, 2-h and 4-h, respectively, $P=0.008$).

Leg stamping was correlated ($P<0.05$) with head swinging in treatments C (CC, correlation coefficient, +0.92) and 2-h (CC +0.79). Sniffing the ground was correlated ($P<0.05$) with leg stamping and repositioning in treatments 2-h (CC +0.59) and 4-h (CC +0.59) and rubbing head against housing in treatment 2-h (CC +0.61).

Post deprivation. The first meal post-treatment was longest for treatment 4-h cows (median 27 min) compared with cows in treatments C (median 20 min) and 2-h (median 19 min) ($P<0.001$). The first lying bout was longer for cows in treatment 4-h (median 79 min), and, to a lesser extent, for cows in treatment 2-h (median 61 min), compared with cows in treatment C (median 51 min) ($P=0.003$). The feeding time in the first 8-h post deprivation was longest for cows in treatment 4-h (median 115 min) and, to a lesser extent, for cows in treatment 2-h (median 98 min) compared with cows in treatment C (median 73 min). The main feeding bout during this first 8-h period occurred between 1620 and 2220 h (Figure 1). The cumulative feeding time was restored for cows in treatment 2-h by 24 h post deprivation (Figure 2). Cows in treatment 4-h did not compensate for lost feeding time during deprivation over the 41-h post-deprivation period (Figure 2).

Neither standing nor lying time were significantly affected by treatment in any of the 8-h periods post-deprivation ($P>0.05$), but there was one recording (at 1820 h), soon after the end of deprivation, when the cows in treatment 4-h tended to increase their lying time (Figure 3). The absence of restoration of lying at other times of the day meant that the cumulative lying curves were parallel for the three treatments (Figure 4).

Table 3 The median frequency or duration of each behaviour recorded continuously in hours 1 to 4 of the deprivation period, and for the 4 h, for cows in control, 2-h and 4-h treatments

Treatment	Hour				Median value for all 4 hours
	1	2	3	4	
Leg stamping (no. per h)					
Control	3.0	2.0	3.0	2.5	3.1
2 h	1.8	1.8	6.2	7.0	5.5
4 h	9.2	9.2	7.3	9.5	9.2
<i>P</i>	0.03	0.12	0.01	0.04	0.01
Reposing (no. per h)					
Control	0	0	0.5	0	0.3
2 h	0.3	0.5	5.2	6.0	3.3
4 h	5.2	7.0	5.3	6.5	6.4
<i>P</i>	0.002	0.01	0.004	0.004	<0.001
Weight shifting (no. per h)					
Control	0	0.2	0.8	1.3	0.75
2 h	0	0	3.5	4.5	1.6
4 h	1.5	4.8	3.2	4.7	3.4
<i>P</i>	0.001	0.01	0.02	0.07	0.008
Head swinging (no. per h)					
Control	2.7	2.5	1.5	2.0	2.0
2 h	2.0	1.3	1.5	2.0	2.5
4 h	3.8	4.7	3.5	2.0	3.8
<i>P</i>	0.41	0.15	0.90	0.89	0.82
Nose ground (no. per h)					
Control	0	0	0.8	0	0.2
2 h	0	0	1.7	1.2	1.1
4 h	3.5	3.0	3.0	1.8	2.9
<i>P</i>	0.001	<0.001	0.05	0.005	0.001
Sniffing ground (no. per h)					
Control	0.7	0.5	0.7	0.5	0.6
2 h	0.5	0.5	0.8	1.0	1.1
4 h	0.8	2.0	1.0	1.5	1.7
<i>P</i>	0.80	0.06	0.97	0.24	0.17
Licking self (licks per h)					
Control	5.2	1.5	0.7	3.2	3.5
2 h	5.0	3.8	3.0	1.0	4.4
4 h	7.8	5.2	1.0	2.3	5.7
<i>P</i>	0.46	0.37	0.67	0.20	0.041
Rubbing head against housing (s/h)					
Control	0.33	1.0	1.2	3.0	1.7
2 h	1.7	1.5	4.3	9.0	3.3
4 h	4.5	7.5	8.0	6.2	8.3
<i>P</i>	0.32	0.06	0.07	0.51	0.03
Butting (no. per h)					
Control	0	0.2	0	0	0.1
2 h	0.3	0	0.5	0.5	0.5
4 h	1.2	0.8	0.5	0	0.9
<i>P</i>	0.002	0.004	0.08	0.07	0.02
Drinking (s/h)					
Control	13.5	0	6.0	2.7	12.1
2 h	1.8	6.5	8.0	1.3	9.5
4 h	0.2	0	0	0	1.7
<i>P</i>	0.03	0.47	0.12	0.07	0.02
Walking (no. per h)					
Control	2.2	1.3	2.0	1.5	1.8
2 h	1.8	1.9	0	0	0.9
4 h	0	0	0	0	0
<i>P</i>	0.002	0.002	<0.001	<0.001	<0.001




Figure 1 The mean time (min/h) spent feeding for all the experimental animals in the control (◆), 2-h deprivation (■) and 4-h deprivation (▲) treatments for the 4-h deprivation period (1020–1420 h) and 40-h post-deprivation period ($l = 1$ s.e.d.).




Figure 2 The accumulated feeding time (min) for cows in the control (◆), 2-h deprivation (■) and 4-h deprivation (▲) treatments for the 4-h deprivation period (h – 4 to – 1) and post-deprivation period (hours 1 to 40), ($l = 1$ s.e.d. between two means) in experiment 1.

Milk production. There were no significant effects on milk yield at any of the four milkings post treatment (Table 4). However, there were positive correlations ($P < 0.05$) between milk yield and the incidence of leg stamping (CC +0.60) and weight shifting (CC +0.70). For cows in treatments C and 2-h, there were negative correlations between milk yield and time spent lying (CC –0.50 and –0.49, respectively, $P < 0.05$), and there were tendencies for positive correlations with the incidence of head swinging (CC +0.45 and +0.41, respectively, $P < 0.10$).

Experiment 2

Mean milk yields were 22.6 kg/day for the 5 days before trimming and 20.8 kg/day for the 5 days after trimming. Milk yield was reduced by about 2 l/day for 2 days after foot

trimming, and on the day of trimming (Figure 5). Subsequently, there was a recovery for 2 days after which milk yield appeared to resume to its normal pattern of decline. There was no difference in the mean \pm s.e. walking rate before and after trimming (before: 0.82 ± 0.096 ; after: 0.82 ± 0.136 m/s) or in the stepping rate of cows before and after trimming (before: 1.44 ± 0.063 ; after: 1.40 ± 0.043 steps per min).

Discussion

Deprivation period

In experiment 1, cows in the control treatment lay and fed for 100 and 59 min, respectively, over the 4 h, which was therefore likely to be similar to the actual amount of lying and feeding that treatment 4-h cows were deprived of.




Figure 3 The mean time (min/h) spent lying for all the experimental animals in the control (◆), 2-h deprivation (■) and 4-h deprivation (▲) treatments for the 4-h deprivation period (1020–1420 h) and 40 h post-deprivation period ($I = 1$ s.e.d.).




Figure 4 The accumulated lying time (min) for cows in the control (◆), 2-h deprivation (■) and 4-h deprivation (▲) treatments ($I = 1$ s.e.d.) for the 4-h deprivation period (hours –4 to –1) and post-deprivation period (hours 1 to 40), ($I = 1$ s.e.d. between two means) in experiment 1.

Similarly, treatment 2-h cows would have been deprived of 45 and 26 min lying and feeding, respectively. Both treatment 2-h and 4-h cows replaced lying and feeding almost equally with standing not ruminating and standing ruminating during the deprivation period. Munksgaard and Simonsen (1996) reported that cows spent the majority of their time standing ruminating when deprived of lying. Overall, the proportion of time spent ruminating was similar for the cows in treatments C, 2-h and 4-h (41%, 45% and 42%, respectively), which contrasts with the results of a similar, previous experiment (Cooper *et al.*, 2007), in which the time spent ruminating by cows when deprived of only lying decreased with increasing deprivation time. In this previous experiment, the cows had access to feed during lying deprivation, and primarily replaced lying with feeding.

Table 4 Mean post-deprivation milk yields (kg) for cows in the control, 2-h and 4-h treatments in experiment 1

Milking	Mean milk yield (kg) per treatment			P
	Control	2-h	4-h	
First (p.m.)	8.0	8.0	8.0	0.78
Second (a.m.)	11.7	12.0	12.3	0.75
Third (p.m.)	6.3	6.0	5.7	0.15
Fourth (a.m.)	12.0	12.0	12.0	0.76
Combined (p.m. + a.m.)	20.0	19.0	20.0	0.52

Cows responded to the deprivation of lying and feeding in this experiment by leg stamping, repositioning and weight shifting, which has been observed previously as a




Figure 5 Milk yield before and after trimming.

response to lying deprivation (Ruckebusch, 1974; Hopster *et al.*, 2002). The frequency of these behaviours remained approximately constant over the course of the deprivation period for both treatments, suggesting that either any discomfort caused did not increase over time, or that they were not effective coping mechanisms. Butting, although rarely observed, was more frequently performed by cows in treatments 2-h and 4-h. Behaviours such as leg stamping, repositioning and weight shifting may enable the cow to cope with forced standing by alleviating strain on the legs and hoofs. Butting is more likely to be a behavioural expression of frustration, representing an attempt to cope with a negative situation (Sandem *et al.*, 2002). Butting has been previously observed in response to food deprivation in cows (Sandem *et al.*, 2002). The ability of the deprived cows to see other cows eating may have increased the frustration, and may not necessarily occur if cows were isolated for treatment. In addition, the increased butting activity may have partly been due to being confined during deprivation.

Deprived cows were observed to nose the ground at the place where food was usually presented, suggesting that they may have been searching for food. In Cooper *et al.* (2007) we reported that lying-deprived cows sniffed the ground more, but this was interpreted as intentional lying behaviour. Nosing the ground was not observed when cows were deprived of lying only (Cooper *et al.*, 2007), as food was available. Nosing the ground in cows may be the equivalent of beak-wiping in chickens, which is performed when they are prevented from feeding (Preston, 1987). It seems likely that the motivation to feed was constant throughout the deprivation period, as the frequency of this behaviour did not differ between hours. The increase in nosing the ground, leg stamping, repositioning, weight shifting and butting over time indicates increased activity with deprivation, which has also been observed in chickens (Webster, 2000), turkeys (Hocking *et al.*, 1999) and pigs (Day *et al.*, 1995). Typical responses are increased pacing and non-nutritive pecking in fowl, and increased aggression and rooting behaviour in pigs. The increased activity, with the exception of nosing the floor, was probably related to frustration, rather than food seeking, since it was also observed when cows were deprived of lying only (Cooper *et al.*, 2007), and food restriction alone does not increase locomotion in pigs (Day *et al.*, 1995).

Sniffing the ground was positively correlated with leg stamping and repositioning when cows were in treatment 2-h and 4-h, suggesting that the inability to lie down led to frustration. Over the entire deprivation period, the cows rubbed their heads against the housing more when they were deprived of lying and feeding for 4 h than for 2 h. Other research has found that cattle deprived of lying interacted with their surroundings more frequently, i.e. licking or chewing stall fixtures (Munksgaard and Simonsen, 1996; Munksgaard *et al.*, 1999), reflecting a possible need to compensate for a lack of stimulation (Munksgaard and Simonsen, 1996).

Cows deprived of lying and feeding for 4 h slept and drank less. Reduced sleep is associated with stress in other species (Abou-Ismael *et al.*, 2007) and drinking has also been observed in food-deprived chickens (Preston, 1987), pigs (Day *et al.*, 1995) and sheep (Cockram *et al.*, 1999). Water is required while eating to prepare a bolus of the correct moisture content for swallowing and after feeding to clear the mouth of debris, but is also required to maintain a constant osmolarity of rumen contents. In treatment C, the cows spent the most time drinking in the first hour, which is also when they spent the most time feeding. There was no treatment effect on self-licking, in contrast to some studies in which this increased during lying deprivation (Munksgaard and Simonsen, 1996; Munksgaard *et al.*, 1999), but not all (Cooper *et al.*, 2007). However, both Munksgaard and Simonsen (1996) and Munksgaard *et al.* (1999) deprived cows of lying for 14 h/day over several weeks, and the oral behaviours observed by Munksgaard *et al.* (1999) did not increase until week 3. Munksgaard and Simonsen (1996) recorded an associated increased concentration of ACTH in the first hour of deprivation, but others have suggested that these orally centred body-care activities are induced by oxytocin (Van Erp *et al.*, 1993), which remains high for several minutes after cessation of milking (Mayer *et al.*, 1984). The increased rumination in deprived cows may have decreased the motivation for grooming.

Post deprivation

During the first 8-h post deprivation, compared with treatment C, cows in treatments 2-h and 4-h increased their feeding time and not the time spent standing or lying, indicating that the motivation to feed was greater than to lie down after being deprived of both. This is in agreement with Ruckebusch (1974), who also found that cattle chose to feed immediately after being deprived of lying and feeding for 14 and 22 h/day. Munksgaard *et al.* (2005) found that cattle experienced less of a proportional reduction in feeding than lying time during deprivation for 12 h/day for 18 days, and that a reduction in feeding time can be ameliorated by an increase in feed intake rate. There was limited evidence that the more time-restricted cows learnt to prioritise feeding during the early stages of this experiment.

The prevention of feeding by cows in treatments 2-h and 4-h resulted in the greatest increase in feeding time, compared to treatment C, during the first 8 h post deprivation,

in comparison to any other 8-h period. During this time the cows in treatments 2-h and 4-h compensated for about 94% (31 min) and 47% (28 min), respectively, of their lost feeding time. This rebound indicates that the motivation to feed builds up significantly after only a few hours of deprivation, but the extent of feeding may be limited by the gastrointestinal capacity, since cows deprived for 4 h did not feed for significantly longer time than those deprived for 2 h, and cows and pigs deprived for longer periods (14 and 20 h, respectively) have also been observed to only increase feeding time by 2 to 3 h immediately post deprivation (Ruckebusch, 1974; Beattie *et al.*, 2002). Besides gastrointestinal capacity, there is the possibility that biting rate or bite mass was increased, as previously observed in cows (Patterson *et al.*, 1998), sheep (Iason *et al.*, 1999) and pigs (Day *et al.*, 1995) after fasting.

Milk production

In experiment 1, the absence of any effect on milk yields suggests that the increase in feeding time post-treatment was successful in maintaining feed intakes to sustain milk production. Any reduction in milk yield following lying deprivation is small and short-lived (Munksgaard and Løvendahl, 1993; Verkerk *et al.*, 1999; Cooper *et al.*, 2007). In experiment 2, the major reduction in milk yield suggests that the foot trimming caused stress, but of lesser magnitude than relocation stress, which was observed to cause a 10 kg/day yield reduction for 3 days (Varner *et al.*, 1983). More minor stressors cause less effect – tuberculin testing has been observed to reduce milk yield by 1 kg/day (King, 1976) and visitor disturbances resulted in reduced yield by 0.5 kg/day (King, 1978), both of these for about 2 days. Other researchers have found that a single hoof trimming either improved milk production in late lactation (Tanaka *et al.*, 1994) or it did not significantly adversely affect the milk yield of a small sample of cows with healthy hooves (Nishimori *et al.*, 2006). The benefits of improved hoof health and frictional properties following trimming (Phillips *et al.*, 1998) may provide long-term improvements in the ability of cows to feed, whereas the short-term impact may be a reduction in milk yield as a result of stress.

Conclusions

Depriving lactating dairy cows of the ability to lie and feed for 2 to 4 h increases the time ruminating, but they will also spend an approximately equal amount of time to ruminating just standing. Behaviours indicative of discomfort, frustration and tiredness: leg stamping, butting, repositioning themselves and weight shifting – quickly develop, and are at a greater frequency in a 4 h deprivation than in a 2 h one. Nosing the ground may develop as a lying intention and rubbing the head against the housing was increasingly frequent as the length of deprivation increased. Following deprivation, cows deprived for just 2 h restored their feeding time but not lying time. Cows deprived for 4 h did not restore feeding or lying time. Milk yield was transiently affected by the deprivation, but only when a 4 h deprivation

was accompanied by an event that is likely to be stressful, foot trimming, and lasted for 3 days. The well-being of dairy cows is therefore likely to be reduced by temporary deprivation of food and lying, and milk production is likely to be affected if deprivation is accompanied by a stressful procedure.

Acknowledgements

The authors acknowledge the provision of a postgraduate studentship for MDC and facilities for the experiment by Moulton College, a Universities Federation for Animal Welfare research grant and formal supervision for MDC by Dr L Sinclair, Harper Adams University College.

References

- Abou-Ismaïl UA, Burman OHP, Nicol CJ and Mendl M 2007. Can sleep behaviour be used as an indicator of stress in group-housed rats (*Rattus norvegicus*)? *Animal Welfare* 16, 185–188.
- Beattie VE, Burrows BW, Moss BW and Weathercup RN 2002. The effect of food deprivation prior to slaughter on performance, behaviour and meat quality. *Meat Science* 62, 413–418.
- Bolinger DJ, Albright JL, Morrow-Tesch J, Kenyon SJ and Cunningham MD 1997. The effects of restraint using self-locking stanchions on dairy cows in relation to behavior, feed intake, physiological parameters, health, and milk yield. *Journal of Dairy Science* 80, 2411–2417.
- Cockram MS, Kent JE, Waran NK, McGilp LM, Jackson RM, Amory JR, Southall EL, O’Riordan TO, McConnell TI and Wilkins BS 1999. Effects of a 15 h journey followed by either 12 h starvation or *ad libitum* hay on the behaviour and blood chemistry of sheep. *Animal Welfare* 8, 135–148.
- Cooper MD, Arney DR and Phillips CJC 2007. Two- or four-hour lying deprivation on the behavior of lactating dairy cows. *Journal of Dairy Science* 90, 1149–1158.
- Day JEL, Kyriazakis I and Lawrence AB 1995. The Effect of food deprivation on the expression of foraging and exploratory behaviour in the growing pig. *Applied Animal Behaviour Science* 42, 193–206.
- Fregonesi JA and Leaver D 2001. Behaviour, performance and health indicators of welfare for dairy cows housed in strawyard and cubicle systems. *Livestock Production Science* 68, 205–216.
- Greenough PR 1997. Lameness in cattle, 3rd edition. WB Saunders and Co., Los Angeles, USA.
- Hocking PM, Maxwell MH and Mitchell MA 1999. Welfare of food restricted male and female turkeys. *British Poultry Science* 40, 19–29.
- Hopster H, Hermans GN, Engel B and Van der Werf JTN 2002. Behavioural and physiological consequences of deprivation from nightly lying in dairy cows. Proceedings of the 36th International Congress of the International Society of Applied Ethology, Zan am Zee, the Netherlands, 143 p.
- Iason GR, Mantecon AR, Sim DA, Gonzalez J, Foreman E, Bermudez FF and Elston DA 1999. Can grazing sheep compensate for a daily foraging time constraint? *Journal of Animal Ecology* 68, 87–93.
- Kamphues J 1998. Problems in feeding food producing animals to meet the energy and nutrient requirements as well as to enable animals’ well-being. *Deutsche Tierärztliche Wochenschrift* 105, 117–123.
- King JOL 1976. The effect of disturbance on milk production in cows. *The Veterinary Record* 98, 41–42.
- King JOL 1978. The influence of visiting parties on milk production in cows. *The Veterinary Record* 22, 361–362.
- Mayer H, Schams D, Prokopp A and Worstoff H 1984. Effects of manual stimulation and delayed milking on secretion of oxytocin and milking characteristics in dairy cows. *Milchwissenschaft* 39, 666–670.
- Metz J 1985. The reaction of cows to a short-term deprivation of lying. *Applied Animal Behaviour Science* 13, 301–307.
- Munksgaard L and Løvendahl P 1993. Effect of social and physical stressors on growth hormone levels in dairy cows. *Canadian Journal of Animal Science* 73, 847–853.

- Munksgaard L and Simonsen H 1996. Behavioural and pituitary-adrenal axis responses of dairy cows to social isolation and deprivation of lying down. *Journal of Animal Science* 74, 769–778.
- Munksgaard L, Ingvarsten L, Pedersen LJ and Nielsen VKM 1999. Deprivation of lying down affects behaviour and pituitary-adrenal axis responses in young bulls. *Acta Agriculturae Scandinavica (Section A)* 49, 172–178.
- Munksgaard L, Jensen MB, Pedersen LJ, Hansen SW and Matthews L 2005. Quantifying behavioural priorities – effects of time constraints on behaviour of dairy cows, *Bos taurus*. *Applied Animal Behaviour Science* 92, 3–14.
- Nishimori K, Okada K and Ikuta K 2006. The effects of one-time hoof trimming on blood biochemical composition, milk yield, and milk composition in dairy cows. *The Journal of Veterinary Medical Science* 68, 267–270.
- Patterson HD and Lucas HL 1962. Change-over Designs. North Carolina Agricultural Experiment Station and US Department of Agriculture. Technical Bulletin No. 147, 53 pp.
- Patterson DM, McGilloway DA, Cushnahan A, Mayne CS and Laidlaw AS 1998. Effect of duration of fasting period on short-term intake rates of lactating dairy cows. *Animal Science* 66, 299–305.
- Phillips CJC, Coe R, Colgan M, Duffus C, Ingoldby L, Pond M and Postlethwaite S 1998. Effect of hoof characteristics on the propensity to slip in cattle. *The Veterinary Record* 142, 242–245.
- Preston AP 1987. Restricted feeding time and the behaviour of laying hens. *British Poultry Science* 28, 387–396.
- Ruckebusch Y 1974. Sleep deprivation in cattle. *Brain Research* 78, 495–499.
- Sandem AI, Braastad BO and Boe KE 2002. Eye white may indicate emotional state on a frustration-contentedness axis in dairy cows. *Applied Animal Behaviour Science* 79, 1–10.
- Tanaka T, Onashi T, Tanida H and Yoshimoto T 1994. Effects of hoof cutting on milk production of dairy cows. *Journal of Japanese Livestock Management* 30, 45–51.
- Van Erp AM, Kruk MR and De Kloet ER 1993. Induction of grooming in resting rats by intracerebroventricular oxytocin but not by adrenocorticotrophic hormone-(1-24) and alpha-melanocyte-stimulating hormone. *European Journal of Pharmacology* 232, 217–221.
- Varner MA, Johnson BH, McDaniel BT and Mochrie RD 1983. Influence of herd relocation upon production and endocrine traits of dairy cows. *Journal of Dairy Science* 66, 466–474.
- Verkerk GA, Fisher AD, Morrow CJ and Mathews LR 1999. The effects of stressors on milk yield and composition of dairy cows. *Proceedings of the New Zealand Society of Animal Production* 59, 192–194.
- Webster AB 2000. Behaviour of white leghorn laying hens after withdrawal of feed. *Poultry Science* 79, 192–200.