

## **The timing of ankle fracture surgery and the effect on infectious complications;**

### **A case series and systematic review of the literature**

T. Schepers<sup>1,2</sup>, M.R. De Vries<sup>3</sup>, E.M.M. Van Lieshout<sup>1</sup>, M. Van der Elst<sup>3</sup>

<sup>1</sup> Department of Surgery-Traumatology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

<sup>2</sup> Trauma Unit, Department of Surgery, Academic Medical Center, Amsterdam, The Netherlands

<sup>3</sup> Department of Surgery and Traumatology, Reinier de Graaf Groep Delft, The Netherlands

#### **Corresponding author:**

T. Schepers, MD PhD

Erasmus MC, University Medical Center Rotterdam

Department of Surgery-Traumatology

Room H-822k

P.O. Box 2040

3000 CA Rotterdam

The Netherlands

Tel: +31-10-7031050

Fax: +31-10-7032396

E-mail: [t.schepers@erasmusmc.nl](mailto:t.schepers@erasmusmc.nl)

**Conflicts of Interest and Source of Funding:** no conflict of interest or funding is declared by any of the authors.

**The timing of ankle fracture surgery and the effect on infectious complications;**

**A case series and systematic review of the literature**

## **Abstract**

**Purpose:** Ambiguous data exists concerning the influence of delaying surgery on infectious wound complications. A clinical audit was performed to test for the hypothesis that early surgery lowers the rate of infectious wound complications. Secondly we looked at the influence of surgical delay and complications on patient reported functional outcome

**Methods:** All consecutive, closed distal fibular fractures treated surgically with a plate were included and retrospectively analyzed for the delay in operation and wound complications. In a second cohort of patients with a AO-Weber B-type ankle fracture outcome was measured using the Olerud-Molander Ankle Score (OMAS), the American Orthopaedic Foot and Ankle Society score (AOFAS) and a Visual Analog Score (VAS) for overall satisfaction.

**Results:** Patients treated within one day experienced no wound complications (zero out of 60), whereas in the delayed group 11% (16/145) ( $p=0.004$ ). A similar significant difference was found for the patients treated within one week (2/98) versus after one week (14/107). A systematic review of the literature showed a difference in wound complications of 3.6% (early) versus 12.9% (late) ( $p<0.0001$ ). After 43 months, the median AOFAS was 11.5 points lower in the complication group, the OMAS 10 points, and the VAS 0.5 points, with all differences being statistically significant.

**Conclusions:** Every effort should be made to operate closed ankle fractures as soon as reasonably possible. A delay in surgery is associated with a significant rise in infectious wound complications; which significantly lowers outcome and patient satisfaction. These fractures should preferably be treated within the first day.

**Keywords:** Ankle Fractures; ORIF, complications, outcome

## Introduction

Ankle fractures represent approximately ten percent of all fractures and are among the most frequently encountered surgically treated fractures [10]. The incidence of ankle fractures has increased up to 300 percent in the last 30 years, especially in the elderly population [3, 16]. With increasing rates of comorbidities a concomitant rise in fragility fractures can be expected, which in turn may increase the risk of complications. Surgical treatment of ankle fractures may be accompanied by several complications such as nonunion, malunion, implant failure, and wound complications [12]. The most frequently encountered complications are wound complications, of which the infectious complication may have the most devastating effect.

Wound complications that may occur include wound edge necrosis, wound dehiscence, superficial infection, and deep infection (osteomyelitis), and are caused by surgeon, fracture, and patient characteristics. The surgeon characteristics included the timing of surgery[13], placement of the incision, soft tissue handling[23], type of implant[19, 26], and the aftertreatment.[31] The fracture characteristics include for example the severity of the fracture and the presence of a breach of skin integrity. The related patient characteristics include among others age [18, 20], obesity [9], smoking habits [4], substance abuse [32], peripheral vascular disease [28], and diabetes [11, 34].

Ambiguous data exists concerning the influence of delaying surgery on infectious wound complications. We therefore performed a clinical audit to test for the hypothesis that early surgery lowers the rate of infectious wound complications. Secondly we looked at the influence of surgical delay and complications on patient reported functional outcome. The present study is a continuation of an investigation in which a series of ankle fractures from the period 2004 to 2009 were studied [26, 33].

## Material and methods

This is a retrospective study of all consecutive patients, between January 1, 2004 and December 31, 2009, in which plating of the fibula in a closed ankle fracture was performed [26]. All patients received antibiotic prophylaxis according to hospital protocol (third generation cephalosporin). The use of a tourniquet was based on surgeon's preferences. Fracture approach, reduction and fixation were according to AO standards as published previously [26]. Closure was performed in two layers, no suction-drains were used.

Delay until operation and wound complications were recorded from the patient files, operation reports, and the picture archiving and communication system (PACS).

Infectious wound complications and the need for elective hardware removal were scored. Wound infection was further separated into superficial or minor and deep or major infectious complications by applying the criteria of the Centers for Disease Control and Prevention (CDC) for defining a surgical site infection[21]. Minor complications were defined as a superficial infection treatable with conservative management like oral antibiotics only, without the need for intervention or re-admission. Major complications were defined as a deep infection in need for intervention or re-admission, like intravenous antibiotics, removal of hardware, wound debridement with or without vacuum assisted closure [26]. The differentiation between superficial and deep infection was made by the surgeon or attending physician.

In a second series of consecutive patients aged between 16-65 years with an AO-Weber B type ankle fracture, treated between January 2004 and July 2009 [33], the influence of infectious wound complications on the overall outcome was investigated. Only the Weber-B type fractures were included in order to obtain a more homogenous population. Minimum follow-up was six months. Patient characteristics (*i.e.*, age, gender, diabetes, and smoking habits), fracture characteristics (*i.e.*, fracture side and number of malleoli involved) were recorded. Outcome was measured using standard questionnaires, which consisted of the

Olerud-Molander Ankle Score (OMAS), the American Orthopaedic Foot and Ankle Society Ankle-hindfoot Score (AOFAS), and a single question Visual Analog Scale (VAS) for patient satisfaction with outcome. Two patients with a complication other than a wound infection were excluded (n=2; one complex regional pain syndrome and one deep venous thrombosis) in order to compare patients with versus without infectious complications.

### *Literature search*

A literature search was conducted in order to identify studies in which ankle fractures were treated surgically. The electronic databases up to December 1, 2011 of the Cochrane Library, Pubmed Medline, EMBase, and Google Scholar were explored using the combination of the following search terms and Boolean operators: ankle OR fibula AND complication OR infection OR timing. No restriction in language and publication date was applied. In addition, a comprehensive search of reference lists of all identified articles was conducted in order to find additional studies. An article was found eligible when it concerned 1) the surgical treatment of an ankle fracture, 2) reporting of the number of wound complications, and 3) reporting on two groups treated early or delayed. Abstracts from scientific meetings were included in the current review, when sufficient data could be extracted on functional outcome or complication rate. The cut-off point for early and delayed surgery, the group sizes, and the number of wound complications were extracted from these publications.

### *Data analysis*

The statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) version 16.0 (SPSS, Chicago, IL). The Kolmogorov-Smirnov test was used to test for normality of the data. The Levene's test was applied to assess homogeneity of variance

between groups. Since most numeric variables did not show normal distribution or equal variance, all items were regarded as nonparametric for the statistical analysis. A Mann-Whitney U-test (numeric data), Chi2 analysis or Fisher exact test (categorical data) was performed in order to assess statistical significance of difference between the number of complications in the early and delayed treated patients, and to test for difference in outcome between groups with and without complications. A p-value <0.05 was taken as level of statistical significance. Numeric data are expressed as medians with P<sub>25</sub>-P<sub>75</sub>; categorical data are shown as numbers with percentages.



## Results

### *Influence of surgical delay on complications*

A total of 205 ankle fractures were treated surgically in the six year study period. All received plating of the fibula. The number of fractured malleoli was as followed: 117 unimalleolar, 57 bimalleolar, and 31 trimalleolar. There were 26 fracture-dislocations. The median surgical delay was 7 days for the entire population ( $P_{25}$ - $P_{75}$  1-11 days).

In the unimalleolar group 22.2 percent was treated within the first day (24 hours) and in the bi- or trimalleolar group this percentage was 38.6% ( $p=0.013$ ). None of the 60 ankle fractures treated within one day developed an infectious wound complications, whereas in the group treated after day one 11% (16 out of 145) did develop an infectious wound complications ( $p=0.004$ ). Ten were minor and six were major complication (Figure 1a).

When taking one week as cut-off point for early treatment 43.6 percent of the unimalleolar fractures were treated early versus 53.4 percent of the bi- or trimalleolar fractures ( $p=0.204$ ). Two percent of the patients treated within six days experienced a complication (two out of 98), which were both major complications. Patients treated from day seven experienced an infectious wound complication in 13.1% of cases (14 out of 107; ten minor and four major), which was a significant difference ( $p=0.003$ ).

The complication rate in the unimalleolar group was 5.1% (3 minor and 3 major out of 107 fractures) and in the bi- or trimalleolar group this number was 11.4% (7 minor and 3 major out of 88 fractures). This difference was not statistically significant (Figure 1b;  $p=0.118$ ). Using a post hoc power analysis the groups needed to be three times the size in order to detect a significant difference (with 80% power, two-sided testing, and an alpha of 0.05).

### *Systematic review influence delay on complications*

Including the results of the current study, eleven publications were included in the literature review (Table 1). Considering wound complications and the effect of early versus delayed surgical treatment six showed a significant increase with delaying surgery, two showed a trend for increase complications, and three studies did not find a difference between early and delayed surgery. In the early treatment group 24 events occurred in 673 patients (3.6%) and in the delayed group 66 events occurred in 513 patients (12.9%). This difference was statistically significant ( $P < 0.0001$ ).

#### *Influence delay and complications on outcome*

The second series was comprised of 101 patients, of which 14 experienced an infectious wound complication. With a median follow-up of 43 ( $P_{25}$ - $P_{75}$ : 29–60) months the overall outcome was significantly influenced by the occurrence of an infectious wound complication. The median AOFAS was 11.5 points lower in the complication group, the OMAS 10 points, and the VAS 0.5 points (Table 2). With the exception of gender the two groups of Weber-B fractures were comparable considering age, BMI, number of fractured malleoli, fracture dislocation, smoking, and diabetes.

Patients with a delay of one day to surgery showed no difference in AOFAS and VAS, but a significant five point reduction in the median OMAS (90 versus 85 points). This five point difference in the OMAS was also present when comparing the groups with a delay of one week, but not significant ( $p=0.057$ ). The AOFAS and VAS did not differ between these groups.

## Discussion

The current study shows a significant difference in infectious wound complication for patients treated delayed for a closed ankle fracture. Independent of the cut-off point of 24 hours or one week the chance of an infection is at least a factor six higher in the delayed group. This difference was also noted in the systematic review where 3.6% infectious wound complications occurred compared with 12.9% in the delayed group. More important this increase in complications has a direct negative effect on functional outcome as measure with three renowned scoring systems.

Previous studies on early versus delayed surgical treatment mainly focus on the length of stay and the concomitant increase in costs (up to 1250 Euro per case) with a longer hospital stay [6, 15, 24]. Others have pointed out an improved rate of anatomical reductions in early management [7-8].

Avoiding wound complications is of paramount importance and should receive a high priority. Timing of the surgical stabilization appears to be a controllable factor to achieve lower complication rates. Another factor that can be influenced is the choice of implant, e.g., thinner plates [26] or fibular nailing [2] in fragility fractures or in patients with comorbidities.

The effect of wound complications in general on outcome has been noted in one previous study with an almost identical duration of follow-up as the current study [14]. Especially the major infectious complications (n=2) in that study were associated with a worse outcome, however the effect on functional outcome was not reported. Surgical delay was not associated with a poorer outcome in this study, which is in concurrence with a more recent publication [5]. Our results on the other hand show a lower OMAS in fractures treated after a surgical delay.

Other factors, other than trauma and surgical characteristics, which have been identified as negatively influencing outcome are obesity, smoking, alcohol abuse, and a lower level of education [5, 29]. These parameters were not assessed in the current study.

A limitation of the current study is its retrospective design, as were the included studies in the systematic review. The reasons for postponing the surgical repair were not recorded. In the current study it was the policy to plan the fractures electively and to wait for approximately a week to let the swelling settle, which is not uncommon practice [22]. In some cases a conservative treatment was initiated, but because of secondary dislocation a change to surgery was made. In these patients the delay was longer than one week. The percentage of bi- and trimalleolar fractures treated early was higher than the unimalleolar fractures. Thus, the more complex fractures were apparently not postponed more frequently because of soft-tissue problems.

Open fractures form a completely different entity as orthopaedic emergency and were not included in this study. In case of a fracture-dislocation (n=26) a closed reduction was tried first, which resulted in an early surgical intervention if not successful. If fracture-dislocations were removed from the analyses the difference in complications with cut-off point 1 day remained (0/46 vs 14/133; p=0.022), as well as the difference with cut-off point 1 week (2/81 vs 12/98; p=0.023).

In conclusion, every effort should be made to operate closed ankle fractures as soon as reasonably possible. A delay of more than one week gives a significant rise in infectious wound complications, which significantly lowers functional outcome and patient satisfaction. The fractures should preferably be treated within the first day. Literature data also support that early operative treatment gives a significant reduction in wound complications and a concomitant improved functional outcome for the patients.

## References

1. Adamson SP, Trickett R, Hodgson P, Mohanty K (2009) Ankle fractures: Impact of timing of surgery. *Injury Extra* 40:224
2. Appleton PT, McQueen MM, Court-Brown CM (2006) The Fibula Nail for Treatment of Ankle Fractures in Elderly and High Risk Patients. *Techniques in Foot and Ankle Surgery* 5:204-208
3. Bengner U, Johnell O, Redlund-Johnell I (1986) Epidemiology of ankle fracture 1950 and 1980. Increasing incidence in elderly women. *Acta Orthop Scand* 57:35-37
4. Berkes M, Obremskey WT, Scannell B, Ellington JK, Hymes RA, Bosse M) Maintenance of hardware after early postoperative infection following fracture internal fixation. *J Bone Joint Surg Am* 92:823-828
5. Bhandari M, Sprague S, Ayeni OR, Hanson BP, Moro JK (2004) A prospective cost analysis following operative treatment of unstable ankle fractures: 30 patients followed for 1 year. *Acta Orthop Scand* 75:100-105
6. Breederveld RS, van Straaten J, Patka P, van Mourik JC (1988) Immediate or delayed operative treatment of fractures of the ankle. *Injury* 19:436-438
7. Carragee EJ, Csongradi JJ (1993) Increased rates of complications in patients with severe ankle fractures following interinstitutional transfers. *J Trauma* 35:767-771
8. Carragee EJ, Csongradi JJ, Bleck EE (1991) Early complications in the operative treatment of ankle fractures. Influence of delay before operation. *J Bone Joint Surg Br* 73:79-82
9. Chaudhry S, Egol KA (2011) Ankle injuries and fractures in the obese patient. *Orthop Clin North Am* 42:45-53, vi
10. Court-Brown CM, Caesar B (2006) Epidemiology of adult fractures: A review. *Injury* 37:691-697
11. Flynn JM, Rodriguez-del Rio F, Piza PA (2000) Closed ankle fractures in the diabetic patient. *Foot Ankle Int* 21:311-319

12. Hess F, Sommer C (2011) Minimally invasive plate osteosynthesis of the distal fibula with the locking compression plate: first experience of 20 cases. *J Orthop Trauma* 25:110-115
13. Hoiness P, Stromsoe K (2000) The influence of the timing of surgery on soft tissue complications and hospital stay. A review of 84 closed ankle fractures. *Ann Chir Gynaecol* 89:6-9
14. Hoiness P, Engebretsen L, Stromsoe K (2001) The influence of perioperative soft tissue complications on the clinical outcome in surgically treated ankle fractures. *Foot Ankle Int* 22:642-648
15. James LA, Sookhan N, Subar D (2001) Timing of operative intervention in the management of acutely fractured ankles and the cost implications. *Injury* 32:469-472
16. Kannus P, Palvanen M, Niemi S, Parkkari J, Jarvinen M (2002) Increasing number and incidence of low-trauma ankle fractures in elderly people: Finnish statistics during 1970-2000 and projections for the future. *Bone* 31:430-433
17. Konrath G, Karges D, Watson JT, Moed BR, Cramer K (1995) Early versus delayed treatment of severe ankle fractures: a comparison of results. *J Orthop Trauma* 9:377-380
18. Koval KJ, Zhou W, Sparks MJ, Cantu RV, Hecht P, Lurie J (2007) Complications after ankle fracture in elderly patients. *Foot Ankle Int* 28:1249-1255
19. Lamontagne J, Blachut PA, Broekhuysen HM, O'Brien PJ, Meek RN (2002) Surgical treatment of a displaced lateral malleolus fracture: the antiglide technique versus lateral plate fixation. *J Orthop Trauma* 16:498-502
20. Lynde MJ, Sautter T, Hamilton GA, Schuberth JM (2012) Complications after open reduction and internal fixation of ankle fractures in the elderly. *Foot Ankle Surg* 18:103-107
21. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR (1999) Guideline for Prevention of Surgical Site Infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control* 27:97-132; quiz 133-134; discussion 196

22. Miller AG, Margules A, Raikin SM (2012) Risk factors for wound complications after ankle fracture surgery. *J Bone Joint Surg Am* 94:2047-2052
23. Ovaska M, Lindahl J, Mäkinen T, Madanat R, Pulliainen L, Kiljunen V, Hirvensalo E, Tukiainen E (2011) Postoperative infection after closed and open ankle fractures. *Suomen Ortopedia ja Traumatologia* 34:30-33
24. Pietzik P, Qureshi I, Langdon J, Molloy S, Solan M (2006) Cost benefit with early operative fixation of unstable ankle fractures. *Ann R Coll Surg Engl* 88:405-407
25. Saithna A, Moody W, Jenkinson E, Almazedi B, Sargeant I (2009) The influence of timing of surgery on soft tissue complications in closed ankle fractures. *Eur J Orthop Surg Traumatol* 19:481-484
26. Schepers T, Van Lieshout EM, De Vries MR, Van der Elst M (2011) Increased rates of wound complications with locking plates in distal fibular fractures. *Injury* 42:1125-1129
27. Singh BI, Balaratnam S, Naidu V (2005) Early versus delayed surgery for ankle fractures: a comparison of results. *Eur J Orthop Surg Traumatol* 15:23-27
28. SooHoo NF, Krenke L, Eagan MJ, Gurbani B, Ko CY, Zingmond DS (2009) Complication rates following open reduction and internal fixation of ankle fractures. *J Bone Joint Surg Am* 91:1042-1049
29. Still GP, Atwood TC (2009) Operative outcome of 41 ankle fractures: a retrospective analysis. *J Foot Ankle Surg* 48:330-339
30. Sukeik M, Qaffaf M, Ferrier G (2010) Ankle fractures: impact of swelling on timing of surgery, length of hospital stay and the economic burden. *Injury Extra* 41:133
31. Thomas G, Whalley H, Modi C (2009) Early mobilization of operatively fixed ankle fractures: a systematic review. *Foot Ankle Int* 30:666-674
32. Tonnesen H, Pedersen A, Jensen MR, Moller A, Madsen JC (1991) Ankle fractures and alcoholism. The influence of alcoholism on morbidity after malleolar fractures. *J Bone Joint Surg Br* 73:511-513

33. Van Schie-Van der Weert EM, Van Lieshout EM, De Vries MR, Van der Elst M, Schepers T (2012)  
Determinants of outcome in operatively and non-operatively treated Weber-B ankle fractures.  
Arch Orthop Trauma Surg 132:257-263
34. Wukich DK, Kline AJ (2008) The management of ankle fractures in patients with diabetes. J  
Bone Joint Surg Am 90:1570-1578



**Table 1.** Literature overview showing influence of timing of ankle fracture surgery and wound complications

Study (year)	Cut-off point	Wound complications		Conclusion
		Early	Delayed	
<b>Breederveld (1988)[6]</b>	24 hours	3/72 (2 in open#)	2/20	No significant difference
<b>Caragee (1991)[8]</b>	4 days*	17.9%	39.5%	P = 0.008 all complications
<b>Caragee (1993)[7]</b>	Same day	3/43	10/34	P < 0.001
<b>Konrath (1995)[17]</b>	5 days	5/105	6/97	No significant difference
<b>Hoiness (2000)[13]</b>	< 8h or >5d**	5/67	7/17	P = 0.0018
<b>Singh (2005)[27]</b>	24 hours	0/24	6/38	Trend (P = 0.07)
<b>Pietzik (2006)[24]</b>	48 hours	0/62	1/21	No significant difference
<b>Adamson (2009)[1]</b>	24 hours	1/67	8/84	P = 0.04
<b>Saithna (2009)[25]</b>	6 days	2/56	6/29	P = 0.01
<b>Sukeik (2010)[30]</b>	24 hours	5/117	4/28	Trend (P = 0.07)
<b>Current study</b>	24 hours	0/60	16/145	P = 0.004
<b>Combined***</b>		<b>24/673</b>	<b>66/513</b>	<b>P &lt; 0.0001</b>
		<b>(3.6%)</b>	<b>(12.9%)</b>	

\* Combination of all types of complications (infection, reduction, other) reported

\*\* Wound infection and wound edge necrosis

\*\*\* Caragee 1991 not included, 2-sided Fisher exact test

**Table 2.** Demographics and outcome of surgically treated Weber-B ankle fracture with or without infectious wound complication

<b>Parameter</b>	<b>Total group (n=101)</b>	<b>Without complication (n=87)</b>	<b>With complications (n=14)</b>	<b>p value</b>
<b>Gender (%male)</b>	40 (39.6%)	30 (34.5%)	19 (71.4)	<b>0.016</b> <sup>1</sup>
<b>Age (years)</b>	50.7 (40.9-60.7)	51.7 (41.5-60.7)	46.7 (22.1-63.3)	0.814 <sup>2</sup>
<b>BMI (kg/m2)</b>	25.9 (23.8-29.2)	25.9 (23.5-28.9)	25.9 (24.1-29.6)	0.734 <sup>2</sup>
<b>Unimalleolar %</b>	69 (68.3%)	60 (69.0)	9 (64.3%)	0.115 <sup>1</sup>
<b>Fracture dislocation (%)</b>	12 (11.9%)	9 (10.3%)	3 (21.4%)	0.366 <sup>1</sup>
<b>Smoking (%)</b>	21 (20.8%)	17 (19.5%)	4 (28.6%)	0.482 <sup>1</sup>
<b>Diabetic (%)</b>	6 (5.9%)	5 (5.7%)	1 (7.1%)	1.000 <sup>1</sup>
<b>AOFAS</b>	97 (82-100)	97 (84-100)	85.5 (65-98.5)	<b>0.036</b> <sup>2</sup>
<b>OMAS</b>	90 (80-100)	90 (80-100)	80 (66.3-90)	<b>0.012</b> <sup>2</sup>
<b>VAS</b>	8 (7-9)	8 (7-9.1)	7.5 (6-8)	<b>0.027</b> <sup>2</sup>


Data are shown as number with percentages, or as median with P<sub>25</sub>-P<sub>75</sub>.

BMI; body mass index, AOFAS; American Orthopaedic Foot Ankle Society hindfoot score, OMAS; Olerud Molander Ankle Score, VAS; Visual Analogue Scale

<sup>1</sup> Fisher exact test

<sup>2</sup> Mann-Whitney U test

**Figure 1.** Kaplan-Meier plot of complications in relation to delay



Cumulative number of patients with an infectious wound complication in relation to surgical delay for the total study sample (A), and the difference in events in unimalleolar and bi- or trimalleolar fractures (B).