

Acta Crystallographica Section E

Structure Reports**Online**

ISSN 1600-5368

(E)-1-[2-(4-Fluoro-2-nitrostyryl)-1-phenylsulfonyl-1H-indol-3-yl]propan-1-one

**M. Umadevi,^a V. Saravanan,^b R. Yamuna,^{c*}
A. K. Mohanakrishnan^b and G. Chakkavarthi^{d*}**

^aResearch Scholar (Chemistry), Bharathiyar University, Coimbatore 641 046, Tamilnadu, India, ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India, ^cDepartment of Sciences, Chemistry and Materials Research Lab, Amrita Vishwa Vidyapeetham University, Ettimadai, Coimbatore 641 112, India, and ^dDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India
Correspondence e-mail: ryamuna1@gmail.com, chakkavarthi_2005@yahoo.com


Received 10 November 2013; accepted 11 November 2013

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.055; wR factor = 0.176; data-to-parameter ratio = 26.6.

In the title compound, $\text{C}_{25}\text{H}_{19}\text{FN}_2\text{O}_5\text{S}$, the substituted phenyl ring makes a dihedral angle of $12.26(9)^\circ$ with the indole ring system. The nitro group is twisted at an angle of $26.92(8)^\circ$ out of the plane of the ring to which it is attached. The molecular structure is stabilized by weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. In the crystal, weak $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{F}$ and $\pi-\pi$ [centroid-centroid distance = $3.6645(11)\text{ \AA}$] interactions link the molecules, forming a three-dimensional network.

Related literature

For the biological activity of indole derivatives, see: Pomarnacka & Kozlarska-Kedra (2003); Srivastava *et al.* (2011). For related structures, see: Chakkavarthi *et al.* (2008, 2010). For details of the configuration at the S atom, see: Bassindale (1984). For details of N-atom hybridization, see: Beddoes *et al.* (1986).

**Experimental***Crystal data*

$\text{C}_{25}\text{H}_{19}\text{FN}_2\text{O}_5\text{S}$
 $M_r = 478.48$
Triclinic, $P\bar{1}$
 $a = 8.2615(3)\text{ \AA}$
 $b = 10.7624(5)\text{ \AA}$
 $c = 13.2432(6)\text{ \AA}$
 $\alpha = 68.606(2)^\circ$
 $\beta = 80.554(3)^\circ$
 $\gamma = 81.012(2)^\circ$
 $V = 1075.53(8)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.20\text{ mm}^{-1}$
 $T = 295\text{ K}$
 $0.30 \times 0.24 \times 0.20\text{ mm}$

Data collection

Bruker APEXII diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.942$, $T_{\max} = 0.961$
27388 measured reflections
8185 independent reflections
5506 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.176$
 $S = 1.03$
308 parameters
8185 reflections
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.96\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.61\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C8—H8 \cdots O1	0.93	2.33	2.913 (3)	120
C11—H11 \cdots O3	0.93	2.40	2.905 (3)	114
C16—H16A \cdots F1 ⁱ	0.97	2.54	3.192 (2)	124
C22—H22 \cdots O4 ⁱⁱ	0.93	2.52	3.438 (2)	170

Symmetry codes: (i) $-x + 1, -y - 1, -z + 2$; (ii) $x + 1, y, z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors wish to acknowledge the SAIF, IIT, Madras, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT6945).

References

- Bassindale, A. (1984). In *The Third Dimension in Organic Chemistry*. New York: John Wiley and Sons.
- Beddoes, R. L., Dalton, L., Joule, T. A., Mills, O. S., Street, J. D. & Watt, C. I. F. (1986). *J. Chem. Soc. Perkin Trans. 2*, pp. 787–797.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chakkavarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2008). *Acta Cryst. E64*, o749.
- Chakkavarthi, G., Panchatcharam, R., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2010). *Acta Cryst. E66*, o2957.
- Pomarnacka, E. & Kozlarska-Kedra, I. (2003). *Il Farmaco*, **58**, 423–429.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Srivastava, Anupam & Pandeya, S. N. (2011). *JCPD*, **1**, 1–17.

supporting information

Acta Cryst. (2013). E69, o1780 [doi:10.1107/S1600536813030961]

(E)-1-[2-(4-Fluoro-2-nitrostyryl)-1-phenylsulfonyl-1*H*-indol-3-yl]propan-1-one

M. Umadevi, V. Saravanan, R. Yamuna, A. K. Mohanakrishnan and G. Chakkavarthi

S1. Comment

Indole derivatives are known to exhibit antimicrobial, antibiotic, analgesic, anticancer and anti-HIV (Pomarnacka & Kozlarska-Kedra, 2003; Srivastava *et al.*, 2011) activities. In continuation of our studies on indole derivatives, we determined the crystal structure of the title compound (I). The geometric parameters of (I) (Fig. 1) are agree well with the reported structures (Chakkavarthi *et al.*, 2008; 2010).

Due to Thorpe-Ignold effect (Bassindale, 1984), bond angles around atom S1 show significant deviation from ideal tetrahedral value, with significant deviations in angles O1—S1—O2 [120.42 (8) $^{\circ}$] and N1—S1—C1 [104.66 (7) $^{\circ}$]. The phenyl ring (C1—C6) makes the dihedral angle of 85.05 (8) $^{\circ}$ with the indole ring system. The phenyl ring (C1—C6) and the benzene ring (C20—C25) are inclined at an angle of 12.26 (9) $^{\circ}$. The nitro group is twisted at an angle of 26.92 (8) $^{\circ}$ with the attached benzene ring (C20—C25). The sum of the bond angles around N1 (358.26 $^{\circ}$) indicates the sp^2 hybridization of N1 atom (Beddoes *et al.*, 1986).


The molecular structure is stabilized by weak intramolecular C—H \cdots O hydrogen bonds (Table 1). The crystal structure exhibit weak intermolecular C—H \cdots O, C—H \cdots F (Table 1 & Fig. 2) and $\pi\cdots\pi$ [$Cg4\cdots Cg4^i = 3.6645$ (11) Å; (i) 1 - x , -1 - y , 2 - z ; $Cg4$ is the centroid of the ring (C20—C25)] interactions.

S2. Experimental


A solution of 1-(2-(bromomethyl)-1-(phenylsulfonyl)-1*H*-indol-3-yl) propan-1-one (5 g, 12.31 mmol) and triphenyl-phosphine (3.5 g, 13.54 mmol) in dry THF (100 ml) was refluxed for 6 h. After consumption of the starting material, the solvent was removed under vacuo and the solid was washed with diethyl ether to give the phosphonium salt. Then, the mixture of phosphonium salt (8 g, 11.97 mmol), 4-fluoro-2-nitrobenzaldehyde (2.24 g, 13.17 mmol) and K_2CO_3 (3.30 g, 23.95 mmol) in DCM (70 ml) was stirred at room temperature for 22 h. After completion of the reaction (monitored by TLC), it was diluted using DCM (30 ml), washed with water (2 x 100 ml) and dried (Na_2SO_4). Removal of solvent *in vacuo* followed by trituration of the crude product with MeOH (20 ml) afforded the title compound suitable for X-ray diffraction quality.

S3. Refinement

The H atoms were positioned geometrically and refined using a riding model with C—H = 0.93–0.97 Å, and with $U_{iso}(H) = 1.2U_{eq}(C)$ (or) $U_{iso}(H) = 1.5U_{eq}(C)$.

**Figure 1**

The molecular structure of the title compound, with atom labels and 30% probability displacement ellipsoids for non-H atoms.

**Figure 2**

The packing of the title compound, view onto the ac plane. Intermolecular hydrogen bonds are shown as dashed lines. H atoms not involving hydrogen bonding have been omitted.

(E)-1-[2-(4-Fluoro-2-nitrostyryl)-1-phenylsulfonyl-1*H*-indol-3-yl]propan-1-one

Crystal data

$C_{25}H_{19}FN_2O_5S$
 $M_r = 478.48$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 8.2615 (3) \text{ \AA}$
 $b = 10.7624 (5) \text{ \AA}$
 $c = 13.2432 (6) \text{ \AA}$
 $\alpha = 68.606 (2)^\circ$
 $\beta = 80.554 (3)^\circ$

$\gamma = 81.012 (2)^\circ$
 $V = 1075.53 (8) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 496$
 $D_x = 1.477 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 5619 reflections
 $\theta = 2.1\text{--}31.1^\circ$
 $\mu = 0.20 \text{ mm}^{-1}$

$T = 295\text{ K}$
Block, colourless

$0.30 \times 0.24 \times 0.20\text{ mm}$

Data collection

Bruker APEXII
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω and φ scan
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.942$, $T_{\max} = 0.961$

27388 measured reflections
8185 independent reflections
5506 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\max} = 35.0^\circ$, $\theta_{\min} = 1.7^\circ$
 $h = -12 \rightarrow 12$
 $k = -16 \rightarrow 17$
 $l = -20 \rightarrow 21$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.176$
 $S = 1.03$
8185 reflections
308 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.089P)^2 + 0.2703P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.96\text{ e }{\text{\AA}}^{-3}$
 $\Delta\rho_{\min} = -0.61\text{ e }{\text{\AA}}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.05337 (18)	0.17906 (15)	0.76350 (12)	0.0387 (3)
C2	-0.0663 (2)	0.06833 (18)	0.85866 (14)	0.0476 (4)
H2	0.0275	0.0203	0.8901	0.057*
C3	-0.2214 (2)	0.0306 (2)	0.90598 (16)	0.0563 (4)
H3	-0.2324	-0.0430	0.9702	0.068*
C4	-0.3602 (2)	0.1021 (2)	0.85825 (16)	0.0540 (4)
H4	-0.4641	0.0762	0.8907	0.065*
C5	-0.3457 (2)	0.2109 (2)	0.76326 (16)	0.0548 (4)
H5	-0.4397	0.2576	0.7314	0.066*
C6	-0.1920 (2)	0.25156 (19)	0.71465 (14)	0.0489 (4)
H6	-0.1817	0.3257	0.6507	0.059*
C7	0.1475 (2)	0.19323 (17)	0.50807 (12)	0.0439 (3)
C8	0.0691 (3)	0.3166 (2)	0.44803 (15)	0.0596 (5)
H8	0.0385	0.3851	0.4765	0.072*

C9	0.0389 (3)	0.3323 (2)	0.34425 (16)	0.0668 (6)
H9	-0.0130	0.4135	0.3020	0.080*
C10	0.0836 (3)	0.2308 (2)	0.30181 (15)	0.0637 (5)
H10	0.0627	0.2457	0.2312	0.076*
C11	0.1583 (2)	0.1080 (2)	0.36119 (13)	0.0544 (4)
H11	0.1858	0.0397	0.3322	0.065*
C12	0.1919 (2)	0.08841 (17)	0.46722 (12)	0.0429 (3)
C13	0.27195 (19)	-0.02364 (15)	0.54893 (12)	0.0403 (3)
C14	0.27322 (18)	0.01380 (14)	0.63737 (11)	0.0367 (3)
C15	0.3524 (3)	-0.14616 (18)	0.52664 (14)	0.0522 (4)
C16	0.4209 (3)	-0.26746 (18)	0.61156 (15)	0.0563 (4)
H16A	0.3365	-0.2957	0.6730	0.068*
H16B	0.5119	-0.2449	0.6373	0.068*
C17	0.4816 (4)	-0.3835 (2)	0.5708 (2)	0.0767 (7)
H17A	0.3896	-0.4137	0.5537	0.115*
H17B	0.5345	-0.4558	0.6265	0.115*
H17C	0.5589	-0.3542	0.5065	0.115*
C18	0.34816 (19)	-0.05675 (15)	0.73895 (11)	0.0380 (3)
H18	0.4365	-0.0220	0.7510	0.046*
C19	0.29602 (19)	-0.16804 (15)	0.81477 (11)	0.0380 (3)
H19	0.2011	-0.1976	0.8059	0.046*
C20	0.37971 (18)	-0.24690 (14)	0.91153 (11)	0.0360 (3)
C21	0.5492 (2)	-0.24341 (16)	0.90850 (14)	0.0454 (3)
H21	0.6053	-0.1874	0.8457	0.054*
C22	0.6361 (2)	-0.31934 (18)	0.99463 (15)	0.0510 (4)
H22	0.7478	-0.3125	0.9913	0.061*
C23	0.5542 (2)	-0.40558 (17)	1.08570 (15)	0.0511 (4)
C24	0.3899 (2)	-0.41643 (16)	1.09468 (13)	0.0462 (4)
H24	0.3370	-0.4760	1.1569	0.055*
C25	0.30441 (19)	-0.33571 (15)	1.00810 (11)	0.0384 (3)
N1	0.20107 (17)	0.14790 (13)	0.61298 (10)	0.0410 (3)
N2	0.12683 (19)	-0.34714 (16)	1.02366 (11)	0.0489 (3)
O1	0.12824 (18)	0.36656 (12)	0.64011 (11)	0.0580 (3)
O2	0.25410 (15)	0.17470 (13)	0.78405 (10)	0.0492 (3)
O3	0.3708 (3)	-0.1422 (2)	0.43278 (14)	0.1065 (8)
O4	0.03576 (17)	-0.25160 (16)	0.97436 (13)	0.0684 (4)
O5	0.0783 (2)	-0.45267 (19)	1.08539 (15)	0.0913 (6)
S1	0.14288 (5)	0.22764 (4)	0.70332 (3)	0.04104 (11)
F1	0.64046 (18)	-0.48260 (14)	1.16881 (11)	0.0778 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0383 (7)	0.0433 (7)	0.0372 (7)	-0.0005 (5)	-0.0030 (5)	-0.0194 (6)
C2	0.0446 (8)	0.0492 (9)	0.0453 (8)	-0.0023 (6)	-0.0044 (6)	-0.0135 (7)
C3	0.0536 (10)	0.0562 (10)	0.0553 (10)	-0.0114 (8)	0.0045 (8)	-0.0171 (8)
C4	0.0426 (8)	0.0676 (11)	0.0620 (11)	-0.0103 (8)	0.0031 (7)	-0.0366 (9)
C5	0.0401 (8)	0.0756 (12)	0.0585 (10)	0.0042 (8)	-0.0096 (7)	-0.0372 (10)

C6	0.0465 (9)	0.0570 (10)	0.0430 (8)	0.0034 (7)	-0.0073 (6)	-0.0199 (7)
C7	0.0439 (8)	0.0475 (8)	0.0315 (6)	-0.0027 (6)	-0.0015 (6)	-0.0055 (6)
C8	0.0646 (11)	0.0569 (11)	0.0416 (9)	0.0094 (9)	-0.0054 (8)	-0.0053 (8)
C9	0.0628 (12)	0.0732 (13)	0.0416 (9)	0.0056 (10)	-0.0100 (8)	0.0033 (9)
C10	0.0587 (11)	0.0875 (15)	0.0335 (8)	-0.0111 (10)	-0.0098 (7)	-0.0047 (9)
C11	0.0584 (10)	0.0704 (12)	0.0337 (7)	-0.0154 (9)	-0.0069 (7)	-0.0131 (7)
C12	0.0430 (8)	0.0509 (8)	0.0304 (6)	-0.0113 (6)	-0.0013 (5)	-0.0078 (6)
C13	0.0456 (8)	0.0412 (7)	0.0322 (6)	-0.0095 (6)	-0.0010 (5)	-0.0100 (5)
C14	0.0390 (7)	0.0370 (7)	0.0303 (6)	-0.0049 (5)	-0.0004 (5)	-0.0082 (5)
C15	0.0694 (11)	0.0485 (9)	0.0438 (8)	-0.0094 (8)	-0.0045 (8)	-0.0216 (7)
C16	0.0739 (12)	0.0455 (9)	0.0468 (9)	0.0002 (8)	0.0042 (8)	-0.0202 (7)
C17	0.1053 (19)	0.0522 (11)	0.0682 (13)	-0.0006 (11)	0.0154 (13)	-0.0298 (10)
C18	0.0422 (7)	0.0379 (7)	0.0326 (6)	-0.0020 (5)	-0.0038 (5)	-0.0118 (5)
C19	0.0397 (7)	0.0407 (7)	0.0315 (6)	-0.0033 (5)	-0.0026 (5)	-0.0110 (5)
C20	0.0412 (7)	0.0340 (6)	0.0318 (6)	-0.0024 (5)	-0.0033 (5)	-0.0116 (5)
C21	0.0423 (8)	0.0421 (8)	0.0453 (8)	-0.0035 (6)	-0.0041 (6)	-0.0085 (6)
C22	0.0444 (8)	0.0480 (9)	0.0570 (10)	-0.0009 (7)	-0.0135 (7)	-0.0127 (7)
C23	0.0628 (11)	0.0431 (8)	0.0458 (8)	0.0028 (7)	-0.0217 (8)	-0.0105 (7)
C24	0.0606 (10)	0.0419 (8)	0.0334 (7)	-0.0055 (7)	-0.0074 (6)	-0.0089 (6)
C25	0.0460 (8)	0.0367 (7)	0.0322 (6)	-0.0055 (5)	-0.0031 (5)	-0.0119 (5)
N1	0.0473 (7)	0.0399 (6)	0.0302 (5)	0.0019 (5)	-0.0018 (5)	-0.0093 (5)
N2	0.0490 (8)	0.0547 (8)	0.0382 (7)	-0.0129 (6)	0.0003 (6)	-0.0096 (6)
O1	0.0655 (8)	0.0389 (6)	0.0634 (8)	-0.0044 (5)	0.0000 (6)	-0.0142 (6)
O2	0.0431 (6)	0.0580 (7)	0.0523 (7)	-0.0025 (5)	-0.0092 (5)	-0.0258 (6)
O3	0.187 (2)	0.0818 (12)	0.0574 (9)	0.0227 (13)	-0.0319 (12)	-0.0396 (9)
O4	0.0455 (7)	0.0704 (9)	0.0690 (9)	-0.0035 (6)	-0.0025 (6)	-0.0031 (7)
O5	0.0707 (10)	0.0820 (11)	0.0877 (12)	-0.0339 (9)	-0.0065 (9)	0.0192 (9)
S1	0.0421 (2)	0.03958 (19)	0.0412 (2)	-0.00283 (14)	-0.00230 (14)	-0.01553 (15)
F1	0.0838 (9)	0.0747 (8)	0.0621 (7)	-0.0007 (7)	-0.0385 (7)	0.0009 (6)

Geometric parameters (\AA , $^{\circ}$)

C1—C2	1.385 (2)	C15—O3	1.213 (2)
C1—C6	1.391 (2)	C15—C16	1.482 (3)
C1—S1	1.7532 (16)	C16—C17	1.520 (3)
C2—C3	1.383 (3)	C16—H16A	0.9700
C2—H2	0.9300	C16—H16B	0.9700
C3—C4	1.383 (3)	C17—H17A	0.9600
C3—H3	0.9300	C17—H17B	0.9600
C4—C5	1.374 (3)	C17—H17C	0.9600
C4—H4	0.9300	C18—C19	1.330 (2)
C5—C6	1.385 (3)	C18—H18	0.9300
C5—H5	0.9300	C19—C20	1.4678 (19)
C6—H6	0.9300	C19—H19	0.9300
C7—C8	1.394 (2)	C20—C21	1.401 (2)
C7—C12	1.397 (2)	C20—C25	1.403 (2)
C7—N1	1.4155 (19)	C21—C22	1.375 (2)
C8—C9	1.382 (3)	C21—H21	0.9300

C8—H8	0.9300	C22—C23	1.373 (3)
C9—C10	1.379 (3)	C22—H22	0.9300
C9—H9	0.9300	C23—F1	1.3465 (19)
C10—C11	1.375 (3)	C23—C24	1.363 (3)
C10—H10	0.9300	C24—C25	1.384 (2)
C11—C12	1.410 (2)	C24—H24	0.9300
C11—H11	0.9300	C25—N2	1.466 (2)
C12—C13	1.448 (2)	N1—S1	1.6824 (13)
C13—C14	1.373 (2)	N2—O5	1.215 (2)
C13—C15	1.488 (2)	N2—O4	1.216 (2)
C14—N1	1.4131 (19)	O1—S1	1.4193 (13)
C14—C18	1.467 (2)	O2—S1	1.4235 (13)
C2—C1—C6	121.56 (16)	C17—C16—H16A	109.0
C2—C1—S1	118.92 (12)	C15—C16—H16B	109.0
C6—C1—S1	119.52 (13)	C17—C16—H16B	109.0
C3—C2—C1	118.70 (16)	H16A—C16—H16B	107.8
C3—C2—H2	120.7	C16—C17—H17A	109.5
C1—C2—H2	120.7	C16—C17—H17B	109.5
C2—C3—C4	120.28 (18)	H17A—C17—H17B	109.5
C2—C3—H3	119.9	C16—C17—H17C	109.5
C4—C3—H3	119.9	H17A—C17—H17C	109.5
C5—C4—C3	120.48 (17)	H17B—C17—H17C	109.5
C5—C4—H4	119.8	C19—C18—C14	123.25 (14)
C3—C4—H4	119.8	C19—C18—H18	118.4
C4—C5—C6	120.46 (17)	C14—C18—H18	118.4
C4—C5—H5	119.8	C18—C19—C20	123.78 (14)
C6—C5—H5	119.8	C18—C19—H19	118.1
C5—C6—C1	118.52 (17)	C20—C19—H19	118.1
C5—C6—H6	120.7	C21—C20—C25	115.36 (14)
C1—C6—H6	120.7	C21—C20—C19	119.86 (13)
C8—C7—C12	122.28 (16)	C25—C20—C19	124.64 (14)
C8—C7—N1	130.77 (17)	C22—C21—C20	122.68 (15)
C12—C7—N1	106.93 (13)	C22—C21—H21	118.7
C9—C8—C7	117.0 (2)	C20—C21—H21	118.7
C9—C8—H8	121.5	C23—C22—C21	118.50 (17)
C7—C8—H8	121.5	C23—C22—H22	120.8
C10—C9—C8	121.67 (19)	C21—C22—H22	120.8
C10—C9—H9	119.2	F1—C23—C24	119.04 (17)
C8—C9—H9	119.2	F1—C23—C22	118.45 (17)
C11—C10—C9	121.72 (18)	C24—C23—C22	122.51 (16)
C11—C10—H10	119.1	C23—C24—C25	117.75 (15)
C9—C10—H10	119.1	C23—C24—H24	121.1
C10—C11—C12	118.23 (19)	C25—C24—H24	121.1
C10—C11—H11	120.9	C24—C25—C20	123.15 (15)
C12—C11—H11	120.9	C24—C25—N2	115.60 (14)
C7—C12—C11	119.09 (16)	C20—C25—N2	121.24 (13)
C7—C12—C13	108.19 (13)	C14—N1—C7	108.72 (12)

C11—C12—C13	132.70 (17)	C14—N1—S1	125.70 (10)
C14—C13—C12	107.60 (14)	C7—N1—S1	123.84 (11)
C14—C13—C15	129.90 (15)	O5—N2—O4	123.33 (17)
C12—C13—C15	122.08 (14)	O5—N2—C25	117.83 (16)
C13—C14—N1	108.51 (13)	O4—N2—C25	118.84 (14)
C13—C14—C18	130.55 (14)	O1—S1—O2	120.42 (8)
N1—C14—C18	120.63 (13)	O1—S1—N1	105.64 (7)
O3—C15—C16	119.04 (18)	O2—S1—N1	106.91 (7)
O3—C15—C13	117.57 (18)	O1—S1—C1	109.25 (8)
C16—C15—C13	123.24 (14)	O2—S1—C1	108.80 (7)
C15—C16—C17	113.04 (17)	N1—S1—C1	104.66 (7)
C15—C16—H16A	109.0		
C6—C1—C2—C3	-0.6 (3)	C25—C20—C21—C22	-1.3 (2)
S1—C1—C2—C3	179.82 (14)	C19—C20—C21—C22	-177.11 (15)
C1—C2—C3—C4	0.5 (3)	C20—C21—C22—C23	2.4 (3)
C2—C3—C4—C5	0.2 (3)	C21—C22—C23—F1	178.20 (16)
C3—C4—C5—C6	-0.7 (3)	C21—C22—C23—C24	-1.5 (3)
C4—C5—C6—C1	0.6 (3)	F1—C23—C24—C25	179.81 (15)
C2—C1—C6—C5	0.0 (2)	C22—C23—C24—C25	-0.5 (3)
S1—C1—C6—C5	179.63 (13)	C23—C24—C25—C20	1.7 (2)
C12—C7—C8—C9	-1.0 (3)	C23—C24—C25—N2	-177.55 (15)
N1—C7—C8—C9	177.10 (18)	C21—C20—C25—C24	-0.8 (2)
C7—C8—C9—C10	0.2 (3)	C19—C20—C25—C24	174.77 (14)
C8—C9—C10—C11	1.0 (3)	C21—C20—C25—N2	178.40 (14)
C9—C10—C11—C12	-1.3 (3)	C19—C20—C25—N2	-6.0 (2)
C8—C7—C12—C11	0.7 (3)	C13—C14—N1—C7	2.37 (17)
N1—C7—C12—C11	-177.79 (14)	C18—C14—N1—C7	176.59 (13)
C8—C7—C12—C13	179.31 (17)	C13—C14—N1—S1	167.73 (11)
N1—C7—C12—C13	0.79 (17)	C18—C14—N1—S1	-18.1 (2)
C10—C11—C12—C7	0.4 (3)	C8—C7—N1—C14	179.72 (18)
C10—C11—C12—C13	-177.72 (17)	C12—C7—N1—C14	-1.92 (17)
C7—C12—C13—C14	0.66 (17)	C8—C7—N1—S1	14.0 (3)
C11—C12—C13—C14	178.97 (17)	C12—C7—N1—S1	-167.62 (11)
C7—C12—C13—C15	-172.65 (15)	C24—C25—N2—O5	-27.0 (2)
C11—C12—C13—C15	5.7 (3)	C20—C25—N2—O5	153.67 (18)
C12—C13—C14—N1	-1.84 (16)	C24—C25—N2—O4	153.21 (16)
C15—C13—C14—N1	170.76 (16)	C20—C25—N2—O4	-26.1 (2)
C12—C13—C14—C18	-175.29 (15)	C14—N1—S1—O1	161.77 (13)
C15—C13—C14—C18	-2.7 (3)	C7—N1—S1—O1	-34.98 (15)
C14—C13—C15—O3	-161.0 (2)	C14—N1—S1—O2	32.38 (15)
C12—C13—C15—O3	10.7 (3)	C7—N1—S1—O2	-164.37 (13)
C14—C13—C15—C16	14.4 (3)	C14—N1—S1—C1	-82.94 (14)
C12—C13—C15—C16	-173.90 (17)	C7—N1—S1—C1	80.31 (14)
O3—C15—C16—C17	-10.1 (3)	C2—C1—S1—O1	-154.93 (13)
C13—C15—C16—C17	174.49 (19)	C6—C1—S1—O1	25.46 (15)
C13—C14—C18—C19	-67.9 (2)	C2—C1—S1—O2	-21.66 (15)
N1—C14—C18—C19	119.31 (17)	C6—C1—S1—O2	158.73 (12)

C14—C18—C19—C20	173.75 (13)	C2—C1—S1—N1	92.33 (13)
C18—C19—C20—C21	−26.3 (2)	C6—C1—S1—N1	−87.28 (13)
C18—C19—C20—C25	158.25 (15)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C8—H8···O1	0.93	2.33	2.913 (3)	120
C11—H11···O3	0.93	2.40	2.905 (3)	114
C16—H16 <i>A</i> ···F1 ⁱ	0.97	2.54	3.192 (2)	124
C22—H22···O4 ⁱⁱ	0.93	2.52	3.438 (2)	170

Symmetry codes: (i) $-x+1, -y-1, -z+2$; (ii) $x+1, y, z$.