

Animal health aspects of adaptation to climate change: beating the heat and parasites in a warming Europe

P. J. Skuce^{1†}, E. R. Morgan², J. van Dijk³ and M. Mitchell⁴

¹Moredun Research Institute, Pentlands Science Park, Midlothian EH26 0PZ, UK; ²School of Biological Sciences, University of Bristol, Langford, Bristol BS40 5DU, UK; ³Institute of Infection and Global Health, University of Liverpool, Leahurst Campus, CH64 7TE, UK; ⁴Scotland's Rural College (SRUC), The Roslin Building, Easter Bush, Midlothian EH25 9RG, UK

(Received 11 February 2013; Accepted 29 March 2013)

Weather patterns in northern European regions have changed noticeably over the past several decades, featuring warmer, wetter weather with more extreme events. The climate is projected to continue on this trajectory for the foreseeable future, even under the most modest warming scenarios. Such changes will have a significant impact on livestock farming, both directly through effects on the animals themselves, and indirectly through changing exposure to pests and pathogens. Adaptation options aimed at taking advantage of new opportunities and/or minimising the risks of negative impacts will, in themselves, have implications for animal health and welfare. In this review, we consider the potential consequences of future intensification of animal production, challenges associated with indoor and outdoor rearing of animals and aspects of animal transportation as key examples. We investigate the direct and indirect effects of climate change on the epidemiology of important livestock pathogens, with a particular focus on parasitic infections, and the likely animal health consequences associated with selected adaptation options. Finally, we attempt to identify key gaps in our knowledge and suggest future research priorities.

Keywords: climate change, livestock, adaptation, parasites

Implications

The climate of northern Europe has changed over the past several decades and will continue to do so for the foreseeable future. These changes will have an impact directly and indirectly on livestock farming systems, the animals themselves and the pests and pathogens that constrain them. We need to remain vigilant to such changes, through enhanced veterinary investigation and disease surveillance services. We also need to implement more sustainable disease control programmes, which are robust and flexible enough to cope with future changes.

Introduction: climate change and animal agriculture

It is now widely accepted that climate change is happening and that it represents one of the greatest threats faced by our planet, its population and economies. Agriculture is in the front line and among the first to feel the effects of a changing climate. Impacts are likely to be most severe in developing countries but no less important in Europe. The 'perfect storm' of real and projected rising atmospheric CO₂ and global temperatures, changes in annual and seasonal

precipitation and extreme weather events will undoubtedly affect the yield, quality and stability of food production. The projected changes offer opportunities and challenges but will undoubtedly lead to risk and uncertainty in agricultural production, food prices and farm incomes. Responses to climate change include 'adaptation' to reduce vulnerability of agricultural systems to change, and 'mitigation' to reduce the extent of the change. Adaptation has always been an integral part of the history of farming. Farmers have always adapted to change, whether climatic, economic or societal; this is based on a sound knowledge and understanding of their crops and livestock and the environment in which they live and work. Many farmers have already begun adapting their farming practices and strategies in response to recent changes in weather patterns and will continue to do so, both to exploit new opportunities, for example, increasing demand for their products, and to avoid or reduce any negative consequences of change. Adaptation options include, for example, production adjustments, diversification, intensification, changing land-use, altering timing of operations and so on. However, it is anticipated that, although projected climatic changes are not expected to severely disturb overall EU agricultural production until the ~2050s, the extent of change in the coming decades may overload individual farmers' capacity

† E-mail: philip.skuce@moredun.ac.uk

to adapt (EU, 2009). That said, strategies to adapt to the potential changes in climate are not without precedent nor do they need to be developed entirely *de novo*. The climatic conditions likely to apply in Europe, even under high emission scenarios, already exist in other countries, for example, South Africa and Australia, thereby providing the basis for the adaptive strategies likely to be appropriate under such conditions.

Therefore, the effects of climate change on livestock production and the health and welfare of animals in production systems, as well as the contribution of farmed animals to climate change through greenhouse gas (GHG) emissions, pose some of the major questions relating to the global issue of 'sustainable food production in a changing world' (Garnett, 2013; Smith and Gregory, 2013). Livestock are also central to the Food Security and 'Sustainable Intensification' agendas (Beddington, 2011), that is, the requirement for increased food production, with less waste and environmental impact to feed a growing world population, estimated to reach 9 billion by the 2050s. Demand for meat and dairy products, in particular, is projected to increase markedly as the developing world becomes more affluent (Gill *et al.*, 2010; Thornton, 2010).

For the purposes of this review, we focus specifically on sheep, cattle, pigs and poultry and restrict our coverage to temperate northern European regions. The direct effects of the predicted changes in climatic conditions, weather, frequency of extreme events and altered thermal environments will potentially have an impact on animal production efficiency and through the imposition of stress, resulting in consequences ranging from increased mortality to stress-induced pathologies, altered disease resistance and poor welfare (Gaughan *et al.*, 2009). Livestock will also be affected indirectly through changing exposure to important livestock diseases. The sectoral reports of both UK and Scottish Climate Change Risk Assessments (CCRA), for example, specifically mention livestock disease risk as a direct (negative) consequence of climate change (e.g. Moran *et al.*, 2009; Knox *et al.*, 2012).

Animal disease issues can have devastating consequences for the livestock sector and the wider economy, as exemplified by the UK BSE crisis in the late 1980s and the foot and mouth disease outbreak in 2001. Many of the diseases considered important in a northern European context have little or no obvious climate dimension, other than that they contribute to loss and waste in the system and increase the carbon footprint of livestock farming. However, others, most notably vector-borne diseases, such as those caused by Bluetongue and Schmallenberg viruses, do have an obvious climate dimension, through an increased range of their biting midge vectors. Incursions of such 'exotic' pathogens have received most attention in the farming and scientific literature (e.g. Baylis, 2006). However, arguably, the greatest constraint on efficient, sustainable livestock production comes from endemic rather than exotic pathogens. All sheep and cattle, in particular, are exposed to helminth parasite (or 'worm') infections and infestations by arthropod ectoparasites while grazing. Losses associated with increased infection intensity can be substantial.

In this review, we explore the complex interactions of direct and indirect effects of climate change on parasites, hosts and farm management practices and the implications for animal health and welfare. Such a review can never be comprehensive, and therefore we have chosen specific examples to help illustrate key points.

Direct effects of climate change on animal health, welfare and production

Predicted changes in climatic conditions, frequency and intensity of extreme events and altered thermal environments will have an impact directly on animal production efficiency and have likely knock-on deleterious effects on animal health and welfare. The severity of these effects may be forecast by examining the available climate change predictions for the appropriate time scales and periods and on the basis of the different emission scenarios and the estimated likelihoods, probabilities and uncertainties currently being modelled. A range of approaches may be used to determine the severity of impacts and their regional distributions (e.g. Rivington *et al.*, 2013).

In response to the changing climate, it will be necessary for agriculture and, indeed, the whole of society to adopt and implement reliable strategies for adaptation. Adaptation strategies are needed at all levels of administration, from the local to the international level (EC – Europa, 2012 and Defra, 2013). It is acknowledged that the currently predicted rates of change in climate variables will exceed the capacity of the traditional spontaneous adaptations by communities and animal species, and new and scientifically based strategies must be developed (IFAD, 2010). Analysing the impacts of climate change on animal well-being is complex. Even more complicated is the prediction of the direct effects on animal health and welfare of any associated adaptations that might be implemented by the animal production industry in response to reduced productivity and efficiency, increased losses, falling profits and changes in animal health. This also applies to concerns relating to the well-being of livestock in systems unable to cope with the climate change challenges or in response to adaptive changes that may alter their welfare status. For the purposes of the current review, we focus on warming scenarios and their potential impacts on livestock production systems in the United Kingdom, as an example broadly representative of northern temperate regions. We further examine the potential adaptations that might be required to minimise the deleterious effects of the stressors imposed by climate change.

Extensive and intensive production systems

In relation to adaptation strategies and their impacts, the important differences between extensive and intensive systems for livestock production need to be considered. Thus, although all livestock production systems are likely to be sensitive to the threats posed by predicted climate change scenarios, intensive animal production, specifically meat

poultry and pig systems, might be of particular concern in terms of direct adverse thermal effects. The size of the pig and meat poultry industries in the United Kingdom alone indicates the importance of reliable analysis of the climate change impacts and their likelihood and the development of effective and efficient adaptations. As indicated above, pigs raised in indoor systems are less vulnerable to changes in outdoor conditions, as long as the magnitude of any imposed thermal challenge does not exceed the capacity of the building design and ventilation system (McGlone and Pond, 2003). Similar arguments may be offered for intensive broiler production. However, as indicated elsewhere, it is unlikely that current climate control systems and production/husbandry practice will be adequate in the face of the challenges presented under some of the climate change projections currently in place.

In addition, animal transportation, including the movement of slaughter meat animals from intensive systems, constitutes a component of the process that may be considered particularly vulnerable to the thermal challenges imposed by all climate change scenarios and, in particular, increased incidence of extreme events. Animal transportation may present opportunities for the design and implementation of a wide range of adaptation options by the industry, to ameliorate or minimise the effects of the impacts resulting from the climate change scenarios.

Intensive production and animal transport systems present some complex challenges in terms of prioritisation of adaptations and assessment of such strategies in relation to the true cost–benefit. Indoor intensive systems require various degrees of environmental control and are thus deemed less sensitive to changes in external environments or meteorological conditions; however, current systems may not have the capacity for effective control in the face of large changes in thermal challenges or during an increased incidence of extreme events. In addition, the adaptations required under such conditions, for example, improved house design and structures, ventilation systems and insulation, water provision and husbandry practices and procedures have economic costs, energy costs, welfare costs and consequences for the carbon footprints of the systems.

Thus, adaptation strategies must be based on factors other than production indices and animal health and welfare, including socio-economic factors and socio-cultural values (Thornton, 2010). In the case of animal transportation, the same issues must be addressed, but it is proposed that the time course of implementation of adaptations may be significantly shorter as the impacts of warming will become apparent even before 2020 (Haskell *et al.*, 2011). Adaptations should be designed to concurrently maximise productive efficiency and ensure animal health and welfare status. However, current policy must also consider other consequences and sequelae of the adaptations; thus, for each system, it is important to model holistically all aspects of the adaptive strategies. A study by Turnpenny *et al.* (2001) recognised the importance of examining the impacts of climate change on intensive systems and has proposed

models that facilitate identification of risks of thermal stress and identifies some adaptations (e.g. changing stocking densities and improved ventilation regimes that might reduce such risks).

Thermal stress – impacts and adaptations

The UKCCRA (Knox *et al.*, 2012) has concluded that there is a significant risk of heat stress affecting dairy production in the United Kingdom under specific prediction scenarios but that major effects would not be encountered until 2050. Referring specifically to the study by Haskell *et al.* (2011), Knox *et al.* (2012) acknowledge that failure to assess the impacts of the predictions on intensive production of pigs and poultry is an important omission, indicating that ‘In particular, pigs and poultry could be more severely affected by heat stress’.

Haskell *et al.* (2011) reviewed the impacts of climate change on livestock and included dairying, beef cattle, pigs and poultry. Their evidence relates primarily to the direct impacts of thermal stress on animal well-being and health, as well as indirect climate impacts, including impacts on feed availability, competition for land-use and climate-induced consequences on animal transport. The study by Haskell *et al.* (2011) reports that the effect of climate change on the welfare of dairy cows may be primarily a consequence of increased temperature, although the magnitude of this effect is difficult to predict from currently available data. Heat stress during summer months, at which time the majority of the UK dairy cattle graze outside on pasture, may not be a concern until 2050 in areas such as the south-west of England, but may be of more significance under the scenarios predicted for 2080. Animals grazed outdoors are expected to suffer some effects from high ambient temperatures, high direct and indirect solar radiation and high humidity, particularly during heat waves, all stressors that will negatively affect welfare. It is interesting to note that, in a more recent comprehensive review of climate change and dairy farming in Central Europe (Gauly *et al.*, 2013), the authors identified feed availability and quality as key constraints. They argue, however, that the effects on farm economy could be neutralised if effective adaptation and mitigation strategies, such as management, nutrition, animal health, plant and animal breeding, are implemented.

In contrast, it is thought that beef cattle will be spared the major effects of heat stress, as the geographical areas in which the majority are reared (west and north) will not see major temperature increases until 2080 under the high emission scenario. In any of the predicted scenarios, providing an appropriate environment can reduce heat stress. Adaptations suggested have been the provision of shade or the use of active cooling mechanisms such as sprinklers, increasing air velocity and provision of cool drinking water. In many cases, the provision of shade may be the most economical solution of reducing high heat load. It is suggested that a well-designed shade structure should reduce the total heat load by 30% to 50%. Shade can be provided in

many different ways, for example, shade can be natural (trees) or artificial. It has been suggested that shade from trees is more effective and is preferred by cattle. Air movement is an important factor in the relief of heat stress. Air movement is critical if cooling is to be effective, whether outside or inside a building. The use of natural ventilation in animal buildings should be maximised through the use of open-sided sheds. Direct access to water such as rivers and pools is effective in cooling animals in a grazing situation, as animals will stand in the water or splash it over themselves. The use of sprinklers, misting and fogging is believed to reduce heat load. Body temperature can be reduced by a few degrees for perhaps 3 to 4 h following a substantial wetting of cattle.

Intensive production systems and transportation – adaptation strategies

Therefore, for present purposes, it may be proposed that, in the United Kingdom, extensively produced (outdoor) livestock will not be at risk of primary thermal stress for most of the currently predicted scenarios and that the focus should be on the requirements, nature and additional implications of adaptations required in the intensive production of pigs and poultry. It is therefore proposed that it is important to consider all aspects of the adaptations required by the EU poultry and pig intensive production sectors both in relation to the production or housing requirements and transportation of animals by road to slaughter.

Animal transportation may be considered to be extremely vulnerable to changes in average and maximum temperatures and to any change in the frequency and/or severity of extreme events. It may be proposed that the thermal micro-environment within the transport containers or vehicles is extremely vulnerable to changes in external thermal conditions. As such, in response to any potential increase in thermal load it poses the greatest threat to the animals' welfare and well-being (Appleby and Lawrence, 1999; Mitchell and Kettlewell, 1998; Cockram and Mitchell, 1999; Mitchell, 2006). Adverse thermal conditions resulting in either heat or cold stress may lead to reduced welfare, overt tissue damage or injury and increases in mortality in transit (Mitchell, 2006; Mitchell and Kettlewell, 2006; Mitchell and Kettlewell, 2008). The thermal micro-environment in transport containers or vehicles may be complex and results from the interactions of several factors. These include the external climatic conditions, heat and water production of the animals, ventilation regimes, the distribution and flow rates of air movement and any additional external sources of heat and/or moisture. The metabolic heat and water production of animals in transit is a major determinant of the 'on-board' thermal environment and this can be of vital importance, if external conditions and/or inadequate ventilation are likely to precipitate heat stress (Kettlewell *et al.*, 2001a, 2001b and 2001c).

We will now examine the direct effects of climate change on intensive pig and poultry production and the transportation of these animals to slaughter, identify potential adaptations, as well as their secondary costs and consequences.

Current animal housing design and operation assures a controlled environment (temperature control) and this in turn involves or influences a number of other variables. Thus, houses may be mechanically ventilated and supplementary heating and/or cooling may be supplied. Some facilities will have natural ventilation where control is very limited. The houses require water supplies, bedding and/or litter, the condition of which is influenced by the hygrothermal environment and thus ventilation regime. The houses accumulate waste and contamination that require removal, storage and disposal. The houses require cleaning and disinfection and disease control measures. All of the factors are intimately linked, and changes in climate necessitating alterations in the operation or specification of the primary control systems will have far-reaching consequences. Thus, if the thermal demands placed on an intensive system (e.g. a broiler production shed) by a 'heat wave' or long-term increase in maximum temperature exceed the ventilation capacity, then, in addition to potential heat stress on the birds, other variables will be affected. Water usage by the stressed birds will increase (May and Lott, 1992; Belay and Teeter, 1993), litter quality may deteriorate, this will result in increased incidence of pathology and losses (Bessai, 2006; Manning *et al.*, 2007; Meluzzi *et al.*, 2008; Alain *et al.*, 2009; Kyvsgaard *et al.*, 2013) and the final wet litter will present problems of emissions and disposal (Topper *et al.*, 2008; Singh *et al.*, 2009). Obviously, during animal transport (catching/handling/loading/in transit) and the associated procedures, similar factors will determine the on-board environment for the animals and their effects. Climate change scenarios involving significant warming (increased average temperature and changes in average, min/max) and increased incidence of extreme events will have direct impacts on pigs/birds in the shed or in transit. These will include:

(1) losses, that is, mortality and 'dead on arrival'; (2) poor welfare; (3) growth penalty/product quality; (4) reduced feed conversion efficiency; (5) increased diseases' susceptibility; and (6) reduced bedding/litter quality (with associated lesions, infections and pathologies).

The potential adaptations necessary to mitigate against the climate change-induced problems will be: (1) improved mechanical ventilation systems/regimes; (2) additional cooling/heating systems; (3) changes in stocking density; (4) slower growing pigs/birds (to reduce thermal loads and incidence of growth-associated pathologies); (5) more heat-tolerant lines/strains (genetic selection/genomic strategies); and (6) nutritional measures.

It is immediately apparent that some of these adaptations to reduce the impact of climate change are themselves likely to contribute to climate change. Thus, the efficacy of each of these approaches requires consideration along with any environmental impact and an assessment of the real costs and therefore definition of the most cost-effective strategies for adaptations in intensive production systems. It is immediately apparent that a major investment in building structures and perhaps an increase in the scale of production are important options and costs. The installation of higher-capacity ventilation

or environmental control systems in new or existing housing will have a very significant cost. The energy (electricity, gas) and water requirements of such higher-capacity/performance systems will be a major concern. The GHG/emissions or carbon footprints and pollution and waste generation potential associated with these systems and perhaps implemented in larger-scale production systems will require estimation and incorporation in holistic, integrative models. Such models must also be applied to animal transportation including costs for improved vehicle design and operation. For housed animals, the engineering solutions must be analysed in the context of management of the animals and their bedding or litter. The short-term alternatives to technological adaptations such as reducing stocking densities and/or growth rates must be carefully evaluated. The modelling of these adaptation strategy costs must also consider the effects on disease and health status of the animals and any likely impacts on zoonoses. Other factors that might be incorporated in a long-term strategy include the likelihood of new regulatory requirements (animal health and welfare) as applied to new and improved production systems and the benefits or otherwise of relocation of production systems or trade-based solutions to avoid the impacts under consideration.

Unfortunately, at present, it is not possible to provide definitive answers to any of the questions raised. There are a number of studies and programmes in place where modelling of the impacts on intensive livestock and the costs of potential adaptations are to be addressed and assessed. The key issue is to recognise that intensive animal production can be a key component of the building of sustainable animal protein supply in the future. Intensive production should offer a degree of protection against changing climates and, in particular, thermal challenges. However, current climate change predictions indicate that additional adaptations may be required and these will have secondary consequences that require very detailed and comprehensive assessment before identification of the most cost-effective policies and strategies.

Climate change and livestock pathogens – some examples, with a focus on parasites

The impacts of climate change on the lifecycle and distribution of insects, and subsequently on vector-borne diseases, are becoming increasingly common in livestock. Climate change also has well-documented effects on the lifecycle of ticks, with warming scenarios generally favouring development and increased spread of tick-borne infections, which threaten both humans and animals (e.g. White *et al.*, 2003; Gray *et al.*, 2009). Extreme weather events, for example, flooding can carry a risk of *Cryptosporidium* parasites and/or enterohaemorrhagic *Escherichia coli* emerging as diffuse pollution in a run-off from agricultural land. This poses an obvious threat to other livestock and is also a zoonotic risk to humans through contamination of water supplies. Future challenges in the control of parasitic zoonoses, including those related to climate change, deserve increasing attention alongside

production-limiting disease (Polley and Thompson, 2009). Extreme weather events can also lead to animal health problems, even in housed animals. For example, infestation of laying hen houses with the poultry red mite, *Dermanyssus gallinae*, is thought to cost the EU poultry industry ~€130 millions/year (Maurer and Baumgartner, 1992). During extreme weather events, red mite populations can become uncontrollable in both laying hen systems and broiler production units and have been implicated in the deaths of large numbers of hens during the summer heat wave of 2003.

Although extreme weather events pose specific challenges, gradual changes in climate can also have a major impact on animal health. The effects of climate change on infectious diseases of farmed animals, and the complex interactions determining whether animal health, production and welfare will be compromised, are arguably best demonstrated by the example of helminth parasites of ruminants. Several different species of gastrointestinal nematodes (roundworms), notably *Ostertagia (Teladorsagia) spp.*, *Haemonchus contortus*, *Nematodirus battus* and *Trichostrongylus spp.*, can be found on most ruminant farms globally. Similarly, the liver fluke, *Fasciola hepatica*, is a highly pathogenic flatworm parasite of sheep and cattle and is responsible for considerable disease, death and economic losses (e.g. Schweizer *et al.*, 2005; Sargison and Scott, 2011). Because it requires a small mud snail intermediate host to complete its lifecycle and disseminate infectious stages onto pasture, fluke has typically been reported in the wetter areas, for example, north-west Europe. Heavy infestations of livestock with some of these parasites can prove fatal but more commonly they cause lower-level infections resulting in sub-acute or chronic disease, for example, anorexia, diarrhoea, weight loss and reduced milk yield in production animals. Associated losses to the meat and milk producing industry are very significant. For example, chronic roundworm infections of sheep, and their prevention, have been estimated to cost the GB sheep industry £84 millions/year (Nieuwhof and Bishop, 2005). In recent years, the diagnostic incidence of both acute and chronic helminth disease in cattle and sheep in the United Kingdom, where surveillance records enable meaningful analysis, has increased threefold to fourfold (van Dijk *et al.*, 2010). Parasite species such as the rumen fluke (or paramphistomes), which traditionally caused no measurable production losses, have recently been diagnosed as the cause of acute disease and death in young sheep and cattle in the United Kingdom (Mason *et al.* 2012; Millar *et al.*, 2012). Rumen fluke have also become a significant problem over recent years in other EU countries, such as Spain, France, Italy and Ireland (González-Warleta *et al.*, 2013). In the United Kingdom, rumen fluke outbreaks have been specifically associated with flooded farm land; hence, infection and transmission may be exacerbated by extreme weather, that is, rainfall events. These most alarming trends are also accompanied by a change in the seasonality of these parasites, with, in general, more diseases being witnessed later in the grazing season (van Dijk *et al.*, 2008). Moreover,




Figure 1 The life-cycle of *Fasciola hepatica*, the sheep/cattle liver fluke, showing the respective environmental stages, that is eggs, miracidia, snails, cercariae and cysts.

the spatial distribution of various species appears to be changing with a northwards spread of *H. contortus*, a parasite originating in the tropics, and west-to-east spread of the liver fluke, *F. hepatica*. Part of the overall increase in the abundance of these parasites has been ascribed to the ever-increasing problem of anthelmintic resistance (as reviewed by Wolstenholme *et al.*, 2004), which, before the recent arrival of the first new commercially available anthelmintic compounds for a generation, was already driving some sheep farmers out of business (Sargison *et al.*, 2005). However, some of the trends described above, and the simultaneous increase in the abundance of so many different worm species, cannot be explained by anthelmintic resistance alone. Climate change has been shown to be a likely significant driver of the problems experienced (van Dijk *et al.*, 2008; Fox *et al.*, 2011); combating this is currently the subject of substantial research efforts across Europe (e.g. <http://www.gloworm.eu/project>).

The lifecycle of the pathogenic helminths, for example, the sheep/cattle liver fluke *F. hepatica* (Figure 1), consists of parasitic and non-parasitic stages. The latter are voided onto pasture as eggs and need to undergo a period of development, to the infective stage larvae, at pasture. The dependence of the

developmental success and survival of these stages on climatic factors (O'Connor *et al.*, 2006), as well as the complexity involved in predicting the overall, often contrasting, effects of climate change on the abundance of infective larvae (van Dijk *et al.*, 2010; Morgan and van Dijk, 2012) have been reviewed in detail. Briefly, increases in environmental temperature will increase the proportion of parasite populations being able to develop, as well as development rates of those populations. Simultaneously, the death rate of the (pre-) infective stages is increased for most pathogens infecting ruminants. In temperate regions, increases in developmental success will generally outweigh the negative effects. Thus, for example, the proportion of *H. contortus* eggs developing successfully to the infective L3 stage in the laboratory rises from around a quarter at 15°C to half at 25°C, falling back to a third at 35° because of elevated mortality (van Dijk, 2008). This magnitude of change in the number of larvae reaching the herbage is likely to have substantial effects on infection pressure for sheep. Eggs and larvae also need a certain amount of moisture to be able to develop and subsequently migrate onto herbage. Moisture present in faeces and soil is normally sufficient to cover early development of nematodes (van Dijk and Morgan, 2011). Soil is also likely to buffer desiccation. However, especially when




Figure 2 Effects of changing temperatures and rainfall patterns on host–parasite interactions: examples of the complex interactive pathways and feedback loops co-determining the effects of climate change on the grazing ruminant host.

development occurred during warm, dry days, free water may subsequently be needed for the larvae to migrate out of the faeces (van Dijk and Morgan, 2011). The overall effect of the total amount of rain falling in fewer rain events is the subject of current mathematical modelling efforts but for now is difficult to predict. These changing rainfall patterns are likely to have a significant effect on the free-living stages of *F. hepatica* (and rumen fluke). On the one hand, if new water bodies are formed, new parasite habitats may be established. On the other hand, dry periods may make parasite development and survival less predictable. Furthermore, negative effects of changing rainfall patterns may be offset by positive influences of increased temperature on the development of both pre-infective stages and the intermediate snail host.

When the overall abundance of pathogenic helminths on farms is increased, a complex network of interactions subsequently determines the extent to which this leads to increases in disease prevalence. Some examples are given in Figure 2. In general, larger worm burdens will require increased efforts of the immune system of the host and, especially in growing animals; apart from reduced growth rates, this may lead to a reduced ability to combat other infective organisms. At the cost of meat and milk production, the host population may be able to dampen the increase in the parasite population (Hudson *et al.*, 2006). As the traditionally seasonal availability of infective stages is altered for all different worm species, co-infection patterns are likely to change, with parasite species competing with others to inhabit the same host (van Dijk *et al.*, 2010). If the host becomes overwhelmed by worms, the negative feedback mechanism of host immunity on the worm population will be diminished. As sheep and cattle hosts become less-efficient production animals, their GHG emission to production ratio, and thereby their contribution to climate change, will increase (O'Mara, 2012). On warm days, temperature increases experienced by hosts, especially sheep, are likely to

decrease their feed intake, as animals congregate in the shade for large parts of the day, although the impact of this may be offset by grazing during cooler periods or at night. Similarly, periods of heavy downpours have been shown to have a negative effect on herbage intake (Champion *et al.*, 1994), although they would rarely be sufficiently prolonged to seriously affect animal production or body condition. A decreased energy and protein intake will, in turn, negatively affect the immune system (Sargison *et al.*, 2002; Houdijk, 2008).

Climate change and farm management factors

Host–parasite interactions will not only be directly affected by climate change but also through alterations to farm management. If farmers are to maintain productivity under future climate change scenarios (Bell *et al.*, 2012), while limiting GHG emissions (Eckhard and Cullen, 2011), plasticity in farm systems is desirable (Rodríguez *et al.*, 2011). For the dairy industry, the need to decrease CH₄ and N₂O emissions is likely to speed up the move towards zero-grazing and diets with a lower forage-to-concentrate ratios, among other mitigation options (Aguerre *et al.*, 2011; Novak and Fiorelli, 2011). This may solve the problem of helminth-associated production losses, at least in adult cattle, although the effects of prolonged housing on the transmission of other infectious diseases, for example, bovine tuberculosis and calf pneumonia, remain to be investigated. However, it needs to be remembered that such cattle are less immune to these parasites, as a result of decreased exposure, and serious diseases can be expected when they are sold to farms where cattle are pastured. There may also be animal welfare considerations of prolonged housing. On those farms that are likely to have seen stock increased in recent years, total GHG emissions may be lowered by the creation of both intensely grazed and animal-free pastures (Schonbach *et al.*, 2012). Intensely grazed

pasture systems will inevitably depend on anthelmintic treatment for worm control. Anthelmintic resistance, the development of which itself may further be enhanced by climatic changes limiting the survival of free-living stages (Papadopoulos *et al.*, 2001; Waghorn *et al.*, 2009), may threaten the sustainability of these farm systems.

Temperature increases will quite likely lengthen the grass growth season and farmers will be tempted to turn out stock earlier and to delay housing in autumn. However, lengthening the grazing season without increasing the hectareage of land grazed may negatively affect sward quality and herbage utilisation (Roca-Fernandez *et al.*, 2011), especially later in the grazing season (Kanneganti and Kaffka, 1995). Climate change may also negatively influence herbage utilisation through increased growth rates of unwanted plants at pasture (Tiley, 2010) and unpredictable rainfall patterns/periods of drought. Unless animals are supplemented at pasture, a relatively lowered energy intake, over lengthened grazing seasons, may constrain the immune system of hosts. Especially in the beef and sheep sector, both direct climatic effects and changing farm management may change the breed composition of national herds and flocks (Aby *et al.*, 2012). Different breeds may be more susceptible but could also be more resilient to parasites.

The effects of changing farm management on parasites

A lengthening grazing season may markedly alter the epidemiology of helminths. The earlier animals are turned out onto pasture, the higher the initial infection with over-wintered larvae will be. These larvae would normally die off rapidly during the late spring in temperate regions (van Dijk *et al.*, 2009). In addition, with grass growth and parasite development occurring above very similar temperature thresholds, more parasite generations become possible per grazing season, potentially leading to sharply increased worm burdens. Bringing lambing forward may appear to be advantageous in that hosts infected in autumn will be older and more 'immunocompetent' (Morgan and van Dijk, 2012). Similarly, longer growing seasons would be expected to offer better quality (or at least cheaper) nutrition, which may, in turn, improve the resilience of stock to helminth infections. However, across the different worm species, more diseases are witnessed in autumn in the United Kingdom (van Dijk *et al.*, 2008), suggesting that increased worm abundance at pasture regularly overwhelms the immune system of hosts. If these trends continue, then autumnal parasite transmission success itself may provide a negative feedback on the length of the grazing season. Changes to farm management also provide opportunities to combat the effects of climate change on parasite populations (Morgan and Wall, 2009). Earlier mowing of pastures ought to greatly enhance the spring die-off of larvae (Charlier *et al.*, 2005; van Dijk *et al.*, 2009), and strategic alterations to the timing of the application of anthelmintic treatment may be able to nullify the effects of the lengthening transmission season (Morgan and Wall, 2009).




Figure 3 Potential for elongated season of challenge to undermine parasite management of disease risk. In the top graph (a), the solid line represents earlier emergence or build-up of parasite challenge in spring under warm conditions relative to the present (broken line). In the bottom graph (b), treatment with persistent or repeated parasiticide (light shaded bar) must commence sooner (dark shaded bar) to control disease, leaving a new window of risk in autumn. Increased duration of coverage is costly and risks strengthening selection for drug resistance. This general schematic does not take into account system-specific parasite population dynamics, but finds theoretical and empirical support from myiasis in sheep (Wall *et al.* 2011).

Adapting future parasite control strategies

Given the complexity and uncertainty of possible effects of climate change on parasite epidemiology, it is difficult to design new strategic treatment or prophylactic regimes that will be robust to future scenarios. Moreover, increased use of chemical protection is almost certainly unsustainable, as resistance typically outpaces the development of new active compounds (Kaplan, 2004). In many cases, longer seasons of parasite challenge will force farmers into difficult choices. For example, countering early emergence of *Lucilia sericata* blowflies by early chemical treatment or shearing can eliminate predicted spring peaks of fly strike in sheep in warmer conditions, but also lead to new unprotected periods of risk in autumn (Figure 3; Wall *et al.*, 2011). Rather than seeking to protect livestock more comprehensively at all times of the year, parasite control strategies that are well adapted to future threats will take a risk-based approach. Although mathematical modelling can help to identify high-risk periods and to explore likely effectiveness of control strategies, uncertainties inherent in their performance on individual farms might encourage farmers to include a monitoring component in future parasite management plans. In some cases, for example, sheep scab or louse infestations, this might be based on appearance of clinical signs, whereas in others, notably gastrointestinal nematodes, this could be routine parasitological and production monitoring. Applied at group

level, such monitoring enables treatment at appropriate times, whereas at an individual level it is the bedrock of targeted selective treatment (TST). Such strategies target anthelmintic treatments to only those individuals or groups in need of treatment (or most likely to benefit from treatment) and are designed to preserve drug efficacy and susceptibility in the parasite population (van Wyk, 2001; Kenyon and Jackson, 2012).

TST approaches have so far been most developed for small ruminants (Kenyon and Jackson, 2012), and uptake is variable because monitoring costs often exceed proximal benefits from treatment savings, whereas economic benefits of prolonging the useful life of individual drugs are hard to quantify. Much work remains to be conducted on the economics of production loss from endemic animal disease, before risk monitoring can be optimised to future scenarios. At present, an increasingly sound evidence base for sustainable general practices is being developed (e.g. Leathwick *et al.*, 2009 and 2012), which will provide a platform for context-specific approaches. Considerable local adaptation to take account of climate and management will be needed. For example, strategic and selective treatments should take account of climate not only in determining when livestock should be treated, but also in how intensively refugia-based approaches should be applied at a given time of the year. This is because the fate of offspring of parasites left *in refugia* will depend strongly on climate. As ever, simple solutions that are based on sound science will be needed by producers, but the current state of the art is unequal to the task of designing such solutions from the top down, not least because system behaviour will be locally specific (Dobson *et al.*, 2011). It is for this very reason that seeking to emulate production and parasite control systems from warmer parts of the world is a simplistic approach to climate change adaptation, and one that is likely to meet with limited success. Drivers and constraints on farm management systems are legion, and their depth and complexity makes it difficult to overturn established practice, which itself might have arisen for sound reasons of local adaptation (de Rancourt *et al.*, 2006; Sturaro *et al.*, 2009). Trade-offs between immediate productivity and long-term sustainability also provide justified disincentives to farmers to change practice too quickly (Waghorn *et al.*, 2008). It is possible that placing better tools in the hands of farmers for local risk assessment and appropriate responses will be more useful in enabling them to quickly and sustainably adapt to unpredictable changes in parasite epidemiology. Options for adaptation, which are more creative than simply altering treatment strategies, are available and should be fully used, for example, altering patterns of reproduction, housing, grazing and even breed selection (White *et al.*, 2003).

The Red Queen – coming soon to a farm near you?

Like farmers, parasites are unlikely to stand still in the face of climate change. Direct effects of climate on development and survival of life-cycle stages outside the definitive

host, as well as indirect effects on fitness acting through host availability and susceptibility, for example, as a result of the management changes discussed above, will exert selective pressure. Therefore, parasite life histories could evolve to become better adapted to their new surroundings. High-effective population sizes, high levels of genetic diversity and short generation times mean that parasites are potentially better equipped than their hosts to adapt rapidly to environmental change, especially when small changes in prevailing conditions could markedly affect the chances of individuals finding a host. Potentially rapid evolution in response to strong selective pressure is abundantly illustrated by the emergence of anthelmintic resistance among gastrointestinal parasites of livestock (Kaplan, 2004; Wolstenholme *et al.*, 2004). Other evolutionary responses of parasites could, although less obvious, have significant practical implications for parasite control in a changing climate.

Direct evidence for the adaptation of parasite life history in response to changes in management or climate is hard to find, but geographic variation in life traits of epidemiological importance suggest that it occurs, and could have a bearing on the success of efforts to adapt future disease control strategies to changing climates. In some cases, this could involve upregulation of mechanisms of protection against environmental noxes (Baker *et al.*, 2012). In others, the targets of selection could be subtle and not the most obvious. Thus, free-living stages of *H. contortus* from sheep in Sweden had very similar characteristics to the populations in Africa, even though improved ability to develop or survive at lower temperatures might have been expected to enhance the ability of northern isolates to colonise cooler areas (Troell *et al.*, 2005 and 2006). However, northern populations made greater use of developmental arrest or hypobiosis within the host to improve the efficiency of overwinter survival. The plasticity of this species therefore allows it to ensure transmission in widely different climates, and we should not be surprised by its ability to adapt to rapidly changing climates in future. Similar variation was found in the propensity of *Ixodes ricinus* ticks to enter diapause in winter, depending on prevailing climate (Dobson *et al.*, 2011). In this case, models predicting the abundance and seasonality of ticks in the United Kingdom had to be altered to take into account presumed differences in diapause in populations in Spain to provide a satisfactory fit to local data. This is likely to be the result of genetic differences between populations that underlie phenotypic variations in life history. Experimental work to elucidate mechanisms and rates of adaptation to climate change in parasites of veterinary importance is limited; in other systems, this suggests that responses at low and high temperatures can differ and affect ability to adapt to environmental conditions in different parts of a parasite's geographic range (Ford and Chintala, 2006) and, by extension, to climate change.

A particularly interesting example of phenotypic variation, which could indicate the potential for rapid parasite adaptation to climate change, is seen in the intestinal nematode of lambs, *N. battus*. In this species, eggs shed by lambs in

spring develop slowly through the summer and typically do not hatch until after they have experienced a period of cold exposure over winter, followed by a rise in temperature in spring. In this way, infection is carried over from year to year, while adult worms are rapidly eliminated from lambs by an effective immune response. However, recent work found that there is an upper temperature threshold above which hatching ceases, such that under rapid spring increases in ambient temperature in southern England, hatching is often interrupted (van Dijk and Morgan, 2008). This removes the advantage to the parasite of delaying hatching until after winter, and in these populations a proportion of eggs does not require chilling in order to hatch (van Dijk *et al.*, 2010). As there is no evidence for genetic segregation of worms whose offspring hatch with or without chilling, this appears to be a form of 'bet hedging', such that parasite genotypes are well equipped to deal with reliable spring conditions for hatching, leading to efficient infection of young, susceptible lambs, while also retaining the ability to hatch in autumn, such that adverse weather conditions in a particular year are not disastrous for transmission (van Dijk and Morgan, 2010). Crucially, management change that seeks to decrease opportunities for parasite transmission, such as alternating grazing of lambs on a particular pasture between years to avoid spring peaks in *N. battus* larvae, are destined to select for parasite phenotypes that are able to cope with those conditions. This example also shows that parasite adaptation strategies can be very refined and flexible, and given that hatching behaviour of *N. battus* appears to have diverged in northern and southern United Kingdom over only a few decades, adaptation can be rapid. Parasite control strategies that are well adapted to anticipate future conditions but that ignore the ability of parasites to change could fail sooner and more comprehensively than expected.

General discussion – knowledge gaps and research priorities

It is clear that climate change has had both direct and indirect impacts on livestock production, animal health and welfare over the past number of decades. If even the most modest climate change projections are to be realised over the coming decades, then further changes will be inevitable. It is, however, important to appreciate that the direct and indirect effects discussed in this review do not operate independently but are inter-related, with many knock-on effects and feedback loops, and often with unforeseen or consequences. For example, climate change may bring increased incidence or intensity of endemic disease, which in turn has an impact on production efficiency, which in turn contributes to climate change through unnecessary GHG emissions from unproductive livestock. Similarly, putting animals outdoors to prevent heat stress may bring a greater risk of pasture-borne diseases and parasites, whereas bringing animals indoors to avoid extreme weather and enhance productivity may bring an increased risk of housing-associated

diseases. More needs to be done to integrate measures aimed at improving animal welfare and productivity under climate change with the animal health implications. To ensure future sustainability of the livestock industry, even in temperate regions, the industry must be able to adapt to the challenges and opportunities presented. This requires, first, a detailed understanding of the interactive network of drivers, and second, a calculation of costs and benefits of adaptation options at the individual farm- and industry level. Such analyses involve sophisticated modelling of the likely climate change scenarios, their effects on individual drivers and their interaction at a local, regional and international level. Given the between-year variability in weather conditions, it will be a great challenge to design strategies that are cost-effective overall. If such strategies are to be adopted, detailed risk analyses, followed by clear communication of risk, will be required from researchers. In the context of livestock disease, our ever-increasing understanding of the epidemiology of important disease-causing pathogens and diagnostics can now be interfaced with very sophisticated geographic information system technology and both statistically and process-based modelling approaches to produce risk maps for disease at an appropriate tempo-spatial scale (e.g. Cornell, 2005; Bergquist and Rinaldi, 2010; Fox *et al.*, 2012). However, there are still gaps in our knowledge in relation to the biology of parasites and pathogens and how they will respond to changing climatic conditions. Much of the published work has focused on responses to increasing temperatures, linking rainfall, evapotranspiration rates, etc. to the ecology of parasites and vectors that remains a challenge. It goes without saying that epidemiological models are only as good as the assumptions they are based on. Assumptions, which may be valid in certain experimental settings, have to be appropriate for all spatial and temporal scales modelled. In the short-medium term, we need to validate models and quantify trends, through improved surveillance, ideally active rather than passive, of animal health disease issues, whether exotic or endemic. This places a requirement on improved diagnostic capabilities, with the potential to be rapid, high-throughput and cost-effective at an appropriate regional/national/international scale. In the context of the endoparasites and ectoparasites specifically, vaccines to protect animals from infection, although not a complete solution, would be highly desirable. However, development of such vaccines has proven to be extremely technically demanding for a number of reasons (e.g. Vercruyse *et al.*, 2007). Until such time as vaccines do become available, we need to reduce the usage of and reliance on chemical treatments and promote more sustainable parasite control strategies. Climate change is an obvious driver in some of these scenarios of change but we also need to gain a better understanding of the impact of other confounding factors, such as global economic change, CAP reform, anthelmintic resistance, changing farm management practices, etc. and their implications for the EU livestock sector. The key to combating the effects of climate change will be embracing change ourselves.

Acknowledgements

The authors acknowledge the financial support of the Scottish Government Strategic Research Programme, the ClimateXChange Centre of Expertise (M.M. & P.J.S.) and the EU FP7 GLOWORM project no. 288975 (E.R.M., J.v.D. & P.J.S.).

This paper was published as part of a supplement to *animal*, publication of which was supported by the Greenhouse Gases & Animal Agriculture Conference 2013. The papers included in this supplement were invited by the Guest Editors and have undergone the standard journal formal review process. They may be cited. The Guest Editors appointed to this supplement are R. J. Dewhurst, D. R. Chadwick, E. Charmley, N. M. Holden, D. A. Kenny, G. Lanigan, D. Moran, C. J. Newbold, P. O'Kiely, and T. Yan. The Guest Editors declare no conflict of interest.

References

- Aby BA, Aass L, Sehested E and Vangen O 2012. Effects of changes in external production conditions on economic values of traits in continental and British beef cattle breeds. *Livestock Science* 150, 80–93.
- Aguerre MJ, Wattiaux MA, Powell JM, Broderick GA and Arndt C 2011. Effect of forage-to-concentrate ratio in dairy cow diets on emission of methane, carbon dioxide, and ammonia, lactation performance, and manure excretion. *Journal of Dairy Science* 94, 3081–3093.
- Alain V, Mirabito L, Arnould C, Colas M, Le Bouquin S, Lupo C and Michel V 2009. Skin lesions in broiler chickens measured at the slaughterhouse: relationships between lesions and between their prevalence and rearing factors. *British Poultry Science* 50, 407–417.
- Appleby MC and Lawrence TL (eds) 1999. *Farm animal welfare – who writes the rules?* Occasional Publication no. 23. British Society of Animal Production, Edinburgh, pp. 43–64.
- Baker RH, Britton C, Roberts B, Loer CM, Matthews JB and Nisbet AJ 2012. Melanisation of *Teladorsagia circumcincta* larvae exposed to sunlight: a role for GTP-cyclohydrolase in nematode survival. *International Journal of Parasitology* 42, 887–891.
- Baylis M 2006. Foresight. Infectious diseases: preparing for the future. T7.3: The effects of climate change on infectious diseases of animals. The Government Office for Science, London. Retrieved from <http://www.foresight.gov.uk>
- Beddington J 2011. Foresight: The future of food and farming. The Government Office for Science, London. Retrieved from <http://www.foresight.gov.uk>
- Belay T and Teeter RG 1993. Broiler water balance and thermo-balance during thermoneutral and high temperature exposure. *Poultry Science* 72, 116–124.
- Bell MJ, Eckhard RJ and Cullen BR 2012. The effect of future climate change scenarios on the balance between productivity and greenhouse gas emissions from sheep grazing systems. *Livestock Science* 147, 126–138.
- Bergquist R and Rinaldi L 2010. Health research based on geospatial tools: a timely approach in a changing environment. *Journal of Helminthology* 84, 1–11.
- Bessai W 2006. Welfare of broilers: a review. *Worlds Poultry Science Journal* 62, 455–466.
- Champion RA, Rutter SM, Penning PD and Rook AJ 1994. Temporal variation in grazing behaviour of sheep and the reliability of sampling periods. *Applied Animal Behaviour Science* 42, 99–108.
- Charlier J, Claerebout E, De Muelenaere E and Vercauteren J 2005. Associations between dairy herd management factors and bulk tank milk antibody levels against *Ostertagia ostertagi*. *Veterinary Parasitology* 133, 91–100.
- Cockram MS and Mitchell MA 1999. Role of research in the formulation of 'rules' to protect the welfare of farm animals during road transportation. In *EC Climate Change Policy* (ed. AJ Russel, CA Morgan, CJ Savory), pp. 43–64. British Association for Animal Science, Edinburgh, UK. Retrieved from http://ec.europa.eu/clima/policies/adaptation/index_en.htm
- Cornell S 2005. Modelling nematode populations: 20 years of progress. *Trends Parasitol* 21, 542–545.
- Defra 2013. Adapting to climate change. Retrieved April 24, 2013, from <http://www.gov.uk/government/policies/adapting-to-climate-change>
- de Rancourt M, Fois N, Lavin M, Tchakerian E and Vallerand F 2006. Mediterranean sheep and goats production: an uncertain future. *Small Ruminant Research* 62, 167–179.
- Dobson ADM, Finnie TJR and Randolph SE 2011. A modified matrix model to describe the seasonal population ecology of the European tick *Ixodes ricinus*. *Journal of Applied Ecology* 48, 1017–1028.
- Dobson RJ, Barnes EH, Tyrrell KL, Hosking BC, Larsen JWA, Besier RB, Love S, Rolfe PF and Bailey JN 2011. A multi-species model to assess the effect of refugia on worm control and anthelmintic resistance in sheep grazing systems. *Australian Veterinary Journal* 89, 200–208.
- Eckard RJ and Cullen BR 2011. Impacts of future climate scenarios on nitrous oxide emissions from pasture based dairy systems in south eastern Australia. *Animal Feed Science and Technology* 166–167, 736–748.
- Ford SE and Chintala MM 2006. Northward expansion of a marine parasite: testing the role of temperature adaptation. *Journal of Experimental Marine Biology and Ecology* 339, 226–235.
- EU 2009. Agriculture and Rural Development. Retrieved January 22, 2013, from http://ec.europa.eu/agriculture/climate-change/pdf/sum2009_en.pdf
- Europa 2012. Climate action: forests and agriculture. Retrieved January 22, 2013, from http://ec.europa.eu/clima/policies/adaptation/index_en.htm
- Fox NJ, Marion G, Davidson RS, White PCL and Hutchings MR 2012. Livestock helminths in a changing climate: approaches and restrictions to meaningful predictions (Review). *Animals* 2, 93–107.
- Fox NJ, White PCL, McClean CJ, Marion G, Evans A and Hutchings M 2011. Predicting impacts of climate change on *Fasciola hepatica* risk. *PLoS One* 6, e16126. doi:10.1371/journal.pone.0016126.
- Garnett T 2013. Food sustainability: problems, perspectives and solutions. *Proceedings of the Nutrition Society* 72, 29–39.
- Gaughan JB, Lacetera N, Valtorta SE, Khalifa HH, Hahn L and Mader T 2009. Response of domestic animals to climate challenges. In *Biometeorology for adaption to climate variability and change* (ed. KL Ebi, I Burton, GR McGregor), vol. 1, pp. 131–170. Springer.
- Gauly M, Bollwein H, Breves G, Brugemann K, Danicke S, Das G, Demeler J, Hansen H, Isselstein J, König S, Loholter M, Martinsohn M, Meyer U, Potthoff M, Sanker C, Schroder B, Wrage N, Meibaum B, von Samson-Himmelstjerna G, Stinshoff H and Wrenzycki C 2013. Future consequences and challenges for dairy cow production systems arising from climate change in Central Europe – a review. *Animal* 7, 843–859.
- Gill M, Smith P and Wilkinson JM 2010. Mitigating climate change: the role of domestic livestock. *Animal* 4, 323–333.
- Gray J, Dautel H, Estrada-Pena A, Kahl O and Lindgren E 2009. Effects of climate change on ticks and tick-borne diseases in Europe. Invited Review. *Interdisciplinary perspectives on infectious diseases 2009*, article ID 593232, 12pp., doi:10.1155/2009/593232.
- González-Warleta M, Lladosa S, Castro-Hermida JA, Martínez-Ibeas AM, Conesa D, Muñoz F, López-Quílez A, Manga-González Y and Mezo M 2013. Bovine paramphistomosis in Galicia (Spain): prevalence, intensity, aetiology and geospatial distribution of the infection. *Veterinary Parasitology* 191, 252–263.
- Haskell M, Kettlewell P, McCloskey E, Wall E, Hutchings Mand and Mitchell M 2011. *Animal Welfare and Climate Change: Impacts, Adaptations, Mitigation and Risks*. Defra Project Report AW0513. Retrieved from <http://www.defra.gov.uk/>
- Houdijk JGM 2008. Influence of periparturient nutritional demand on resistance to parasites in livestock. *Parasite Immunology* 30, 113–121.
- Hudson PJ, Cattadori IM, Boag B and Dobson AP 2006. Climate disruption and parasite-host dynamics: patterns and processes associated with warming and the frequency of extreme climatic events. *Journal of Helminthology* 80, 175–182.
- IFAD 2010. *Livestock and Climate Change – Livestock Thematic Papers* – Retrieved from <http://www.ifad.org/climate/>
- Kanneganti VR and Kaffka SR 1995. Forage availability from a temperate pasture managed with intensive rotational grazing. *Grass and Forage Science* 50, 55–62.
- Kaplan RM 2004. Drug resistance in nematodes of veterinary importance: a status report. *Trends in Parasitology* 20, 477–481.
- Kenyon F and Jackson F 2012. Targeted flock/herd and individual ruminant treatment approaches. *Veterinary Parasitology* 186, 10–17.

- Kettlewell PJ, Hampson CJ, Green NR, Teer NJ, Veale BM and Mitchell MA 2001a. Heat and moisture generation of livestock during transportation. In Proceedings of 6th International livestock environment symposium, 21–23 May, Louisville, Kentucky (ed. RR Stowell, R Bucklin, RW Bottcher), pp. 519–526. American Society of Agricultural Engineers, St Joseph, Missouri.
- Kettlewell PJ and Mitchell MA 2001b. Mechanical ventilation, improving the welfare of boiler chickens in transit. *Journal of the Royal Agricultural Society* 162, 175–184.
- Kettlewell PJ, Hoxey RP, Hampson CJ, Green NR, Veale BM and Mitchell MA 2001c. Design and operation of a prototype mechanical ventilation system for livestock transport vehicles. *Journal of Agricultural Engineering Research* 79, 429–439.
- Knox JW, Hurford A, Hargreaves L and Wall E 2012. Climate Change Risk Assessment for the Agriculture Sector – (Defra Project Code–GA0204) January 2012.
- Kyvsgaard N-C, Jensen HB, Ambrosen T and Toft N 2013. Temporal changes and risk factors for foot-pad dermatitis in Danish broilers. *Poultry Science* 92, 26–32.
- Leathwick DM, Hosking BC, Bisset SA and McKay CH 2009. Managing anthelmintic resistance: is it feasible in New Zealand to delay the emergence of resistance to a new anthelmintic class? *New Zealand Veterinary Journal* 57, 181–192.
- Leathwick DM, Waghorn TS, Miller CM, Candy PM and Oliver AMB 2012. Managing anthelmintic resistance – use of a combination anthelmintic and leaving some lambs untreated to slow the development of resistance to ivermectin. *Veterinary Parasitology* 187, 285–294.
- Manning L, Chadd SA and Baines RN 2007. Water consumption in broiler chicken: a welfare indicator. *Worlds Poultry Science Journal* 63, 63–71.
- Mason C, Stevenson H, Cox A, Dick I and Rodger C 2012. Disease associated with immature paramphistome infection in sheep. *Veterinary Record* 170, 343–344.
- Maurer V and Baumgärtner J 1992. Temperature influence on life table statistics of the chicken mite *Dermanyssus gallinae* (Acari: Dermanyssidae). *Experimental and Applied Acarology* 15, 27–40.
- May JD and Lott BD 1992. Feed and water consumption patterns of broilers at high environmental temperatures. *Poultry Science* 71, 331–336.
- McGlone J and Pond WG 2003. Pig production: biological principles and applications. Delmar Publishers, Clifton Park, NY.
- Meluzzi A, Fabbri C, Folegatti E and Sirri F 2008. Survey of chicken rearing conditions: effects of litter quality and stocking density on productivity, foot dermatitis and carcass injuries. *British Poultry Science* 49, 257–264.
- Millar M, Colloff A and Scholes S 2012. Disease associated with immature paramphistome infection. *Veterinary Record* 171, 509–510.
- Mitchell MA 2006. Using physiological models to define environmental control strategies. In *Mechanistic modelling in pig and poultry production* (ed. RM Gous, TR Morris, C Fisher), pp. 209–228. CABI International, Wallingford, Oxfordshire, UK.
- Mitchell MA and Kettlewell PJ 1998. Physiological stress and welfare of broiler chickens in transit, solutions not problems!. *Poultry Science* 77, 1803–1814.
- Mitchell MA and Kettlewell PJ 2006. Minimización del stress en el transporte de ganado porcino. Plenary Lecture, Proceedings of the III Congreso Mundial de Jamon, Sobre Ciencia, Tecnología y Commercialization, Teruel, Spain, May 17–20, 2005, 11pp.
- Mitchell MA and Kettlewell PJ 2008. Engineering and design of vehicles for long distance road transport of livestock (ruminants, pigs and poultry). An invited review. *Veterinaria Italiana* 44, 213–225.
- Moran D, Topp K, Wall E, Wreford A and Mitchell M. 2009. Climate change impacts on the livestock sector. Defra Project Report AC0307. Retrieved from <http://www.defra.gov.uk/>
- Morgan ER and Wall R 2009. Climate change and parasitic disease: farmer mitigation? *Trends in Parasitology* 25, 308–313.
- Morgan ER and van Dijk J 2012. Climate and the epidemiology of gastrointestinal nematode infections of sheep in Europe. *Veterinary Parasitology* 189, 8–14.
- Nieuwhof GJ and Bishop SC 2005. Costs of the major endemic diseases in Great Britain and the potential benefits of reduction in disease impact. *Animal Science* 81, 23–29.
- Novak SM and Fiorelli JL 2011. Greenhouse gases and ammonia emissions from organic mixed crop-dairy systems: a critical review of mitigation options. *Sustainable Agriculture* 2, 529–556.
- O'Connor LJ, Walkden-Brown SW and Kahn LP 2006. Ecology of the free-living stages of major trichostrongylid parasites of sheep. *Veterinary Parasitology* 142, 1–15.
- O'Mara FP 2012. The role of grasslands in food security and climate change. *Annals of Botany* 6, 1263–1270.
- Papadopoulos E, Himonas C and Coles GC 2001. Drought and flock isolation may enhance the development of anthelmintic resistance in nematodes. *Veterinary Parasitology* 97, 253–259.
- Polley L and Thompson RCA 2009. Parasite zoonoses and climate change: molecular tools for tracking shifting boundaries. *Trends in Parasitology* 25, 285–291.
- Rivington M, Matthews KB, Buchan K, Miller DG, Bellocchi G and Russel G 2013. Climate change impacts and adaptation, scope for agriculture indicated by agro-meteorological metrics. *Agricultural Systems* 114, 15–31.
- Roca-Fernandez AI, O'Donovan MA, Curran J and Gonzalez-Rodriguez A 2011. Effect of pre-grazing herbage mass and daily herbage allowance on perennial ryegrass swards structure, pasture dry matter intake and milk performance of Holstein-Friesian dairy cows. *Spanish Journal of Agricultural Research* 9, 86–99.
- Rodriguez D, de Voil P, Power B, Cox H, Crimp S and Meinke H 2011. The intrinsic plasticity of farm businesses and their resilience to change. An Australian example. *Field Crop Research* 124, 157–170.
- Sargison ND and Scott PR 2011. Diagnosis and economic consequences of triclabendazole resistance in *Fasciola hepatica* in a sheep flock in south-east Scotland. *Veterinary Record* 168, 159.
- Sargison ND, Scott PR and Jackson F 2002. Teladorsagiosis in young lambs and extended periparturient susceptibility in moxidectin-treated ewes grazing heavily contaminated pastures. *Veterinary Record* 151, 353–355.
- Sargison ND, Jackson F, Bartley DJ and Moir A 2005. Failure of moxidectin to control benzimidazole-, levamisole- and ivermectin-resistant teladorsagia circumcincta in a sheep flock. *Veterinary Record* 156, 105–109.
- Schonbach P, Wolf B, Dickhofer U, Wiesmeier M, Chen W, Hongwei W, Gierus M, Butterbach-Bahl K, Kogel-Knabner I, Susenbeth A, Zheng X and Taube F 2012. Grazing effects on the greenhouse gas balance of a temperate steppe ecosystem. *Nutrient Cycling in Agroecosystems* 93, 357–371.
- Schweizer G, Braun W, Deplazes D and Torgerson P 2005. The economic effects of bovine fasciolosis in Switzerland. *Veterinary Record* 157, 188–193.
- Singh A, Casey KD, King WD, Pescatore AJ, Gates RS and Ford MJ 2009. Efficacy of urease inhibitor to reduce ammonia emission from poultry houses. *Journal of Applied Poultry Science* 18, 34–42.
- Smith P and Gregory PJ 2013. Climate change and sustainable food production. *Proceedings of the Nutrition Society* 72, 21–28.
- Sturaro E, Cocca G, Gallo L, Mrad M and Ramanzin M 2009. Livestock systems and farming styles in Eastern Italian Alps: an on-farm survey. *Italian Journal of Animal Science* 8, 541–554.
- Thornton PK 2010. Livestock production: recent trends, future prospects. *Philosophical Transactions of the Royal Society B* 365, 2853–2867.
- Tiley GED 2010. Biological flora of the British Isles: *Cirsium arvense* (L.). *Journal of Ecology* 98, 938–983.
- Topper PA, Wheeler EF, Zajackowski JS, Gates RS, Xin H, Liang Y and Casey KD 2008. Ammonia emissions from two empty broiler houses with built up litter. *Transactions of the ASABE* 51, 219–225.
- Troell K, Waller P and Hoglund J 2005. The development and overwintering survival of free-living larvae of *Haemonchus contortus* in Sweden. *Journal of Helminthology* 79, 373–379.
- Troell K, Tingstedt C and Hoglund J 2006. Phenotypic characterization of *Haemonchus contortus*: a study of isolates from Sweden and Kenya in experimentally infected sheep. *Parasitology* 132, 403–409.
- Turnpenney JR, Parsons DJ, Armstrong AC, Clark JA, Cooper K and Matthews AM 2001. Integrated models of livestock systems for climate change studies. 2. Intensive systems. *Global Change Biology* 7, 163–170.
- van Dijk J 2008. Climate change and the epidemiology of gastrointestinal nematodes of sheep. Unpublished PhD thesis, University of Bristol, UK.
- van Dijk J and Morgan ER 2008. The influence of temperature on the development, hatching and survival of *Nematodirus battus* larvae. *Parasitology* 135, 269–283.
- van Dijk J and Morgan ER 2010. Variation in the hatching behavior of *Nematodirus battus*: polymorphic bet hedging? *International Journal of Parasitology* 40, 675–681.

- van Dijk J and Morgan ER 2011. The influence of water on the migration of infective trichostrongyloid larvae onto grass. *Parasitology* 138, 780–788.
- van Dijk J, David GP, Baird G and Morgan ER 2008. Back to the future: developing hypotheses on the effects of climate change on ovine parasitic gastroenteritis from historical data. *Veterinary Parasitology* 158, 73–84.
- van Dijk J, De Louw MDE, Kalis LPA and Morgan ER 2009. Ultraviolet light increases mortality of nematode larvae and can explain patterns of larval availability at pasture. *International Journal for Parasitology* 39, 1151–1156.
- van Dijk J, Sargison ND, Kenyon F and Skuce PJ 2010. Climate change and infectious disease: helminthological challenges to farmed ruminants in temperate regions. *Animal* 4, 377–392.
- Van Wyk JA 2001. Refugia – overlooked as perhaps the most potent factor concerning the development of anthelmintic resistance. *Onderstepoort Journal of Veterinary Research* 68, 55–67.
- Waghorn TS, Leathwick DM, Miller CM and Atkinson DS 2008. Brave or gullible: testing the concept that leaving susceptible parasites in refugia will slow the development of anthelmintic resistance. *New Zealand Veterinary Journal* 56, 158–163.
- Waghorn TS, Miller CM, Oliver AM and Leathwick DM 2009. Drench-and-shift is a high risk practice in the absence of refugia. *New Zealand Veterinary Journal* 57, 359–363.
- Wall R, Rose H, Elise L and Morgan E 2011. Livestock ectoparasites: integrated management in a changing climate. *Veterinary Parasitology* 180, 82–89.
- White N, Sutherst RW, Hall N and Whish-Wilson P 2003. The vulnerability of the Australian beef industry to impacts of the cattle tick (*Boophilus microplus*) under climate change. *Climatic Change* 61, 157–190.
- Vercruyse J, Schetters T, Knox DP, Willadsen P and Claerebout E 2007. Control of parasitic disease by vaccination – an answer to drug resistance? *Reviews in Science and Technology* 26, 105–115.
- Wolstenholme AJ, Fairweather I, Prichard RK, von Samson-Himmelstjerna G and Sangster NC 2004. Drug resistance in veterinary helminths. *Trends in Parasitology* 20, 469–476.