

Advances in microbial ecosystem concepts and their consequences for ruminant agriculture

J. E. Edwards[†], S. A. Huws, E. J. Kim, M. R. F. Lee, A. H. Kingston-Smith and N. D. Scollan

Institute of Grassland & Environmental Research, Plas Gogerddan, Aberystwyth, Ceredigion, Wales SY23 3EB, UK

(Received 4 December 2007; Accepted 6 December 2007)

Microbial transformations in the rumen ecosystem have a major impact on our ability to meet the challenge of reducing the environmental footprint of ruminant livestock agriculture, as well as enhancing product quality. Current understanding of the rumen microbial ecosystem is limited, and affects our ability to manipulate rumen output. The view of ruminal fermentation as the sum of activities of the dominant rumen microbiota is no longer adequate, with a more holistic approach required. This paper reviews rumen functionality in the context of the microbiota of the rumen ecosystem, addressing ruminal fermentation as the product of an ecosystem while highlighting the consequences of this for ruminant agriculture. Microbial diversity in the rumen ecosystem enhances the resistance of the network of metabolic pathways present, as well as increasing the potential number of new pathways available. The resulting stability of rumen function is further promoted by the existence of rumen microbiota within biofilms. These protected, structured communities offer potential advantages, but very little is currently known about how ruminal microorganisms interact on feed-surfaces and how these communities develop. The temporal and spatial development of biofilms is strongly linked to the availability of dietary nutrients, the dynamics of which must also be given consideration, particularly in fresh-forage-based production systems. Nutrient dynamics, however, impact not only on pathway inputs but also the turnover and output of the whole ecosystem. Knowledge of the optimal balance of metabolic processes and the corresponding microbial taxa required to provide a stable, balanced ecosystem will enable a more holistic understanding of the rumen. Future studies should aim to identify key ecosystem processes and components within the rumen, including microbial taxa, metabolites and plant-based traits amenable to breeding-based modification. As well as gaining valuable insights into the biology of the rumen ecosystem, this will deliver realistic and appropriate novel targets for beneficial manipulation of rumen function.

Keywords: ecosystem, metabolism, microbiota, nutrient, rumen

Introduction

The purpose of the rumen is to convert dietary material into microbial biomass and end products that can be utilised by the animal. Rumen microorganisms anaerobically ferment complex lignocellulosic plant material, which otherwise would be unable to be utilised by ruminants. Fermentation of dietary material generates microbial biomass and valuable microbial end products, the outflow/diffusion of which provides the ruminant with usable forms of protein and energy. The efficiency and sustainability of ruminant production relies on driving the ecosystem carrying out this conversion to follow an optimal network of metabolic processes. Microbial protein from the rumen accounts for more than half of the total protein entering the intestines

(Broudiscou and Jouany, 1995), with three main taxonomic groups accounting for the majority of this microbial biomass (bacteria, protozoa and fungi) (Hobson and Stewart, 1997). Despite this symbiotic host–microbe relationship, conversion of dietary material by ruminants into saleable product is an inefficient process.

A large part of this inefficiency is due to ruminal processes, resulting in ruminants typically converting only ~20% of dietary protein into meat and/or milk (MacRae *et al.*, 1975; Dewhurst *et al.*, 2000). Ruminal recycling of microbial protein and excessive production of ammonia result in poor nitrogen use efficiency (Ulyatt *et al.*, 1980) while the excretion of methane (Monteny *et al.*, 2006) and unincorporated dietary protein (Marini and Van Amburgh, 2005) pose significant environmental problems. These issues form the basis of one of the main challenges that face this industry, namely to reduce the environmental

[†] E-mail: joan.edwards@bbsrc.ac.uk

impact of livestock agriculture. Similarly, ruminal processes are also important in terms of improving the quality of animal products. Consumers are becoming increasingly aware of the relationships between diet and health, resulting in a greater impetus to enhance the nutritional value of ruminant meat and milk. In response to current dietary guidelines (WHO, 2003), research into functional food components has focused on modifying the quantities of those nutrients present naturally in ruminant products, which are considered to play important roles in health. Current targets include increasing the n-3 polyunsaturated fatty acid content of meat (Scollan *et al.*, 2006) and milk (Vlaeminck *et al.*, 2006). Manipulation of microbial processes in the rumen could have a major impact on our ability to meet these challenges, however recent methodological advances have highlighted that our knowledge of ruminal microorganisms is limited. Current understanding of rumen microbial fermentation is based on the sum of the metabolic activities of the dominant species cultivated from this ecosystem (Krause and Russell, 1996) and not of the hundreds of species now acknowledged to exist (Edwards *et al.*, 2004). However, understanding a microbial ecosystem requires more than just comprehending the sum of the activities of the microorganisms present. Consideration must also be given to what is known about the range of microbial taxa present within the ecosystem.

Improving the efficiency and sustainability of ruminant agriculture relies on the rumen ecosystem following an optimal network of metabolic activities. This review aims to broaden our current thinking of rumen function by considering the central properties of the rumen in the context of an ecosystem in which diversity, stability, structure, dynamics (microbial and nutrient) and biomass (dietary and microbial) are the key elements. Enhancing our understanding of the rumen ecosystem will enable targeted and sustainable manipulation of rumen function to be more effectively realised in the future, including exploiting approaches to manipulate rumen function with dietary forage-based approaches.

Current understanding of rumen microbial diversity

Historically, cultivation-based techniques have enabled identification of a large number of species present in the rumen (Hungate, 1966; Mackie and White, 1997). More recently, application of cultivation-independent methodologies to complex ecosystems has highlighted that the number of bacterial species present is underestimated by these approaches (Rappe and Giovannoni, 2003). Reports of analogous findings relating to the main microbial taxa resident in the rumen (Moon-van der Staay *et al.*, 2002; Edwards *et al.*, 2004; Tuckwell *et al.*, 2005; Wright *et al.*, 2006) confirms that the rumen ecosystem is no exception.

The ability of very simple methodological steps, such as incubation time and inoculum size (Davis *et al.*, 2005), to limit cultivation of certain microorganisms goes some way in explaining why only 11% of the bacterial taxa present




Figure 1 The uncultivated bacterial diversity of the rumen. The percentage of operational taxonomic units (OTU) represented by uncultivated (light grey) and cultivated bacteria (dark grey) for the ^atotal rumen bacterial biota is shown (Edwards *et al.*, 2004). The phylum distribution of these are shown for the two predominant phyla, ^b*Bacteroidetes* (*Cytophaga-Flexibacter-Bacteroides* phylum) and ^c*Firmicutes* (low G + C Gram-positive bacteria), along with ^dother minor phyla and OTU of unknown phylum association.

within the rumen has been recovered in culture (Figure 1) (Edwards *et al.*, 2004). Based on 16S rDNA-based analysis it has been suggested that 300 to 400 different bacterial species actually exist within the rumen (Edwards *et al.*, 2004; Yu *et al.*, 2006), as well as other mammalian gut ecosystems (Eckburg *et al.*, 2005). Use of novel and informed cultivation techniques together with molecular-based approaches that can encompass the whole ecosystem (i.e. metagenomics) have highlighted the potential to overcome some of these limitations (Hugenholtz, 2002). Only by utilising novel 'omic' approaches in tandem with new cultivation techniques (Connon and Giovannoni, 2002; Kaerberlein *et al.*, 2002; Zengler *et al.*, 2002), can we effectively start to comprehend the functional role of microorganisms present within the rumen.

There is a need to define how important a diverse range of microorganisms really is to rumen function. Diversity is not only the 'variety' of individuals present within an ecosystem but also their abundance (Magurran, 1988). Numbers have consequences; e.g. the presence of rumen lactic acid producing bacteria (i.e. *Streptococcus bovis* and *Lactobacillus* spp.) does not by itself induce acute acidosis, but a tenfold increase in their number as a result of rapid dietary change does (Coe *et al.*, 1999). Similarly, numbers of individual species do not necessarily equate to importance. High specific activities can result in functional significance despite low numbers, as is the case with hyper-ammonia-producing bacteria, e.g. *Peptostreptococcus anaerobius* and *Clostridium sticklandii* (Russell *et al.*, 1992; Russell and Rychlik, 2000). Hence, our comprehension of rumen microbial diversity needs to move beyond a species inventory towards considering the significance of diversity, and its impact on ecosystem stability.

Ecosystem stability

Stability in microbial ecosystems is a trait derived from two key system properties, resistance and resilience. Resistance

is the ability of a system to withstand change. Diverse ecosystems are thought to buffer perturbations through the capacity of multiple components to carry out related functions. Microbial diversity in the rumen can be predicted to enhance the resistance of the network of metabolic pathways by increasing the number of genes encoding the pathway, as has been observed in other systems (Cohen *et al.*, 2000; Buchanan, 2002). By increasing the gene pool, alternative metabolic pathways also become available, enabling the ecosystem to stabilise more rapidly after change to a new equilibrium (e.g. change in diet). The more resistant the metabolic pathways, and the more diverse the resource of novel pathways, the more resilient the microbial ecosystem will become.

Resilience is the speed with which a system can return to its original or a new equilibrium following perturbation. System resilience, however, is a complex concept. It is widely accepted that diversity plays a key role in determining resilience, as diversity has been demonstrated to stabilise community and ecosystem processes (Girvan *et al.*, 2005). However, it is also apparent that systems of equivalent diversity do not always have the same stability (Griffiths *et al.*, 2004), and that diversity itself can also destabilise population processes (Tilman, 1996). System resilience is therefore concluded not to be a consequence of diversity alone.

Describing the rumen in terms of system stability should help to increase our ability to comprehend the ecosystem and its fundamental structure. Stability, as a function of resistance and resilience, has been previously related to specific components within a community (Griffiths *et al.*, 2004). However, complex microbial ecosystems often contain more than one species capable of carrying out the same role. Due to this redundancy, it is clear that linking resilience to individual system components is of limited value in complex microbial ecosystems like the rumen, particularly as no role has been assigned to the majority of the taxa present.

Structural organisation of rumen microbiota

Together with increased understanding of the metabolic pathways that occur within the rumen ecosystem, we must also consider the physical location of the microbial consortia resident within the rumen, which are responsible for these processes. Traditionally, the structure of the rumen bacterial population is considered to be the sum of three pools of microorganisms; those that are solid associated, liquid associated and tissue associated. It is clear that these populations are different from each other (LaRue *et al.*, 2005; Cho *et al.*, 2006; Sadet *et al.*, 2007), and that, of these, the solid-associated microbial biomass is the largest (Craig *et al.*, 1987) and plays the most important role in feed digestion (McAllister *et al.*, 1994).

Recent microscopical evidence has also shown that attached bacterial communities are commonly enclosed in an extracellular polymeric matrix, thus defining them as biofilm communities (Figure 2) (Mayorga *et al.*, 2007). Organisation of bacteria into biofilms has two main




Figure 2 Microbial feed-associated biofilms. Scanning electron microscopy image of a biofilm community present on the adaxial surface of *Lolium perenne* following 2 h of incubation in the presence of rumen fluid and under *in vitro* conditions mimicking the rumen ecosystem (Mayorga *et al.*, 2007). The plant waxy cuticle (PWC) and the scale bar of 10 μm are also indicated on the image.

advantages. Firstly, the self-produced extracellular polymeric substances (EPS) that coat biofilm communities trap and concentrate nutrients and enzymes, enabling microbes within the biofilm to readily exchange substrates (Minato *et al.*, 1966; Wolin *et al.*, 1997; Michalet-Doreau *et al.*, 2001). Secondly, mature 'stable' multi-species biofilms are resistant to detachment (McAllister *et al.*, 1994; Harrison *et al.*, 2005; LaRue *et al.*, 2005) and so microbes are structurally protected from a range of forms of attack including antibodies (Costerton *et al.*, 1981), antimicrobial agents (Gilbert and Brown, 1995; Allison *et al.*, 2000) and bacteriophage (Costerton *et al.*, 1987). It has also been considered that biofilms offer protection from predation (Costerton *et al.*, 1987), although data now suggests that many protozoa have mechanisms of efficiently preying on these attached communities (Lawrence and Snyder, 1998; Pederson, 1990; Huws *et al.*, 2005). Therefore, due to the structural stability of biofilm communities, it is important to consider this phenotype in the context of the rumen microbial ecosystem, particularly when developing novel strategies to manipulate rumen microbiota.

Microorganisms attach rapidly to dietary substrates in the rumen (Bonhomme, 1990; McAllister *et al.*, 1990; Edwards *et al.*, 2007). Association of microbes with feed particles occurs either randomly or through response to chemo-attractants such as soluble carbohydrates (McAllister *et al.*, 1994; Dehority and Orpin, 1997). Attachment itself can occur in a range of ways, from specific mechanisms requiring binding proteins and receptors to non-specific mechanisms that rely on physico-chemical forces such as van-der Waals forces (McAllister *et al.*, 1994; Miron *et al.*, 2001).

The specific order in which various microorganisms colonise the surface of feed is thought to influence the spatial organisation within a biofilm community (McAllister *et al.*, 1994), although the rate of initial attachment of rumen bacteria to forage in the rumen is thought to be similar between colonising species (Koike *et al.*, 2003; Edwards *et al.*, 2007). Primary colonising microorganisms

attached in favourable nutrient niches proliferate to form functional micro-colonies, where high concentrations of specific digestive enzymes are produced (Cheng *et al.*, 1995; Saginala *et al.*, 1997). Release of nutrients or end products by the primary colonisers attract secondary microorganisms to the site of digestion, with the EPS produced facilitating their attachment (McAllister *et al.*, 1994; McAllister and Cheng, 1996). Utilisation of nutrients by primary and secondary colonisers stimulates further proliferation and subsequent development and maturation of the biofilm into a structured consortium (Costerton *et al.*, 1987). Formation of mature biofilms on fresh forage has recently been shown to be maximal after 2 h of incubation, under conditions mimicking the rumen (Mayorga *et al.*, 2007).

The diverse range of ruminant dietary material available provides a wide variety of substrates for colonisation in the rumen. This will affect the development of biofilm communities, as will the physical structure of forage. Structurally complex lignocellulosic substrates undergo a complex and sequential microbial attack (Cheng *et al.*, 1980), involving both bacteria and anaerobic fungi (Theodorou *et al.*, 1996; Gordon and Phillips, 1998). Furthermore, the mechanical process of ingestive mastication increases the accessibility of dietary nutrients and increases the surface area available for microbial attachment and enzymatic attack (Bowman and Firkins, 1993; McAllister *et al.*, 1994; Pan *et al.*, 2003). *In sacco* studies with fresh forage have shown that mastication has an effect on microbial colonisation. Early rumen bacterial populations (<4 h) colonising on intact perennial ryegrass were observed to differ from those colonising on masticated perennial ryegrass (Figure 3) (Kim *et al.*, 2006).

Impact of diet on rumen microbial biomass

Rumen microbial fermentation essentially occurs under continuous-culture conditions, whereby a frequent input of dietary material and removal of microbial biomass and end products enables the continual growth of rumen microbes. Diets influence ruminal turnover through differences in the relative rates of fermentation and energy/nutrient density, e.g. readily fermentable processed barley grains *v.* bulky barley straw. However, the greater the flow rate of nutrients through the rumen, the smaller the window of opportunity for microbial digestion and growth (Robinson *et al.*, 1985). On this basis it could be suggested that decreasing the ruminal passage of dietary material may improve the efficiency of its conversion into rumen microbial biomass. In addition to the type of diet, however, differences in intake, feeding regimes and the way the diet is presented (i.e. ground *v.* chopped) also influence the rate of passage (Russell *et al.*, 1992), making the practical optimisation of ruminal flow rate complex.

As protein is one of the most expensive dietary components, much work has focussed on maximising the amount of protein reaching the ruminant hindgut (Russell *et al.*, 1992). Rumen microbial biomass can account for more than




Figure 3 Effect of mastication on rumen bacteria colonising fresh perennial ryegrass. Bacterial 16S rDNA denaturing gradient gel electrophoresis profiles of rumen *in sacco* incubated samples of intact (F) and masticated (B) perennial ryegrass were analysed (Kim *et al.*, 2006). Branch labels indicate sample type and incubation time (h), with the scale indicating the percentage of similarity between bacterial populations which have colonised the bag residues. White shading highlights the early incubation intact forage samples (0.25 to 2 h), which formed a separate and distinct cluster.

50% of the protein entering the intestines (Broudicou and Jouany, 1995) with the remainder derived from endogenous nitrogen sources (8% to 10%) (Bartram, 1987) and undegraded protein. Generally two main approaches have been used to increase the supply of utilisable protein entering the intestines: enhancing the efficiency of rumen microbial biomass production and protecting dietary protein from ruminal degradation. Efforts to increase the efficiency of microbial growth have centred on improving the supply of nutrients to the rumen microbial population through a variety of different approaches, e.g. synchrony/balance (Rooke *et al.*, 1987; Dewhurst *et al.*, 2000), forage quality (Beever and Siddons, 1986; Lee *et al.*, 2002; Merry *et al.*, 2006) and concentrate supplementation (Thomas *et al.*, 1980; Harstad and Vik-Mo, 1985). To increase the amount of undegraded dietary protein entering the intestines, diets high in non-degradable protein sources have been used, as well as processing (i.e. heat or chemical treatment) and the addition of protective agents such as tannins, which complex with protein to decrease degradation. In some situations this leads to a deficiency of fermentable energy for rumen microorganisms and a consequent reduction in microbial growth (Siddons *et al.*, 1985; Rooke *et al.*, 1986; Christensen *et al.*, 1993). Conversely though, it has been noted that responses of microbial protein synthesis to protein supplements differ according to the type of energy supplied in the diet (Dewhurst *et al.*, 1999).

In addition to the passage and fermentation rates of the feed, consideration should be paid to the temporal availability of feed nutrients to the rumen microbiota. For example, in fresh-forage-based systems not all of the

soluble nutrients are immediately available as a result of mastication (Boudon and Peyraud, 2001; Kim *et al.*, 2006) or as a result of exposure of ingested viable plant cells to the rumen environment (Kingston-Smith *et al.*, 2003b). Although it has been clearly demonstrated that dietary protein can be directly utilised or broken down in the rumen by microbial taxa (Wallace *et al.*, 1997), *in situ* the plant protein may be protected from degradation by complexing with phenolic compounds (Theodorou *et al.*, 2000; Winters and Minchin, 2001) or can be inaccessible due to being compartmented within plant cells (Kingston-Smith and Theodorou, 2000).

Plant cells are complex, containing numerous single- and double-membrane-bound organelles (e.g. plant chloroplast, mitochondria and vacuoles). Spatial separation has an important role in coordinating plant metabolism. The majority of plant protease activity exists in lytic vacuoles away from the cytosolic proteins, which are potential substrates, thereby minimising the futile cycling of proteins (Thomas and Stoddart, 1980; Feller, 1986). During stress or herbivory, proteins can be degraded in response to cellular signals or physical loss of compartmentation. As recent estimates suggest that approximately half of the plant cells are intact and viable in ingested fresh forage (Gay, Kim and Kingston-Smith, unpublished), it is clear that plant structure plays an important role in determining nutrient availability (Kingston-Smith *et al.*, 2008). Furthermore, the ability of the plant cell's own enzymatic complement to access the nutrients in intact plant cells before they can be accessed by ruminal microorganisms also gives fresh forage a dynamic and temporal metabolite fingerprint distinct from that of conserved forages or concentrate feeds (Zhu *et al.*, 1999; Kingston-Smith *et al.*, 2003a and 2005). Therefore, the role of dietary material type can have an impact on the rumen microbial biomass beyond the chemical analysis of the nutrients it contains.

Manipulation of rumen microbial processes

The complexities of rumen function indicated above make manipulation of rumen microbial processes a significant challenge. Some goals for the ruminant livestock industry target the optimisation of metabolic processes within the rumen ecosystem as a whole, whereas others are focussed on particular metabolic pathways within the ecosystem. Improvements in the balance of rumen microbial metabolic processes to date have been achieved either by indirectly perturbing the microbial populations by exploiting specific characteristics of feed or supplements, or by intervening directly to modulate numbers of ruminal microorganisms with desirable traits.

Differences in the chemical composition of plants mean that different feeds provide distinct opportunities for microbial fermentation because of the presence of specific niches. Understanding of substrate-based microbial population shifts, and why they occur, will enable us to apply the appropriate techniques to modify ruminal microbial

populations and fermentation end products. For example, significant shifts in the microbial population can be achieved according to silage type (grass *v.* red clover) combined with dietary supplementation with fish oil at 2% and 3% of dry matter intake (Huws *et al.*, 2006). The latter result could explain how increased substrate supply and altered microbial lipid metabolism resulting from dietary fish oil supplementation increases the deposition of beneficial n-3 polyunsaturated fatty acids into ruminant products (Shingfield *et al.*, 2003; Lee *et al.*, 2005).

Direct manipulation of ruminal microorganisms can be achieved by use of feed additives such as ionophores, antibiotics and methane inhibitors (Nagaraja *et al.*, 1997). The modes of action of these additives vary but in general they result in beneficial changes in ruminal metabolism mediated through their effects on a wide or specific range of taxa. Ionophores act on a wide range of Gram-positive bacteria and generally cause an increase in the ruminal production of the glucogenic precursor propionate relative to a decrease in acetate, as well as decreasing the ruminal production of lactate, methane and ammonia (Russell and Strobel, 1989). Some antibiotics change ruminal fermentation parameters more specifically, e.g. virginiamycin decreases lactate production (Coe *et al.*, 1999). Other antibiotics act by inhibiting bacteria that are detrimental to ruminant health, such as *Fusobacterium necrophorum* (Nagaraja *et al.*, 1999; Wallace *et al.*, 2001; Edwards *et al.*, 2005). Other direct approaches to modify ruminal microbiota have involved either the removal or addition of microorganisms, e.g. by defaunation (Demeyer and Van Nevel, 1979) or introduction of probiotic microorganisms with and without genetically enhanced metabolic capabilities (Gregg, 1995; Kamra and Pathak, 2005). The practical implementation of these latter approaches, however, has had limited success to date within the livestock industry (Kobayashi *et al.*, 2000 and 2004).

Since the ban on the use of antimicrobial feed additives for growth promotion within the EU, there has been a move to the development and use of plant secondary metabolites as alternative manipulators of rumen function (Wallace, 2004). Plant secondary metabolites can exert their effect directly on microbial taxa and their metabolism. Saponins (Shi *et al.*, 2004) and catecholamines (Lafontan *et al.*, 2002) directly affect ruminal lipid metabolism by inhibiting the action of lipolytic bacteria. Indirect effects can also occur when the availability of dietary metabolites is modified, e.g. by condensed tannins (Theodorou *et al.*, 2000; Min *et al.*, 2003). Harnessing the plant enzyme polyphenol oxidase (PPO) decreases the extent of red clover proteolysis both *in silo* (Sullivan and Hatfield, 2006) and in the rumen (Albrecht and Broderick, 1992) through the complexing of leaf proteins (Winters and Minchin, 2001) and/or the denaturation of plant proteases (Jones *et al.*, 1995; Lee *et al.*, 2004). Similar mechanisms have also been proposed for observed beneficial effects on rumen lipolysis and biohydrogenation (Lee *et al.*, 2003, 2004 and 2007). Modifying plant enzymatic capabilities is another promising area, with

recent advances resulting in the generation of forage that can breakdown its own cell-wall structure upon cell death (Buanafina *et al.*, 2006). Clear potential exists for sustainable improvements in nutrient use efficiency and product quality through the further exploitation of plant secondary metabolites and metabolism.

Summary and conclusions

Comprehension of the rumen microbial ecosystem based on the sum of activities of dominant rumen microbiota is limited, particularly in light of the lack of phenotypic knowledge of the wealth of microbial taxa now acknowledged to populate this ecosystem. Microbial diversity and its implications for metabolic pathways in the rumen should be carefully considered if targeted modification of ruminal metabolism is to be effectively realised, especially when processes occur within biofilms. Understanding of rumen metabolism needs to incorporate temporal and spatial dynamics of ruminal microbiota, together with the dynamics associated with dietary material and nutrient availability. Combined knowledge of the optimal balance of microbial taxa, their metabolic processes and factors affecting the properties of the rumen ecosystem is limited. A more holistic approach to comprehending the rumen microbial ecosystem is urgently required.

Methodological advances in metabolite, microbial and sequence-based analyses provide the potential for identification of key ecosystem components. However, the power of this increasing toolbox of techniques can only be realised when they are combined with a strong experimental base to allow ecosystem changes with a significant impact to be identified. This will generate novel targets requiring the ability to manipulate key processes through modifying the relevant microbial species, metabolites and related ecosystem properties. Further identification of key metabolites and plant-based traits/activities that can be modified in important agronomical forage crops will offer novel means to manipulate dietary material through selective plant breeding and modification. By comprehending more fully the ecosystem processes that occur within the rumen, targeted manipulation of rumen function will be more effectively realised, while gaining valuable insight into the biology of this system.

Acknowledgements

The authors gratefully acknowledge BBSRC, DEFRA and the EU for funding their research. Thanks are also given to Mike Theodorou for his support and encouragement of this work. The scanning electron microscopy expertise of Barry Thomas is also gratefully acknowledged.

References

Albrecht KA and Broderick GA 1992. Ruminal *in vitro* degradation of protein from different legume species. US Dairy Forage Research Centre, Madison, USA. Research Summaries, pp. 92–94.

Allison DG, McBain AJ and Gilbert P 2000. Biofilms: problems of control. In Community structure and co-operation in biofilms (ed. DG Allison, P Gilbert, HM Lappin-Scott and M Wilson). Cambridge University Press, Cambridge, UK.

Bartram CJ 1987. The endogenous protein content of ruminant proximal duodenal digests. PhD Thesis, University of Nottingham, Sutton Bonington, Leicestershire, UK.

Beever DE and Siddons RC 1986. Digestion and metabolism in the grazing ruminant. In Control of digestion and metabolism in the ruminant (ed. IW McDonald and ACI Warner), pp. 432–447. University of New England Publishing Unit, Armidale, Australia.

Bonhomme A 1990. Rumen ciliates: their metabolism and relationships with bacteria and their hosts. *Animal Feed Science and Technology* 30, 203–212.

Boudon A and Peyraud JL 2001. The release of intracellular constituents from fresh ryegrass (*Lolium perenne*) during ingestive mastication in dairy cows: effect of intracellular constituent, season and stage of maturity. *Animal Feed Science and Technology* 93, 229–245.

Bowman JGP and Firkins JL 1993. Effects of forage species and particle size on bacterial cellulolytic activity and colonisation *in situ*. *Journal of Animal Science* 71, 1623–1633.

Broudicou L and Jouany JP 1995. Reassessing the manipulation of protein synthesis by rumen microbes. *Reproduction, Nutrition, Development* 35, 517–535.

Buanafina MM, de O Langdon T, Hauck BD, Dalton SJ and Morris P 2006. Manipulating the phenolic acid content and digestibility of Italian ryegrass (*Lolium multiflorum*) by vacuolar targeted expression of a fungal ferulic acid esterase. *Applied Biochemistry and Biotechnology* 130, 415–426.

Buchanan M 2002. Nexus: small worlds and the groundbreaking science of networks. W. W. Norton & Company Ltd., London, UK.

Cheng KJ, Fay JP, Howarth RE and Costerton JW 1980. Sequence of events in the digestion of fresh legume leaves by rumen bacteria. *Applied and Environmental Microbiology* 40, 613–625.

Cheng KJ, McAllister TA and Costerton JW 1995. Biofilm of the ruminant digestive tract. In *Microbial biofilms* (ed. H Lappin-Scott and JM Costerton), pp. 221–232. Cambridge University Press, Cambridge, UK.

Cho SJ, Cho KM, Shin EC, Lim WJ, Hong SY, Choi BR, Kang JM, Lee SM, Kim YH, Kim H and Yun HD 2006. 16S rDNA analysis of bacterial diversity in three fractions of cow rumen. *Journal of Microbiology and Biotechnology* 16, 92–101.

Christensen RA, Cameron MR, Klusmeyer TH, Elliot JP, Clark JH, Nelson DR and Yu Y 1993. Influence of the amount and degradability of dietary protein on nitrogen utilization by dairy cows. *Journal of Dairy Science* 76, 3497–3513.

Coe ML, Nagaraja TG, Sun YD, Wallace N, Towne EG, Kemp KE and Hutcheson JP 1999. Effect of virginiamycin on ruminal fermentation in cattle during adaptation to a high concentrate diet and during an induced acidosis. *Journal of Animal Science* 77, 2259–2268.

Cohen R, Erez K, ben-Avraham D and Havlin S 2000. Resilience of the internet to random breakdowns. *Physical Review Letters* 85, 4626–4628.

Connon SA and Giovannoni SJ 2002. High-throughput methods for culturing microorganisms in very-low nutrient media yield diverse new marine isolates. *Applied and Environmental Microbiology* 68, 3878–3885.

Costerton JW, Irvin RT and Cheng KJ 1981. The bacterial glycocalyx in nature and disease. *Annual Review Microbiology* 35, 299–324.

Costerton JW, Cheng KJ, Geesey GG, Ladd TI, Nickel JC, Dasgupta M and Marrie TJ 1987. Bacterial biofilms in nature and disease. *Annual Review Microbiology* 41, 435–464.

Craig WM, Broderick GA and Ricker DB 1987. Quantitation of microorganisms associated with the particulate phase of ruminal ingesta. *Journal of Nutrition* 117, 56–62.

Davis KE, Joseph SJ and Janssen PH 2005. Effects of growth medium, inoculum size, and incubation time on culturability and isolation of soil bacteria. *Applied and Environmental Microbiology* 71, 826–834.

Dehority BA and Orpin GC 1997. Development of and natural fluctuations in rumen microbial populations. In *The rumen microbial ecosystem* (ed. PH Hobson and CS Stewart), pp. 196–245. Chapman & Hall, London, UK.

Demeyer DI and Van Nevel CJ 1979. Effect of defaunation on the metabolism of rumen micro-organisms. *British Journal of Nutrition* 42, 515–524.

Dewhurst RJ, Aston K, Fisher WJ, Evans RT, Dhanoa MS and McAllan AB 1999. Comparison of energy and protein sources offered at low levels in grass-silage based diets for dairy cows. *Animal Science* 68, 789–799.

- Dewhurst RJ, Davies DR and Merry RJ 2000. Microbial protein supply in the rumen. *Animal Feed Science and Technology* 85, 1–21.
- Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, Gill SR, Nelson KE and Relman DA 2005. Diversity of the human intestinal microbial flora. *Science* 308, 1635–1638.
- Edwards JE, McEwan NR, Travis AJ and Wallace RJ 2004. 16s rDNA library-based analysis of ruminal bacterial diversity. *Antonie van Leeuwenhoek* 86, 263–281.
- Edwards JE, McEwan NR, McKain N, Walker N and Wallace RJ 2005. Influence of flavomycin on ruminal fermentation and microbial populations in sheep. *Microbiology* 151, 717–725.
- Edwards JE, Huws SA, Kim EJ and Kingston-Smith AH 2007. Characterisation of the dynamics of initial bacterial colonisation of non-conserved forage in the rumen. *FEMS Microbiology Ecology* 62, 323–335.
- Feller U 1986. Plant proteolytic enzymes in relation to leaf senescence. In *Plant proteolytic enzymes* (ed. MJ Dalling), pp. 49–68. CRC Press, Boca Raton, FL, USA.
- Gilbert P and Brown MRW 1995. Mechanisms of the protection of bacterial biofilms from antimicrobial agents. In *Microbial biofilms – plant microbial biotechnology research 5* (ed. HM Lappin-Scott and JW Costerton), Cambridge University Press, Cambridge, UK.
- Girvan MS, Campbell CD, Killham K, Prosser JI and Glover LA 2005. Bacterial diversity promotes community stability and functional resilience after perturbation. *Environmental Microbiology* 7, 301–313.
- Gordon GLR and Phillips MW 1998. The role of anaerobic gut fungi in ruminants. *Nutrition Research Reviews* 11, 133–168.
- Gregg K 1995. Engineering gut flora of ruminant livestock to reduce forage toxicity: progress and problems. *Trends in Biotechnology* 13, 418–421.
- Griffiths BS, Kuan HL, Ritz K, Glover LA, McCaig AE and Fenwick C 2004. The relationship between microbial community structure and functional stability, tested experimentally in an upland pasture soil. *Microbial Ecology* 47, 104–113.
- Harrison JJ, Turner RJ, Marques LLR and Ceri H 2005. Biofilms: a new understanding of these microbial communities is driving a revolution that may transform the science of microbiology. *American Scientist* 93, 508.
- Harstad OM and Vik-Mo L 1985. Estimation of microbial and undegraded protein in sheep on grass silage based diets. *Acta Agriculturae Scandinavica* 25, 37–48.
- Hobson PN and Stewart CS 1997. *The rumen microbial ecosystem*. Chapman & Hall, London, UK.
- Hugenholtz P 2002. Exploring prokaryotic diversity in the genomic era. *Genome Biology* 3, 1–8.
- Hungate RE 1966. *The rumen and its microbes*. Academic Press Inc., New York, USA.
- Huws SA, McBain AJ and Gilbert P 2005. Protozoan grazing and its impact upon population dynamics in biofilm communities. *Journal Applied Microbiology* 98, 238–244.
- Huws SA, Lee MRF, Muetzel S, Wallace RJ and Scollan ND 2006. Effect of forage type and level of fish oil inclusion on bacterial diversity in the rumen. *Reproduction, Nutrition, Development* 46, S99.
- Jones BA, Muck RE and Hartfield RD 1995. Red clover extracts inhibit legume proteolysis. *Journal of the Science of Food and Agriculture* 67, 329–333.
- Kaerberlein T, Lewis K and Epstein SS 2002. Isolating “uncultivable” microorganisms in pure culture in a simulated natural environment. *Science* 296, 1127–1129.
- Kamra DN and Pathak NN 2005. Improvement in livestock productivity by use of probiotics: a review. *Indian Journal Animal Science* 75, 128–134.
- Kim EJ, Edwards JE, Sanderson R, Kingston-Smith AH, Scollan ND and Theodorou MK 2006. Effects of mastication on temporal bacterial colonisation and degradation of fresh perennial ryegrass in the rumen. *Reproduction, Nutrition, Development* 46, S19.
- Kingston-Smith AH and Theodorou MK 2000. Post-ingestion metabolism of fresh forage. *New Phytologist* 148, 37–55.
- Kingston-Smith AH, Bollard AL, Armstead IP, Thomas BJ and Theodorou MK 2003a. Proteolysis and cell death in clover leaves is induced by grazing. *Protoplasma* 220, 119–129.
- Kingston-Smith AH, Bollard AL, Thomas BJ, Brooks AE and Theodorou MK 2003b. Nutrient availability during the early stages of colonization of fresh forage by rumen micro-organisms. *New Phytologist* 158, 119–130.
- Kingston-Smith A, Merry RJ, Leemans DK, Thomas H and Theodorou MK 2005. Evidence in support of a role of plant-mediated proteolysis in the rumens of grazing animals. *British Journal of Nutrition* 93, 73–79.
- Kingston-Smith AH, Davies TE, Edwards JE and Theodorou MK 2008. From plants to animals: the role of plant cell death in ruminant herbivores. *Journal of Experimental Botany*, doi: 10.1093/jxb/erm326.
- Kobayashi Y, Forster RJ and Teather RM 2000. Development of a competitive polymerase chain reaction assay for the ruminal bacterium *Butyrivibrio fibrisolvens* OB156-derived recombinant. *FEMS Microbiology Letters* 188, 185–190.
- Kobayashi Y, Koike S, Taguchi H, Itabashi H, Kam DK and Ha JK 2004. Recent advances in gut microbiology and their possible contribution to animal health and production – A review. *Asian-Australasian Journal of Animal Science* 17, 877–884.
- Koike S, Pan J, Kobayashi Y and Tanaka K 2003. Kinetics of *in sacco* fiber-attachment of representative ruminal cellulolytic bacteria monitored by competitive PCR. *Journal of Dairy Science* 86, 1429–1435.
- Krause DO and Russell JB 1996. How many ruminal bacteria are there? *Journal of Dairy Science* 79, 1467–1475.
- Lafontan M, Berlan M, Stich V, Crampes F, Riviere D, De Glisezinski I, Sengenès C and Galitzky J 2002. Recent data on the regulation of lipolysis by catecholamines and natriuretic peptides. *Annales D’Endocrinologie* 63, 86–90.
- LaRue R, Yu Z, Parisi VA, Egan AR and Morrison M 2005. Novel microbial diversity adherent to plant biomass in the herbivore gastrointestinal tract, as revealed by ribosomal intergenic spacer analysis and *rrs* gene sequencing. *Environmental Microbiology* 7, 530–543.
- Lawrence JR and Snyder RA 1998. Feeding behaviour and grazing impacts of a *Euplotes* sp. on attached bacteria. *Canadian Journal of Microbiology* 44, 623–629.
- Lee MRF, Harris LJ, Moorby JM, Humphreys MO, Theodorou MK, MacRae JC and Scollan ND 2002. Rumen metabolism and nitrogen flow to the small intestine in steers offered *Lolium perenne* containing different levels of water-soluble carbohydrate. *Animal Science* 74, 587–596.
- Lee MRF, Martinez EM and Scollan ND 2003. Plant enzyme mediated lipolysis of *Lolium perenne* and *Trifolium pratense* in an *in vitro* simulated rumen environment. *Aspects of Applied Biology* 70, 115–120.
- Lee MRF, Winters AL, Scollan ND, Dewhurst RJ, Theodorou MK and Minchin FR 2004. Plant mediated lipolysis and proteolysis in red clover with different polyphenol oxidase activities. *Journal of the Science of Food and Agriculture* 84, 1639–1645.
- Lee MRF, Tweed JKS, Moloney AP and Scollan ND 2005. The effects of fish oil supplementation on rumen metabolism and the biohydrogenation of unsaturated fatty acids in beef steers given diets containing sunflower oil. *Animal Science* 80, 361–367.
- Lee MRF, Parfitt L, Scollan ND and Minchin FR 2007. Lipolysis in red clover with different polyphenol oxidase activities in the presence of rumen fluid. *Journal of the Science of Food and Agriculture* 87, 1308–1314.
- Mackie RI and White BA 1997. *Gastrointestinal ecosystems and fermentations*. Chapman & Hall, London, UK.
- MacRae JC, Campbell DR and Eadie J 1975. Changes in the biochemical composition of herbage upon freezing and thawing. *The Journal of Agricultural Science* 84, 125–131.
- Magurran AE 1988. *Ecological diversity and its measurement*. Croom Helm, London, UK.
- Marini JC and Van Amburgh ME 2005. Partition of nitrogen excretion in urine and the feces of Holstein replacement heifers. *Journal of Dairy Science* 88, 1778–1784.
- Mayorga O, Huws SA, Kim EJ, Kingston-Smith AH, Newbold CJ and Theodorou MK 2007. Microbial colonization and subsequent biofilm formation by ruminal microorganisms on fresh perennial ryegrass. *Microbial Ecology in Health and Disease* 19, 26.
- McAllister TA and Cheng KJ 1996. Microbial strategies in the ruminal digestion of cereal grains. *Animal Feed Science and Technology* 62, 29–36.
- McAllister TA, Rode LM, Major DJ, Cheng KJ and Buchanan-Smith JG 1990. The effect of ruminal microbial colonization on cereal grain digestion. *Canadian Journal of Animal Science* 70, 571–579.
- McAllister TA, Bae HD, Jones GA and Cheng KJ 1994. Microbial attachment and feed digestion in the rumen. *Journal of Animal Science* 72, 3004–3018.

- Merry RJ, Davies DR, Lee MRF, Dewhurst RJ, Moorby JM, Leemans DK, Scollan ND and Theodorou MK 2006. Effects of high-sugar ryegrass silage and mixtures with red clover silage on ruminant digestion. 1. *In vitro* and *in vivo* studies of nitrogen utilization. *Journal of Animal Science* 84, 3049–3060.
- Michalet-Doreau B, Fernandez I, Peyron C, Millet L and Fonty G 2001. Fibrolytic activities and cellulolytic bacterial community structure in the solid and liquid phases of rumen contents. *Reproduction, Nutrition, Development* 41, 187–194.
- Min BR, Barry TN, Attwood GT and McNabb WC 2003. The effect of condensed tannins on the nutrition and health of ruminants fed fresh temperate forages: a review. *Animal Feed Science and Technology* 106, 3–19.
- Minato H, Endo A, Ootomo Y and Uemura T 1966. Ecological treatise on the rumen fermentation. II. The amylolytic and cellulolytic activities of fractionated bacterial portions attached to the rumen solids. *The Journal of General and Applied Microbiology* 12, 53–69.
- Miron J, Ben-Ghedalia D and Morrison M 2001. Adhesion mechanisms of rumen cellulolytic bacteria. *Journal of Dairy Science* 84, 1294–1309.
- Monteny GJ, Bannink A and Chadwick D 2006. Greenhouse gas abatement strategies for animal husbandry. *Agriculture, Ecosystems and Environment* 112, 163–170.
- Moon-van der Staay SY, van der Staay GWM, Javorský P, Jouany JP, Michalowski T, Nsabimana E, Machebeuf D, Kišidayová S, Varadyova Z, McEwan NR, Newbold CJ and Hackstein JHP 2002. Diversity of rumen ciliates revealed by 18S ribosomal DNA analysis. *Reproduction, Nutrition, Development* 42, 576.
- Nagaraja TG, Newbold CJ, Van Nevel CJ and Demeyer DI 1997. Manipulation of ruminal fermentation. In *The rumen microbial ecosystem* (ed. PH Hobson and CS Stewart), pp. 523–632. Chapman & Hall, London, UK.
- Nagaraja TG, Sun Y, Wallace N, Kemp KE and Parrott CJ 1999. Effects of tylosin on concentrations of *Fusobacterium necrophorum* and fermentation products in the rumen of cattle fed a high-concentrate diet. *American Journal of Veterinary Research* 60, 1061–1065.
- Pan J, Koike S, Suzuki T, Ueda K, Kobayashi Y, Tanaka K and Okubo M 2003. Effect of mastication on degradation of orchardgrass hay stem by rumen microbes: fibrolytic enzyme activities and microbial attachment. *Animal Feed Science and Technology* 106, 69–79.
- Pederson K 1990. Biofilm development on stainless steel and PVC surfaces in drinking water. *Water Research* 24, 239–243.
- Rappe MS and Giovannoni SJ 2003. The uncultured microbial majority. *Annual Review of Microbiology* 57, 369–394.
- Robinson PH, Sniffen CJ and Van Soest PJ 1985. Influence of level of feed intake on digestion and bacterial yield in the forestomachs of dairy cattle. *Canadian Journal of Animal Science* 65, 437–444.
- Rooke JA, Alvarez P and Armstrong DG 1986. The digestion by cattle of barley and silage diets containing increasing quantities of soya-bean meal. *The Journal of Agricultural Science* 107, 263–272.
- Rooke JA, Lee NH and Armstrong DG 1987. The effects of intraruminal infusions of urea, casein, glucose syrup, and a mixture of casein and glucose syrup on nitrogen digestion in the rumen of cattle receiving grass-silage diets. *British Journal of Nutrition* 32, 89–94.
- Russell JB and Rychlik JL 2000. The isolation, characterisation and enumeration of hyper-ammonia producing ruminal bacteria. *Asian-Australasian Journal of Animal Sciences* 13, 121–127.
- Russell JB and Strobel HJ 1989. Effect of ionophores on ruminal fermentation. *Applied and Environmental Microbiology* 55, 1–6.
- Russell JB, O'Connor JD, Fox DG, Van Soest PJ and Sniffen CJ 1992. A net carbohydrate and protein system for evaluating cattle diets: 1. Ruminal fermentation. *Journal of Animal Science* 70, 3551–3561.
- Sadet S, Martin C, Meunier B and Morgavi DP 2007. PCR-DGGE analysis reveals a distinct diversity in the bacterial population attached to the rumen epithelium. *Animal* 1, 939–944.
- Saginala S, Nagaraja TG, Lechtenberg KF, Chengappa MM, Kemp KE and Hine PM 1997. Effect of *Fusobacterium necrophorum* leukotoxin vaccine on susceptibility to experimentally induced liver abscesses in cattle. *Journal of Animal Science* 75, 1160–1166.
- Scollan ND, Hocquette JF, Nuernberg K, Dannenberger D, Richardson RI and Moloney AP 2006. Innovations in beef production systems that enhance the nutritional and health value of beef lipids and their relationship with meat quality. *Meat Science* 74, 17–33.
- Shi J, Arunasalam K, Yeung D, Kakuda Y, Mittal G and Jiang Y 2004. Saponins from edible legumes: chemistry, processing and health benefits. *Journal of Medicinal Food* 7, 67–78.
- Shingfield KJ, Ahvenjarvi S, Toivonen V, Arola A, Nurmela KVV, Huhtanen P and Griinari JM 2003. Effect of dietary fish oil on biohydrogenation of fatty acids and milk fatty acid content in cows. *Animal Science* 77, 165–179.
- Siddons RC, Paradine J, Gale DL and Evans RT 1985. Estimation of the degradability of dietary protein in the sheep rumen by *in vivo* and *in vitro* procedures. *British Journal of Nutrition* 54, 545–561.
- Sullivan ML and Hatfield RD 2006. Polyphenol oxidase and o-diphenols inhibit post harvest proteolysis in red clover and alfalfa. *Crop Science* 46, 662–670.
- Theodorou MK, Mennim G, Davies DR, Zhu WY, Trinci APJ and Brookman JL 1996. Anaerobic fungi in the digestive tract of mammalian herbivores and their potential for exploitation. *Proceedings of the Nutrition Society* 55, 913–926.
- Theodorou MK, Barahona R, Kingston-Smith A, Sanchez S, Lascano C, Owen E and Morris P 2000. New perspectives on the degradation of plant biomass in the rumen in the absence and presence of condensed tannins. In *Tannins in livestock and human nutrition* (ed. JD Brooker), pp. 44–51. Australian Centre for International Agricultural Research Proceedings Series, 92, Adelaide, Australia.
- Thomas H and Stoddart JL 1980. Leaf senescence. *Annual Review of Plant Physiology* 31, 83–111.
- Thomas PC, Chamberlain DG, Kelly NC and Wait MK 1980. The nutritive value of silages. Digestion of nitrogenous constituents in sheep receiving diets of grass silage and barley. *British Journal of Nutrition* 43, 469–479.
- Tilman D 1996. Biodiversity: population versus ecosystem stability. *Ecology* 77, 350–363.
- Tuckwell DS, Nicholson MJ, McSweeney CS, Theodorou MK and Brookman JL 2005. The rapid assignment of ruminal fungi to presumptive genera using ITS1 and ITS2 RNA secondary structures to produce group-specific fingerprints. *Microbiology* 151, 1557–1567.
- Ulyatt MJ, Beever DE, Thomson DJ, Evans RT and Haines MJ 1980. Measurement of nutrient supply at pasture. *Proceedings of the Nutrition Society* 39, 67A.
- Vlaeminck B, Fievez V, Tamminga S, Dewhurst RJ, van Vuuren AM, De Brabander D and Demeyer D 2006. Milk odd- and branched-chain fatty acids in relation to the rumen fermentation pattern. *Journal of Dairy Science* 89, 3954–3964.
- Wallace RJ 2004. Antimicrobial properties of plant secondary metabolites. *Proceedings of the Nutrition Society* 63, 621–629.
- Wallace RJ, Onodera R and Cotta MA 1997. Metabolism of nitrogen-containing compounds. In *The rumen microbial ecosystem* (ed. PN Hobson and CS Stewart), pp. 283–328. Chapman & Hall, London, UK.
- Wallace RJ, Newbold CJ, Bequette BJ, MacRae JC and Lobley GE 2001. Increasing the flow of protein from ruminal fermentation – review. *Asian-Australasian Journal of Animal Science* 14, 885–893.
- WHO 2003. Diet, nutrition and prevention of chronic diseases. Report of a joint WHO/FAO expert consultation. WHO Technical Report Series, 916. WHO, Geneva.
- Winters AL and Minchin FR 2001. Red clover and the future for pasture legumes as an alternative protein source for ruminants. *IGER Innovations* 5, 30–33.
- Wolin MJ, Miller TL and Stewart CS 1997. Microbe–microbe interactions. In *The rumen microbial ecosystem* (ed. PH Hobson and CS Stewart), pp. 467–491. Chapman & Hall, London, UK.
- Wright AG, Toovey AF and Pimm CL 2006. Molecular identification of methanogenic archaea from sheep in Queensland, Australia reveals more uncultured novel archaea. *Anaerobe* 12, 134–139.
- Yu Z, Yu M and Morrison M 2006. Improved serial analysis of V1 ribosomal sequence tags (SARST-V1) provides a rapid, comprehensive, sequence-based characterization of bacterial diversity and community composition. *Environmental Microbiology* 8, 603–611.
- Zengler K, Toledo G, Rappe M, Elkins J, Mathur EJ, Short JM and Keller M 2002. Cultivating the uncultured. *Proceedings of the National Academy of Science* 99, 15681–15686.
- Zhu WY, Kingston-Smith AH, Troncoso D, Merry RJ, Davies DR, Pichard G, Thomas H and Theodorou MK 1999. Evidence of a role for plant proteases in the degradation of herbage proteins in the rumen of grazing cattle. *Journal of Dairy Science* 82, 2651–2658.