

## Reproductive capacity of Merino ewes fed a high-salt diet

S. N. Digby<sup>1,4†</sup>, D. G. Masters<sup>2,4</sup>, D. Blache<sup>3,4</sup>, M. A. Blackberry<sup>3</sup>, P. I. Hynd<sup>1</sup> and D. K. Revell<sup>1,2,4</sup>

<sup>1</sup>Discipline of Agricultural and Animal Science, School of Agriculture, Food and Wine, The University of Adelaide, Roseworthy, SA 5371, Australia; <sup>2</sup>CSIRO Livestock Industries, Private Bag 5, Wembley, WA 6913, Australia; <sup>3</sup>School of Animal Biology M085, Faculty of Natural and Agricultural Sciences, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia; <sup>4</sup>Future Farm Industries Cooperative Research Centre M081, Faculty of Natural and Agricultural Sciences, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

(Received 19 October 2007; Accepted 6 April 2008; First published online 8 July 2008)

An option to increase the productivity of saline land is to graze sheep on salt-tolerant plants, which, during the summer/autumn period, can contain 20% to 25% of their dry matter as salt. This study assessed the impact of coping with high dietary salt loads on the reproductive performance of grazing ewes. From the time of artificial insemination until parturition, 2-year-old maiden Merino ewes were fed either a high-salt diet (NaCl 13% of dry matter) or control diet (NaCl 0.5% of dry matter). Pregnancy rates, lamb birth weights, milk composition and the plasma concentrations of hormones related to salt and water balance, and energy metabolism were measured. Leptin and insulin concentrations were lower ( $1.4 \pm 0.09$  v.  $1.5 \pm 0.12$  ng/ml;  $P < 0.05$ ) and  $7.2 \pm 0.55$  v.  $8.2 \pm 0.83$  ng/ml;  $P < 0.02$ ) in response to high-salt ingestion as was aldosterone concentration ( $27 \pm 2.7$  v.  $49 \pm 5.4$  pg/ml;  $P < 0.05$ ), presumably to achieve salt and water homeostasis. Arginine vasopressin concentration was not significantly affected by the diets, but plasma concentration of  $T_3$  differed during gestation ( $P < 0.02$ ), resulting in lower concentrations in the high-salt group in the first third of gestation ( $1.2 \pm 0.18$  v.  $1.3 \pm 0.14$  pmol/ml) and higher concentrations in the final third of gestation ( $0.8 \pm 0.16$  v.  $0.6 \pm 0.06$  pmol/ml).  $T_4$  concentration was lower in ewes ingesting high salt for the first two-thirds of pregnancy ( $162 \pm 8.6$  v.  $212 \pm 13$  ng/ml;  $P < 0.001$ ). No substantial effects of high salt ingestion on pregnancy rates, lamb birth weights or milk composition were detected.

**Keywords:** aldosterone, high salt, insulin, leptin, reproduction

### Introduction

Salinity is an increasing problem in agriculture worldwide (Ghassemi *et al.*, 1995) and the use of halophytic plants such as saltbush represents one of the few options available to revegetate salinised landscapes and re-establish grazing systems (Masters *et al.*, 2007). Some landholders in Australia are grazing sheep on saltbush to fill a summer/autumn feed gap (Masters *et al.*, 2006), a period that coincides with the greater demands of late pregnancy for autumn- or winter-lambing ewes. However, feeding saltbush may possibly have a negative impact on reproductive performance of the ewes as high salt intake may reduce intake and cause physiological changes associated with adaptation to the salt load.

High salt intake has been shown to reduce voluntary feed intake (Masters *et al.*, 2005; Blache *et al.*, 2007) as well as

the efficiency of energy use for production (Arieli *et al.*, 1989) in sheep. A decrease either in voluntary feed intake or in fat reserves is usually associated with a decrease in the concentration of metabolic hormones such as insulin and leptin (Chilliard *et al.*, 2005) and recently a high salt ingestion has been shown by Blache *et al.* (2007) to affect energy metabolism through changes in insulin concentrations in sheep fed high salt (20% NaCl) diets. Thus, the ingestion of large amount of salt may impact energy availability and consequently reproductive performance.

There are also physiological 'conflicts' for pregnant ewes fed high-salt diets. Pregnancy is characterised by sodium retention and increased extracellular volume necessary for the maintenance of the mother and growth of the foetus (Davison and Lindheimer, 1989). The renin-angiotensin system (RAS) is responsible for the maintenance of water and salt balance, aldosterone-controlling sodium retention and arginine vasopressin (AVP)-controlling water re-absorption by the kidney. Therefore high salt consumption leads to changes in the RAS and decreased aldosterone

<sup>†</sup> Present address: Liggins Institute, The University of Auckland, Ngapouri Research Farm, Reporoa, 2379 State Highway 5 RD2, Reporoa 3083, New Zealand. E-mail: s.digby@auckland.ac.nz

concentration, which reduces sodium re-absorption and increases sodium excretion. In contrast to aldosterone, high salt intake requires no change in AVP plasma concentration if the intake of fresh water is sufficient to maintain a salt and water balance (Cowley *et al.*, 1986). No change in AVP concentration or plasma osmolality was observed in pregnant sheep compared to non-pregnant sheep (Bell *et al.*, 1986).

Due to the potential impact on feed intake, energy availability and the potential challenge to water and salt balance induced by high salt intake, we hypothesised that high-salt diets will reduce the reproductive capacity of Merino ewes through a reduction in pregnancy rates and foetal growth (birth weight). To investigate the effect of high-salt diets on water and sodium balance, concentrations of aldosterone and AVP were measured, as these hormones are the end products of the regulatory RAS. The concentrations of leptin and insulin were also measured as indicators of changes in energy metabolism.

## Material and methods

### Experimental design

From the time of artificial insemination until parturition, Merino ewes were fed either a high-salt diet or a control diet containing low salt. Ewes receiving the high-salt diet were fed *ad libitum*, while the ewes on the control diet were fed the same amount of organic matter as their paired high-salt counterparts. The reproductive capacities of the ewes were assessed by measuring pregnancy rates, lamb birth weights and milk volume and composition. The concentrations of hormones relating to salt and water balance and energy metabolism were measured to assess the metabolic changes experienced by the ewes.

### Animals

A total of 76 Merino maiden ewes, aged 2 years, and weighing on average 45 kg, were held at The University of Adelaide, Roseworthy Campus. They were allocated randomly to four small paddocks (approximately 6 m × 12 m), with 19 in each paddock. Ovulation was synchronised through the use of intravaginal progesterone sponges (40 mg; Intervet Australia Pty Ltd, Bendigo, Victoria), and 48 h after progesterone sponges were removed, ewes were artificially inseminated via laparoscopic insemination using a small volume (0.3 ml/uterine horn) of diluted fresh mixed semen from two sires (day 0). The experiment was approved by the Animal Ethics Committee of The University of Adelaide.

### Experimental diets

Two pelleted diets were used based on barley and lupin grain, and their ingredient composition and estimated nutrient specifications are shown in Table 1. The high-salt diet was formulated to contain NaCl at a concentration of 130 g/kg dry matter to mimic the concentration of dietary salt in animals grazing saltbush-based pastures in summer and autumn, given that saltbush contains 20% to 25% NaCl (Norman *et al.*, 2002) and that over 50% of the

**Table 1** Ingredient composition and nutrient specifications of the high-salt and control diet

	Control	High salt
Ingredients (% air-dry basis)		
Lupins	38.9	33.8
Barley	17	14.8
Mill mix	25	21.7
Oat offal	5	4.3
Rice hulls	10	8.8
Canola oil	3	2.61
Limestone	1	0.9
Mineral mix	0.1	0.09
Added NaCl	0	13
Nutrient composition (% of dry matter)		
<i>In vitro</i> digestibility of dry matter	70.9	74.9
CP	18.8	18.1
ADF	18.8	14.1
Na content	0.19	4.7

selected diet at this time of year is saltbush (Thomas, 2006; H. C. Norman, personal communication). The diets were prepared by a commercial manufacturer (Lauckes, Murray Bridge, South Australia) and analysed for nutritive value by Feedtest<sup>®</sup>, Hamilton, Victoria. Feed was analysed for sodium concentration by inductively coupled plasma atomic emission spectrometry (Dahlquist and Knoll, 1978) using a Spectro CIROS ICPAES machine (Waite Analytical Services, The University of Adelaide). The daily intake of metabolisable energy (ME) was calculated from organic matter intake and from the estimated ME content of the organic matter fraction of the two diets (9.6 MJ ME/kg OM).

### Timeline

From day 0 through to day 44, ewes were grouped in paddocks. Two groups were fed the control diet and two groups were fed the high-salt diet. Each of the control groups were pair-fed with a group fed the high-salt diet (see below for pair-feeding methodology). During this time, water consumption was determined on a group basis by fitting a flow meter to the water line supplying each small holding paddock. Pregnancy diagnosis was conducted by ultrasound examination on day 44 of gestation, after which 40 single-bearing ewes (20 fed the high-salt diet and 20 fed the control diet) were placed into individual pens in an animal house. Water intake was not measured when animals were in individual pens due to a technical difficulty with flow meters. All ewes offered the high-salt diet were fed *ad libitum* and each day between 0800 and 0900 h, and the amount offered and refused was recorded. If the daily residue exceeded 10% of the total offered for ewes fed the high-salt diet, the amount fed the following day was reduced by 100 g; if no residue was present, the amount fed the following day was increased by 100 g. Ewes in the group fed the control diet were paired with individuals in the high-salt group based on body weight, and their organic matter intake was restricted to match that of the voluntary

intake of their high-salt pair from the previous day. Feed refusals were bulked together over 1 week for analysis and dried at 65°C for 24 h. Dry matter and organic matter contents of the refusals were determined as described elsewhere (Masters *et al.*, 2005). A second ultrasound scan was conducted on day 78 to confirm pregnancy in all ewes. Blood samples by venipuncture of the jugular vein were taken on day 0, 21, 51, 79, 115 and 140 of gestation for hormone analysis, haematocrit and glucose concentrations. Animals were also weighed on these days. At parturition, lambs were weighed within 4 h of birth and the crown-to-rump length measured as an indication of foetal bone growth. Udder volume was determined within 1 week of birth using measurements taken longitudinally over the left and right teats (Bencini and Purvis, 1990). A 5 ml milk sample from each teat was also taken by injecting 1 IU of synthetic oxytocin (Lyppards, Thebarton, South Australia) intramuscularly. Samples were frozen for later analysis of fat and protein using a Milko Scan 133 (Foss Electric, Hillerod, Denmark) calibrated for sheep as described by Bencini (1999).

#### Hormone assays

Plasma leptin was measured in duplicate by a double-antibody radioimmunoassay (Blache *et al.*, 2000). All samples were processed in a single assay and the limit of detection was 0.06 ng/ml. The assay included six replicates of three control samples containing 0.43, 0.98 and 1.68 ng/ml, which were used to estimate the intra-assay coefficients of variation of 6.1%, 6.0% and 6.5%, respectively.


Plasma insulin was assayed in duplicate by a double-antibody radioimmunoassay (Tindal *et al.*, 1978) that had been validated for sheep plasma (Miller *et al.*, 1995). All samples were processed in a single assay and the limit of detection was 0.78  $\mu$ U/ml. Six replicates of three control samples containing 2.45, 3.63 and 8.70  $\mu$ U/ml were included in the assay and were used to estimate the intra-assay coefficients of variation of 7.5%, 2.2% and 3.5%, respectively.

Concentrations of tri-iodothyronine ( $T_3$ ) and thyroxine ( $T_4$ ) were measured using a double-antibody radioimmunoassay (Dawson *et al.*, 1996) and validated for sheep plasma (Zhang *et al.*, 2005). The samples for  $T_3$  were assayed as duplicate 20  $\mu$ l aliquots with the limit of detection 0.3 ng/ml, whereas those for  $T_4$  were assayed as duplicate 20  $\mu$ l (1:40 dilution) aliquots with the limit of detection 0.25 nM/l. Four replicates of two control samples containing 0.87 and 1.34 ng/ml were included in the  $T_3$  assay and were used to estimate the intra-assay coefficients of variation of 4.6% and 7.1%, respectively. Five replicates of two control samples containing 10.8 and 18.7 nM/l were included in the  $T_4$  assay and were used to estimate the intra-assay coefficients of variation of 6.9% and 3.7%.

Concentration of progesterone in the plasma was measured in a single assay using a double-antibody radioimmunoassay after extraction with hexane (Gales *et al.*, 1997). Intra-assay variation was 6.7% at 1.03 ng/ml, 5.3% at 2.18 ng/ml and 5.2% at 4.10 ng/ml. The limit of detection was 0.10 ng/ml.

Concentration of AVP in the plasma was measured using a double-antibody radioimmunoassay after extraction following the assay procedure for arginine vasotocin (Rice, 1982). AVP (Sigma-Aldrich Pty. Ltd, Sydney, Australia) was labelled with  $^{125}$ I (ANSTO, Menai, NSW, Australia) using the chloramine-T method (Greenwood and Hunter, 1963). The first antibody, raised in a rabbit against AVP, reacts with rat, mouse, human, sheep and rabbit and has less than 1% reactivity to oxytocin (Fitzgerald Industries International, Concord, MA, USA). Plasma samples (1 ml) were extracted by reverse-phase liquid chromatography using Maxi-clean C18, 300 mg cartridges (Alltech, Dandenong South, Victoria, Australia) eluted with acetonitrile and 4% acetic acid (75/25). The extract was dried under compressed air at 37°C and reconstituted in 500  $\mu$ l of 0.1 M phosphate buffer (pH 7.5). Duplicates of 200  $\mu$ l were added with 100  $\mu$ l of 0.1 M phosphate buffer and 100  $\mu$ l of 0.1 M phosphate + 0.5% bovine serum albumin (BSA) (pH 7.5). After addition of 50  $\mu$ l of the first antibody the sample was diluted 1 in 30,000 in 0.1 M phosphate and incubated overnight at 4°C. Iodinated AVP (50  $\mu$ l) was then diluted in 0.1 M phosphate to create approximately 5000 cpm/tube was added and incubated overnight at 4°C. After addition of 100  $\mu$ l of donkey anti-rabbit serum (one in 15 in 0.1 M phosphate) containing one in 800 normal rabbit serum and 500  $\mu$ l of 10% polyethylene glycol, the tubes were incubated overnight at 4°C and spun to separate free from bound AVP. The extraction efficiency was 80% as determined from pooled samples spiked with  $^{125}$ I-AVP and incubated for 60 min at 37°C before extraction. The assay was validated for sheep plasma by checking for parallelism using a serial dilution of pooled samples of sheep plasma and measuring AVP in sheep with access to water or with no access to water for 24 h (Figure 1).

Concentration of aldosterone in plasma was measured using a radioimmunoassay described previously by James and Wilson (1976) and modified at The University of Western Australia. Briefly, aldosterone was extracted from 200  $\mu$ l of plasma and from aldosterone standards (Sigma-Aldrich Pty. Ltd) using 2 ml dichloromethane (BDH, Australia) by vortexing for 5 min. After freezing the plasma using a dry ice/acetone bath, the solvent was transferred in a new tube and dried with compressed air at 37°C. The dried extracts were reconstituted in 200  $\mu$ l phosphate buffered saline (PBS) + 0.5% BSA (pH 7.5) and added with 100  $\mu$ l of rabbit anti-aldosterone-3-BSA (1:30 000 in PBS + 0.5% BSA, Fitzgerald Industries International). The antisera cross reactivity was 1.1% with 11-deoxycorticosterone, 0.01% with testosterone and less than 0.001% with estradiol, 11-deoxycortisol, androstosterone, 21-deoxycortisol, oestriol, cortisone, oestrone and dihydrotestosterone (Fitzgerald Industries International). Following addition of antisera, 100  $\mu$ l of tracer (approximately 10 000 dpm/tube in PBS, (1,2,6,7- $^3$ H)-aldosterone; Amersham, Australia) in PBS + 0.5% BSA was added. The tubes were incubated at 4°C for 48 h and then the free and bound aldosterone were separated using 200  $\mu$ l of Dextran-coated charcoal method (1% charcoal, 0.1% Dextran, Sigma, USA).



**Figure 1** Standard curves for radioimmunoassay for (a) aldosterone and (b) arginine vasopressin (AVP) showing parallelism with serial dilution of sheep plasma.

After addition of 2 ml of scintillant (Starcint; Packard Chemical Operations, USA), the radioactivity of the supernatant was counted in a liquid scintillation counter (Tri Carb 1500; Packard, Downers Grove, IL, USA) for 3 min. The extraction efficiency was greater than 95%, as determined from pooled samples spiked with  $^3\text{H}$ -aldosterone and incubated for 60 min at 37°C before extraction. The assay was validated for sheep plasma by checking for parallelism using a serial dilution of pooled samples of sheep plasma and measuring aldosterone in sheep with access to water or with no access to water for 24 h (Figure 1).

*Data analysis*

The data were analysed using Genstat 8th Edition (Lawes Agricultural Trust, Rothamsted, UK). Pregnancy rates were analysed using the  $\chi^2$  test. Hormone concentrations, live

weight and feed intake were analysed by repeated measure analysis of variance, using conservative *F*-tests, with the AVP and aldosterone concentration log transformed and progesterone transformed to the fourth root in order to better approximate the assumption of homogeneity of variance. Lamb birth weight, crown-to-rump measurement and milk composition were analysed by analysis of variance.

**Results**

*Feed and water intake*

Pair-feeding was successfully achieved, as feed intake did not differ between the two dietary treatments (Table 2). The estimated average energy ME intake for ewes fed the control and high-salt diets was  $8.9 \pm 0.47$  MJ ME/head per day and  $9.4 \pm 0.49$  MJ ME/head per day, respectively. Sodium intake was  $1.9 \pm 0.096$  g Na/head per day and  $55.1 \pm 2.89$  g Na/head per day for the control and high-salt groups, respectively. The ewes fed the high-salt diet weighed 3 kg more ( $P < 0.05$ ) than their counterparts from the second month of pregnancy to parturition (Table 2). By 25 days from insemination, ewes fed the high-salt diet drank  $4.0 \pm 0.01$  l of water/head per day, while ewes fed the control diet drank  $2.43 \pm 0.145$  l of water/head per day.

*Reproductive performance*

Pregnancy rates were not significantly affected by feeding the high-salt diet throughout pregnancy (Table 3). Approximately 60% of the original group of ewes were single-bearing and 17% were twin-bearing. There were no significant differences in gestation length (overall mean 150.1 days), lamb weight (mean  $4.85 \pm 0.16$  kg) and crown-to-rump measurements (mean  $45.5 \pm 1.01$  cm) between treatment groups (Table 3).

Udder volume was not different between the two treatments (high-salt;  $1487 \pm 121$  ml, control;  $1670 \pm 232$  ml) and there was no difference between fat and protein content in the milk samples between the two treatments (fat: high-salt,  $8.1 \pm 0.38\%$ ; control,  $8.8 \pm 0.41\%$  and protein: high-salt,  $4.7 \pm 0.13\%$ ; control,  $4.8 \pm 0.22\%$ ).

*Endocrine and metabolic changes*

Plasma progesterone concentration was not different between the two treatments (Figure 2). Prior to pregnancy, progesterone was low ( $0.69 \pm 0.13$  ng/ml and  $0.47 \pm 0.08$  ng/ml for control and high-salt ewes, respectively), before rising slowly to 2.6 ng/ml by day 79. Progesterone concentration then increased more rapidly, reaching  $5.6 \pm 0.45$  ng/ml on day 115 and  $12.4 \pm 1.06$  ng/ml on day 140.

Prior to insemination, plasma concentrations of aldosterone were not different between ewes fed the high-salt diet ( $43 \pm 4.1$  pg/ml) and ewes fed the control diet ( $38 \pm 3.8$  pg/ml). Concentrations of aldosterone of ewes fed the high-salt diet were lower than that of ewes fed the control diet during pregnancy ( $27 \pm 2.7$  v.  $49 \pm 5.4$  pg/ml  $P < 0.05$ ; Figure 2). In the last month of pregnancy, plasma concentrations of aldosterone increased by approximately 20 pg/ml in ewes fed both diets but were still higher

**Table 2** Organic matter intake, sodium intake and live weight (mean  $\pm$  s.e.)

Day of pregnancy	Organic matter intake (kg OM/head per day)	Sodium intake (g Na/head per day)	Estimated ME intake (MJ ME/head per day)	Day of pregnancy	Live weight (kg)
Control					
0 to 44	0.93 $\pm$ 0.074	1.83 $\pm$ 0.146	8.9 $\pm$ 0.71	0	44.8 $\pm$ 0.90
45 to 100	0.98 $\pm$ 0.030	1.98 $\pm$ 0.061	9.4 $\pm$ 0.29	31	43.7 $\pm$ 1.04
				62	46.9 <sup>a</sup> $\pm$ 1.05
101 to term	0.88 $\pm$ 0.044	1.76 $\pm$ 0.082	8.4 $\pm$ 0.41	94	50.8 <sup>a</sup> $\pm$ 0.85
				126	53.4 <sup>a</sup> $\pm$ 1.08
				150	58.9 <sup>a</sup> $\pm$ 1.23
High salt					
0 to 44	0.98 $\pm$ 0.077	55.0 $\pm$ 4.28	9.4 $\pm$ 0.73	0	45.0 $\pm$ 0.87
45 to 100	1.03 $\pm$ 0.033	57.8 $\pm$ 1.84	9.9 $\pm$ 0.31	31	42.9 $\pm$ 0.67
				62	49.0 <sup>b</sup> $\pm$ 0.87
101 to term	0.94 $\pm$ 0.046	52.6 $\pm$ 2.55	9.0 $\pm$ 0.44	94	53.0 <sup>b</sup> $\pm$ 0.82
				126	56.7 <sup>b</sup> $\pm$ 1.13
				150	62.5 <sup>b</sup> $\pm$ 1.40

ME = metabolisable energy; OM = organic matter.

<sup>a,b</sup>Differences ( $P < 0.05$ ) between the two groups at a given stage of pregnancy are shown by different superscripts.

**Table 3** Reproductive performance of ewes fed a high-salt or control diet

	Control	High salt
Pregnancy rates (%)	82	73
Gestation length (days)	150 $\pm$ 0.45	150 $\pm$ 0.57
Singleton lamb characteristics		
Birth weight (kg)	4.81 $\pm$ 0.15	4.90 $\pm$ 0.17
Crown-to-rump length (cm)	44.9 $\pm$ 0.96	46.1 $\pm$ 1.06
Survival rate (%)	97	97

Pregnancy rate was measured on day 44 of gestation. Characteristic of the singleton lambs used in the study were measured at birth. Survival rate was measured at weaning.

( $P < 0.05$ ) in the ewes fed the high-salt diet than in ewes fed the control diet (74  $\pm$  7.9 v. 45  $\pm$  4.6 pg/ml).

Prior to pregnancy, plasma concentrations of AVP were high in both groups of ewes and then decreased during early and mid pregnancy by approximately 18 pg/ml for ewes fed the high-salt diet and 10 pg/ml for the ewes fed the control diet; however, AVP concentrations were not different between the two dietary treatments during pregnancy (Figure 2). AVP concentration remained relatively constant until the last month of pregnancy where there was an increase of approximately 5 pg/ml in both groups (to 7 pg/ml).

Glucose concentrations did not differ between ewes fed the high-salt diet or ewes fed the controls. At day 21 of gestation, glucose concentration averaged 4.2  $\pm$  0.14 mmol/l and at day 106 it averaged 3.2  $\pm$  0.10 mmol/l. There were no differences in haematocrit values between groups at any stage of pregnancy, ranging from 28% to 34%.


Overall, ewes that were fed the high-salt diet had a lower concentration of leptin ( $P < 0.05$ ) and insulin ( $P < 0.02$ ) than the control ewes (1.4  $\pm$  0.09 v. 1.5  $\pm$  0.12 ng/ml; and 7.2  $\pm$  0.55 v. 8.2  $\pm$  0.83 ng/ml). Insulin concentrations for control ewes were higher in the first half of gestation before

declining to the same concentration as the ewes fed high salt by day 115 of gestation (Figure 3). Leptin concentration on the other hand was similar between the two treatment groups during the first third of gestation (at 1.19 ng/ml) but at day 51 of gestation leptin concentrations of control ewes increased above that of ewes fed the high-salt diet (1.8  $\pm$  0.1 v. 1.7  $\pm$  0.1 ng/ml; Figure 3). Both leptin and insulin decreased between days 115 and 140 of gestation for both control and high-salt ewes (from 1.7  $\pm$  0.1 to 1.19  $\pm$  0.09 ng/ml; and from 8.5  $\pm$  0.9 to 5.6  $\pm$  0.5 ng/ml).

There was an interaction between time and dietary treatment on concentration of T<sub>3</sub> ( $P < 0.02$ ). In the first third of gestation (day 0 to 50), the plasma concentration of T<sub>3</sub> was lower in the ewes fed the high-salt diet than that in the ewes fed the control diet (1.2  $\pm$  0.18 v. 1.3  $\pm$  0.14 pmol/ml; Figure 3). During the second third of gestation (day 51 to 115), the concentration of T<sub>3</sub> was similar in both treatment groups (0.92  $\pm$  0.16 v. 0.87  $\pm$  0.1 pmol/ml). However, in the last third of gestation (day 116 to 140) the plasma concentration of T<sub>3</sub> was higher in the ewes fed the high-salt diet than that in the ewes fed the control diet (0.8  $\pm$  0.16 v. 0.6  $\pm$  0.06 pmol/ml). The concentration of T<sub>4</sub> in ewes fed the high-salt diet was lower than that of ewes fed the control diet up to day 115 (162  $\pm$  8.6 v. 212  $\pm$  13 ng/ml;  $P < 0.001$ ) but, by day 140, the T<sub>4</sub> concentrations were similar for both groups (166 ng/ml).


## Discussion

Feeding Merino ewes a high-salt (13% NaCl) diet from the time of conception until parturition did not compromise reproductive efficiency. The ewes fed the high-salt diet produced lambs with normal crown-to-rump length, birth weight and milk with similar composition, indicating they were able to adequately regulate salt and water balance and adapt metabolically to the challenges associated with high salt ingestion.



**Figure 2** Plasma concentrations of progesterone, aldosterone and arginine vasopressin (AVP) during pregnancy in ewes fed a control or high-salt diet. Asterisks denote differences ( $P < 0.05$ ) between groups at a given stage of pregnancy. Open circles: control group; closed circles: ewes fed high-salt diet.

The high-salt ewes were heavier, probably as a consequence of water retention as feed intake of the ewes did not differ between groups (they were pair-fed). An increase in water retention could be associated with an increase of 10% to 15% (3 to 5 kg) in total body water normally associated with osmotic regulation in sheep consuming high-salt feed (Warren *et al.*, 1995) or an increase in muscle deposition associated with a higher flow of protein to the small intestine in sheep fed high salt (Hemsley *et al.*, 1975; Masters *et al.*, 2005), as changes in body water, protein and fat do occur during pregnancy (Masters and Mata, 1996). It is possible that the live weight of the high-salt ewes could



**Figure 3** Plasma concentrations of insulin, leptin, T3 and T4 during pregnancy in ewes fed a control or high-salt diet. Asterisks denote differences ( $P < 0.05$ ) between groups at a given stage of pregnancy. Open circles: control group; closed circles: ewes fed high-salt diet.

reflect an increase in the weight of the fetuses and placental components, but it is unlikely as the mean birth weights of the lambs from different dietary treatments did not differ.

Ingestion of high salt had a specific effect on both insulin and leptin concentrations that could not be attributed to

differences in feed intake or body weight. A strong direct effect of salt (dietary intake or infused) on insulin concentration has been reported in non-pregnant humans (Goodfriend *et al.*, 1991) but the mechanism is not known. An earlier study with wethers found that plasma concentrations of insulin concentration decreased in wethers fed 20% NaCl for 14 days (Blache *et al.*, 2007) and our results extend the work by Blache *et al.* (2007) by showing a persistent (i.e. longer term) effect of salt ingestion on insulin secretion in sheep. In the study of Blache *et al.* (2007), the drop in insulin concentration with high salt ingestion with wethers was associated with lower glucose concentration, but this was not observed in the current study. However, this may have been due to a different physiological regulation during pregnancy whereby a higher maternal glucose concentration was maintained in order to meet the demands of the foetus. It would appear that the ewes were consuming sufficient energy to meet the energy requirements for maternal tissue and foetal growth (McDonald *et al.*, 1995), and thus were able to maintain glucose concentrations. Therefore, the lower insulin concentrations in pregnant ewes that consumed a high-salt diet did not lead to an increase in maternal glucose concentration.

In contrast to observations obtained with wethers (Blache *et al.*, 2007), high salt consumption decreased leptin concentrations in pregnant ewes independent of both the level of food intake (because they were pair fed) or body weight (similar between the two groups). Our study illustrates that another mechanism must have been operating to reduce leptin concentrations rather than the classical regulation of plasma leptin by voluntary feed intake (e.g. Woods *et al.*, 2004) or fat deposition (e.g. Chilliard *et al.*, 2005). In humans, concentrations of circulating progesterone have been correlated to plasma leptin during the menstrual cycle (Hardie *et al.*, 1997) but, in the current study, progesterone concentrations were not different between the ewes fed the high-salt diet and the ewes fed the control diet. Another factor that decreases leptin secretion is an increase in energy expenditure (Zimmermann-Belsing *et al.*, 2003). However, in the present study there was not a large difference in plasma  $T_3$  concentration between ewes fed high salt and control animals, suggesting that energy expenditure was similar between the two groups. However, plasma  $T_4$  concentrations were lower in ewes fed high salt, possibly reflecting a higher conversion of  $T_4$  to  $T_3$ , which could reflect a higher metabolic rate in the ewes fed high salt.

Despite the low levels of leptin and insulin, udder volume and milk composition in week 1 of lactation were not affected. This result is consistent with Chagas *et al.* (2006) where dairy cows during early lactation have extremely low insulin and leptin concentrations when milk production is at its peak. Furthermore, a study by Abu-Zanat and Tabbaa (2006) found no significant effect on milk production when pregnant Awassi ewes were fed a diet containing either 40% or 60% of saltbush, and Olsson *et al.* (2006) found that milk secretory mechanisms in lactating goats were unaffected by plasma hyperosmolality.

High salt consumption during pregnancy required a lowering of aldosterone concentration, but this did not involve a concomitant effect on AVP. Plasma concentrations of AVP did not increase despite high salt ingestion and the water-retaining requirements of pregnancy possibly because the ewes had free access to fresh water so the ewes were able to increase their water consumption above normal levels (Hamilton and Webster, 1987; Meintjes and Olivier, 1992). Water consumption increased nearly two-fold in early pregnancy in ewes fed the high-salt diet. We were not able to quantify water consumption during the rest of pregnancy because of technical problems. The high intake of water could also explain the limited response in aldosterone because of the dilution of sodium in the rumen.

Further to the consequences of increased water intake, the lack of a large change in AVP concentration during pregnancy is likely to be also due to changes in osmolality during pregnancy and continual adjustments in the volume-sensing AVP release mechanisms (Lindheimer *et al.*, 1989). AVP is not suppressed at the usual levels of body tonicity (at least in humans; Lindheimer and Davison, 1995), as part of the normal water-retaining mechanisms of pregnancy.

In terms of integrating saltbush into a productive system, the data show that high salt intake does not decrease the reproductive capacity of the ewe. However, this may be so only if the ewes have free access to water. In addition, there is a need to investigate the long-term effect on the offspring. The manipulation of high-salt diets in rats and cattle during pregnancy or postnatally has shown behavioural and/or physiological changes in the offspring (da Silva *et al.*, 2003; Mohamed and Phillips, 2003; Curtis *et al.*, 2004). The occurrence and potential value of such foetal programming needs to be determined with sheep.

### Acknowledgements

SND was supported by an Australian Postgraduate Award and CSIRO Livestock Industries scholarship. We thank Dr Julian Taylor at Biometrics SA, University of Adelaide, Waite, South Australia for the statistical analysis and Dr Roberta Bencini (University of Western Australia) for milk analysis.

### References

- Abu-Zanat MMW and Tabbaa MJ 2006. Effect of feeding *Atriplex* browse to lactating ewes on milk yield and growth rate of their lambs. *Small Ruminant Research* 64, 152–161.
- Arieli ML, Naim E, Benjamin RW and Pasternak D 1989. The effect of feeding saltbush and sodium chloride on energy metabolism in sheep. *Animal Production* 49, 451–457.
- Bell RJ, Laurence BM, Meehan PJ, Congiu M, Scoggins BA and Wintour EM 1986. Regulation and function of arginine vasopressin in pregnant sheep. *American Journal of Physiology – Renal Physiology* 250, F777–F780.
- Bencini R 1999. Development of specialty sheep milk dairy products – increasing the market scope. A report for The Rural Industries Research and Development Corporation. RIRDC publ. No 99/69.
- Bencini R and Purvis IW 1990. The yield and composition of milk from Merino sheep. *Wool Technology and Sheep Breeding* 38, 71–73.
- Blache D, Tellam RL, Chagas LM, Blackberry MA, Vercoe PE and Martin GB 2000. Level of nutrition affects leptin concentrations in plasma and cerebrospinal fluid in sheep. *Journal of Endocrinology* 165, 625–637.

- Blache D, Grandison MJ, Masters DG, Dynes RA, Blackberry MA and Martin GA 2007. Relationships between metabolic endocrine systems and voluntary feed intake in Merino sheep fed a high salt diet. *Australian Journal of Experimental Agriculture* 47, 544–550.
- Chagas LM, Rhodes FM, Blache D, Gore PJS, Macdonald KA and Verkerk GA 2006. Precalving effects on metabolic responses and postpartum anestrus in grazing primiparous dairy cows. *Journal of Dairy Science* 89, 1981–1989.
- Chilliard Y, Delavaud C and Bonnet M 2005. Leptin expression in ruminants: nutritional and physiological regulations in relation with energy metabolism. *Domestic Animal Endocrinology* 29, 3–22.
- Cowley Jr AW, Skelton MM and Merrill DC 1986. Osmoregulation during high salt intake: relative importance of drinking and vasopressin secretion. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 251, R878–R886.
- Curtis KS, Krause EG, Wong DL and Contreras RJ 2004. Gestational and early postnatal dietary NaCl levels affect NaCl intake, but not stimulated water intake, by adult rats. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 286, R1043–R1050.
- da Silva AA, de Noronha IL, de Oliveira IB, Malheiros DM and Heimann JC 2003. Renin angiotensin system function and blood pressure in adult rats after perinatal salt overload. *Nutrition, Metabolism and Cardiovascular Disease* 13, 133–139.
- Dahlquist RL and Knoll JW 1978. Inductively coupled plasma-atomic emission spectrometry: analysis of biological material and soils for major, trace and ultra-trace elements. *Applied Spectroscopy* 32, 1–29.
- Davison JM and Lindheimer MD 1989. Volume homeostasis and osmoregulation in human pregnancy. *Baillieres Clinical Endocrinology and Metabolism* 3, 451–472.
- Dawson A, Deeming DC, Dick AC and Sharp PJ 1996. Plasma thyroxine concentrations in farmed ostriches in relation to age, body weight, and growth hormone. *General and Comparative Endocrinology* 103, 308–315.
- Gales NJ, Williamson P, Higgins LV, Blackberry MA and James I 1997. Evidence for a prolonged postimplantation period in the Australian sea lion (*Neophoca cinerea*). *Journal of Reproduction and Fertility* 111, 159–163.
- Ghassemi F, Jakeman AJ and Nix HA 1995. *Salinisation of Land and Water Resources*. CAB International Publishing, Wallingford, UK.
- Goodfriend TL, Ball DL, Weinberger MH, Moore TJ, Weder AB and Egan BM 1991. Salt loads raise plasma fatty acids and lower insulin. *Hypertension* 17, 958–964.
- Greenwood FC and Hunter WH 1963. The preparation of <sup>131</sup>I-labelled human growth hormone of high specific radioactivity. *Biochemical Journal* 89, 114–123.
- Hamilton JA and Webster MED 1987. Food intake, water intake, urine output, growth rate and wool growth of lambs accustomed to high or low intake of sodium chloride. *Australian Journal of Agricultural Research* 38, 187–194.
- Hardie L, Trayhurn P, Abramovich D and Fowler P 1997. Circulating leptin in women: a longitudinal study in the menstrual cycle and during pregnancy. *Clinical Endocrinology* 47, 101–106.
- Hemsley JA, Hogan JP and Weston RH 1975. Effect of high intakes of sodium chloride on the utilisation of a protein concentrate by sheep. II. Digestion and absorption of organic matter and electrolytes. *Australian Journal of Agricultural Research* 26, 715–727.
- James VHT and Wilson GA 1976. Determination of aldosterone in biological fluids. In *Assay of Drugs and Other Trace Compounds in Biological Fluids* (ed. E Reid), pp. 149–158. Elsevier Publishing, Amsterdam.
- Lindheimer MD and Davison JM 1995. Osmoregulation, the secretion of arginine vasopressin and its metabolism during pregnancy. *European Journal of Endocrinology* 132, 133–143.
- Lindheimer MD, Barron WM and Davison JM 1989. Osmoregulation of thirst and vasopressin release in pregnancy. *American Journal of Physiology – Renal Physiology* 257, F159–F169.
- Masters DG and Mata G 1996. Responses to feeding canola meal or lupin seed to pregnant, lactating, and dry ewes. *Australian Journal of Agricultural Research* 47, 1291–1303.
- Masters DG, Rintoul AJ, Dynes RA, Pearce KL and Norman HC 2005. Feed intake and production in sheep fed diets high in sodium and potassium. *Australian Journal of Agricultural Research* 56, 427–434.
- Masters DG, Edwards N, Sillence M, Avery A, Revell D, Friend M, Sanford P, Saul G, Beverly C and Young J 2006. The role of livestock in the management of dryland salinity. *Australian Journal of Experimental Agriculture* 46, 733–741.
- Masters DG, Benes SE and Norman HC 2007. Biosaline agriculture for forage and livestock production. *Agriculture, Ecosystems and Environment* 119, 234–248.
- McDonald P, Edwards RA, Greenhalgh JFD and Morgan CA 1995. *Animal Nutrition*. Addison Wesley Longman, Singapore.
- Meintjes RA and Olivier R 1992. The effects of salt loading via two different routes on feed intake, body water turnover rate and electrolyte excretion in sheep. *Onderstepoort Journal of Veterinary Research* 59, 91–96.
- Miller DW, Blache D and Martin GB 1995. The role of intracerebral insulin in the effect of nutrition on gonadotrophin secretion in mature male sheep. *Journal of Endocrinology* 147, 321–329.
- Mohamed MO and Phillips CJC 2003. The effect of increasing salt intake of pregnant dairy cows on the salt appetite and growth of their calves. *Animal Science* 77, 181–185.
- Norman HC, Dynes RA and Masters DG 2002. Nutritive value of plants growing on saline land. In *Productive use and rehabilitation of saline land*, 8th National Conference, pp. 59–69. Promaco Conventions Pty Ltd, Fremantle, Australia.
- Olsson K, Mengistu U, Stein J, Bekele T and Bruckmaier RM 2006. Hypertonic NaCl infusions affect milk composition in goats. *Journal of Dairy Research* 73, 306–311.
- Rice GE 1982. Plasma arginine vasotocin concentration in the lizard *Varanus gouldii* (Gray) following water loading, salt loading, and dehydration. *General and Comparative Endocrinology* 47, 1–6.
- Thomas DT, 2006. Using stable carbon isotopes to measure diet selection in sheep grazing saltland pastures. 6th Biennial Conference of the Australian Society of Animal Production in Australia, Short communication number 61.
- Tindal JS, Knaggs GS, Hart IC and Blake LA 1978. Release of growth hormone in lactating and non-lactating goats in relation to behaviour, stages of sleep, electroencephalograms, environmental stimuli and levels of prolactin, insulin, glucose and free fatty acids in the circulation. *Journal of Endocrinology* 76, 333–346.
- Warren BE, Casson T and Barrett-Lennard E 1995. Value of saltbush questioned. *Western Australia Journal of Agriculture* 36, 24–27.
- Woods SC, Benoit SC, Clegg DJ and Seeley RJ 2004. Regulation of energy homeostasis by peripheral signals. *Best Practice and Research Clinical Endocrinology and Metabolism* 18, 497–515.
- Zhang S, Blache D, Blackberry MA and Martin GB 2005. Body reserves affect the reproductive endocrine responses to an acute change in nutrition in mature male sheep. *Animal Reproduction Science* 88, 257–269.
- Zimmermann-Belsing T, Brabant G, Holst JJ and Feldt-Rasmussen U 2003. Circulating leptin and thyroid dysfunction. *European Journal of Endocrinology* 149, 257–271.