

RESEARCH

Open Access

Some fixed and coincidence point theorems for expansive maps in cone metric spaces

Wasfi Shatanawi and Fadi Awawdeh*

* Correspondence: awawdeh@hu.edu.jo
Department of Mathematics,
Hashemite University, Zarqa 13115,
Jordan**Abstract**

In this article, we establish some common fixed and common coincidence point theorems for expansive type mappings in the setting of cone metric spaces. Our results extend some known results in metric spaces to cone metric spaces. Also, we introduce some examples the support the validity of our results.

Mathematics Subject Classification: 54H25; 47H10; 54E50.

Keywords: common fixed point, cone metric space, coincidence fixed point

1. Introduction

Huang and Zhang [1] introduced the notion of cone metric spaces as a generalization of metric spaces. They replacing the set of real numbers by an ordered Banach space. Huang and Zhang [1] presented the notion of convergence of sequences in cone metric spaces and proved some fixed point theorems. Then after, many authors established many fixed point theorems in cone metric spaces. For some fixed point theorems in cone metric spaces we refer the reader to [2-30].

In the present article, E stands for a real Banach space.

Definition 1.1. Let P be a subset of E with $\text{Int}(P) \neq \emptyset$. Then P is called a cone if the following conditions are satisfied:

- (1) P is closed and $P \neq \{\theta\}$.
- (2) $a, b \in \mathbf{R}^+$, $x, y \in P$ implies $ax + by \in P$.
- (3) $x \in P \cap -P$ implies $x = \theta$.

For a cone P , define a partial ordering \preceq with respect to P by $x \preceq y$ if and only if $y - x \in P$. We shall write $x \prec y$ to indicate that $x \preceq y$ but $x \neq y$, while $x \ll y$ will stand for $y - x \in \text{Int} P$. It can be easily shown that $\lambda \text{Int}(P) \subseteq \text{Int}(P)$ for all positive scalar λ .

Definition 1.2. [1] Let X be a nonempty set. Suppose the mapping $d : X \times X \rightarrow E$ satisfies

- (1) $\theta \prec d(x, y)$ for all $x, y \in X$ and $d(x, y) = \theta$ if and only if $x = y$.
- (2) $d(x, y) = d(y, x)$ for all $x, y \in X$.
- (3) $d(x, y) \preceq d(x, z) + d(y, z)$ for all $x, y \in X$.

Then d is called a cone metric on X , and (X, d) is called a cone metric space.

Definition 1.3. [1] Let (X, d) be a cone metric space. Let (x_n) be a sequence in X and $x \in X$. If for every $c \in E$ with $\theta \ll c$, there is an $N \in \mathbf{N}$ such that $d(x_n, x) \ll c$ for all $n \geq N$, then (x_n) is said to be convergent and (x_n) converges to x and x is the limit of (x_n) . We denote this by $\lim_{n \rightarrow +\infty} x_n = x$ or $x_n \rightarrow x$ as $n \rightarrow +\infty$. If for every $c \in E$ with $\theta \ll c$ there is an $N \in \mathbf{N}$ such that $d(x_n, x_m) \ll c$ for all $n, m \geq N$, then (x_n) is called a Cauchy sequence in X . The space (X, d) is called a complete cone metric space if every Cauchy sequence is convergent.

The cone P in a real Banach space E is called normal if there is a number $\lambda > 0$ such that for all $x, y \in E$,

$$\theta \preceq x \preceq y \text{ implies } \|x\| \leq \lambda \|y\|.$$

Iker Ćahin and Mustafa Telci [30] studied a theorem on common fixed points of expansive type mappings in cone metric spaces.

Definition 1.4. [30] Let E and F be real Banach spaces and P and Q be cones in E and F , respectively. Let (X, d) and (Y, ρ) be cone metric spaces, where $d : X \times X \rightarrow E$ and $\rho : Y \times Y \rightarrow F$. A function $f : X \rightarrow Y$ is said to be continuous at $x_0 \in X$, if for every $c \in F$ with $0 \ll c$, there exists $b \in E$ with $0 \ll b$ such that $\rho(fx, fx_0) \ll c$ whenever $x \in X$ and $d(x, x_0) \ll b$.

Lemma 1.1. [30] Let (X, d) and (Y, ρ) be cone metric spaces. A function $f : X \rightarrow Y$ is continuous at a point $x_0 \in X$ if and only if whenever a sequence (x_n) in X converges to x_0 , the sequence (fx_n) converges to fx_0 .

Theorem 1.1. [30] Let (X, d) be a complete cone metric space and P be a cone. Let f and g be surjective self-mappings of X satisfying the following inequalities

$$\begin{aligned} d(gfx, fx) &\succeq ad(fx, x), \\ d(fgx, gx) &\succeq bd(gx, x) \end{aligned}$$

for all $x \in X$, where $a, b > 1$. If either f or g is continuous, then f and g have a common fixed point.

The aim of this article is to study new theorems of common fixed and coincidence point for expansive mappings in cone metric spaces under a set of conditions. Our results generalize several well known comparable results in the literature. Also, we introduced some examples to support the validity of our results.

2. Main results

We start with the following theorem

Theorem 2.1. Let (X, d) be a cone metric space with a solid cone P . Let $T, f : X \rightarrow X$ be mappings satisfying:

$$d(Tx, Ty) \succeq ad(fx, fy) + bd(fx, Tx) + cd(fy, Ty) \tag{1}$$

for all $x, y \in X$ where $a, b, c \geq 0$ with $a + b + c > 1$. Suppose the following hypotheses:

- (1) $b < 1$ or $c < 1$.
- (2) $fX \subseteq TX$.
- (3) TX is a complete subspace of X .

Then T and f have a coincidence point.

Proof. Let $x_0 \in X$. Since $fX \subseteq TX$, we choose $x_1 \in X$ such that $Tx_1 = fx_0$. Again, we can choose $x_2 \in X$ such that $Tx_2 = fx_1$. Continuing in the same way, we construct a sequence (x_n) in X such that $Tx_{n+1} = fx_n$ for all $n \in \mathbf{N} \cup \{0\}$.

If $fx_{m-1} = fx_m$ for some $m \in \mathbf{N}$, then $Tx_m = fx_m$. Thus x_m is a coincidence point of T and f .

Now, assume that $x_{n-1} \neq x_n$ for all $n \in \mathbf{N}$.

Case (1): Suppose $b < 1$.

By (1), we have

$$\begin{aligned} d(fx_{n-1}, fx_n) &= d(Tx_n, Tx_{n+1}) \\ &\succcurlyeq ad(fx_n, fx_{n+1}) + bd(fx_n, Tx_n) + cd(fx_{n+1}, Tx_{n+1}) \\ &= ad(fx_n, fx_{n+1}) + bd(fx_n, fx_{n-1}) + cd(fx_{n+1}, fx_n). \end{aligned}$$

Thus, we have

$$(1 - b)d(fx_{n-1}, fx_n) \succcurlyeq (a + c)d(fx_{n+1}, fx_n).$$

Hence

$$d(fx_{n+1}, fx_n) \preccurlyeq \frac{1 - b}{a + c}d(fx_{n-1}, fx_n). \tag{2}$$

Case (2): Suppose $c < 1$.

By (1), we have

$$\begin{aligned} d(fx_n, fx_{n-1}) &= d(Tx_{n+1}, Tx_n) \\ &\succcurlyeq ad(fx_{n+1}, fx_n) + bd(fx_{n+1}, Tx_{n+1}) + cd(fx_n, Tx_n) \\ &= ad(fx_n, fx_{n+1}) + bd(fx_{n+1}, fx_n) + cd(fx_n, fx_{n-1}). \end{aligned}$$

Thus, we have

$$(1 - c)d(fx_{n-1}, fx_n) \succcurlyeq (a + b)d(fx_{n+1}, fx_n).$$

Hence

$$d(fx_{n+1}, fx_n) \preccurlyeq \frac{1 - c}{a + b}d(fx_{n-1}, fx_n). \tag{3}$$

In Case (1), we let

$$\lambda = \frac{1 - b}{a + c}$$

and in Case (2), we let

$$\lambda = \frac{1 - c}{a + b}.$$

Thus in both cases, we have $\lambda < 1$ and

$$d(fx_{n+1}, fx_n) \preccurlyeq \lambda d(fx_{n-1}, fx_n). \tag{4}$$

By (4), we have

$$\begin{aligned} d(fx_{n+1}, fx_n) &\preceq \lambda d(fx_{n-1}, fx_n) \\ &\preceq \lambda^2 d(fx_{n-2}, fx_{n-1}) \\ &\vdots \\ &\preceq \lambda^n d(fx_0, fx_1). \end{aligned}$$

So for $m > n$, we have

$$\begin{aligned} d(fx_n, fx_m) &\preceq d(fx_n, fx_{n+1}) + d(fx_{n+1}, fx_{n+2}) + \dots + d(fx_{m-1}, fx_m) \\ &\preceq (\lambda^n + \lambda^{n+1} + \dots + \lambda^{m-1})d(fx_0, fx_1) \\ &\preceq \lambda^n \sum_{i=0}^{m-n} \lambda^i d(fx_0, fx_1) \\ &= \frac{\lambda^n}{1-\lambda} d(fx_0, fx_1). \end{aligned}$$

Let $\theta \ll c$ be given, choose $\delta > 0$ such that $c + N_\delta(0) \subseteq P$, where

$$N_\delta(0) = \{y \in E : \|y\| < \delta\}.$$

Also, choose a natural number N_1 such that

$$\frac{\lambda^n}{1-\lambda} d(fx_0, fx_1) \in N_\delta(0),$$

for $m \geq N_1$. Then

$$\frac{\lambda^n}{1-\lambda} d(fx_0, fx_1) \ll c,$$

for all $m \geq N_1$. Thus,

$$d(fx_n, fx_m) \preceq \frac{\lambda^n}{1-\lambda} d(fx_0, fx_1) \ll c,$$

for all $m > n$. Therefore (Tx_n) is a Cauchy sequence in (TX, d) . Since (TX, d) is a complete cone metric space, there is $u \in X$ such that (Tx_n) converges to Tu as $n \rightarrow +\infty$. Hence fx_n converges to Tu as $n \rightarrow +\infty$. Since $a + b + c > 1$, we have a, b and c are not all 0. So we have the following cases.

Case 1: If $a \neq 0$, then

$$d(Tx_n, Tu) \succcurlyeq ad(fx_n, fu) + bd(fx_n, Tx_n) + cd(fu, Tu) \succcurlyeq ad(fx_n, fu).$$

Hence

$$d(fx_n, fu) \preceq \frac{1}{a} d(Tx_n, Tu).$$

Let $\theta \ll c$ be given, choose $\delta > 0$ such that $c + N_\delta(0) \subseteq P$, where

$$N_\delta(0) = \{y \in E : \|y\| < \delta\}.$$

Since $\frac{1}{a}d(Tx_n, Tu) \rightarrow \theta$. We choose a natural number $n_0 \in \mathbf{N}$ such that

$$\frac{1}{a}d(Tx_n, Tu) \in N_\delta(0),$$

for $n \geq n_0$. Then

$$\frac{1}{a}d(Tx_n, Tu) \ll c,$$

for all $n \geq n_0$. Thus,

$$d(fx_n, fu) \preceq \frac{1}{a}d(Tx_n, Tu) \ll c,$$

for all $n \geq n_0$. Thus $fx_n \rightarrow fu$ as $n \rightarrow +\infty$. By uniqueness of limit, we have $Tu = fu$. Therefore T and f have a coincidence point.

Case 2: If $b \neq 0$, then

$$d(Tu, Tx_n) \succcurlyeq ad(fx_n, fu) + bd(fu, Tu) + cd(fx_n, Tx_n) \succcurlyeq bd(fu, Tu).$$

Hence

$$d(fu, Tu) \preceq \frac{1}{b}d(Tx_n, Tu).$$

As similar proof of Case (1), we can show that $fu = Tu$. Thus f and T have a coincidence point.

Case 3: If $c \neq 0$, then

$$d(Tx_n, Tu) \succcurlyeq ad(fx_n, fu) + bd(fx_n, Tx_n) + cd(Tu, fu) \succcurlyeq cd(fu, Tu).$$

Hence

$$d(fu, Tu) \preceq \frac{1}{c}d(Tx_n, Tu).$$

As similar proof of Case (1), we can show that $fu = Tu$. Thus f and T have a coincidence point.

Corollary 2.1. Let (X, d) be a cone metric space with a solid cone P . Let $T, f: X \rightarrow X$ be mappings satisfying:

$$d(Tx, Ty) \succcurlyeq ad(fx, fy) + bd(fx, Tx)$$

for all $x, y \in X$ where $a, b \geq 0$ with $a + b > 1$ and $b < 1$. Suppose the following hypotheses:

- (1) $fX \subseteq TX$.
- (2) TX is a complete subspace of X .

Then T and f have a coincidence point.

Corollary 2.2. Let (X, d) be a complete cone metric space with a solid cone P . Let $T, f: X \rightarrow X$ be mappings satisfying:

$$d(Tx, Ty) \succcurlyeq ad(fx, fy)$$

for all $x, y \in X$ where $a > 1$. Suppose the following hypotheses:

- (1) $fX \subseteq TX$.
- (2) TX is a complete subspace of X .

Then T and f have a coincidence point.

Corollary 2.3. Let (X, d) be a complete cone metric space with a solid cone P . Let $T : X \rightarrow X$ be a surjective mapping satisfying:

$$d(Tx, Ty) \succcurlyeq ad(x, y) + bd(x, Tx) + cd(y, Ty)$$

for all $x, y \in X$ where $a, b, c \geq 0$ with $a + b + c > 1$. Suppose $b < 1$ or $c < 1$. Then T has a fixed point.

Proof. Follows from Theorem 2.1 by taking $f = I$, the identity map.

Corollary 2.4. Let (X, d) be a complete cone metric space with a solid cone P . Let $T : X \rightarrow X$ be a surjective mapping satisfying:

$$d(Tx, Ty) \succcurlyeq ad(x, y)$$

for all $x, y \in X$ where with $a > 1$. Then T has a fixed point.

Putting $E = \mathbf{R}$, $P = \{x \in \mathbf{R} : x \geq 0\}$ and $d : X \times X \rightarrow \mathbf{R}$ in Corollaries 2.1 and 2.2, we have the following results:

Corollary 2.5. Let (X, d) be a complete metric space. Let $T : X \rightarrow X$ be a surjective mapping satisfying:

$$d(Tx, Ty) \geq ad(x, y) + bd(x, Tx)$$

for all $x, y \in X$ where $a, b \geq 0$ with $a + b > 1$ and $b < 1$. Then T has a fixed point.

Corollary 2.6. Let (X, d) be a complete metric space. Let $T : X \rightarrow X$ be a surjective mapping satisfying:

$$d(Tx, Ty) \geq ad(x, y) + bd(y, Ty)$$

for all $x, y \in X$ where $a, b \geq 0$ with $a + b > 1$ and $b < 1$. Then T has a fixed point.

Now, we present a fixed point theorem for two maps.

Theorem 2.2. Let $T, S : X \rightarrow X$ be two surjective mappings of a complete cone metric space (X, d) with a solid cone P . Suppose that T and S satisfying the following inequalities

$$d(T(Sx), Sx) + kd(T(Sx), x) \succcurlyeq ad(Sx, x) \tag{5}$$

and

$$d(S(Tx), Tx) + kd(S(Tx), x) \succcurlyeq bd(Tx, x) \tag{6}$$

for all $x \in X$ and some nonnegative real numbers a, b and k with $a > 1 + 2k$ and $b > 1 + 2k$. If T or S is continuous, then T and S have a common fixed point

Proof. Let x_0 be an arbitrary point in X . Since T is surjective, there exists $x_1 \in X$ such that $x_0 = Tx_1$. Also, since S is surjective, there exists $x_2 \in X$ such that $x_2 = Sx_1$. Continuing this process, we construct a sequence (x_n) in X such that $x_{2n} = Tx_{2n+1}$ and $x_{2n+1} = Sx_{2n+2}$ for all $n \in \mathbf{N} \cup \{0\}$. Now, for $n \in \mathbf{N} \cup \{0\}$, we have

$$d(T(Sx_{2n+2}), Sx_{2n+2}) + kd(T(Sx_{2n+2}), x_{2n+2}) \succcurlyeq ad(Sx_{2n+2}, x_{2n+2}).$$

Thus, we have

$$d(x_{2n}, x_{2n+1}) + kd(x_{2n}, x_{2n+2}) \succcurlyeq ad(x_{2n+1}, x_{2n+2}),$$

which implies that

$$d(x_{2n}, x_{2n+1}) + kd(x_{2n}, x_{2n+1}) + kd(x_{2n+1}, x_{2n+2}) \succcurlyeq ad(x_{2n+1}, x_{2n+2}).$$

Hence

$$d(x_{2n+1}, x_{2n+2}) \preccurlyeq \frac{1+k}{a-k}d(x_{2n}, x_{2n+1}) \tag{7}$$

On other hand, we have

$$d(S(Tx_{2n+1}), Tx_{2n+1}) + kd(S(Tx_{2n+1}), x_{2n+1}) \succcurlyeq bd(Tx_{2n+1}, x_{2n+1}).$$

Thus, we have

$$d(x_{2n-1}, x_{2n}) + kd(x_{2n-1}, x_{2n+1}) \succcurlyeq bd(x_{2n}, x_{2n+1}).$$

Since $d(x_{2n-1}, x_{2n}) + d(x_{2n}, x_{2n+1}) \succcurlyeq d(x_{2n-1}, x_{2n+1})$, we have

$$d(x_{2n-1}, x_{2n}) + kd(x_{2n-1}, x_{2n}) + kd(x_{2n}, x_{2n+1}) \succcurlyeq bd(x_{2n}, x_{2n+1}).$$

Hence

$$d(x_{2n}, x_{2n+1}) \preccurlyeq \frac{1+k}{b-k}d(x_{2n-1}, x_{2n}) \tag{8}$$

Let

$$\lambda = \max \left\{ \frac{1+k}{a-k}, \frac{1+k}{b-k} \right\}.$$

Then by combining (7) and (8), we have

$$d(x_n, x_{n+1}) \leq \lambda d(x_{n-1}, x_n) \quad \forall n \in \mathbf{N} \cup \{0\}. \tag{9}$$

Repeating (9) n -times, we get

$$d(x_n, x_{n+1}) \preccurlyeq \lambda^n d(x_0, x_1).$$

Thus, for $m > n$, we have

$$\begin{aligned} d(x_n, x_m) &\preccurlyeq d(x_n, x_{n+1}) \cdots + d(x_{m-1}, x_m) \\ &\preccurlyeq (\lambda^n + \cdots + \lambda^{m-1})d(x_0, x_1) \\ &\preccurlyeq \frac{\lambda^n}{1-\lambda}d(x_0, x_1). \end{aligned}$$

As similar arguments to proof of Theorem 2.1, we can show that (x_n) is a Cauchy sequence in the complete cone metric space (X, d) . Then there exists $v \in X$ such that $x_n \rightarrow v$ as $n \rightarrow +\infty$. Therefore $x_{2n+1} \rightarrow v$ and $x_{2n+2} \rightarrow v$ as $n \rightarrow +\infty$. Without loss of generality, we may assume that T is continuous, then $Tx_{2n+1} \rightarrow Tv$ as $n \rightarrow +\infty$. But $Tx_{2n+1} = x_{2n} \rightarrow v$ as $n \rightarrow +\infty$. Thus, we have $Tv = v$. Since S is surjective, there exists $w \in X$ such that $Sw = v$. Now,

$$d(T(Sw), Sw) + kd(T(Sw), w) \succcurlyeq ad(Sw, w),$$

implies that $kd(v,w) \succcurlyeq ad(v,w)$. Thus

$$d(v,w) \preccurlyeq \frac{k}{a}d(v,w).$$

Since $a > k$, we conclude that $d(v,w) = \theta$. So $v = w$. Hence $Tv = Sv = v$. Therefore v is a common fixed point of T and S .

By taking $b = a$ in Theorem 2.2, we have the following result.

Corollary 2.7. *Let $T, S : X \rightarrow X$ be two surjective mappings of a complete cone metric space (X, d) with a solid cone P . Suppose that T and S satisfying the following inequalities*

$$d(T(Sx), Sx) + kd(T(Sx), x) \succcurlyeq ad(Sx, x) \tag{10}$$

and

$$d(S(Tx), Tx) + kd(S(Tx), x) \succcurlyeq ad(Tx, x) \tag{11}$$

for all $x \in X$ and some nonnegative real numbers a and k with $a > 1 + 2k$. If T or S is continuous, then T and f have a common fixed point

By taking $S = T$ in Corollary 2.7, we have the following corollary.

Corollary 2.8. *Let $T : X \rightarrow X$ be a surjective mapping of a complete cone metric space (X, d) with a solid cone P . Suppose that T satisfying*

$$d(T(Tx), Tx) + kd(T(Tx), x) \succcurlyeq ad(Tx, x) \tag{12}$$

for all $x \in X$ and some nonnegative real number a and k with $a > 1 + 2k$. If T is continuous, then T has a fixed point.

Now, we present some examples to illustrate the useability of our results.

Example 2.1. (The case of normal cone) *Let $X = [0, +\infty)$, $E = \mathbf{R}^2$. Let $P = \{(a, b) : a \geq 0, b \geq 0\}$ be the cone with $d(x, y) = (|x - y|, |x - y|)$. Then (X, d) is a complete cone metric space. Define $T : X \rightarrow X$ by $Tx = 2x$. Then T has a fixed point.*

Proof. Note that

$$d(T(Tx), Tx) + d(T(Tx), x) \geq 4d(Tx, x)$$

for all $x \in X$. Thus T satisfies all the hypotheses of Corollary 2.8 and hence T has a fixed point. Here 0 is the fixed point of T .

Example 2.2. (The case of non-normal cone) *Let $X = [0, 1]$, $E = C_{\mathbf{R}}^1([0, 1])$. Let $P = \{\varphi \in E : \varphi(t) \geq 0, t \in [0, 1]\}$. Define the mapping $d : X \times X \rightarrow E$ by*

$$d(x, y)(t) := |x - y| \phi(t),$$

where $\varphi \in P$ is a fixed function, for example $\varphi(t) = e^t$. Define $T, f : X \rightarrow X$ by $fx = \frac{1}{16}x$ and $Tx = \frac{1}{16}x$. Then T and f have a coincidence point.

Proof. Note that

$$\begin{aligned} d(Tx, Ty)(t) &= \left| \frac{1}{4}x - \frac{1}{4}y \right| e^t \\ &= 4 \left| \frac{1}{16}x - \frac{1}{16}y \right| e^t \\ &= 4d(fx, fy)(t) \end{aligned}$$

for all $x, y \in X$ and $t \in [0, 1]$. Thus T and f satisfy all the hypotheses of Corollary 2.2 and hence T and f have a coincidence point. Here 0 is the coincidence point of T and f .

Remarks:

- (1) Theorem 4.1 of [29] is a special case of Theorem 2.2.
- (2) Corollary 4.1 of [29] is a special case of Corollary 2.8.
- (3) Theorem 4 of [31] is a special case of Corollary 2.8.

Acknowledgements

The authors thank the editor and the referees for their useful comments and suggestions.

Authors' contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 11 October 2011 Accepted: 20 February 2012 Published: 20 February 2012

References

1. Huang, LG, Zhang, X: Cone metric spaces and fixed point theorems of contractive mappings. *J Math Anal Appl.* **332**, 1468–1476 (2007). doi:10.1016/j.jmaa.2005.03.087
2. Arshad, M, Azam, A, Beg, I: Common fixed points of two maps in cone metric spaces. *Rend Circ Mat Palermo.* **57**, 433–441 (2008). doi:10.1007/s12215-008-0032-5
3. Arshad, M, Azam, A, Vetro, P: Some common fixed point results in cone metric spaces. *Fixed Point Theory Appl.* **2009**, 11 (2009). (Article ID 493965)
4. Abbas, M, Ali Khan, M, Radenović, S: Common coupled fixed point theorems in cone metric spaces for w -compatible mappings. *Appl Math Comput.* (2010)
5. Abbas, M, Rhoades, BE: Fixed and periodic point results in cone metric spaces. *Appl Math Lett.* **22**, 511–515 (2009). doi:10.1016/j.aml.2008.07.001
6. Abbas, M, Jungck, G: Common fixed point results for noncommuting mappings without continuity in cone metric spaces. *J Math Anal Appl.* **341**, 416–420 (2008). doi:10.1016/j.jmaa.2007.09.070
7. Altun, I, Durmaz, G: Some fixed point theorems on ordered cone metric spaces. *Rend Circ Mat Palermo.* **58**, 319–325 (2009). doi:10.1007/s12215-009-0026-y
8. Altun, I, Damjanović, B, Djorić, D: Fixed point and common fixed point theorems on ordered cone metric spaces. *Appl Math Lett.* **23**, 310–316 (2010). doi:10.1016/j.aml.2009.09.016
9. Aydi, H, Nashine, HK, Samet, B, Yazidi, H: Coincidence and common fixed point results in partially ordered cone metric spaces and applications to integral equations. *Nonlinear Anal.* **74**, 6814–6825 (2011). doi:10.1016/j.na.2011.07.006
10. Aydi, H, Samet, B, Vetro, C: Coupled fixed point results in cone metric spaces for compatible mappings. *Fixed Point Theory Appl.* **2011**, 27 (2011). doi:10.1186/1687-1812-2011-27
11. Azam, A, Beg, I, Arshad, M: Fixed point in topological vector space-valued cone metric spaces. *Fixed Point Theory Appl.* **2010**, 9 (2010). (Article ID 604084)
12. Choudhury, BS, Metiya, N: Fixed points of weak contractions in cone metric spaces. *Nonlinear Anal.* **72**, 1589–1593 (2010). doi:10.1016/j.na.2009.08.040
13. Choudhury, BS, Metiya, N: The point of coincidence and common fixed point for a pair of mappings in cone metric spaces. *Comput Math Appl.* **60**, 1686–1695 (2010). doi:10.1016/j.camwa.2010.06.048
14. Dordivić, M, Dorić, D, Kadelburg, Z, Radenović, S, Spasić, D: Fixed point results under c -distance in tvs-cone metric spaces. *Fixed Point Theory Appl.* **2011**, 29 (2011). doi:10.1186/1687-1812-2011-29
15. Di Bari, C, Vetro, P: Weakly ϕ -pairs and common fixed points in cone metric spaces. *Rend Circ Mat Palermo.* **58**, 125–132 (2009). doi:10.1007/s12215-009-0012-4
16. Di Bari, D, Vetro, P: ϕ -pairs and common fixed points in cone metric spaces. *Rend Circolo Matematico di Palermo.* **57**, 279–285 (2008). doi:10.1007/s12215-008-0020-9
17. Wei-Shih, Du: A note on cone metric fixed point theory and its equivalence. *Nonlinear Anal.* **72**, 2259–2261 (2010). doi:10.1016/j.na.2009.10.026
18. Ilić, D, Rakočević, V: Quasi-contraction on a cone metric space. *Appl Math Lett.* **22**, 728–731 (2009). doi:10.1016/j.aml.2008.08.011
19. Ilić, D, Rakočević, V: Common fixed points for maps on cone metric space. *J Math Anal Appl.* **341**, 876–882 (2008). doi:10.1016/j.jmaa.2007.10.065
20. Janković, S, Kadelburg, Z, Radenović, S, Rhoades, BE: Assad-Kirk-Type fixed point theorems for a pair of nonself mappings on cone metric spaces. *Fixed Point Theory Appl.* **2009** (2009). (Article ID 761086)
21. Janković, S, Golubović, Z, Radenović, S: Compatible and weakly compatible mappings in cone metric spaces. *Math Comput Model.* **52**, 1728–1738 (2010). doi:10.1016/j.mcm.2010.06.043
22. Kadelburg, Z, Pavlović, M, Radenović, S: Common fixed point theorems for ordered contractions and quasicontractions in ordered cone metric spaces. *Comput Math Appl.* **59**, 3148–3159 (2010). doi:10.1016/j.camwa.2010.02.039

23. Khojasteh, F, Razani, A, Moradi, S: A fixed point of generalized T_F -contraction mappings in cone metric spaces. *Fixed Point Theory Appl.* **2011**, 14 (2011). doi:10.1186/1687-1812-2011-14
24. Sumitra, R, Rhymend Uthariaraj, R, Hemavathy, R, Vijayaraju, P: Common fixed point theorem for non-self mappings satisfying generalized Ćirić type contraction condition in cone metric space. *Fixed Point Theory Appl.* **2010**, 17 (2010). (Article ID 408086)
25. Shatanawi, W: Some common coupled fixed point results in cone metric spaces. *Int J Math Anal.* **4**, 2381–2388 (2010)
26. Shatanawi, W: Partially ordered cone metric spaces and coupled fixed point results. *Comput Math Appl.* **60**, 2508–2515 (2010). doi:10.1016/j.camwa.2010.08.074
27. D Turkoglu, D, Abuloha, M: Cone metric spaces and fixed point theorems in diametrically contractive mappings. *Acta Mathematica Sinica (English Series)*. **26**, 489–496 (2010)
28. Vetro, P: Common fixed points in cone metric spaces. *Rend Circ Mat Palermo.* **56**, 464–468 (2010)
29. Şahin, İ, Telci, M: Fixed points of contractive mappings on complete cone metric spaces. *Hacet J Math Stat.* **38**, 59–67 (2009)
30. Şahin, İ, Telci, M: A theorem on common fixed points of expansion type mappings in cone metric spaces. *An Şt Univ Ovidius Constanţa.* **18**, 329–336 (2010)
31. Wang, SZ, Li, BY, Gao, ZM, Iseki, K: Some fixed point theorems on expansion mappings. *Math Japon.* **29**, 631–636 (1984)

doi:10.1186/1687-1812-2012-19

Cite this article as: Shatanawi and Awawdeh: Some fixed and coincidence point theorems for expansive maps in cone metric spaces. *Fixed Point Theory and Applications* 2012 **2012**:19.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
