

RESEARCH

Open Access



On Appell-type Changhee polynomials and numbers

Jeong Gon Lee¹, Lee-Chae Jang^{2*}, Jong-Jin Seo³, Sang-Ki Choi⁴ and Hyuck In Kwon⁵

*Correspondence:

lcjang@konkuk.ac.kr

²Graduate School of Education,
Konkuk University, Seoul, 143-701,
Republic of KoreaFull list of author information is
available at the end of the article**Abstract**

In this paper, we consider the Appell-type Changhee polynomials and derive some properties of these polynomials. Furthermore, we investigate certain identities for these polynomials.

MSC: 05A10; 11B68; 11S80; 05A19**Keywords:** Changhee polynomials; Appell-type Changhee polynomials; degenerate Bernoulli polynomials; beta functions**1 Introduction**

Let p be a fixed odd prime number. Throughout this paper, we denote by \mathbb{Z}_p , \mathbb{Q}_p , and \mathbb{C}_p the ring of p -adic integers, the field of p -adic numbers, and the completion of algebraic closure of \mathbb{Q}_p . The p -adic norm $|\cdot|_p$ is normalized as $|p|_p = \frac{1}{p}$. Let $C(\mathbb{Z}_p)$ be the space of continuous functions on \mathbb{Z}_p . For $f \in C(\mathbb{Z}_p)$, the fermionic p -adic integral on \mathbb{Z}_p is defined by Kim to be

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x)(-1)^x \quad (1)$$

(see [1–19]). For $f_1(x) = f(x+1)$, we have

$$I_{-1}(f_1) + I_{-1}(f) = 2f(0). \quad (2)$$

As is well known, the Changhee polynomials are defined by the generating function

$$\int_{\mathbb{Z}_p} (1+t)^{x+y} d\mu_{-1}(y) = \frac{2}{2+t}(1+t)^x = \sum_{n=0}^{\infty} \text{Ch}_n(x) \frac{t^n}{n!}. \quad (3)$$

When $x=0$, $\text{Ch}_n = \text{Ch}_n(0)$ are called the Changhee numbers (see [17, 18, 20]). The gamma and beta functions are defined by the following definite integrals: for $\alpha > 0$, $\beta > 0$,

$$\Gamma(\alpha) = \int_0^{\infty} e^{-t} t^{\alpha-1} dt \quad (4)$$

and

$$\begin{aligned}
 B(\alpha, \beta) &= \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} dt \\
 &= \int_0^\infty \frac{t^{\alpha-1}}{(1+t)^{\alpha+\beta}} dt
 \end{aligned}
 \tag{5}$$

(see[20, 21]). Thus, by (4) and (5) we have

$$\Gamma(\alpha + 1) = \alpha\Gamma(\alpha), \quad B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)}.
 \tag{6}$$

Stirling numbers of the first kind are defined by

$$(\log(1 + t))^n = n! \sum_{m=n}^\infty S_1(m, n) \frac{t^m}{m!},
 \tag{7}$$

and the Stirling numbers of the second kind are defined by

$$(e^t - 1)^n = n! \sum_{l=n}^\infty S_2(n, l) \frac{t^l}{l!} \quad (n \geq 0).
 \tag{8}$$

Recently, Lim and Qi [20] have derived integral identities for Appell-type λ -Changhee numbers from the fermionic integral equation. The degenerate Bernoulli polynomials, a degenerate version of the well-known family of polynomials, were introduced by Carlitz, and after that, many researchers have studied the degenerate special polynomials (see [1–3, 20, 22–28]).

The goal of this paper is to consider the Appell-type Changhee polynomials, another version of the Changhee polynomials in (3), and derive some properties of these polynomials. Furthermore, we investigate certain identities for these polynomials.

2 Some identities for Appell-type Changhee polynomials

Now we define the Appell-type Changhee polynomials $Ch_n^*(x)$ by

$$\frac{2}{2+t} e^{xt} = \sum_{n=0}^\infty Ch_n^*(x) \frac{t^n}{n!}.
 \tag{9}$$

When $x = 0$, the Changhee numbers $Ch_n^* = Ch_n^*(0)$ are equal to the Changhee numbers $Ch_n = Ch_n(0)$. From (9) we have

$$\begin{aligned}
 \frac{2}{2+t} e^{xt} &= \left(\sum_{m=0}^\infty Ch_m^* \frac{t^m}{m!} \right) \left(\sum_{l=0}^\infty x^l \frac{t^l}{l!} \right) \\
 &= \sum_{n=0}^\infty \left(\sum_{m=0}^n \binom{n}{m} Ch_m^* x^{n-m} \right) \frac{t^n}{n!}.
 \end{aligned}
 \tag{10}$$

By (10) we have the following theorem.

Theorem 1 For $n \in \mathbb{N}$, we have

$$\text{Ch}_n^*(x) = \sum_{m=0}^n \binom{n}{m} \text{Ch}_m^* x^{n-m}. \tag{11}$$

By (9), replacing t by $e^t - 1$, we get

$$\frac{2}{2 + e^t - 1} e^{x(e^t-1)} = \sum_{n=0}^{\infty} \text{Ch}_n^*(x) \frac{(e^t - 1)^n}{n!}. \tag{12}$$

Then we have

$$\begin{aligned} \text{RHS} &= \sum_{n=0}^{\infty} \text{Ch}_n^*(x) \frac{(e^t - 1)^n}{n!} \\ &= \sum_{n=0}^{\infty} \text{Ch}_n^*(x) \frac{1}{n!} \sum_{l=n}^{\infty} S_2(l, n) \frac{t^l}{l!} \\ &= \sum_{l=0}^{\infty} \sum_{n=0}^l \text{Ch}_n^*(x) S_2(l, n) \frac{t^l}{l!}, \end{aligned} \tag{13}$$

where $S_2(l, n)$ are the Stirling numbers of the second kind, and

$$\begin{aligned} \text{LHS} &= \frac{2}{1 + e^t} e^{x(e^t-1)} \\ &= \sum_{m=0}^{\infty} E_m \frac{t^m}{m!} \sum_{n=0}^{\infty} \text{Bel}_n(x) \frac{t^n}{n!} \\ &= \sum_{l=0}^{\infty} \sum_{n=0}^l \binom{l}{n} E_n \text{Bel}_{l-n}(x) \frac{t^l}{l!}. \end{aligned} \tag{14}$$

It is well known that the Bell polynomials are defined by the generating function

$$e^{x(e^t-1)} = \sum_{n=0}^{\infty} \text{Bel}_n(x) \frac{t^n}{n!}$$

(see [8]). By (13) and (14) we have the following theorem.

Theorem 2 For $l \in \mathbb{N}$, we have

$$\sum_{n=0}^l \text{Ch}_n^*(x) S_2(l, n) = \sum_{n=0}^l \binom{l}{n} E_n \text{Bel}_{l-n}(x). \tag{15}$$

By (11) we can derive the following equation:

$$\begin{aligned} \frac{d}{dx} \text{Ch}_n^*(x) &= \sum_{m=0}^{n-1} \binom{n}{m} \text{Ch}_m^*(n-m) x^{n-m-1} \\ &= n \text{Ch}_{n-1}^*(x). \end{aligned} \tag{16}$$

From (16) we get

$$\begin{aligned}
 n \int_0^x \text{Ch}_{n-1}^*(s) ds &= \int_0^x \frac{d}{ds} \text{Ch}_n^*(s) ds \\
 &= \text{Ch}_n^*(s) \Big|_{s=0}^x \\
 &= \text{Ch}_n^*(x) - \text{Ch}_n^*.
 \end{aligned}
 \tag{17}$$

By (17) we can derive the following theorem.

Theorem 3 For $n \in \mathbb{N}$, we have

$$\frac{\text{Ch}_{n+1}^*(x) - \text{Ch}_{n+1}^*}{n + 1} = \int_0^x \text{Ch}_n^*(s) ds.
 \tag{18}$$

By (4) we note that

$$\begin{aligned}
 2 &= \left(\sum_{n=0}^{\infty} \text{Ch}_n^* \frac{t^n}{n!} \right) (2 + t) \\
 &= \left(\sum_{n=0}^{\infty} 2 \text{Ch}_n^* \frac{t^n}{n!} \right) + t \sum_{n=0}^{\infty} \text{Ch}_n^* \frac{t^n}{n!} \\
 &= \left(\sum_{n=0}^{\infty} 2 \text{Ch}_n^* \frac{t^n}{n!} \right) + \sum_{n=1}^{\infty} n \text{Ch}_{n-1}^* \frac{t^n}{n!} \\
 &= 2 \text{Ch}_0^* + \sum_{n=1}^{\infty} (2 \text{Ch}_n^* + n \text{Ch}_{n-1}^*) \frac{t^n}{n!}.
 \end{aligned}
 \tag{19}$$

By (19) we have the following theorem.

Theorem 4 For $n \in \mathbb{N}$, we have

$$\text{Ch}_0^* = 1, \quad 2 \text{Ch}_n^* + n \text{Ch}_{n-1}^* = 0 \quad \text{if } n \geq 1.
 \tag{20}$$

Now we observe that

$$\begin{aligned}
 \sum_{n=0}^{\infty} \text{Ch}_n^*(1-x) \frac{t^n}{n!} &= \frac{2}{2+t} e^{(1-x)t} \\
 &= \frac{2}{2+t} e^t e^{-xt} \\
 &= \left(\sum_{l=0}^{\infty} \text{Ch}_l^*(1) \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} (-x)^m \frac{t^m}{m!} \right) \\
 &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \binom{n}{m} \text{Ch}_{n-m}^*(1) (-x)^m \right) \frac{t^n}{n!}.
 \end{aligned}
 \tag{21}$$

From (21) we obtain the following theorem.

Theorem 5 For $n \in \mathbb{N}$, we have

$$\text{Ch}_n^*(1-x) = \sum_{m=0}^n \binom{n}{m} \text{Ch}_{n-m}^*(1)(-x)^m. \tag{22}$$

By (22) we get

$$\begin{aligned} \int_0^1 \text{Ch}_n^*(1-x)x^n dx &= \sum_{m=0}^n \binom{n}{m} \text{Ch}_{n-m}^*(1)(-1)^m \int_0^1 x^{n+m} dx \\ &= \sum_{m=0}^n \binom{n}{m} (-1)^m \frac{\text{Ch}_{n-m}^*(1)}{n+m+1}. \end{aligned} \tag{23}$$

From (16) we note that

$$\begin{aligned} &\int_0^1 y^n \text{Ch}_n^*(x+y) dy \\ &= \frac{y^{n+1}}{n+1} \text{Ch}_n^*(x+y) \Big|_{y=0}^1 - \frac{1}{n+1} \int_0^1 y^{n+1} \frac{d}{dy} \text{Ch}_n^*(x+y) dy \\ &= \frac{\text{Ch}_n^*(x+1)}{n+1} - \frac{n}{n+1} \int_0^1 y^{n+1} \text{Ch}_{n-1}^*(x+y) dy \\ &= \frac{\text{Ch}_n^*(x+1)}{n+1} - \frac{n}{n+1} \left(\frac{\text{Ch}_{n-1}^*(x+y)}{n+2} y^{n+2} \Big|_{y=0}^1 \right) \\ &\quad + (-1)^2 \frac{n}{n+1} \frac{1}{n+2} (n-1) \int_0^1 y^{n+2} \text{Ch}_{n-2}^*(x+y) dy \\ &= \frac{\text{Ch}_n^*(x+1)}{n+1} - \frac{n}{n+1} \frac{\text{Ch}_{n-1}^*(x+1)}{n+2} + (-1)^2 \frac{n}{n+1} \frac{n-1}{n+2} \int_0^1 y^{n+2} \text{Ch}_{n-2}^*(x+y) dy \\ &= \frac{\text{Ch}_n^*(x+1)}{n+1} - \frac{n}{n+1} \frac{\text{Ch}_{n-1}^*(x+1)}{n+2} + (-1)^2 \frac{n}{n+1} \frac{n-1}{n+2} \frac{\text{Ch}_{n-2}^*(x+1)}{n+3} \\ &\quad + (-1)^3 \frac{n}{n+1} \frac{n-1}{n+2} \frac{n-2}{n+3} \int_0^1 y^{n+3} \text{Ch}_{n-3}^*(x+y) dy. \end{aligned} \tag{24}$$

Also, we get

$$\int_0^1 y^{2n-1} \text{Ch}_1^*(x+y) dy = \frac{\text{Ch}_1^*(x+y)}{2n} y^{2n} \Big|_{y=0}^1 - \frac{1}{2n} \int_0^1 y^{2n} \text{Ch}_0^*(x+y) dy. \tag{25}$$

From (11) we get

$$\text{Ch}_0^*(x) = 1, \tag{26}$$

and hence

$$\begin{aligned} \int_0^1 y^{2n-1} \text{Ch}_1^*(x+y) dy &= \frac{\text{Ch}_1^*(x)}{2n} - \frac{1}{2n} \int_0^1 y^{2n} dy \\ &= \frac{\text{Ch}_1^*(x)}{2n} - \frac{1}{2n(2n+1)}. \end{aligned} \tag{27}$$

By (27), continuing the process in (24), we have

$$\begin{aligned} & \int_0^1 y^n \text{Ch}_n^*(x+y) dy \\ &= \frac{\text{Ch}_n^*(x+1)}{n+1} + \sum_{m=1}^n (-1)^m \text{Ch}_{n-m}^*(x+1) \frac{n(n-1)\cdots(n-m+1)}{(n+1)(n+2)\cdots(n+m+1)}. \end{aligned} \tag{28}$$

We note that

$$\begin{aligned} \text{Ch}_n^*(x+y) &= \text{Ch}_n^*(x+1+y-1) \\ &= \sum_{l=1}^n \binom{n}{l} \text{Ch}_l^*(x+1) (-1)^{n-l} (1-y)^{n-l}. \end{aligned} \tag{29}$$

By (29) we get

$$\begin{aligned} & \int_0^1 y^n \text{Ch}_n^*(x+y) dy \\ &= \sum_{l=1}^n \binom{n}{l} \text{Ch}_l^*(x+1) (-1)^{n-l} \int_0^1 y^n (1-y)^{n-l} dy \\ &= \sum_{l=1}^n \binom{n}{l} \text{Ch}_l^*(x+1) (-1)^{n-l} B(n+1, n-l+1) \\ &= \sum_{l=0}^n \binom{n}{l} \text{Ch}_l^*(x+1) (-1)^{n-l} \frac{\Gamma(n+1)\Gamma(n-l+1)}{\Gamma(2n-l+2)} \\ &= \sum_{l=0}^n (-1)^{n-l} \binom{n}{l} \frac{n!(n-l)!}{(2n-l+1)!} \text{Ch}_l^*(x+1) \\ &= \sum_{l=0}^n (-1)^{n-l} \frac{n \binom{n}{l}}{(2n-l+1) \binom{2n-l}{n}} \text{Ch}_l^*(x+1). \end{aligned} \tag{30}$$

By (28) and (30) we have the following theorem.

Theorem 6 For $n \in \mathbb{N}$, we have

$$\begin{aligned} & \sum_{l=0}^n (-1)^{n-l} \frac{n \binom{n}{l}}{(2n-l+1) \binom{2n-l}{n}} \text{Ch}_l^*(x+1) \\ &= \frac{\text{Ch}_n^*(x+1)}{n+1} + \sum_{m=1}^n (-1)^m \text{Ch}_{n-m}^*(x+1) \frac{n(n-1)\cdots(n-m+1)}{(n+1)(n+2)\cdots(n+m+1)}. \end{aligned} \tag{31}$$

From (16) we note that

$$\begin{aligned} & \int_0^1 y^n \text{Ch}_n^*(x+y) dy \\ &= \frac{\text{Ch}_{n+1}^*(x+y)}{n+1} \Big|_{y=0}^1 - \frac{1}{n+1} n \int_0^1 y^{n-1} \text{Ch}_{n+1}^*(x+y) dy \end{aligned}$$

$$\begin{aligned}
 &= \frac{\text{Ch}_{n+1}^*(x+1)}{n+1} - \frac{n}{n+1} \int_0^1 y^{n-1} \text{Ch}_{n+1}^*(x+y) dy \\
 &= \frac{\text{Ch}_{n+1}^*(x+1)}{n+1} - \frac{n}{n+1} \frac{\text{Ch}_{n+2}^*(x+1)}{n+2} + \frac{n(n-1)}{(n+1)(n+2)} \int_0^1 y^{n-2} \text{Ch}_{n+2}^*(x+y) dy \\
 &= \frac{\text{Ch}_{n+1}^*(x+1)}{n+1} - \frac{n}{n+1} \frac{\text{Ch}_{n+2}^*(x+1)}{n+2} + \frac{n(n-1)}{(n+1)(n+2)} \frac{\text{Ch}_{n+3}^*(x+1)}{n+3} \\
 &\quad - \frac{n(n-1)(n-2)}{(n+1)(n+2)(n+3)} \int_0^1 y^{n-3} \text{Ch}_{n+3}^*(x+y) dy. \tag{32}
 \end{aligned}$$

Also, we have

$$\begin{aligned}
 &\int_0^1 y \text{Ch}_{2n-1}^*(x+y) dy \\
 &= \frac{\text{Ch}_{2n}^*(x+y)}{2n} y \Big|_{y=0}^1 - \frac{1}{2n} \int_0^1 1 \cdot \text{Ch}_{2n}^*(x+y) dy \\
 &= \frac{\text{Ch}_{2n}^*(x+1)}{2n} - \frac{1}{2n} \frac{1}{2n+1} \text{Ch}_{2n+1}^*(x+y) \Big|_{y=0}^1 \\
 &= \frac{\text{Ch}_{2n}^*(x+1)}{2n} - \frac{\text{Ch}_{2n+1}^*(x+1) - \text{Ch}_{2n+1}^*(x)}{2n(2n+1)}. \tag{33}
 \end{aligned}$$

By (30), continuing the process in (28), we obtain the following theorem.

Theorem 7 For $n \in \mathbb{N}$, we have

$$\begin{aligned}
 &\sum_{l=0}^n (-1)^{n-l} \frac{n^{(n)}}{(2n-l+1) \binom{2n-l}{n}} \text{Ch}_l^*(x+1) \\
 &= \frac{\text{Ch}_{n+1}^*(x+1)}{n+1} + \sum_{m=1}^{n-1} (-1)^m \text{Ch}_{n+m+1}^*(x+1) \frac{n(n-1) \cdots (n-m+1)}{(n+1)(n+2) \cdots (n+m+1)} \\
 &\quad + (-1)^n \frac{n!}{(2n+1)_{n+1}} (\text{Ch}_{2n+1}^*(x+1) - \text{Ch}_{2n+1}^*(1)). \tag{34}
 \end{aligned}$$

Now, we have

$$\begin{aligned}
 &\int_0^1 \text{Ch}_n^*(x) \text{Ch}_m^*(x) dx \\
 &= \frac{\text{Ch}_{n+1}^*(x) \text{Ch}_m^*(x)}{n+1} \Big|_0^1 - \frac{1}{n+1} m \int_0^1 \text{Ch}_{n+1}^*(x) \text{Ch}_{m-1}^*(x) dx \\
 &= \frac{1}{n+1} (\text{Ch}_{n+1}^*(1) \text{Ch}_m^*(1) - \text{Ch}_{n+1}^*(0) \text{Ch}_m^*(0)) \\
 &\quad - \frac{m}{n+1} \int_0^1 \text{Ch}_{n+1}^*(x) \text{Ch}_{m-1}^*(x) dx \\
 &= \frac{\text{Ch}_{n+1}^*(1) \text{Ch}_m^*(1) - \text{Ch}_{n+1}^* \text{Ch}_m^*}{n+1} - \frac{m}{n+1} \frac{\text{Ch}_{n+2}^*(1) \text{Ch}_{m-1}^*(1) - \text{Ch}_{n+2}^* \text{Ch}_{m-1}^*}{n+2} \\
 &\quad + (-1)^2 \frac{m}{n+1} \frac{m-1}{n+2} \int_0^1 \text{Ch}_{n+2}^*(x) \text{Ch}_{m-2}^*(x) dx \tag{35}
 \end{aligned}$$

and

$$\begin{aligned} & \int_0^1 \text{Ch}_{n+m-1}^*(x) \text{Ch}_1^*(x) dx \\ &= \frac{\text{Ch}_{n+m}^*(1) \text{Ch}_1^*(1) - \text{Ch}_{n+m}^* \text{Ch}_1^*}{n+m} - \frac{1}{n+m} \int_0^1 \text{Ch}_{n+m}^*(x) \text{Ch}_0^*(x) dx \\ &= \frac{\text{Ch}_{n+m}^*(1) \text{Ch}_1^*(1) - \text{Ch}_{n+m}^* \text{Ch}_1^*}{n+m} - \frac{1}{n+m} \frac{\text{Ch}_{n+m+1}^*(1) - \text{Ch}_{n+m+1}^*}{n+m+1}. \end{aligned} \tag{36}$$

By (30) with $x = 0$ we get

$$\begin{aligned} & \int_0^1 \text{Ch}_n^*(x) \text{Ch}_m^*(x) dx \\ &= \sum_{j=0}^m \binom{m}{j} \text{Ch}_j^* \int_0^1 x^{m-j} \text{Ch}_m^*(x) dx \\ &= \sum_{j=0}^m \binom{m}{j} \text{Ch}_j^* \sum_{l=0}^{m-j} (-1)^{m-j-l} \frac{(m-j) \binom{m-j}{l}}{(2(m-j)-l+1) \binom{2(m-j)-l}{m-j}} \text{Ch}_l^*(1) \\ &= \sum_{j=0}^m \sum_{l=0}^{m-j} \binom{m}{j} (-1)^{m-j-l} \frac{(m-j) \binom{m-j}{l}}{(2(m-j)-l+1) \binom{2(m-j)-l}{m-j}} \text{Ch}_j^* \text{Ch}_l^*(1). \end{aligned} \tag{37}$$

By (37), continuing the process in (35), we obtain the following theorem.

Theorem 8 For $n \in \mathbb{N}$, we have

$$\begin{aligned} & \sum_{j=0}^m \sum_{l=0}^{m-j} \binom{m}{j} (-1)^{m-j-l} \frac{(m-j) \binom{m-j}{l}}{(2(m-j)-l+1) \binom{2(m-j)-l}{m-j}} \text{Ch}_j^* \text{Ch}_l^*(1) \\ &= \frac{\text{Ch}_{n+1}^*(1) \text{Ch}_m^*(1) - \text{Ch}_{n+1}^* \text{Ch}_m^*}{n+1} \\ &+ \sum_{k=1}^{m-1} (-1)^k \frac{m(m-1) \cdots (m-k+1)}{(n+1)(n+2) \cdots (n+k+1)} \\ &\times (\text{Ch}_{n+k+1}^*(1) \text{Ch}_{m-k}^*(1) - \text{Ch}_{n+k+1}^* \text{Ch}_{m-k}^*) \\ &+ (-1)^m \frac{m!}{(n+m+1)_{m+1}} (\text{Ch}_{n+m+1}^*(1) - \text{Ch}_{n+m+1}^*). \end{aligned} \tag{38}$$

3 Remarks

In this section, by using the fermionic p -adic integral on \mathbb{Z}_p , we derive some identities for Changhee polynomials, Stirling numbers of the first kind, and Euler numbers. By (2) we note that

$$\begin{aligned} \frac{2}{2+t} e^{xt} &= \int_{\mathbb{Z}_p} (1+t)^y e^{xt} d\mu_{-1}(y) \\ &= \int_{\mathbb{Z}_p} e^{y \log(1+t) + xt} d\mu_{-1}(y) \end{aligned} \tag{39}$$

and

$$\begin{aligned}
 e^{xt} e^{y \log(1+t)} &= \left(\sum_{m=0}^{\infty} x^m \frac{t^m}{m!} \right) \left(\sum_{l=0}^{\infty} \frac{y^l (\log(1+t))^l}{l!} \right) \\
 &= \left(\sum_{m=0}^{\infty} x^m \frac{t^m}{m!} \right) \left(\sum_{l=0}^{\infty} y^l \sum_{k=l}^{\infty} S_1(k, l) \frac{t^k}{k!} \right) \\
 &= \left(\sum_{m=0}^{\infty} x^m \frac{t^m}{m!} \right) \left(\sum_{k=0}^{\infty} \sum_{l=0}^k y^l S_1(k, l) \frac{t^k}{k!} \right) \\
 &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \sum_{l=0}^k \binom{n}{k} x^{n-k} y^l S_1(k, l) \right) \frac{t^n}{n!}.
 \end{aligned} \tag{40}$$

Thus, by (39) and (40) we have

$$\begin{aligned}
 \sum_{n=0}^{\infty} \text{Ch}_n^*(x) \frac{t^n}{n!} &= \int_{\mathbb{Z}_p} e^{y \log(1+t)} e^{xt} d\mu_{-1}(y) \\
 &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \sum_{l=0}^k \binom{n}{k} x^{n-k} \int_{\mathbb{Z}_p} y^l d\mu_{-1}(y) S_1(k, l) \right) \frac{t^n}{n!} \\
 &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \sum_{l=0}^k \binom{n}{k} x^{n-k} E_l S_1(k, l) \right) \frac{t^n}{n!}.
 \end{aligned} \tag{41}$$

From (41) we have the following theorem.

Theorem 9 For $n \in \mathbb{N}$, we have

$$\text{Ch}_n^*(x) = \sum_{k=0}^n \sum_{l=0}^k \binom{n}{k} x^{n-k} E_l S_1(k, l). \tag{42}$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to this work. All authors read and approved the final manuscript.

Author details

¹Division of Mathematics and Informational Statistics, Nanoscale Science and Technology Institute, Wonkwang University, Iksan, 570-749, Republic of Korea. ²Graduate School of Education, Konkuk University, Seoul, 143-701, Republic of Korea. ³Department of Applied Mathematics, Pukyong National University, Busan, 608-737, Republic of Korea. ⁴Department of Mathematics Education, Konkuk University, Seoul, 143-701, Republic of Korea. ⁵Department of Mathematics, Kwangwoon University, Seoul, 139-701, Republic of Korea.

Acknowledgements

This paper was supported by Wonkwang University in 2015.

Received: 10 February 2016 Accepted: 17 May 2016 Published online: 21 June 2016

References

1. Bayad, A, Kim, T: Identities for the Bernoulli, the Euler and the Genocchi numbers and polynomials. *Adv. Stud. Contemp. Math. (Kyungshang)* **20**(2), 247-253 (2010)
2. Bayad, A, Kim, T: Identities involving values of Bernstein, q -Bernoulli, and q -Euler polynomials. *Russ. J. Math. Phys.* **18**(2), 133-143 (2011)
3. Carlitz, L: Degenerate Stirling, Bernoulli and Eulerian numbers. *Util. Math.* **15**, 51-88 (1979)

4. Kim, BM, Jang, LC: A note on the von Staudt-Clausen's theorem for the weighted q -Genocchi numbers. *Adv. Differ. Equ.* **2015**, 4 (2015)
5. Kim, DS, Kim, T: Some identities of degenerate Euler polynomials arising from p -adic fermionic integrals on \mathbb{Z}_p . *Integral Transforms Spec. Funct.* **26**(4), 295-302 (2015)
6. Kim, DS, Kim, T: Some identities of degenerate special polynomials. *Open Math.* **13**, 380-389 (2015)
7. Kim, DS, Kim, T, Dolgy, DV: Degenerate q -Euler polynomials. *Adv. Differ. Equ.* **2015**, 246 (2015)
8. Kim, DS, Kim, T: Some identities of Bell polynomials. *Sci. China Math.* **58**(10), 1-10 (2015)
9. Kim, T: Note on the Euler numbers and polynomials. *Adv. Stud. Contemp. Math. (Kyungshang)* **17**(2), 131-136 (2008)
10. Kim, T: Some properties on the integral of the product of several Euler polynomials. *Quaest. Math.* **38**(4), 553-562 (2015)
11. Kim, T: Degenerate Euler zeta function. *Russ. J. Math. Phys.* **22**(4), 469-472 (2015)
12. Kim, T: On the multiple q -Genocchi and Euler numbers. *Russ. J. Math. Phys.* **15**(4), 481-486 (2008)
13. Kim, T, Mansour, T: Umbral calculus associated with Frobenius-type Eulerian polynomials. *Russ. J. Math. Phys.* **21**(4), 484-493 (2008)
14. Kim, T: New approach to q -Euler, Genocchi numbers and their interpolation functions. *Adv. Stud. Contemp. Math. (Kyungshang)* **18**(2), 105-112 (2009)
15. Kim, T: On Euler-Barnes multiple zeta functions. *Russ. J. Math. Phys.* **10**(3), 261-267 (2003)
16. Kim, T: A study on the q -Euler numbers and the fermionic q -integral of the product of several type q -Bernstein polynomials on \mathbb{Z}_p . *Adv. Stud. Contemp. Math. (Kyungshang)* **23**(1), 5-11 (2013)
17. Kim, T, Kim, DS: A note on nonlinear Changhee differential equations. *Russ. J. Math. Phys.* **23**(1), 1-5 (2016)
18. Wang, NL, Li, H: Some identities on the higher-order Daehee and Changhee numbers. *Pure Appl. Math. J.* **5**, 33-37 (2015)
19. Yilmaz Yasar, B, Özarslan, MA: Frobenius-Euler and Frobenius-Genocchi polynomials and their differential equations. *New Trends Math. Sci.* **3**(2), 172-180 (2015)
20. Lim, D, Qi, F: On the Appell type λ -Changhee polynomials. *J. Nonlinear Sci. Appl.* **9**, 1872-1876 (2016)
21. Kim, T, Park, JW, Seo, JJ: A note on λ -zeta function. *Glob. J. Pure Appl. Math.* **11**(5), 3501-3506 (2015)
22. Adelberg, A: A finite difference approach to degenerate Bernoulli and Stirling polynomials. *Discrete Math.* **140**(1-3), 1-21 (1995)
23. Carlitz, L: A degenerate Staudt-Clausen theorem. *Arch. Math. (Basel)* **7**, 28-33 (1956)
24. Gaboury, S, Tremblay, R, Fugère, B-J: Some explicit formulas for certain new classes of Bernoulli, Euler and Genocchi polynomials. *Proc. Jangjeon Math. Soc.* **17**(1), 115-123 (2014)
25. Howard, FT: Explicit formulas for degenerate Bernoulli numbers. *Discrete Math.* **162**(1-3), 175-185 (1996)
26. Liu, GD: Degenerate Bernoulli numbers and polynomials of higher order. *J. Math. (Wuhan)* **25**(3), 283-288 (2005) (in Chinese)
27. Mahmudov, NI, Akkeles, A, Öneren, A: On two dimensional q -Bernoulli and q -Genocchi polynomials: properties and location of zeros. *J. Comput. Anal. Appl.* **18**(5), 834-843 (2015)
28. Kwon, JK: A note on weighted Boole polynomials. *Glob. J. Pure Appl. Math.* **11**(4), 2055-2063 (2015)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
