

## ERROR ESTIMATES FOR THE FINITE ELEMENT SOLUTIONS OF VARIATIONAL INEQUALITIES

**M. ASLAM NOOR**

Mathematics Department  
Islamia University  
Bahawalpur, Pakistan

(Received September 11, 1979 and in revised form January 10, 1980)

ABSTRACT. For piecewise linear approximation of variational inequalities associated with the mildly nonlinear elliptic boundary value problems having auxiliary constraint conditions, we prove that the error estimate for  $u-u_h$  in the  $W^{1,2}$ -norm is of order  $h$ .

KEY WORDS AND PHRASES. *Finite Element, Variational Inequalities, Approximation, Mildly nonlinear.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. *Primary 35J20, 65N30, 41A15.*

### 1. INTRODUCTION.

In this paper, we derive the finite element error estimates for the approximate solution of mildly nonlinear boundary value problems having auxiliary constraint conditions. A much used approach with any elliptic problem is to reformulate it in a weak or variational form. It has been shown by Noor and Whiteman

[1] that in the presence of a constraint, such an approach leads to a variational inequality which is the weak formulation. An approximate formulation of the variational inequality is then defined, and the error estimates involving the difference between the solutions of the exact and approximate formulation in the  $W^{1,2}$ -norm is obtained, which is in fact of order  $h$ . This result is an extension of that obtained by Falk [2] and Mosco and Strang [3] for the constrained linear problem.

## 2. MAIN RESULTS

For simplicity, we consider the problem of the following type:

$$\left. \begin{aligned} -\Delta u(\underline{x}) &= F(\underline{x}, u), & \underline{x} \in \Omega \\ u(\underline{x}) &= 0, & \underline{x} \in \partial\Omega \end{aligned} \right\} \quad (2.1)$$

where  $\Omega$  is a convex polygon domain in  $\mathbb{R}^n$  with boundary  $\partial\Omega$  and  $\bar{\Omega} = \Omega \cup \partial\Omega$ , its closure. The given function  $f(u) = F(\underline{x}, u) \in C^0(\bar{\Omega} \times \mathbb{R})$  is a real-value function involving the unknown  $u$ . If  $f(u)$  is both antimonotone and Lipschitz continuous, then it is known that there does exist a unique solution of (2.1); see Noor [4, p. 57-62]. We study this problem in the usual Sobolev space  $W_2^1(\Omega) = H^1$ , the space of functions which together with their generalized derivatives of order one are in  $L_2(\Omega)$ . The subspace of functions from  $H^1$ , which in a generalized sense satisfy the homogeneous boundary conditions on  $\partial\Omega$ , is  $W_1^0(\Omega) = H_0^1$ .

It has been shown by Tonti [5] that, in its direct variational formulation, (2.1) is equivalent to finding  $u \in H_0^1$  such that

$$I[u] \leq I[v] \quad \text{for all } v \in H_0^1,$$

where

$$\begin{aligned} I[v] &= \int_{\Omega} \left( \frac{\partial v}{\partial \underline{x}} \right)^2 d\Omega - 2 \int_{\Omega} \int_0^v f(\eta) d\eta d\Omega \\ &= a(v, v) - 2F(v) \end{aligned} \quad (2.2)$$

is the energy functional associated with (2.1).

We now consider the case when the solution  $u$  of (2.1) is required to satisfy the condition  $u \geq \psi$ , where  $\psi \in C^0(\bar{\Omega}) \cap H^1_0$ ,  $\psi \leq 0$  on  $\partial\Omega$ ; (see Glowinski [6, p. IV. 2]). In this situation, our problem is to find

$$u \in K \stackrel{\text{def}}{=} \{v; v \in H^1_0, v \geq \psi \text{ on } \Omega\},$$

a closed convex subset of  $H^1_0$  ( see Mosco [7] ), such that  $u$  minimizes  $I[v]$  on  $K$ . It has been shown by Noor and Whiteman [1] that the minimum of  $I[v]$  on  $K$  can be characterized by a class of variational inequalities

$$a(u, v-u) \geq \langle F'(u), v-u \rangle \quad \text{for all } v \in K, \tag{2.3}$$

where  $F'(u)$  is the Frechet differential of  $F(u)$  and is in fact,

$$\langle F'(u), v \rangle = \int_{\Omega} f(u) v \, d\Omega. \tag{2.4}$$

The finite dimensional form of (2.3) is to find  $u_h \in K_h$  such that

$$a(u_h, v_h - u_h) \geq \langle F'(u_h), v_h - u_h \rangle \quad \text{for all } v_h \in K_h. \tag{2.5}$$

Here  $K_h$  is a finite dimensional convex subset of  $H^1_0$ ; for the construction of  $K_h$ , see Mosco [3]. Let  $\Omega$  be the convex polygon. We partition it into triangles of side length less than  $h$ . We consider  $S_h \subset H^1_0$ , the subspace of continuous piecewise linear functions on the triangulation of  $\Omega$ , vanishing on the boundary  $\partial\Omega$ . Let  $\psi_h$  be the interpolant of  $\psi$  such that  $\psi_h$  agrees with  $\psi$  at all the vertices of the triangulation. For our purpose, it is enough to choose the finite dimensional convex subset  $K_h = S_h \cap \{v_h \geq \psi_h \text{ on } \Omega\}$ . For other choices of convex subsets  $K_h$ , see Nitsche [8], where he has chosen  $K_h = K \cap S_h$ .

We also want to know the regularity of the function  $u \in K$  satisfying (2.3). In this case Brézis and Stampacchia [9] have shown that, if  $\psi$  lies in both  $H^1_0$  and  $H^2$ , then the solution  $u \in K$  satisfying (2.3) also lies in  $H^2$ .

Its norm can be estimated from the data:

$$\|u\|_2 \leq e \|\psi\|_2.$$

Moreover, if  $\tilde{u}$  is the interpolant of  $u$ , which agrees with  $u$  at every vertex of  $\Omega$ , then  $\tilde{u}$  lies in  $K_h$ . It is well known from approximation theory (see Strang and Fix [10]) that

$$\|u - \tilde{u}\| \leq eh \|u\|_2. \tag{2.6}$$

We also note that in certain cases, the equality holds instead of the inequality in (2.3). This happens when  $v$ , together with  $2u-v$ , also lies in  $K$ . In this case, we get

$$a(u, v-u) = \langle F'(u), v-u \rangle. \tag{2.7}$$

Finally let  $C$  and  $C_h$  be the cones composed of non-negative functions on  $H_0^1$  and its subspace  $S_h$ . Thus, it is clear that

$$U = u - \psi \text{ is in } C$$

$$U_h = u_h - \psi_h \text{ is in } C_h.$$

From these relations, it follows that

$$u - u_h = U - U_h + \psi - \psi_h. \tag{2.8}$$

**DEFINITION.** An operator  $T$  on  $H_0^1$  is said to be quasi-monotone, if for all  $z, u, v, w, \in H_0^1$ ,

$$\langle Tz - Tv, w - z \rangle \geq 0. \tag{2.9}$$

We also need the following result of Mosco and Strang [3].

**THEOREM 1:** Suppose that  $U \geq 0$  in the plane polygon  $\Omega$  and that  $U$  lies in both  $H_0^1$  and  $H^2$ . Then, there exists a  $V_h$  in  $S_h$  such that

$$0 \leq V_h \leq U \text{ in } \Omega$$

and

$$\|U - V_h\| \leq eh \|U\|_2. \tag{2.10}$$

Now we state and prove the main result.

THEOREM 2. Let  $a(u,v)$  be a continuous coercive bilinear form and  $F'(u)$  be a quasi-monotone operator on  $H_0^1$ . If  $V_h \in C_h$  and  $2U-V_h \in C$ , then

$$|| u-u_h || = O(h),$$

where  $u$  and  $u_h$  are the solutions of (2.3) and (2.5) respectively.

PROOF. Since both  $v = \psi + V_h$  and  $2u-v = \psi + (2U-V_h)$  are in  $K$ , we have from (2.3) and (2.7) that

$$a(u, V_h-U) = \langle F'(u), V_h-U \rangle. \tag{2.11}$$

Letting  $v_h = \psi_h + V_h$  and  $u_h = \psi_h + U_h$  in (2.5), we have

$$a(u_h, V_h-U_h) \geq \langle F'(u_h), V_h-U_h \rangle. \tag{2.12}$$

Using  $v = \psi + U_h$  in (2.3), we get

$$a(u, U_h-U) \geq \langle F'(u), U_h-U \rangle. \tag{2.13}$$

From (2.11) and (2.13), we obtain

$$a(u, U_h-V_h) \geq \langle F'(u), U_h-V_h \rangle, \tag{2.14}$$

and from (2.12) and (2.14), we get

$$a(u-u_h, U_h-V_h) \geq \langle F'(u) - F'(u_h), U_h-V_h \rangle.$$

Thus, using the quasi-monotonicity of  $F'(u)$ , we have

$$a(u-u_h, U_h-V_h) \geq 0,$$

which can be written as

$$a(u-u_h, U-U_h) \leq a(u-u_h, U-V_h). \tag{2.15}$$

Now by coercivity of  $a(u,v)$ , it follows that there exists a constant  $\alpha > 0$  such that

$$\begin{aligned} \alpha ||u-u_h||^2 &\leq a(u-u_h, u-u_h) \\ &= a(u-u_h, \psi - \psi_h) + a(u-u_h, U-U_h), \text{ from (2.8)} \\ &\leq a(u-u_h, \psi - \psi_h) + a(u-u_h, U-V_h) \\ &\leq \beta ||u-u_h|| \{ ||\psi - \psi_h|| + ||U-V_h|| \}, \end{aligned}$$

where  $\beta > 0$  is a continuity constant of the bilinear form  $a(u,v)$ .

Hence, it follows that

$$\begin{aligned} \|u-u_h\| &\leq \frac{\beta}{\alpha} \{ \|\psi-\psi_h\| + \|u-v_h\| \} \\ &\leq \frac{\beta}{\alpha} Ch \{ \|\psi\|_2 + \|u\|_2 \}, \text{ by (2.6)} \end{aligned}$$

and (2.10), from which the required estimate follows.

Remark: The problem of deriving the  $L^\infty$ -norm estimates for the mildly nonlinear problems having constraint conditions is still open.

Acknowledgement: The author would like to thank Professor J.R. Whiteman for his helpful criticism and encouragement.

#### REFERENCES

1. Noor, M. A. and Whiteman, J.R.; Finite element error analysis of nonlinear variational inequalities, Appl. Math. and Opt., (to appear).
2. Falk, R. S.; Error estimates for the approximation of a class of variational inequalities. Math. Comp. 28(1974), 963-971.
3. Mosco, U. and Strang, G.; One sided approximation and variational inequalities, Bull. Amer. Math. Soc., 80 (1974), 308-312.
4. Noor, M. A.; On variational inequalities, Ph. D. Thesis, Brunel University, U. K. 1975.
5. Tonti, E.; Variational formulation of nonlinear differential equations, Bull. Acad. Royale de Belgique, (1969), 137-165 & 262-278.
6. Glowinski, R.; Introduction to the approximation of elliptic variational inequalities, T/R. 76006, Universite Paris VI, 1976.
7. Mosco, U.; An introduction to the approximate solution of variational inequalities in Constructive Aspects of Functional Analysis, Edizione Cremonese, Roma, (1973), 499-685.
8. Nitsche, J.;  $L_\infty$ -convergence of finite element approximation in Mathematical Aspects of Finite Element Method, Rome, 1975.
9. Brezis, H., and Stampacchia, G.; Sur la regularite de la solution d'inequations elliptiques. Bull. Soc. Math. France. 96(1968). 153-180.
10. Strang, G. and Fix, G.; An analysis of the finite element method, Prentice Englewood Cliff., N. J., 1973.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

