

Hindawi Publishing Corporation
Journal of Applied Mathematics
Volume 2012, Article ID 329391, 13 pages
doi:10.1155/2012/329391

Research Article

Almost Sure Central Limit Theory for Self-Normalized Products of Sums of Partial Sums

Qunying Wu^{1,2}

¹ College of Science, Guilin University of Technology, Guilin 541004, China

² Guangxi Key Laboratory of Spatial Information and Geomatics, Guilin University of Technology, Guilin 541004, China

Correspondence should be addressed to Qunying Wu, wqy666@glite.edu.cn

Received 18 January 2012; Accepted 7 February 2012

Academic Editor: Naseer Shahzad

Copyright © 2012 Qunying Wu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let X, X_1, X_2, \dots be a sequence of independent and identically distributed random variables in the domain of attraction of a normal law. An almost sure limit theorem for the self-normalized products of sums of partial sums is established.

1. Introduction

Let $\{X, X_n\}_{n \in \mathbb{N}}$ be a sequence of independent and identically distributed (i.i.d.) positive random variables with a nondegenerate distribution function and $\mathbb{E}X = \mu > 0$. For each $n \geq 1$, the symbol S_n/V_n denotes self-normalized partial sums, where $S_n = \sum_{i=1}^n X_i$ and $V_n^2 = \sum_{i=1}^n (X_i - \mu)^2$. We say that the random variable X belongs to the domain of attraction of the normal law if there exist constants $a_n > 0$, $b_n \in \mathbb{R}$ such that

$$\frac{S_n - b_n}{a_n} \xrightarrow{d} \mathcal{N}, \quad \text{as } n \rightarrow \infty, \quad (1.1)$$

here and in the sequel \mathcal{N} is a standard normal random variable. We say that $\{X, X_n\}_{n \in \mathbb{N}}$ satisfies the central limit theorem (CLT).

It is known that (1.1) holds if and only if

$$\lim_{x \rightarrow \infty} \frac{x^2 \mathbb{P}(|X| > x)}{\mathbb{E}X^2 I(|X| \leq x)} = 0. \quad (1.2)$$

In contrast to the well-known classical central limit theorem, Gine et al. [1] obtained the following self-normalized version of the central limit theorem: $(S_n - \mathbb{E}S_n)/V_n \xrightarrow{d} \mathcal{N}$ as $n \rightarrow \infty$ if and only if (1.2) holds.

Brosamler [2] and Schatte [3] obtained the following almost sure central limit theorem (ASCLT): let $\{X_n\}_{n \in \mathbb{N}}$ be i.i.d. random variables with mean 0, variance $\sigma^2 > 0$, and partial sums S_n . Then

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I \left\{ \frac{S_k}{\sigma \sqrt{k}} < x \right\} = \Phi(x) \quad \text{a.s. } \forall x \in \mathbb{R}, \quad (1.3)$$

with $d_k = 1/k$ and $D_n = \sum_{k=1}^n d_k$; here and in the sequel I denotes an indicator function, and $\Phi(x)$ is the standard normal distribution function. Some ASCLT results for partial sums were obtained by Lacey and Philipp [4], Ibragimov and Lifshits [5], Miao [6], Berkes and Csáki [7], Hörmann [8], Wu [9, 10], and Ye and Wu [11]. Huang and Pang [12] and Zhang and Yang [13] obtained ASCLT results for self-normalized version. However, ASCLT results for self-normalized products of sums of partial sums have not been reported yet.

Under mild moment conditions, ASCLT follows from the ordinary CLT, but in general the validity of ASCLT is a delicate question of a totally different character as CLT. The difference between CLT and ASCLT lies in the weight in ASCLT. The terminology of summation procedures (see e.g., Chandrasekharan and Minakshisundaram [14], page 35) shows that the larger the weight sequence $\{d_k; k \geq 1\}$ in (1.3) is, the stronger the relation becomes. By this argument, one should also expect to get stronger results if we use larger weights. And it would be of considerable interest to determine the optimal weights.

On the other hand, by the Theorem 1 of Schatte [3], (1.3) fails for weight $d_k = 1$. The optimal weight sequence remains unknown.

The purpose of this paper is to study and establish the ASCLT for self-normalized products of sums of partial sums of random variables in the domain of attraction of the normal law, we will show that the ASCLT holds under a fairly general growth condition on $d_k = k^{-1} \exp((\ln k)^\alpha)$, $0 \leq \alpha < 1/2$.

In the following, we assume that $\{X, X_n\}_{n \in \mathbb{N}}$ is a sequence of i.i.d. positive random variables in the domain of attraction of the normal law with $\mathbb{E}X = \mu > 0$ and define $S_k = \sum_{i=1}^k X_i$, $V_k = \sum_{i=1}^k (X_i - \mu)^2$, $T_k = \sum_{i=1}^k S_i$. Let $b_{n,k} = \sum_{j=k}^n 1/j$, $c_{n,k} = 2 \sum_{j=k}^n (j+1-k)/(j(j+1))$, and $d_{n,k} = (n+1-k)/(n+1)$ for $1 \leq k \leq n$. $I(A)$ denotes the indicator function of set A , and $a_n \sim b_n$ denotes $a_n/b_n \rightarrow 1, n \rightarrow \infty$. The symbol c stands for a generic positive constant, which may differ from one place to another. Let

$$\begin{aligned} l(x) &= \mathbb{E}(X - \mu)^2 I\{|X - \mu| \leq x\}, & b &= \inf\{x \geq 1; l(x) > 0\}, \\ \eta_j &= \inf \left\{ s; s \geq b + 1, \frac{l(s)}{s^2} \leq \frac{1}{j} \right\} & & \text{for } j \geq 1. \end{aligned} \quad (1.4)$$

By the definition of η_j , we have $jl(\eta_j) \leq \eta_j^2$ and $jl(\eta_j - \varepsilon) > (\eta_j - \varepsilon)^2$ for any $\varepsilon > 0$. It implies that

$$nl(\eta_n) \sim \eta_n^2, \quad \text{as } n \rightarrow \infty, \eta_n < n + 1. \quad (1.5)$$

Our theorem is formulated in a more general setting.

Theorem 1.1. Let $\{X, X_n\}_{n \in \mathbb{N}}$ be a sequence of i.i.d. positive random variables in the domain of attraction of the normal law with $\mathbb{E}X = \mu > 0$. Suppose

$$l(\eta_n) \sim l\left(\frac{\eta_n}{\ln n}\right). \quad (1.6)$$

For $0 \leq \alpha < 1/2$, set

$$d_k = \frac{\exp(\ln^\alpha k)}{k}, \quad D_n = \sum_{k=1}^n d_k. \quad (1.7)$$

Then

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I\left(\left(\prod_{j=1}^k \left(\frac{T_j}{j(j+1)\mu/2}\right)\right)^{\mu/V_k} \leq x\right) = F(x) \quad a.s., \quad (1.8)$$

for any $x \in \mathbb{R}$, where F is the distribution function of the random variable $\exp(\sqrt{10/3}\mathcal{N})$.

By the terminology of summation procedures, we have the following corollary.

Corollary 1.2. Theorem 1.1 remains valid if we replace the weight sequence $\{d_k\}_{k \in \mathbb{N}}$ by $\{d_k^*\}_{k \in \mathbb{N}}$ such that $0 \leq d_k^* \leq d_k$, $\sum_{k=1}^{\infty} d_k^* = \infty$.

Remark 1.3. If $\mathbb{E}X^2 = \sigma^2 < \infty$, then X is in the domain of attraction of the normal law and $l(x) \rightarrow \sigma^2$, $\eta_n^2 \sim \sigma^2 n$, thus (1.6) holds. Therefore, the class of random variables in Theorems 1.1 is of very broad range.

Remark 1.4. Whether Theorem 1.1 holds for $1/2 \leq \alpha < 1$ remains open.

2. Proofs

Furthermore, the following four lemmas will be useful in the proof, and the first is due to Csörgo et al. [15].

Lemma 2.1. Let X be a random variable with $\mathbb{E}X = \mu$, and denote $l(x) = \mathbb{E}(X - \mu)^2 I\{|X - \mu| \leq x\}$. The following statements are equivalent:

- (i) $l(x)$ is a slowly varying function at ∞ ;
- (ii) X is in the domain of attraction of the normal law;
- (iii) $x^2 \mathbb{P}(|X - \mu| > x) = o(l(x))$;
- (iv) $x \mathbb{E}(|X - \mu| I(|X - \mu| > x)) = o(l(x))$;
- (v) $\mathbb{E}(|X - \mu|^\alpha I(|X - \mu| \leq x)) = o(x^{\alpha-2} l(x))$ for $\alpha > 2$.

Lemma 2.2. Let $\{\xi, \xi_n\}_{n \in \mathbb{N}}$ be a sequence of uniformly bounded random variables. If there exist constants $c > 0$ and $\delta > 0$ such that

$$|\mathbb{E}\xi_k \xi_j| \leq c \left(\frac{k}{j}\right)^\delta, \quad \text{for } 1 \leq k < j, \quad (2.1)$$

then

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k \xi_k = 0 \quad \text{a.s.}, \quad (2.2)$$

where d_k and D_n are defined by (1.7).

Proof. We can easily apply the similar arguments of (2.1) in Wu [16] to get Lemma 2.2, and we omit the details here. \square

The following Lemma 2.3 can be directly verified.

Lemma 2.3. (i) $c_{n,i} = 2(b_{n,i} - d_{n,i})$;

$$(ii) \sum_{i=1}^n b_{n,i}^2 = 2n - b_{n,1} \sim 2n;$$

$$(iii) \sum_{i=1}^n d_{n,i}^2 = \frac{n}{3} - \frac{n}{6(n+1)} \sim \frac{n}{3};$$

$$(iv) \sum_{i=1}^n c_{n,i}^2 = \frac{10n}{3} - 4b_{n,1} + \frac{10n}{3(n+1)} \sim \frac{10n}{3}.$$

For every $1 \leq i \leq k \leq n$, let

$$\bar{X}_{ki} = (X_i - \mu)I(|X_i - \mu| \leq \eta_k), \quad \bar{S}_{k,i} = \sum_{j=1}^i c_{k,j} \bar{X}_{kj}, \quad \bar{V}_k^2 = \sum_{j=1}^k \bar{X}_{kj}^2. \quad (2.3)$$

Lemma 2.4. Suppose that the assumptions of Theorem 1.1 hold. Then

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I \left\{ \frac{\bar{S}_{k,k} - \mathbb{E}\bar{S}_{k,k}}{\sqrt{10kl(\eta_k)/3}} \leq x \right\} = \Phi(x) \quad \text{a.s. for any } x \in \mathbb{R}, \quad (2.4)$$

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k \left(I \left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k) \right) - \mathbb{E} I \left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k) \right) \right) = 0 \quad \text{a.s.}, \quad (2.5)$$

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k \left(f \left(\frac{\bar{V}_k^2}{kl(\eta_k)} \right) - \mathbb{E} f \left(\frac{\bar{V}_k^2}{kl(\eta_k)} \right) \right) = 0 \quad \text{a.s.}, \quad (2.6)$$

where d_k and D_n are defined by (1.7) and f is a nonnegative, bounded Lipschitz function.

Proof. By $\mathbb{E}(X - \mu) = 0$, Lemma 2.1 (iv), we have

$$\left| \mathbb{E} \bar{X}_{ni} \right| \leq \mathbb{E} |X - \mu| I(|X - \mu| > \eta_n) = \frac{o(l(\eta_n))}{\eta_n}. \quad (2.7)$$

Thus, by (1.5) and Lemma 2.3 (iv),

$$\text{Var} \bar{X}_{ni} = \mathbb{E} \bar{X}_{ni}^2 - \left(\mathbb{E} \bar{X}_{ni} \right)^2 \sim l(\eta_n), \quad \sum_{i=1}^n \text{Var} \left(c_{n,i} \bar{X}_{ni} \right) \sim \frac{10n}{3} l(\eta_n) := B_n^2. \quad (2.8)$$

By (1.5) and Lemma 2.3 (i),

$$\max_{1 \leq i \leq n} c_{n,i} \leq 2 \max_{1 \leq i \leq n} b_{n,i} \leq 2b_{n,1} \sim 2 \ln n, \quad \frac{\ln n \max_{1 \leq i \leq n} \left| \mathbb{E} \bar{X}_{ni} \right|}{B_n} \rightarrow 0. \quad (2.9)$$

Thus by combining Lemma 2.3 (iv), (1.6), and (2.8), Lindeberg condition

$$\begin{aligned} \frac{1}{B_n^2} \sum_{i=1}^n \mathbb{E} \left(c_{n,i} \bar{X}_{ni} \right)^2 I_{(|c_{n,i} \bar{X}_{ni} - \mathbb{E} c_{n,i} \bar{X}_{ni}| > \varepsilon B_n)} &\sim \frac{c}{nl(\eta_n)} \sum_{i=1}^n c_{n,i}^2 \mathbb{E} \bar{X}_{ni}^2 I_{(|\bar{X}_{ni} - \mathbb{E} \bar{X}_{ni}| > \varepsilon B_n / 2 \ln n)} \\ &\leq \frac{c}{nl(\eta_n)} \sum_{i=1}^n c_{n,i}^2 \mathbb{E} \bar{X}_{ni}^2 I_{(|\bar{X}_{ni}| > \varepsilon B_n / 4 \ln n)} \\ &= \frac{c}{nl(\eta_n)} \sum_{i=1}^n c_{n,i}^2 \mathbb{E} (X - \mu)^2 I_{(c\eta_n / \ln n < |X - \mu| \leq \eta_n)} \\ &= c \frac{l(\eta_n) - l(c\eta_n / \ln n)}{l(\eta_n)} \rightarrow 0, \quad \text{as } n \rightarrow \infty \end{aligned} \quad (2.10)$$

hold.

Hence, it follows that

$$\frac{\bar{S}_{n,n} - \mathbb{E} \bar{S}_{n,n}}{B_n} \xrightarrow{d} \mathcal{N}, \quad \text{as } n \rightarrow \infty. \quad (2.11)$$

This implies that for any $g(x)$, which is a nonnegative, bounded Lipschitz function,

$$\mathbb{E} g \left(\frac{\bar{S}_{n,n} - \mathbb{E} \bar{S}_{n,n}}{B_n} \right) \rightarrow \mathbb{E} g(\mathcal{N}), \quad \text{as } n \rightarrow \infty. \quad (2.12)$$

Hence, we obtain

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k \mathbb{E} g \left(\frac{\bar{S}_{k,k} - \mathbb{E} \bar{S}_{k,k}}{B_k} \right) = \mathbb{E} g(\mathcal{N}) \quad (2.13)$$

from the Toeplitz lemma.

On the other hand, note that (2.4) is equivalent to

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k g \left(\frac{\bar{S}_{k,k} - \mathbb{E} \bar{S}_{k,k}}{B_k} \right) = \mathbb{E} g(\mathcal{N}) \quad \text{a.s.}, \quad (2.14)$$

from Theorem 7.1 of Billingsley [17] and Section 2 of Peligrad and Shao [18]. Hence, to prove (2.4), it suffices to prove

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k \left(g \left(\frac{\bar{S}_{k,k} - \mathbb{E} \bar{S}_{k,k}}{B_k} \right) - \mathbb{E} g \left(\frac{\bar{S}_{k,k} - \mathbb{E} \bar{S}_{k,k}}{B_k} \right) \right) = 0 \quad \text{a.s.}, \quad (2.15)$$

for any $g(x)$ which is a nonnegative, bounded Lipschitz function.

Let

$$\xi_k = g \left(\frac{\bar{S}_{k,k} - \mathbb{E} \bar{S}_{k,k}}{B_k} \right) - \mathbb{E} g \left(\frac{\bar{S}_{k,k} - \mathbb{E} \bar{S}_{k,k}}{B_k} \right), \quad \text{for } k \geq 1. \quad (2.16)$$

Clearly, there is a constant $c > 0$ such that

$$|g(x)| \leq c, \quad |g(x) - g(y)| \leq c|x - y| \quad \text{for any } x, y \in \mathbb{R}, \quad |\xi_k| \leq 2c, \quad \text{for any } k. \quad (2.17)$$

For any $1 \leq k < l$, note that

$$\bar{S}_{l,l} - \bar{S}_{l,k} = \sum_{i=1}^l c_{l,i} \bar{X}_{li} - \sum_{i=1}^k c_{l,i} \bar{X}_{li} = \sum_{i=k+1}^l c_{l,i} \bar{X}_{li}. \quad (2.18)$$

For any $1 \leq k < j$, note that $g((\bar{S}_{k,k} - \mathbb{E}\bar{S}_{k,k})/B_k)$ and $g((\bar{S}_{j,j} - \mathbb{E}\bar{S}_{j,j} - (\bar{S}_{j,k} - \mathbb{E}\bar{S}_{j,k}))/B_j)$ are independent, $g(x)$ is a nonnegative, bounded Lipschitz function, $\sum_{i=1}^k c_{k,i}^2 \sim 10k/3, c_{j,i}^2 \leq 4b_{j,i}^2, \sum_{i=1}^k b_{k,i}^2 \sim 2k$, and $\ln x \leq 4x^{1/4}, x \geq 1$. By the definition of η_j , we get

$$\begin{aligned}
|\mathbb{E}\xi_k \xi_j| &= \left| \text{Cov} \left(g \left(\frac{\bar{S}_{k,k} - \mathbb{E}\bar{S}_{k,k}}{B_k} \right), g \left(\frac{\bar{S}_{j,j} - \mathbb{E}\bar{S}_{j,j}}{B_j} \right) \right) \right| \\
&= \left| \text{Cov} \left(g \left(\frac{\bar{S}_{k,k} - \mathbb{E}\bar{S}_{k,k}}{B_k} \right), g \left(\frac{\bar{S}_{j,j} - \mathbb{E}\bar{S}_{j,j}}{B_j} \right) - g \left(\frac{\bar{S}_{j,j} - \mathbb{E}\bar{S}_{j,j} - (\bar{S}_{j,k} - \mathbb{E}\bar{S}_{j,k})}{B_j} \right) \right) \right| \\
&\leq c \frac{\mathbb{E} \left| \sum_{i=1}^k c_{j,i} (\bar{X}_{ji} - \mathbb{E}\bar{X}_{ji}) \right|}{\sqrt{j}l(\eta_j)} \leq c \frac{\sqrt{\mathbb{E} \left(\sum_{i=1}^k c_{j,i} (\bar{X}_{ji} - \mathbb{E}\bar{X}_{ji}) \right)^2}}{\sqrt{j}l(\eta_j)} \\
&\leq c \frac{\sqrt{\sum_{i=1}^k b_{j,i}^2 \mathbb{E}\bar{X}_{ji}^2}}{\sqrt{j}l(\eta_j)} \leq c \frac{\sqrt{\sum_{i=1}^k (b_{k,i} + b_{j,k+1})^2 l(\eta_j)}}{\sqrt{j}l(\eta_j)} \\
&= c \frac{\sqrt{\sum_{i=1}^k b_{k,i}^2 + \sum_{i=1}^k b_{j,k+1}^2}}{\sqrt{j}} \leq c \frac{\sqrt{k + k \ln^2(j/k)}}{\sqrt{j}} \\
&\leq c \left(\frac{k}{j} \right)^{1/4}.
\end{aligned} \tag{2.19}$$

By Lemma 2.2, (2.15) holds.

Now we prove (2.5). Let

$$Z_k = I \left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k) \right) - \mathbb{E} I \left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k) \right) \quad \text{for any } k \geq 1. \tag{2.20}$$

It is known that $I(A \cup B) - I(B) \leq I(A)$ for any sets A and B ; then for $1 \leq k < j$, by Lemma 2.1 (iii) and (1.5), we get

$$\mathbb{P}(|X - \mu| > \eta_j) = o(1) \frac{l(\eta_j)}{\eta_j^2} = \frac{o(1)}{j}. \tag{2.21}$$

Hence for $1 \leq k < j$,

$$\begin{aligned}
|\mathbb{E}Z_k Z_j| &= \left| \text{Cov} \left(I \left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k) \right), I \left(\bigcup_{i=1}^j (|X_i - \mu| > \eta_j) \right) \right) \right| \\
&= \left| \text{Cov} \left(I \left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k) \right), I \left(\bigcup_{i=1}^j (|X_i - \mu| > \eta_j) \right) \right. \right. \\
&\quad \left. \left. - I \left(\bigcup_{i=k+1}^j (|X_i - \mu| > \eta_j) \right) \right) \right| \\
&\leq \mathbb{E} \left| I \left(\bigcup_{i=1}^j (|X_i - \mu| > \eta_j) \right) - I \left(\bigcup_{i=k+1}^j (|X_i - \mu| > \eta_j) \right) \right| \\
&\leq \mathbb{E} I \left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_j) \right) \leq k \mathbb{P}(|X - \mu| > \eta_j) \\
&\leq \frac{k}{j}.
\end{aligned} \tag{2.22}$$

By Lemma 2.2, (2.5) holds.

Finally, we prove (2.6). Let

$$\zeta_k = f \left(\frac{\bar{V}_k^2}{kl(\eta_k)} \right) - \mathbb{E} f \left(\frac{\bar{V}_k^2}{kl(\eta_k)} \right) \quad \text{for any } k \geq 1. \tag{2.23}$$

For $1 \leq k < j$,

$$\begin{aligned}
|\mathbb{E}\zeta_k \zeta_j| &= \left| \text{Cov} \left(f \left(\frac{\bar{V}_k^2}{kl(\eta_k)} \right), f \left(\frac{\bar{V}_j^2}{jl(\eta_j)} \right) \right) \right| \\
&= \left| \text{Cov} \left(f \left(\frac{\bar{V}_k^2}{kl(\eta_k)} \right), f \left(\frac{\bar{V}_j^2}{jl(\eta_j)} \right) - f \left(\frac{\bar{V}_j^2 - \sum_{i=1}^k (X_i - \mu)^2 I(|X_i - \mu| \leq \eta_j)}{jl(\eta_j)} \right) \right) \right| \\
&\leq c \frac{\mathbb{E} \left(\sum_{i=1}^k (X_i - \mu)^2 I(|X_i - \mu| \leq \eta_j) \right)}{jl(\eta_j)} = c \frac{k \mathbb{E} (X - \mu)^2 I(|X - \mu| \leq \eta_j)}{jl(\eta_j)} = c \frac{kl(\eta_j)}{jl(\eta_j)} \\
&= c \frac{k}{j}.
\end{aligned} \tag{2.24}$$

By Lemma 2.2, (2.6) holds. This completes the proof of Lemma 2.4. \square

Proof of Theorem 1.1. Let $Z_j = T_j / (j(j+1)\mu/2)$; then (1.8) is equivalent to

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I \left(\frac{\sqrt{3}\mu}{\sqrt{10}V_k} \sum_{j=1}^k \ln Z_j \leq x \right) = \Phi(x), \quad \text{a.s. for any } x, \quad (2.25)$$

where $\Phi(x)$ is the distribution function of the standard normal random variable \mathcal{N} .

Let $q \in (4/3, 2)$, then $\mathbb{E}|X|^q < \infty$. Using Marcinkiewicz-Zygmund strong large number law, we have

$$S_k - \mu k = o(k^{1/q}) \quad \text{a.s.} \quad (2.26)$$

Thus,

$$\begin{aligned} |Z_i - 1| &= \left| \frac{\sum_{j=1}^i S_j - i(i+1)\mu/2}{i(i+1)\mu/2} \right| \leq \frac{\sum_{j=1}^i |S_j - \mu j|}{i(i+1)\mu/2} \leq \frac{\sum_{j=1}^i j^{1/q}}{i(i+1)\mu/2} \leq c \frac{i^{1/q+1}}{i^2} \\ &= i^{1/q-1} \rightarrow 0, \quad \text{a.s.} \end{aligned} \quad (2.27)$$

Hence by $|\ln(1+x) - x| = O(x^2)$ for $|x| < 1/2$, for any $0 < \varepsilon < 1$,

$$\begin{aligned} \left| \frac{1}{\sqrt{1 \pm \varepsilon B_k}} \sum_{i=1}^k \ln Z_i - \frac{1}{\sqrt{1 \pm \varepsilon B_k}} \sum_{i=1}^k (Z_i - 1) \right| &\leq c \frac{1}{\sqrt{kl(\eta_k)}} \sum_{i=1}^k (Z_i - 1)^2 \leq \frac{c}{\sqrt{kl(\eta_k)}} \sum_{i=1}^k i^{2(1/q-1)} \\ &\leq c \frac{1}{k^{3/2-2/q} \sqrt{l(\eta_k)}} \rightarrow 0 \quad \text{a.s. } k \rightarrow \infty, \end{aligned} \quad (2.28)$$

from $3/2 - 2/q > 0$, $l(x)$ is a slowly varying function at ∞ , and $\eta_k \leq k + 1$.

Hence for almost every event ω and any $\delta > 0$, there exists $k_0 = k_0(\omega, \delta, x)$ such that for $k > k_0$,

$$\begin{aligned} I \left(\frac{\mu}{\sqrt{1 \pm \varepsilon B_k}} \sum_{i=1}^k (Z_i - 1) \leq x - \delta \right) &\leq I \left(\frac{\mu}{\sqrt{1 \pm \varepsilon B_k}} \sum_{i=1}^k \ln Z_i \leq x \right) \\ &\leq I \left(\frac{\mu}{\sqrt{1 \pm \varepsilon B_k}} \sum_{i=1}^k (Z_i - 1) \leq x + \delta \right). \end{aligned} \quad (2.29)$$

Note that under condition $|X_j - \mu| \leq \eta_k, 1 \leq j \leq k$,

$$\begin{aligned}
\mu \sum_{i=1}^k (Z_i - 1) &= \sum_{i=1}^k \frac{\sum_{l=1}^i S_l - \mu \sum_{l=1}^i l}{i(i+1)/2} = \sum_{i=1}^k \frac{1}{i(i+1)/2} \sum_{l=1}^i \sum_{j=1}^l (X_j - \mu) \\
&= \sum_{i=1}^k \frac{1}{i(i+1)/2} \sum_{j=1}^i \sum_{l=j}^i (X_j - \mu) = \sum_{i=1}^k \sum_{j=1}^i \frac{2(i+1-j)}{i(i+1)} (X_j - \mu) \\
&= \sum_{j=1}^k \sum_{i=j}^k \frac{2(i+1-j)}{i(i+1)} \bar{X}_{kj} = \sum_{j=1}^k c_{k,j} \bar{X}_{kj} \\
&= \bar{S}_{k,k}.
\end{aligned} \tag{2.30}$$

Thus, for any given $0 < \varepsilon < 1, \delta > 0$, combining (2.29), we have for $k > k_0$

$$\begin{aligned}
I\left(\frac{\sqrt{3}\mu \sum_{i=1}^k \ln Z_i}{\sqrt{10}V_k} \leq x\right) &\leq I\left(\frac{\sqrt{3}\mu \sum_{i=1}^k \ln Z_i}{\sqrt{10(1+\varepsilon)kl(\eta_k)}} \leq x\right) + I(\bar{V}_k^2 > (1+\varepsilon)kl(\eta_k)) \\
&\quad + I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right) \\
&\leq I\left(\frac{\mu \sum_{i=1}^k (Z_i - 1)}{\sqrt{1+\varepsilon}B_k} \leq x + \delta\right) + I(\bar{V}_k^2 > (1+\varepsilon)kl(\eta_k)) \\
&\quad + I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right) \\
&\leq I\left(\frac{\bar{S}_{k,k}}{\sqrt{1+\varepsilon}B_k} \leq x + \delta\right) + I(\bar{V}_k^2 > (1+\varepsilon)kl(\eta_k)) \\
&\quad + 2I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right) \quad \text{for } x \geq 0, \\
I\left(\frac{\sqrt{3}\mu \sum_{i=1}^k \ln Z_i}{\sqrt{10}V_k} \leq x\right) &\leq I\left(\frac{\bar{S}_{k,k}}{\sqrt{1-\varepsilon}B_k} \leq x + \delta\right) + I(\bar{V}_k^2 < (1-\varepsilon)kl(\eta_k)) \\
&\quad + 2I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right) \quad \text{for } x < 0, \\
I\left(\frac{\sqrt{3}\mu \sum_{i=1}^k \ln Z_i}{\sqrt{10}V_k} \leq x\right) &\geq I\left(\frac{\bar{S}_{k,k}}{\sqrt{1-\varepsilon}B_k} \leq x - \delta\right) - I(\bar{V}_k^2 < (1-\varepsilon)kl(\eta_k)) \\
&\quad - 2I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right), \quad \text{for } x \geq 0,
\end{aligned}$$

$$\begin{aligned}
I\left(\frac{\sqrt{3}\mu \sum_{i=1}^k \ln Z_i}{\sqrt{10}V_k} \leq x\right) &\geq I\left(\frac{\bar{S}_{k,k}}{\sqrt{1+\varepsilon}B_k} \leq x - \delta\right) - I(\bar{V}_k^2 > (1+\varepsilon)kl(\eta_k)) \\
&\quad - 2I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right), \quad \text{for } x < 0.
\end{aligned} \tag{2.31}$$

Therefore, to prove (2.25), it suffices to prove

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I\left(\frac{\bar{S}_{k,k}}{B_k} \leq \sqrt{1 \pm \varepsilon}x \pm \delta_1\right) = \Phi(\sqrt{1 \pm \varepsilon}x \pm \delta_1) \quad \text{a.s.}, \tag{2.32}$$

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right) = 0 \quad \text{a.s.}, \tag{2.33}$$

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I(\bar{V}_k^2 > (1+\varepsilon)kl(\eta_k)) = 0 \quad \text{a.s.}, \tag{2.34}$$

$$\lim_{n \rightarrow \infty} \frac{1}{D_n} \sum_{k=1}^n d_k I(\bar{V}_k^2 < (1-\varepsilon)kl(\eta_k)) = 0 \quad \text{a.s.}, \tag{2.35}$$

for any $0 < \varepsilon < 1$ and $\delta_1 > 0$.

Firstly, we prove (2.32). Let $0 < \beta < 1/2$ and $h(\cdot)$ be a real function, such that for any given $x \in \mathbb{R}$,

$$I(y \leq \sqrt{1 \pm \varepsilon}x \pm \delta_1 - \beta) \leq h(y) \leq I(y \leq \sqrt{1 \pm \varepsilon}x \pm \delta_1 + \beta). \tag{2.36}$$

By $\mathbb{E}(X_i - \mu) = 0$, Lemma 2.1(iv) and (1.5), we have

$$\begin{aligned}
|\mathbb{E}\bar{S}_{k,k}| &= \left| \mathbb{E} \sum_{i=1}^k c_{k,i} (X_i - \mu) I(|X_i - \mu| \leq \eta_k) \right| \leq 2 \sum_{i=1}^k b_{k,i} \mathbb{E}|X_i - \mu| I(|X_i - \mu| > \eta_k) \\
&= 2 \sum_{i=1}^k \sum_{j=i}^k \frac{1}{j} \mathbb{E}|X - \mu| I(|X - \mu| > \eta_k) = \sum_{j=1}^k \sum_{i=1}^j \frac{1}{j} \frac{o(l(\eta_k))}{\eta_k} \\
&= o\left(\sqrt{kl(\eta_k)}\right).
\end{aligned} \tag{2.37}$$

This, combining with (2.4), (2.36) and the arbitrariness of β in (2.36), (2.32) holds.

By (2.5), (2.21) and the Toeplitz lemma,

$$\begin{aligned} 0 &\leq \frac{1}{D_n} \sum_{k=1}^n d_k I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right) \sim \frac{1}{D_n} \sum_{k=1}^n d_k \mathbb{E} I\left(\bigcup_{i=1}^k (|X_i - \mu| > \eta_k)\right) \\ &\leq \frac{1}{D_n} \sum_{k=1}^n d_k k \mathbb{P}(|X - \mu| > \eta_k) \longrightarrow 0 \quad \text{a.s.} \end{aligned} \quad (2.38)$$

That is, (2.33) holds.

Now we prove (2.34). For any given $\varepsilon > 0$, let f be a nonnegative, bounded Lipschitz function such that

$$I(x > 1 + \varepsilon) \leq f(x) \leq I\left(x > 1 + \frac{\varepsilon}{2}\right). \quad (2.39)$$

From $\mathbb{E}\bar{V}_k^2 = kl(\eta_k)$, \bar{X}_{ki} is i.i.d., $\mathbb{E}(\bar{X}_{ki}^2 - \mathbb{E}\bar{X}_{ki}^2) = 0$, Lemma 2.1 (v), and (1.5),

$$\begin{aligned} \mathbb{P}\left(\bar{V}_k^2 > \left(1 + \frac{\varepsilon}{2}\right)kl(\eta_k)\right) &= \mathbb{P}\left(\bar{V}_k^2 - \mathbb{E}\bar{V}_k^2 > \frac{\varepsilon}{2}kl(\eta_k)\right) \\ &\leq c \frac{\mathbb{E}\left(\bar{V}_k^2 - \mathbb{E}\bar{V}_k^2\right)^2}{k^2 l^2(\eta_k)} \leq c \frac{\mathbb{E}|X - \mu|^4 I(|X - \mu| \leq \eta_k)}{kl^2(\eta_k)} \\ &= \frac{o(1)\eta_k^2}{kl(\eta_k)} = o(1) \longrightarrow 0. \end{aligned} \quad (2.40)$$

Therefore, combining (2.6) and the Toeplitz lemma,

$$\begin{aligned} 0 &\leq \frac{1}{D_n} \sum_{k=1}^n d_k I\left(\bar{V}_k^2 > (1 + \varepsilon)kl(\eta_k)\right) \leq \frac{1}{D_n} \sum_{k=1}^n d_k f\left(\frac{\bar{V}_k^2}{kl(\eta_k)}\right) \\ &\sim \frac{1}{D_n} \sum_{k=1}^n d_k \mathbb{E} f\left(\frac{\bar{V}_k^2}{kl(\eta_k)}\right) \leq \frac{1}{D_n} \sum_{k=1}^n d_k \mathbb{E} I\left(\bar{V}_k^2 > \left(1 + \frac{\varepsilon}{2}\right)kl(\eta_k)\right) \\ &= \frac{1}{D_n} \sum_{k=1}^n d_k \mathbb{P}\left(\bar{V}_k^2 > \left(1 + \frac{\varepsilon}{2}\right)kl(\eta_k)\right) \\ &\longrightarrow 0 \quad \text{a.s.} \end{aligned} \quad (2.41)$$

Hence, (2.34) holds. By similar methods used to prove (2.34), we can prove (2.35). This completes the proof of Theorem 1.1. \square

Acknowledgments

The author is very grateful to the referees and the Editors for their valuable comments and some helpful suggestions that improved the clarity and readability of the paper. This paper is supported by the National Natural Science Foundation of China (11061012), a project supported by Program to Sponsor Teams for Innovation in the Construction of Talent Highlands in Guangxi Institutions of Higher Learning ([2011] 47), and the support program of Key Laboratory of Spatial Information and Geomatics (1103108-08).

References

- [1] E. Giné, F. Götze, and D. M. Mason, "When is the Student t-statistic asymptotically standard normal?" *The Annals of Probability*, vol. 25, no. 3, pp. 1514–1531, 1997.
- [2] G. A. Brosamler, "An almost everywhere central limit theorem," *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 104, no. 3, pp. 561–574, 1988.
- [3] P. Schatte, "On strong versions of the central limit theorem," *Mathematische Nachrichten*, vol. 137, pp. 249–256, 1988.
- [4] M. T. Lacey and W. Philipp, "A note on the almost sure central limit theorem," *Statistics & Probability Letters*, vol. 9, no. 3, pp. 201–205, 1990.
- [5] I. Ibragimov and M. Lifshits, "On the convergence of generalized moments in almost sure central limit theorem," *Statistics & Probability Letters*, vol. 40, no. 4, pp. 343–351, 1998.
- [6] Y. Miao, "Central limit theorem and almost sure central limit theorem for the product of some partial sums," *Indian Academy of Sciences Proceedings C*, vol. 118, no. 2, pp. 289–294, 2008.
- [7] I. Berkes and E. Csáki, "A universal result in almost sure central limit theory," *Stochastic Processes and their Applications*, vol. 94, no. 1, pp. 105–134, 2001.
- [8] S. Hörmann, "Critical behavior in almost sure central limit theory," *Journal of Theoretical Probability*, vol. 20, no. 3, pp. 613–636, 2007.
- [9] Q. Y. Wu, "Almost sure limit theorems for stable distributions," *Statistics & Probability Letters*, vol. 81, no. 6, pp. 662–672, 2011.
- [10] Q. Y. Wu, "A note on the almost sure limit theorem for self normalized partial sums of random variables in the domain of attraction of the normal law," *Journal of Inequalities and Applications*, vol. 2012, p. 17, 2012.
- [11] D. X. Ye and Q. Y. Wu, "Almost sure central limit theorem for product of partial sums of strongly mixing random variables," *Journal of Inequalities and Applications*, Article ID 576301, 9 pages, 2011.
- [12] S.-H. Huang and T.-X. Pang, "An almost sure central limit theorem for self-normalized partial sums," *Computers & Mathematics with Applications*, vol. 60, no. 9, pp. 2639–2644, 2010.
- [13] Y. Zhang and X.-Y. Yang, "An almost sure central limit theorem for self-normalized products of sums of i.i.d. random variables," *Journal of Mathematical Analysis and Applications*, vol. 376, no. 1, pp. 29–41, 2011.
- [14] K. Chandrasekharan and S. Minakshisundaram, *Typical Means*, Oxford University Press, Oxford, UK, 1952.
- [15] M. Csörgő, B. Szyszkowicz, and Q. Wang, "Donsker's theorem for self-normalized partial sums processes," *The Annals of Probability*, vol. 31, no. 3, pp. 1228–1240, 2003.
- [16] Q. Y. Wu, "An almost sure central limit theorem for the weight function sequences of NA random variables," *Proceedings-Mathematical Sciences*, vol. 121, no. 3, pp. 369–377, 2011.
- [17] P. Billingsley, *Convergence of Probability Measures*, John Wiley & Sons, New York, NY, USA, 1968.
- [18] M. Peligrad and Q. M. Shao, "A note on the almost sure central limit theorem for weakly dependent random variables," *Statistics & Probability Letters*, vol. 22, no. 2, pp. 131–136, 1995.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

