

Research Article

Approximate Solutions of Fractional Riccati Equations Using the Adomian Decomposition Method

Fei Wu¹ and Lan-Lan Huang²

¹Key Laboratory of Energy Engineering Safety and Disaster Mechanics, Ministry of Education, School of Architecture and Environment, Sichuan University, Chengdu 610065, China

²Institute of Applied Nonlinear Science, College of Mathematics and Information Science, Neijiang Normal University, Neijiang 641112, China

Correspondence should be addressed to Lan-Lan Huang; mathlan@126.com

Received 8 June 2014; Accepted 29 August 2014; Published 14 December 2014

Academic Editor: Guotao Wang

Copyright © 2014 F. Wu and L.-L. Huang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The fractional derivative equation has extensively appeared in various applied nonlinear problems and methods for finding the model become a popular topic. Very recently, a novel way was proposed by Duan (2010) to calculate the Adomian series which is a crucial step of the Adomian decomposition method. In this paper, it was used to solve some fractional nonlinear differential equations.

1. Introduction

The fractional derivative has good memory effects compared with the ordinary calculus. In view of this point, it has been proven as a good tool in nonlinear science, for example, the anomalous diffusion [1, 2], the material's viscoelasticity [3, 4], the chaotic behaviors of biology population [5, 6], and so forth.

However, everything has two sides. The fractional derivative's memory effects also lead to the numerical solutions' accumulative errors. Many nonlinear techniques cannot perform the same role as those in ordinary differential equation. For example, the variational iteration method cannot be applied due to the fact that the integral by parts cannot hold and the Lagrange multipliers there are not easily identified; in the Adomian decomposition method, the Adomian series cannot be expanded large enough which greatly affects the solutions' accuracies and even five- or six-order approximation becomes impossible (see the analysis in [7]).

Very recently and fortunately, for the ADM, Duan [8, 9] proposed a convenient way to calculate Adomian series which is the main and crucial step of the classical ADM

developed by Adomian. This way can rapidly decompose the nonlinear terms and some new high order approximation schemes for nonlinear differential equations are proposed [10]. The technique has been successfully extended to fractional differential equations and boundary value problems [11, 12].

In this paper, we investigate the following fractional nonlinear differential equation:

$${}_0^C D_t^\alpha x = 1 + 2x(t) - x(t)^2, \quad 0 < \alpha \leq 2, \quad (1)$$

where ${}_0^C D_t^\alpha x$ is the Caputo derivative with respect to $x(t)$.

The paper is organized as follows: Section 2 introduces some basics of the ADM and the fractional calculus; Section 3 considers the differential equation from the case $\alpha = 1$ in (1)

$$\frac{dx}{dt} = 1 + 2x(t) - x(t)^2, \quad x(0) = 0 \quad (2)$$

and gives the analytical formula.

2. Preliminaries

Definition 1 (see [13]). The Caputo derivative is defined as

$${}_0^C D_t^\alpha u = \frac{1}{\Gamma(m-\alpha)} \int_0^t \frac{1}{(t-\tau)^{\alpha-m+1}} \frac{d^m}{d\tau^m} u(\tau) d\tau, \tag{3}$$

$$0 < t, \quad 0 < \alpha, \quad m = [\alpha] + 1,$$

where Γ is the Gamma function.

Definition 2 (see [14]). The R-L integration of α order is defined by

$${}_0^I_t^\alpha u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} u(\tau) d\tau, \tag{4}$$

$$0 < t, \quad 0 < \alpha.$$

Generally, consider the following nonlinear equation:

$$L[u] + R[u] + N[u] = g(t), \tag{5}$$

where L is the highest derivative, R is the remaining linear part containing the lower order derivatives, and N is the nonlinear operator.

Apply the inverse L^{-1} of the linear operator L in (5) and we can obtain

$$u = u(0) + u'(0)t + \dots + u^{(m-1)}(0) \frac{t^{m-1}}{(m-1)!} + L^{-1}(g(t) - R[u] - N[u]). \tag{6}$$

Assume that

$$u = \sum_{i=0}^{\infty} u_i \tag{7}$$

and expand the term $N[u]$ approximately as

$$N[u] = \sum_{n=0}^{\infty} A_n, \tag{8}$$

where the A_n is calculated by

$$A_n = \frac{1}{n} \sum_{k=0}^{n-1} (k+1) u_{k+1} \frac{dA_{n-1-k}}{du_0}. \tag{9}$$

As a result, one can obtain the analytical iteration scheme as

$$u_{n+1} = -L^{-1}(R[u_n]) - L^{-1}(A_n[u_0, u_1, \dots, u_n]),$$

$$u_0 = u(0) + u'(0)t + \dots + u^{(m-1)}(0) \frac{t^{m-1}}{(m-1)!} + L^{-1}(g(t)). \tag{10}$$

Here we conclude the application of the ADM: firstly, one needs to have equivalent integral form of the original governing equations; second, decompose the nonlinear terms




FIGURE 1: “*” the 20th order approximate solution; “-” the explicit solution.

into linear ones and determine the iteration schemes; third, obtain the solutions successively. For the other applications and modified versions, we do not introduce any more here. The readers who feel interested in the development of the method are referred to [7, 12, 15–19].

3. Numerical Schemes of Integer Equations

According to the ADM, we first establish the integral equation as the one

$$x(t) = x(0) + \int_0^t (1 + 2x(\tau) - x(\tau)^2) d\tau. \tag{11}$$

If we directly use the classical ADM’s idea we can obtain the formula

$$x_{n+1} = 2L^{-1}(x_n) + L^{-1}(A_n[x_0, x_1, \dots, x_n]),$$

$$x(t) = \sum_{i=0}^{\infty} x_i. \tag{12}$$

The explicit solution of (2) was found to be [20]

$$x(t) = 1 + \sqrt{2} \tanh\left(\sqrt{2}t + \frac{1}{2} \log\left(\frac{\sqrt{2}-1}{\sqrt{2}+1}\right)\right). \tag{13}$$

We can assume that

$$A_0 = -x_0^2,$$

$$x_0 = x(0) + t. \tag{14}$$

As a result, from the iteration equations (10) and (12), we can obtain the approximate solutions in comparison with (13).

We find that the approximate solution has a good agreement with the explicit solution in Figure 1. So the scheme is efficient and useful.




FIGURE 2: “-” the residual function.




FIGURE 3: “-” the residual function.

4. Numerical Schemes of Fractional Equations

Case 1 ($0 < \alpha \leq 1$). Similarly, we can have the integral equation for (1):

$$x(t) = x(0) + \int_0^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} (1 + 2x(\tau) - x(\tau)^2) d\tau, \quad (15)$$

$0 < \alpha \leq 1.$

If we directly use the classical ADM’s idea we can obtain the formula

$$x_{n+1} = \int_0^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} A_n [x_0, x_1, \dots, x_n] d\tau, \quad (16)$$

$$x(t) = \sum_{i=0}^{\infty} x_i.$$

Here $x_0 = x(0)$, we give the first solutions as

$$x_0(t) = t + \frac{t^\alpha}{\Gamma(\alpha)\alpha}. \quad (17)$$

For simplicity, we define the residual function as

$$\text{err} = \ln \left| {}^C_0 D_t^\alpha x_n = 1 + 2x_n(t) - x_n(t)^2 \right| \quad (18)$$

and illustrate the errors for $n = 30$ and $\alpha = 0.9$ in Figure 2.

Case 2 ($1 < \alpha \leq 2$). Consider the following.

$$x(t) = x(0) + x'(0)t + \int_0^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} (1 + 2x(\tau) - x(\tau)^2) d\tau, \quad (19)$$

$1 < \alpha \leq 2.$

Here we need two initial conditions $x(0)$ and $x'(0)$. We set

$$x(0) = 0, \quad x'(0) = 1. \quad (20)$$

Similarly, we can have the iteration formula as

$$x_{n+1} = \int_0^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} A_n [x_0, x_1, \dots, x_n] d\tau,$$

$$x_0 = x(0) + x'(0)t = t + \frac{t^\alpha}{\Gamma(\alpha)\alpha}, \quad (21)$$

$$x(t) = \sum_{i=0}^{\infty} x_i,$$

whose error analysis is given in Figure 3. We can see that the approximate solution is reliable.

We plot the approximate solution in Figure 4 in the interval $[0, 1]$.

5. Conclusions

This paper applied the famous Adomian decomposition method to FDEs whose fractional order varied from 0 to 2. First, we revisit the method for integer equation. We conclude the general steps and use the method to solve FDEs analytically. Through the analysis of the residual function, it can be concluded that the ADM using Duan’s way to calculate the Adomian series is very suitable and efficient for obtaining analytical solutions of fractional nonlinear differential equations.




FIGURE 4: “*” the 30th order approximate solution of (19).

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

This work is financially supported by the National Natural Science Foundation (Grant nos. 51120145001, 51374148, and 51104101).

References

- [1] R. Metzler and J. Klafter, “The random walk’s guide to anomalous diffusion: a fractional dynamics approach,” *Physics Reports A: Review Section of Physics Letters*, vol. 339, no. 1, pp. 1–77, 2000.
- [2] H. Sun, W. Chen, and Y. Chen, “Variable-order fractional differential operators in anomalous diffusion modeling,” *Physica A: Statistical Mechanics and its Applications*, vol. 388, no. 21, pp. 4586–4592, 2009.
- [3] R. L. Bagley and P. J. Torvik, “A theoretical basis for the application of fractional calculus to viscoelasticity,” *Journal of Rheology*, vol. 27, no. 3, pp. 201–210, 1983.
- [4] R. L. Bagley and P. J. Torvik, “Fractional calculus—a different approach to the analysis of viscoelastically damped structures,” *AIAA Journal*, vol. 21, no. 5, pp. 741–748, 1983.
- [5] G.-C. Wu and D. Baleanu, “Discrete fractional logistic map and its chaos,” *Nonlinear Dynamics*, vol. 75, no. 1–2, pp. 283–287, 2014.
- [6] G.-C. Wu and D. Baleanu, “Chaos synchronization of the discrete fractional logistic map,” *Signal Processing*, vol. 102, pp. 96–99, 2014.
- [7] D. Baleanu, G.-C. Wu, and J.-S. Duan, “Some analytical techniques in fractional calculus: realities and challenges,” in *Discontinuity and Complexity in Nonlinear Physical Systems*, D. B. J. A. Tenreiro Machado and A. C. J. Luo, Eds., pp. 35–62, Springer, New York, NY, USA, 2014.
- [8] J.-S. Duan, “Recurrence triangle for Adomian polynomials,” *Applied Mathematics and Computation*, vol. 216, no. 4, pp. 1235–1241, 2010.
- [9] J.-S. Duan, “Convenient analytic recurrence algorithms for the Adomian polynomials,” *Applied Mathematics and Computation*, vol. 217, no. 13, pp. 6337–6348, 2011.
- [10] J.-S. Duan and R. Rach, “New higher-order numerical one-step methods based on the Adomian and the modified decomposition methods,” *Applied Mathematics and Computation*, vol. 218, no. 6, pp. 2810–2828, 2011.
- [11] J.-S. Duan, T. Chaolu, and R. Rach, “Solutions of the initial value problem for nonlinear fractional ordinary differential equations by the Rach-Adomian-Meyers modified decomposition method,” *Applied Mathematics and Computation*, vol. 218, no. 17, pp. 8370–8392, 2012.
- [12] J. S. Duan, R. Rach, D. Baleanu, and A.-M. Wazwaz, “A review of the Adomian decomposition method and its applications to fractional differential equations,” *Communications in Fractional Calculus*, vol. 3, no. 2, pp. 73–99, 2012.
- [13] M. Caputo, “Linear models of dissipation whose Q is almost frequency independent—II,” *Geophysical Journal International*, vol. 13, no. 5, pp. 529–539, 1967.
- [14] I. Podlubny, *Fractional Differential Equations*, Academic Press, 1999.
- [15] V. Daftardar-Gejji and H. Jafari, “Adomian decomposition: a tool for solving a system of fractional differential equations,” *Journal of Mathematical Analysis and Applications*, vol. 301, no. 2, pp. 508–518, 2005.
- [16] S. S. Ray and R. K. Bera, “An approximate solution of a nonlinear fractional differential equation by Adomian decomposition method,” *Applied Mathematics and Computation*, vol. 167, no. 1, pp. 561–571, 2005.
- [17] M. G. Sakar and F. Erdogan, “The homotopy analysis method for solving the time-fractional Fornberg-Whitham equation and comparison with Adomian’s decomposition method,” *Applied Mathematical Modelling*, vol. 37, no. 20–21, pp. 8876–8885, 2013.
- [18] V. Daftardar-Gejji and H. Jafari, “Solving a multi-order fractional differential equation using Adomian decomposition,” *Applied Mathematics and Computation*, vol. 189, no. 1, pp. 541–548, 2007.
- [19] C. Li and Y. Wang, “Numerical algorithm based on Adomian decomposition for fractional differential equations,” *Computers and Mathematics with Applications*, vol. 57, no. 10, pp. 1672–1681, 2009.
- [20] M. A. El-Tawil, A. A. Bahnasawi, and A. Abdel-Naby, “Solving Riccati differential equation using Adomian’s decomposition method,” *Applied Mathematics and Computation*, vol. 157, no. 2, pp. 503–514, 2004.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

