

## INTUITIONISTIC FUZZY IDEALS OF BCK-ALGEBRAS

YOUNG BAE JUN and KYUNG HO KIM

(Received 16 February 2000)

**ABSTRACT.** We consider the intuitionistic fuzzification of the concept of subalgebras and ideals in BCK-algebras, and investigate some of their properties. We introduce the notion of equivalence relations on the family of all intuitionistic fuzzy ideals of a BCK-algebra and investigate some related properties.

**Keywords and phrases.** (Intuitionistic) fuzzy subalgebra, (intuitionistic) fuzzy ideal, upper (respectively, lower)  $t$ -level cut, homomorphism.

2000 Mathematics Subject Classification. Primary 06F35, 03G25, 03E72.

**1. Introduction.** After the introduction of the concept of fuzzy sets by Zadeh [9] several researches were conducted on the generalizations of the notion of fuzzy sets. The idea of “intuitionistic fuzzy set” was first published by Atanassov [1, 2], as a generalization of the notion of fuzzy set. The first author (together with Hong, Kim, Kim, Meng, Roh, and Song) considered the fuzzification of ideals and subalgebras in BCK-algebras (cf. [3, 4, 5, 6, 7, 8]). In this paper, using the Atanassov’s idea, we establish the intuitionistic fuzzification of the concept of subalgebras and ideals in BCK-algebras, and investigate some of their properties. We introduce the notion of equivalence relations on the family of all intuitionistic fuzzy ideals of a BCK-algebra and investigate some related properties.

**2. Preliminaries.** First we present the fundamental definitions. By a *BCK-algebra* we mean a nonempty set  $X$  with a binary operation  $*$  and a constant  $0$  satisfying the following conditions:

- (I)  $((x * y) * (x * z)) * (z * y) = 0$ ,
- (II)  $(x * (x * y)) * y = 0$ ,
- (III)  $x * x = 0$ ,
- (IV)  $0 * x = 0$ ,
- (V)  $x * y = 0$  and  $y * x = 0$  imply that  $x = y$

for all  $x, y, z \in X$ .

A partial ordering “ $\leq$ ” on  $X$  can be defined by  $x \leq y$  if and only if  $x * y = 0$ . A nonempty subset  $S$  of a BCK-algebra  $X$  is called a *subalgebra* of  $X$  if  $x * y \in S$  whenever  $x, y \in S$ . A nonempty subset  $I$  of a BCK-algebra  $X$  is called an *ideal* of  $X$  if

- (i)  $0 \in I$ ,
- (ii)  $x * y \in I$  and  $y \in I$  imply that  $x \in I$  for all  $x, y \in X$ .

By a *fuzzy set*  $\mu$  in a nonempty set  $X$  we mean a function  $\mu : X \rightarrow [0, 1]$ , and the complement of  $\mu$ , denoted by  $\bar{\mu}$ , is the fuzzy set in  $X$  given by  $\bar{\mu}(x) = 1 - \mu(x)$  for all  $x \in X$ . A fuzzy set  $\mu$  in a BCK-algebra  $X$  is called a *fuzzy subalgebra* of  $X$  if  $\mu(x * y) \geq$

$\min\{\mu(x), \mu(y)\}$  for all  $x, y \in X$ . A fuzzy set  $\mu$  in a BCK-algebra  $X$  is called a *fuzzy ideal* of  $X$  if

- (i)  $\mu(0) \geq \mu(x)$  for all  $x \in X$ ,
- (ii)  $\mu(x) \geq \min\{\mu(x * y), \mu(y)\}$  for all  $x, y \in X$ .

An intuitionistic fuzzy set (briefly, IFS)  $A$  in a nonempty set  $X$  is an object having the form

$$A = \{(x, \alpha_A(x), \beta_A(x)) \mid x \in X\}, \tag{2.1}$$

where the functions  $\alpha_A : X \rightarrow [0, 1]$  and  $\beta_A : X \rightarrow [0, 1]$  denote the degree of membership and the degree of nonmembership, respectively, and

$$0 \leq \alpha_A(x) + \beta_A(x) \leq 1 \quad \forall x \in X. \tag{2.2}$$

An intuitionistic fuzzy set  $A = \{(x, \alpha_A(x), \beta_A(x)) \mid x \in X\}$  in  $X$  can be identified to an ordered pair  $(\alpha_A, \beta_A)$  in  $I^X \times I^X$ . For the sake of simplicity, we shall use the symbol  $A = (\alpha_A, \beta_A)$  for the IFS  $A = \{(x, \alpha_A(x), \beta_A(x)) \mid x \in X\}$ .

**3. Intuitionistic fuzzy ideals.** In what follows, let  $X$  denote a BCK-algebra unless otherwise specified.

**DEFINITION 3.1.** An IFS  $A = (\alpha_A, \beta_A)$  in  $X$  is called an *intuitionistic fuzzy subalgebra* of  $X$  if it satisfies:

- (IS1)  $\alpha_A(x * y) \geq \min\{\alpha_A(x), \alpha_A(y)\}$ ,
- (IS2)  $\beta_A(x * y) \leq \max\{\beta_A(x), \beta_A(y)\}$ ,

for all  $x, y \in X$ .

**EXAMPLE 3.2.** Consider a BCK-algebra  $X = \{0, a, b, c\}$  with the following Cayley table:

*	0	a	b	c
0	0	0	0	0
a	a	0	0	a
b	b	a	0	b
c	c	c	c	0

Let  $A = (\alpha_A, \beta_A)$  be an IFS in  $X$  defined by

$$\begin{aligned} \alpha_A(0) = \alpha_A(a) = \alpha_A(c) = 0.7 > 0.3 = \alpha_A(b), \\ \beta_A(0) = \beta_A(a) = \beta_A(c) = 0.2 < 0.5 = \beta_A(b). \end{aligned} \tag{3.1}$$

Then  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy subalgebra of  $X$ .

**PROPOSITION 3.3.** Every intuitionistic fuzzy subalgebra  $A = (\alpha_A, \beta_A)$  of  $X$  satisfies the inequalities  $\alpha_A(0) \geq \alpha_A(x)$  and  $\beta_A(0) \leq \beta_A(x)$  for all  $x \in X$ .

**PROOF.** For any  $x \in X$ , we have

$$\begin{aligned} \alpha_A(0) = \alpha_A(x * x) &\geq \min\{\alpha_A(x), \alpha_A(x)\} = \alpha_A(x), \\ \beta_A(0) = \beta_A(x * x) &\leq \max\{\beta_A(x), \beta_A(x)\} = \beta_A(x). \end{aligned} \tag{3.2}$$

This completes the proof. □

**DEFINITION 3.4.** An IFS  $A = (\alpha_A, \beta_A)$  in  $X$  is called an *intuitionistic fuzzy ideal* of  $X$  if it satisfies the following inequalities:

(IF1)  $\alpha_A(0) \geq \alpha_A(x)$  and  $\beta_A(0) \leq \beta_A(x)$ ,

(IF2)  $\alpha_A(x) \geq \min\{\alpha_A(x * y), \alpha_A(y)\}$ ,

(IF3)  $\beta_A(x) \leq \max\{\beta_A(x * y), \beta_A(y)\}$ ,

for all  $x, y \in X$ .

**EXAMPLE 3.5.** Let  $X = \{0, 1, 2, 3, 4\}$  be a BCK-algebra with the following Cayley table:

*	0	1	2	3	4
0	0	0	0	0	0
1	1	0	1	0	0
2	2	2	0	0	0
3	3	3	3	0	0
4	4	3	4	1	0

Define an IFS  $A = (\alpha_A, \beta_A)$  in  $X$  as follows:

$$\begin{aligned} \alpha_A(0) = \alpha_A(2) = 1, & \quad \alpha_A(1) = \alpha_A(3) = \alpha_A(4) = t, \\ \beta_A(0) = \beta_A(2) = 0, & \quad \beta_A(1) = \beta_A(3) = \beta_A(4) = s, \end{aligned} \tag{3.3}$$

where  $t \in [0, 1]$ ,  $s \in [0, 1]$ , and  $t + s \leq 1$ . By routine calculation we know that  $A = (\alpha_A, \beta_A)$  is an *intuitionistic fuzzy ideal* of  $X$ .

**LEMMA 3.6.** Let an IFS  $A = (\alpha_A, \beta_A)$  in  $X$  be an intuitionistic fuzzy ideal of  $X$ . If the inequality  $x * y \leq z$  holds in  $X$ , then

$$\alpha_A(x) \geq \min\{\alpha_A(y), \alpha_A(z)\}, \quad \beta_A(x) \leq \max\{\beta_A(y), \beta_A(z)\}. \tag{3.4}$$

**PROOF.** Let  $x, y, z \in X$  be such that  $x * y \leq z$ . Then  $(x * y) * z = 0$ , and thus

$$\begin{aligned} \alpha_A(x) &\geq \min\{\alpha_A(x * y), \alpha_A(y)\} \\ &\geq \min\{\min\{\alpha_A((x * y) * z), \alpha_A(z)\}, \alpha_A(y)\} \\ &= \min\{\min\{\alpha_A(0), \alpha_A(z)\}, \alpha_A(y)\} \\ &= \min\{\alpha_A(y), \alpha_A(z)\}, \\ \beta_A(x) &\leq \max\{\beta_A(x * y), \beta_A(y)\} \\ &\leq \max\{\max\{\beta_A((x * y) * z), \beta_A(z)\}, \beta_A(y)\} \\ &= \max\{\max\{\beta_A(0), \beta_A(z)\}, \beta_A(y)\} \\ &= \max\{\beta_A(y), \beta_A(z)\}, \end{aligned} \tag{3.5}$$

this completes the proof. □

**LEMMA 3.7.** Let  $A = (\alpha_A, \beta_A)$  be an intuitionistic fuzzy ideal of  $X$ . If  $x \leq y$  in  $X$ , then

$$\alpha_A(x) \geq \alpha_A(y), \quad \beta_A(x) \leq \beta_A(y), \tag{3.6}$$

that is,  $\alpha_A$  is order-reserving and  $\beta_A$  is order-preserving.

**PROOF.** Let  $x, y \in X$  be such that  $x \leq y$ . Then  $x * y = 0$  and so

$$\begin{aligned} \alpha_A(x) &\geq \min \{ \alpha_A(x * y), \alpha_A(y) \} = \min \{ \alpha_A(0), \alpha_A(y) \} = \alpha_A(y), \\ \beta_A(x) &\leq \max \{ \beta_A(x * y), \beta_A(y) \} = \max \{ \beta_A(0), \beta_A(y) \} = \beta_A(y). \end{aligned} \tag{3.7}$$

This completes the proof. □

**THEOREM 3.8.** *If  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$ , then for any  $x, a_1, a_2, \dots, a_n \in X$ ,  $(\dots((x * a_1) * a_2) * \dots) * a_n = 0$  implies*

$$\begin{aligned} \alpha_A(x) &\geq \min \{ \alpha_A(a_1), \alpha_A(a_2), \dots, \alpha_A(a_n) \}, \\ \beta_A(x) &\leq \max \{ \beta_A(a_1), \beta_A(a_2), \dots, \beta_A(a_n) \}. \end{aligned} \tag{3.8}$$

**PROOF.** Using induction on  $n$  and Lemmas 3.6 and 3.7, the proof is straightforward. □

**THEOREM 3.9.** *Every intuitionistic fuzzy ideal of  $X$  is an intuitionistic fuzzy subalgebra of  $X$ .*

**PROOF.** Let  $A = (\alpha_A, \beta_A)$  be an intuitionistic fuzzy ideal of  $X$ . Since  $x * y \leq x$  for all  $x, y \in X$ , it follows from Lemma 3.7 that

$$\alpha_A(x * y) \geq \alpha_A(x), \quad \beta_A(x * y) \leq \beta_A(x), \tag{3.9}$$

so by (IF2) and (IF3),

$$\begin{aligned} \alpha_A(x * y) &\geq \alpha_A(x) \geq \min \{ \alpha_A(x * y), \alpha_A(y) \} \geq \min \{ \alpha_A(x), \alpha_A(y) \}, \\ \beta_A(x * y) &\leq \beta_A(x) \leq \max \{ \beta_A(x * y), \beta_A(y) \} \leq \max \{ \beta_A(x), \beta_A(y) \}. \end{aligned} \tag{3.10}$$

This shows that  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy subalgebra of  $X$ . □

The converse of Theorem 3.9 may not be true. For example, the intuitionistic fuzzy subalgebra  $A = (\alpha_A, \beta_A)$  in Example 3.2 is not an intuitionistic fuzzy ideal of  $X$  since

$$\beta_A(b) = 0.5 > 0.2 = \min \{ \beta_A(b * a), \beta_A(a) \}. \tag{3.11}$$

We now give a condition for an intuitionistic fuzzy subalgebra to be an intuitionistic fuzzy ideal.

**THEOREM 3.10.** *Let  $A = (\alpha_A, \beta_A)$  be an intuitionistic fuzzy subalgebra of  $X$  such that*

$$\alpha_A(x) \geq \min \{ \alpha_A(y), \alpha_A(z) \}, \quad \beta_A(x) \leq \max \{ \beta_A(y), \beta_A(z) \} \tag{3.12}$$

*for all  $x, y, z \in X$  satisfying the inequality  $x * y \leq z$ . Then  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$ .*

**PROOF.** Let  $A = (\alpha_A, \beta_A)$  be an intuitionistic fuzzy subalgebra of  $X$ . Recall that  $\alpha_A(0) \geq \alpha_A(x)$  and  $\beta_A(0) \leq \beta_A(x)$  for all  $X$ . Since  $x * (x * y) \leq y$ , it follows from the hypothesis that

$$\alpha_A(x) \geq \min \{ \alpha_A(x * y), \alpha_A(y) \}, \quad \beta_A(x) \leq \max \{ \beta_A(x * y), \beta_A(y) \}. \tag{3.13}$$

Hence  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$ . □

**LEMMA 3.11.** *An IFS  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$  if and only if the fuzzy sets  $\alpha_A$  and  $\tilde{\beta}_A$  are fuzzy ideals of  $X$ .*

**PROOF.** Let  $A = (\alpha_A, \beta_A)$  be an intuitionistic fuzzy ideal of  $X$ . Clearly,  $\alpha_A$  is a fuzzy ideal of  $X$ . For every  $x, y \in X$ , we have

$$\begin{aligned} \tilde{\beta}_A(0) &= 1 - \beta_A(0) \geq 1 - \beta_A(x) = \tilde{\beta}_A(x), \\ \tilde{\beta}_A(x) &= 1 - \beta_A(x) \geq 1 - \max\{\beta_A(x * y), \beta_A(y)\} \\ &= \min\{1 - \beta_A(x * y), 1 - \beta_A(y)\} \\ &= \min\{\tilde{\beta}_A(x * y), \tilde{\beta}_A(y)\}. \end{aligned} \tag{3.14}$$

Hence  $\tilde{\beta}_A$  is a fuzzy ideal of  $X$ .

Conversely, assume that  $\alpha_A$  and  $\tilde{\beta}_A$  are fuzzy ideals of  $X$ . For every  $x, y \in X$ , we get

$$\alpha_A(0) \geq \alpha_A(x), \quad 1 - \beta_A(0) = \tilde{\beta}_A(0) \geq \tilde{\beta}_A(x) = 1 - \beta_A(x), \tag{3.15}$$

that is,  $\beta_A(0) \leq \beta_A(x)$ ;  $\alpha_A(x) \geq \min\{\alpha_A(x * y), \alpha_A(y)\}$  and

$$\begin{aligned} 1 - \beta_A(x) &= \tilde{\beta}_A(x) \geq \min\{\tilde{\beta}_A(x * y), \tilde{\beta}_A(y)\} \\ &= \min\{1 - \beta_A(x * y), 1 - \beta_A(y)\} \\ &= 1 - \max\{\beta_A(x * y), \beta_A(y)\}, \end{aligned} \tag{3.16}$$

that is,  $\beta_A(x) \leq \max\{\beta_A(x * y), \beta_A(y)\}$ . Hence  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$ . □

**THEOREM 3.12.** *Let  $A = (\alpha_A, \beta_A)$  be an IFS in  $X$ . Then  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$  if and only if  $\square A = (\alpha_A, \tilde{\alpha}_A)$  and  $\diamond A = (\tilde{\beta}_A, \beta_A)$  are intuitionistic fuzzy ideals of  $X$ .*

**PROOF.** If  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$ , then  $\alpha_A = \tilde{\alpha}_A$  and  $\beta_A$  are fuzzy ideals of  $X$  from Lemma 3.11, hence  $\square A = (\alpha_A, \tilde{\alpha}_A)$  and  $\diamond A = (\tilde{\beta}_A, \beta_A)$  are intuitionistic fuzzy ideals of  $X$ . Conversely, if  $\square A = (\alpha_A, \tilde{\alpha}_A)$  and  $\diamond A = (\tilde{\beta}_A, \beta_A)$  are intuitionistic fuzzy ideals of  $X$ , then the fuzzy sets  $\alpha_A$  and  $\tilde{\beta}_A$  are fuzzy ideals of  $X$ , hence  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$ . □

For any  $t \in [0, 1]$  and a fuzzy set  $\mu$  in a nonempty set  $X$ , the set

$$U(\mu; t) = \{x \in X \mid \mu(x) \geq t\} \tag{3.17}$$

is called an *upper  $t$ -level cut* of  $\mu$  and the set

$$L(\mu; t) = \{x \in X \mid \mu(x) \leq t\} \tag{3.18}$$

is called a *lower  $t$ -level cut* of  $\mu$ .

**THEOREM 3.13.** *An IFS  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $X$  if and only if for all  $s, t \in [0, 1]$ , the sets  $U(\alpha_A; t)$  and  $L(\beta_A; s)$  are either empty or ideals of  $X$ .*

**PROOF.** Let  $A = (\alpha_A, \beta_A)$  be an intuitionistic fuzzy ideal of  $X$  and  $U(\alpha_A; t) \neq \emptyset \neq L(\beta_A; s)$  for any  $s, t \in [0, 1]$ . It is clear that  $0 \in U(\alpha_A; t) \cap L(\beta_A; s)$  since  $\alpha_A(0) \geq t$  and  $\beta_A(0) \leq s$ . Let  $x, y \in X$  be such that  $x * y \in U(\alpha_A; t)$  and  $y \in U(\alpha_A; t)$ . Then  $\alpha_A(x * y) \geq t$  and  $\alpha_A(y) \geq t$ . It follows that

$$\alpha_A(x) \geq \min \{ \alpha_A(x * y), \alpha_A(y) \} \geq t \tag{3.19}$$

so that  $x \in U(\alpha_A; t)$ . Hence  $U(\alpha_A; t)$  is an ideal of  $X$ . Now let  $x, y \in X$  be such that  $x * y \in L(\beta_A; s)$  and  $y \in L(\beta_A; s)$ . Then  $\beta_A(x * y) \leq s$  and  $\beta_A(y) \leq s$ , which imply that

$$\beta_A(x) \leq \max \{ \beta_A(x * y), \beta_A(y) \} \leq s. \tag{3.20}$$

Thus  $x \in L(\beta_A; s)$ , and therefore  $L(\beta_A; s)$  is an ideal of  $X$ . Conversely, assume that for each  $t, s \in [0, 1]$ , the sets  $U(\alpha_A; t)$  and  $L(\beta_A; s)$  are either empty or ideals of  $X$ . For any  $x \in X$ , let  $\alpha_A(x) = t$  and  $\beta_A(x) = s$ . Then  $x \in U(\alpha_A; t) \cap L(\beta_A; s)$ , and so  $U(\alpha_A; t) \neq \emptyset \neq L(\beta_A; s)$ . Since  $U(\alpha_A; t)$  and  $L(\beta_A; s)$  are ideals of  $X$ , therefore  $0 \in U(\alpha_A; t) \cap L(\beta_A; s)$ . Hence  $\alpha_A(0) \geq t = \alpha_A(x)$  and  $\beta_A(0) \leq s = \beta_A(x)$  for all  $x \in X$ . If there exist  $x', y' \in X$  such that  $\alpha_A(x') < \min \{ \alpha_A(x' * y'), \alpha_A(y') \}$ , then by taking

$$t_0 = \frac{1}{2} (\alpha_A(x') + \min \{ \alpha_A(x' * y'), \alpha_A(y') \}), \tag{3.21}$$

we have

$$\alpha_A(x') < t_0 < \min \{ \alpha_A(x' * y'), \alpha_A(y') \}. \tag{3.22}$$

Hence  $x' \notin U(\alpha_A; t_0)$ ,  $x' * y' \in U(\alpha_A; t_0)$  and  $y' \in U(\alpha_A; t_0)$ , that is,  $U(\alpha_A; t_0)$  is not an ideal of  $X$ , which is a contradiction. Finally, assume that there exist  $a, b \in X$  such that

$$\beta_A(a) > \max \{ \beta_A(a * b), \beta_A(b) \}. \tag{3.23}$$

Taking  $s_0 := (1/2)(\beta_A(a) + \max \{ \beta_A(a * b), \beta_A(b) \})$ , then

$$\max \{ \beta_A(a * b), \beta_A(b) \} < s_0 < \beta_A(a). \tag{3.24}$$

Therefore  $a * b \in L(\beta_A; s_0)$  and  $b \in L(\beta_A; s_0)$ , but  $a \notin L(\beta_A; s_0)$ , which is a contradiction, this completes the proof. □

Let  $\Lambda$  be a nonempty subset of  $[0, 1]$ .

**THEOREM 3.14.** *Let  $\{I_t \mid t \in \Lambda\}$  be a collection of ideals of  $X$  such that*

- (i)  $X = \cup_{t \in \Lambda} I_t$ ,
- (ii)  $s > t$  if and only if  $I_s \subset I_t$  for all  $s, t \in \Lambda$ .

*Then an IFSA  $A = (\alpha_A, \beta_A)$  in  $X$  defined by*

$$\alpha_A(x) := \sup \{ t \in \Lambda \mid x \in I_t \}, \quad \beta_A(x) := \inf \{ t \in \Lambda \mid x \in I_t \} \tag{3.25}$$

*for all  $x \in X$  is an intuitionistic fuzzy ideal of  $X$ .*

**PROOF.** According to Theorem 3.13, it is sufficient to show that  $U(\alpha_A; t)$  and  $L(\beta_A; s)$  are ideals of  $X$  for every  $t \in [0, \alpha_A(0)]$  and  $s \in [\beta_A(0), 1]$ . In order to prove

that  $U(\alpha_A; t)$  is an ideal of  $X$ , we divide the proof into the following two cases:

- (i)  $t = \sup\{q \in \Lambda \mid q < t\}$ ,
- (ii)  $t \neq \sup\{q \in \Lambda \mid q < t\}$ .

Case (i) implies that

$$x \in U(\alpha_A; t) \iff x \in I_q \quad \forall q < t \iff x \in \bigcap_{q < t} I_q, \tag{3.26}$$

so that  $U(\alpha_A; t) = \bigcap_{q < t} I_q$ , which is an ideal of  $X$ . For the case (ii), we claim that  $U(\alpha_A; t) = \bigcup_{q \geq t} I_q$ . If  $x \in \bigcup_{q \geq t} I_q$ , then  $x \in I_q$  for some  $q \geq t$ . It follows that  $\alpha_A(x) \geq q \geq t$ , so that  $x \in U(\alpha_A; t)$ . This shows that  $\bigcup_{q \geq t} I_q \subseteq U(\alpha_A; t)$ . Now assume that  $x \notin \bigcup_{q \geq t} I_q$ . Then  $x \notin I_q$  for all  $q \geq t$ . Since  $t \neq \sup\{q \in \Lambda \mid q < t\}$ , there exists  $\varepsilon > 0$  such that  $(t - \varepsilon, t) \cap \Lambda = \emptyset$ . Hence  $x \notin I_q$  for all  $q > t - \varepsilon$ , which means that if  $x \in I_q$ , then  $q \leq t - \varepsilon$ . Thus  $\alpha_A(x) \leq t - \varepsilon < t$ , and so  $x \notin U(\alpha_A; t)$ . Therefore  $U(\alpha_A; t) \subseteq \bigcup_{q \geq t} I_q$ , and thus  $U(\alpha_A; t) = \bigcup_{q \geq t} I_q$  which is an ideal of  $X$ . Next we prove that  $L(\beta_A; s)$  is an ideal of  $X$ . We consider the following two cases:

- (iii)  $s = \inf\{r \in \Lambda \mid s < r\}$ ,
- (iv)  $s \neq \inf\{r \in \Lambda \mid s < r\}$ .

For the case (iii), we have

$$x \in L(\beta_A; s) \iff x \in I_r \quad \forall s < r \iff x \in \bigcap_{s < r} I_r, \tag{3.27}$$

and hence  $L(\beta_A; s) = \bigcap_{s < r} I_r$  which is an ideal of  $X$ . For the case (iv) there exists  $\varepsilon > 0$  such that  $(s, s + \varepsilon) \cap \Lambda = \emptyset$ . We will show that  $L(\beta_A; s) = \bigcup_{s \geq r} I_r$ . If  $x \in \bigcup_{s \geq r} I_r$ , then  $x \in I_r$  for some  $r \leq s$ . It follows that  $\beta_A(x) \leq r \leq s$  so that  $x \in L(\beta_A; s)$ . Hence  $\bigcup_{s \geq r} I_r \subseteq L(\beta_A; s)$ . Conversely, if  $x \notin \bigcup_{s \geq r} I_r$ , then  $x \notin I_r$  for all  $r \leq s$ , which implies that  $x \notin I_r$  for all  $r < s + \varepsilon$ , that is, if  $x \in I_r$ , then  $r \geq s + \varepsilon$ . Thus  $\beta_A(x) \geq s + \varepsilon > s$ , that is,  $x \notin L(\beta_A; s)$ . Therefore  $L(\beta_A; s) \subseteq \bigcup_{s \geq r} I_r$  and consequently  $L(\beta_A; s) = \bigcup_{s \geq r} I_r$  which is an ideal of  $X$ . This completes the proof.  $\square$

A mapping  $f : X \rightarrow Y$  of BCK-algebras is called a *homomorphism* if  $f(x * y) = f(x) * f(y)$  for all  $x, y \in X$ . Note that if  $f : X \rightarrow Y$  is a homomorphism of BCK-algebras, then  $f(0) = 0$ . Let  $f : X \rightarrow Y$  be a homomorphism of BCK-algebras. For any IFS  $A = (\alpha_A, \beta_A)$  in  $Y$ , we define a new IFS  $A^f = (\alpha_A^f, \beta_A^f)$  in  $X$  by

$$\alpha_A^f(x) := \alpha_A(f(x)), \quad \beta_A^f(x) := \beta_A(f(x)) \quad \forall x \in X. \tag{3.28}$$

**THEOREM 3.15.** *Let  $f : X \rightarrow Y$  be a homomorphism of BCK-algebras. If an IFS  $A = (\alpha_A, \beta_A)$  in  $Y$  is an intuitionistic fuzzy ideal of  $Y$ , then an IFS  $A^f = (\alpha_A^f, \beta_A^f)$  in  $X$  is an intuitionistic fuzzy ideal of  $X$ .*

**PROOF.** We first have that

$$\begin{aligned} \alpha_A^f(x) &= \alpha_A(f(x)) \leq \alpha_A(0) = \alpha_A(f(0)) = \alpha_A^f(0), \\ \beta_A^f(x) &= \beta_A(f(x)) \geq \beta_A(0) = \beta_A(f(0)) = \beta_A^f(0) \end{aligned} \tag{3.29}$$

for all  $x \in X$ . Let  $x, y \in X$ . Then

$$\begin{aligned}
\min \{ \alpha_A^f(x * y), \alpha_A^f(y) \} &= \min \{ \alpha_A(f(x * y)), \alpha_A(f(y)) \} \\
&= \min \{ \alpha_A(f(x) * f(y)), \alpha_A(f(y)) \} \\
&\leq \alpha_A(f(x)) = \alpha_A^f(x), \\
\max \{ \beta_A^f(x * y), \beta_A^f(y) \} &= \max \{ \beta_A(f(x * y)), \beta_A(f(y)) \} \\
&= \max \{ \beta_A(f(x) * f(y)), \beta_A(f(y)) \} \\
&\geq \beta_A(f(x)) = \beta_A^f(x).
\end{aligned} \tag{3.30}$$

Hence  $A^f = (\alpha_A^f, \beta_A^f)$  is an intuitionistic fuzzy ideal of  $X$ . □

If we strengthen the condition of  $f$ , then we can construct the converse of Theorem 3.15 as follows.

**THEOREM 3.16.** *Let  $f : X \rightarrow Y$  be an epimorphism of BCK-algebras and let  $A = (\alpha_A, \beta_A)$  be an IFS in  $Y$ . If  $A^f = (\alpha_A^f, \beta_A^f)$  is an intuitionistic fuzzy ideal of  $X$ , then  $A = (\alpha_A, \beta_A)$  is an intuitionistic fuzzy ideal of  $Y$ .*

**PROOF.** For any  $x \in Y$ , there exists  $a \in X$  such that  $f(a) = x$ . Then

$$\begin{aligned}
\alpha_A(x) &= \alpha_A(f(a)) = \alpha_A^f(a) \leq \alpha_A^f(0) = \alpha_A(f(0)) = \alpha_A(0), \\
\beta_A(x) &= \beta_A(f(a)) = \beta_A^f(a) \geq \beta_A^f(0) = \beta_A(f(0)) = \beta_A(0).
\end{aligned} \tag{3.31}$$

Let  $x, y \in Y$ . Then  $f(a) = x$  and  $f(b) = y$  for some  $a, b \in X$ . It follows that

$$\begin{aligned}
\alpha_A(x) &= \alpha_A(f(a)) = \alpha_A^f(a) \\
&\geq \min \{ \alpha_A^f(a * b), \alpha_A^f(b) \} \\
&= \min \{ \alpha_A(f(a * b)), \alpha_A(f(b)) \} \\
&= \min \{ \alpha_A(f(a) * f(b)), \alpha_A(f(b)) \} \\
&= \min \{ \alpha_A(x * y), \alpha_A(y) \}, \\
\beta_A(x) &= \beta_A(f(a)) = \beta_A^f(a) \\
&\leq \max \{ \beta_A^f(a * b), \beta_A^f(b) \} \\
&= \max \{ \beta_A(f(a * b)), \beta_A(f(b)) \} \\
&= \max \{ \beta_A(f(a) * f(b)), \beta_A(f(b)) \} \\
&= \max \{ \beta_A(x * y), \beta_A(y) \}.
\end{aligned} \tag{3.32}$$

This completes the proof. □

Let  $\text{IF}(X)$  be the family of all intuitionistic fuzzy ideals of  $X$  and let  $t \in [0, 1]$ . Define binary relations  $U^t$  and  $L^t$  on  $\text{IF}(X)$  as follows:

$$(A, B) \in U^t \iff U(\alpha_A; t) = U(\alpha_B; t), \quad (A, B) \in L^t \iff L(\beta_A; t) = L(\beta_B; t), \tag{3.33}$$

respectively, for  $A = (\alpha_A, \beta_A)$  and  $B = (\alpha_B, \beta_B)$  in  $\text{IF}(X)$ . Then clearly  $U^t$  and  $L^t$  are



equivalence relations on  $\text{IF}(X)$ . For any  $A = (\alpha_A, \beta_A) \in \text{IF}(X)$ , let  $[A]_{U^t}$  (respectively,  $[A]_{L^t}$ ) denote the equivalence class of  $A$  modulo  $U^t$  (respectively,  $L^t$ ), and denote by  $\text{IF}(X)/U^t$  (respectively,  $\text{IF}(X)/L^t$ ) the system of all equivalence classes modulo  $U^t$  (respectively,  $L^t$ ); so

$$\text{IF}(X)/U^t := \{[A]_{U^t} \mid A = (\alpha_A, \beta_A) \in \text{IF}(X)\}, \quad (3.34)$$

respectively,

$$\text{IF}(X)/L^t := \{[A]_{L^t} \mid A = (\alpha_A, \beta_A) \in \text{IF}(X)\}. \quad (3.35)$$

Now let  $I(X)$  denote the family of all ideals of  $X$  and let  $t \in [0, 1]$ . Define maps  $f_t$  and  $g_t$  from  $\text{IF}(X)$  to  $I(X) \cup \{\emptyset\}$  by  $f_t(A) = U(\alpha_A; t)$  and  $g_t(A) = L(\beta_A; t)$ , respectively, for all  $A = (\alpha_A, \beta_A) \in \text{IF}(X)$ . Then  $f_t$  and  $g_t$  are clearly well defined.

**THEOREM 3.17.** *For any  $t \in (0, 1)$  the maps  $f_t$  and  $g_t$  are surjective from  $\text{IF}(X)$  to  $I(X) \cup \{\emptyset\}$ .*

**PROOF.** Let  $t \in (0, 1)$ . Note that  $\mathbf{0}_\sim = (\mathbf{0}, \mathbf{1})$  is in  $\text{IF}(X)$ , where  $\mathbf{0}$  and  $\mathbf{1}$  are fuzzy sets in  $X$  defined by  $\mathbf{0}(x) = 0$  and  $\mathbf{1}(x) = 1$  for all  $x \in X$ . Obviously  $f_t(\mathbf{0}_\sim) = U(\mathbf{0}; t) = \emptyset = L(\mathbf{1}; t) = g_t(\mathbf{0}_\sim)$ . Let  $G (\neq \emptyset) \in I(X)$ . For  $G_\sim = (\chi_G, \bar{\chi}_G) \in \text{IF}(X)$ , we have  $f_t(G_\sim) = U(\chi_G; t) = G$  and  $g_t(G_\sim) = L(\bar{\chi}_G; t) = G$ . Hence  $f_t$  and  $g_t$  are surjective.  $\square$

**THEOREM 3.18.** *The quotient sets  $\text{IF}(X)/U^t$  and  $\text{IF}(X)/L^t$  are equipotent to  $I(X) \cup \{\emptyset\}$  for every  $t \in (0, 1)$ .*

**PROOF.** For  $t \in (0, 1)$  let  $f_t^*$  (respectively,  $g_t^*$ ) be a map from  $\text{IF}(X)/U^t$  (respectively,  $\text{IF}(X)/L^t$ ) to  $I(X) \cup \{\emptyset\}$  defined by  $f_t^*([A]_{U^t}) = f_t(A)$  (respectively,  $g_t^*([A]_{L^t}) = g_t(A)$ ) for all  $A = (\alpha_A, \beta_A) \in \text{IF}(X)$ . If  $U(\alpha_A; t) = U(\alpha_B; t)$  and  $L(\beta_A; t) = L(\beta_B; t)$  for  $A = (\alpha_A, \beta_A)$  and  $B = (\alpha_B, \beta_B)$  in  $\text{IF}(X)$ , then  $(A, B) \in U^t$  and  $(A, B) \in L^t$ ; hence  $[A]_{U^t} = [B]_{U^t}$  and  $[A]_{L^t} = [B]_{L^t}$ . Therefore the maps  $f_t^*$  and  $g_t^*$  are injective. Now let  $G (\neq \emptyset) \in I(X)$ . For  $G_\sim = (\chi_G, \bar{\chi}_G) \in \text{IF}(X)$ , we have

$$\begin{aligned} f_t^*([G_\sim]_{U^t}) &= f_t(G_\sim) = U(\chi_G; t) = G, \\ g_t^*([G_\sim]_{L^t}) &= g_t(G_\sim) = L(\bar{\chi}_G; t) = G. \end{aligned} \quad (3.36)$$

Finally, for  $\mathbf{0}_\sim = (\mathbf{0}, \mathbf{1}) \in \text{IF}(X)$  we get

$$\begin{aligned} f_t^*([\mathbf{0}_\sim]_{U^t}) &= f_t(\mathbf{0}_\sim) = U(\mathbf{0}; t) = \emptyset, \\ g_t^*([\mathbf{0}_\sim]_{L^t}) &= g_t(\mathbf{0}_\sim) = L(\mathbf{0}; t) = \emptyset. \end{aligned} \quad (3.37)$$

This shows that  $f_t^*$  and  $g_t^*$  are surjective. This completes the proof.  $\square$

For any  $t \in [0, 1]$ , we define another relation  $R^t$  on  $\text{IF}(X)$  as follows:

$$(A, B) \in R^t \iff U(\alpha_A; t) \cap L(\beta_A; t) = U(\alpha_B; t) \cap L(\beta_B; t) \quad (3.38)$$

for any  $A = (\alpha_A, \beta_A), B = (\alpha_B, \beta_B) \in \text{IF}(X)$ . Then the relation  $R^t$  is also an equivalence relation on  $\text{IF}(X)$ .

**THEOREM 3.19.** *For any  $t \in (0, 1)$ , the map  $\phi_t : \text{IF}(X) \rightarrow I(X) \cup \{\emptyset\}$  defined by  $\phi_t(A) = f_t(A) \cap g_t(A)$  for each  $A = (\alpha_A, \beta_A) \in \text{IF}(X)$  is surjective.*

**PROOF.** Let  $t \in (0, 1)$ . For  $\mathbf{0}_\sim = (\mathbf{0}, \mathbf{1}) \in \text{IF}(X)$ ,

$$\phi_t(\mathbf{0}_\sim) = f_t(\mathbf{0}_\sim) \cap g_t(\mathbf{0}_\sim) = U(\mathbf{0}; t) \cap L(\mathbf{1}; t) = \emptyset. \quad (3.39)$$

For any  $H \in \text{IF}(X)$ , there exists  $H_\sim = (\chi_H, \bar{\chi}_H) \in \text{IF}(X)$  such that

$$\phi_t(H_\sim) = f_t(H_\sim) \cap g_t(H_\sim) = U(\chi_H; t) \cap L(\bar{\chi}_H; t) = H. \quad (3.40)$$

This completes the proof.  $\square$

**THEOREM 3.20.** *For any  $t \in (0, 1)$ , the quotient set  $\text{IF}(X)/R^t$  is equipotent to  $I(X) \cup \{\emptyset\}$ .*

**PROOF.** Let  $t \in (0, 1)$  and let  $\phi_t^* : \text{IF}(X)/R^t \rightarrow I(X) \cup \{\emptyset\}$  be a map defined by  $\phi_t^*([A]_{R^t}) = \phi_t(A)$  for all  $[A]_{R^t} \in \text{IF}(X)/R^t$ . If  $\phi_t^*([A]_{R^t}) = \phi_t^*([B]_{R^t})$  for any  $[A]_{R^t}, [B]_{R^t} \in \text{IF}(X)/R^t$ , then  $f_t(A) \cap g_t(A) = f_t(B) \cap g_t(B)$ , that is,  $U(\alpha_A; t) \cap L(\beta_A; t) = U(\alpha_B; t) \cap L(\beta_B; t)$ , hence  $(A, B) \in R^t$ . It follows that  $[A]_{R^t} = [B]_{R^t}$  so that  $\phi_t^*$  is injective. For  $\mathbf{0}_\sim = (\mathbf{0}, \mathbf{1}) \in \text{IF}(X)$ ,

$$\phi_t^*([\mathbf{0}_\sim]_{R^t}) = \phi_t(\mathbf{0}_\sim) = f_t(\mathbf{0}_\sim) \cap g_t(\mathbf{0}_\sim) = U(\mathbf{0}; t) \cap L(\mathbf{1}; t) = \emptyset. \quad (3.41)$$

If  $H \in \text{IF}(X)$ , then for  $H_\sim = (\chi_H, \bar{\chi}_H) \in \text{IF}(X)$ , we have

$$\phi_t^*([H_\sim]_{R^t}) = \phi_t(H_\sim) = f_t(H_\sim) \cap g_t(H_\sim) = U(\chi_H; t) \cap L(\bar{\chi}_H; t) = H. \quad (3.42)$$

Hence  $\phi_t^*$  is surjective, this completes the proof.  $\square$

**ACKNOWLEDGEMENT.** The first author was supported by Korea Research Foundation Grant (KRF-99-005-D00003).

## REFERENCES

- [1] K. T. Atanassov, *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems **20** (1986), no. 1, 87–96. MR 87f:03151. Zbl 631.03040.
- [2] ———, *New operations defined over the intuitionistic fuzzy sets*, Fuzzy Sets and Systems **61** (1994), no. 2, 137–142. CMP 1 262 464. Zbl 824.04004.
- [3] Y. B. Jun, *A note on fuzzy ideals in BCK-algebras*, Math. Japon. **42** (1995), no. 2, 333–335. CMP 1 356 395. Zbl 834.06018.
- [4] ———, *Finite valued fuzzy ideals in BCK-algebras*, J. Fuzzy Math. **5** (1997), no. 1, 111–114. CMP 1 441 020. Zbl 868.06010.
- [5] ———, *Characterizations of Noetherian BCK-algebras via fuzzy ideals*, Fuzzy Sets and Systems **108** (1999), no. 2, 231–234. CMP 1 720 432. Zbl 940.06014.
- [6] Y. B. Jun, S. M. Hong, S. J. Kim, and S. Z. Song, *Fuzzy ideals and fuzzy subalgebras of BCK-algebras*, J. Fuzzy Math. **7** (1999), no. 2, 411–418. MR 2000c:06040. Zbl 943.06010.
- [7] Y. B. Jun and E. H. Roh, *Fuzzy commutative ideals of BCK-algebras*, Fuzzy Sets and Systems **64** (1994), no. 3, 401–405. MR 95e:06051. Zbl 846.06011.

- [8] J. Meng, Y. B. Jun, and H. S. Kim, *Fuzzy implicative ideals of BCK-algebras*, *Fuzzy Sets and Systems* **89** (1997), no. 2, 243-248. MR 98a:06033. Zbl 914.06009.
- [9] L. A. Zadeh, *Fuzzy sets*, *Information and Control* **8** (1965), 338-353. MR 36#2509. Zbl 139.24606.

YOUNG BAE JUN: DEPARTMENT OF MATHEMATICS EDUCATION, GYEONGSANG NATIONAL UNIVERSITY, CHINJU 660-701, KOREA

*E-mail address:* ybjun@nongae.gsnu.ac.kr

KYUNG HO KIM: DEPARTMENT OF MATHEMATICS, CHUNGJU NATIONAL UNIVERSITY, CHUNGJU 380-702, KOREA

*E-mail address:* ghkim@gukwon.chungju.ac.kr



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

