

RESEARCH

Open Access

On the sub-supersolution method for $p(x)$ -Kirchhoff type equations

Xiaoling Han and Guowei Dai*

*Correspondence:
daiguowei@nwnu.edu.cn
Department of Mathematics,
Northwest Normal University,
Lanzhou, 730070, P.R. China**Abstract**

This paper deals with the sub-supersolution method for the $p(x)$ -Kirchhoff type equations. A sub-supersolution principle for the Dirichlet problems involving $p(x)$ -Kirchhoff is established. A strong comparison theorem for the $p(x)$ -Kirchhoff type equations is presented. We also give some applications of the abstract theorems obtained in this paper to the eigenvalue problems for the $p(x)$ -Kirchhoff type equation.

MSC: 35D05; 35D10; 35J60**Keywords:** subsolution; supersolution; nonlocal problems; comparison theorem

1 Introduction

In this paper, we study the following problem:

$$\begin{cases} -M(t) \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) = f(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where Ω is a bounded smooth domain in \mathbb{R}^N with $N \geq 1$, $p = p(x) \in C(\overline{\Omega})$ with $1 < p^- := \inf_{\Omega} p(x) \leq p^+ := \sup_{\Omega} p(x) < +\infty$, $f \in C(\overline{\Omega} \times \mathbb{R}, \mathbb{R})$, $M(t)$ is a continuous function with $t := \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx$ and satisfies the following condition:

(M_0) $M : [0, +\infty) \rightarrow (m_0, +\infty)$ is a continuous and increasing function with $m_0 > 0$.

The operator $-\operatorname{div}(|\nabla u|^{p(x)-2} \nabla u)$ is said to be $p(x)$ -Laplacian. The study of various mathematical problems with the variable exponent growth condition has received considerable attention in recent years. These problems are interesting in applications and raise many difficult mathematical problems. We refer the reader to [1] for an overview of and references on this subject.

The solvability of the problem (1.1) can be studied by several approaches; for example, the variational method (see, e.g., [2]). It is well known that, compared with other methods, the sub-supersolution method, or the order method, when it is applicable, has some distinctive advantages. For example, it usually gives some order properties of the solutions. For the applications of the sub-supersolution method to semilinear and quasilinear elliptic problems, we refer to [3, 4] and the references therein. In [3], Fan established a sub-supersolution principle for Dirichlet problems involving $p(x)$ -Laplacian and a strong comparison theorem for $p(x)$ -Laplacian equations. The goal of this paper is to study the sub-supersolution method for (1.1), which is a new research topic.

The problem (1.1) is related to the stationary problem of a model introduced by Kirchhoff [5]. We refer the reader to [6] for an overview of and references on this subject.

In [3], the sub-supersolution principle for $p(x)$ -Laplacian equations established by Fan is based on the properties of $p(x)$ -Laplace, the regularity results and the comparison principle. The aim of the present paper is to establish a sub-supersolution principle for $p(x)$ -Kirchhoff equations.

The rest of this paper is organized as follows. In Section 2, we establish a general principle of the sub-supersolution method for the problem (1.1) based on the regularity results and the comparison principle. In Section 3, we give a special strong comparison principle for the $p(x)$ -Kirchhoff. In Section 4, we give an application of our abstract theorems.

2 Sub-supersolution principle

In this section, we give a general principle of sub-supersolution method for the problem (1.1) based on the regularity results and the comparison principle. We would like to point out that the comparison principle in this section (see Theorem 2.2) is a generalization of Proposition 2.3 of [3]. In addition to the principle of sub-supersolution, we shall establish also a generalization of Theorem 2.1 of [3]. For simplicity, we write $X = W_0^{1,p(x)}(\Omega)$.

Definition 2.1 (1) We say that $u \in X$ is a weak solution of (1.1) if

$$M\left(\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx\right) \int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla \varphi dx = \int_{\Omega} f(x, u) \varphi dx$$

for any $\varphi \in X$.

(2) $u \in W^{1,p(x)}$ is called a subsolution (respectively a supersolution) of (1.1) if $u \leq$ (respectively \geq) 0 on $\partial\Omega$ and, for all $\varphi \in X$ with $\varphi \geq 0$,

$$M\left(\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx\right) \int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla \varphi dx \leq (\text{respectively } \geq) \int_{\Omega} f(x, u) \varphi dx.$$

Regularity results and comparison principles are the basis of the sub-supersolution method. For the regularity results in the variable exponent case, see [7–9]. More precisely, for the L^∞ and $C^{0,\alpha}$ regularity, see [8]; for the local $C^{1,\alpha}$ regularity of the minimizers of the corresponding integral functional, see [7]; for the global $C^{1,\alpha}$ regularity, see [9].

If f is independent of u , we have

Theorem 2.1 *If (M_0) holds and $f(x, u) = f(x)$, $f \in L^{\frac{q(x)}{q(x)-1}}(\Omega)$, then (1.1) has a unique weak solution.*

Proof Clearly, $(f, v) := \int_{\Omega} f(x)v dx$ (for any $v \in X$) defines a continuous linear functional on X . According to Theorem 4.1 of [10], Φ' is a homeomorphism. So, (1.1) has a unique solution, where $\Phi(u) = \widehat{M}\left(\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx\right)$. \square

From Theorem 2.1 we know that, for a given $h \in L^{\frac{q(x)}{q(x)-1}}(\Omega)$, where $q \in C_+(\overline{\Omega})$ and

$$1 < q(x) < p^*(x), \quad \forall x \in \overline{\Omega}, \tag{2.1}$$

the problem

$$\begin{cases} -M(t) \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) = h(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (2.2)$$

has a unique solution $u \in X$ under the condition (M_0) . We denote by $K(h) := u$ the unique solution. K is called a solution operator for (2.2).

From the regularity results and the embedding theorems, we can obtain the properties of the solution operator K as follows.

Proposition 2.1 (1) *If (M_0) holds, the mapping $K : L^{\frac{q(x)}{q(x)-1}}(\Omega) \rightarrow X$ is continuous and bounded. Moreover, the mapping $K : L^{\frac{q(x)}{q(x)-1}}(\Omega) \hookrightarrow L^{q(x)}(\Omega)$ is completely continuous since the embedding $X \hookrightarrow L^{q(x)}(\Omega)$ is compact.*

(2) *If (M_0) holds and p is log-Hölder continuous on $\overline{\Omega}$, then the mapping $K : L^\infty(\Omega) \rightarrow C^{0,\alpha}(\overline{\Omega})$ is bounded, and hence the mapping $K : L^\infty(\Omega) \rightarrow C(\overline{\Omega})$ is completely continuous.*

(3) *If (M_0) holds and p is Hölder continuous on $\overline{\Omega}$, then the mapping $K : L^\infty(\Omega) \rightarrow C^{1,\alpha}(\overline{\Omega})$ is bounded, and hence the mapping $K : L^\infty(\Omega) \rightarrow C^1(\overline{\Omega})$ is completely continuous.*

Definition 2.2 Let $u, v \in W^{1,p(x)}(\Omega)$. We say that $-M(I_0(u))\Delta_{p(x)}(u) \leq -M(I_0(v))\Delta_{p(x)}(v)$ if for all $\varphi \in X$ with $\varphi \geq 0$,

$$M(I_0(u)) \int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla \varphi \, dx \leq M(I_0(v)) \int_{\Omega} |\nabla v|^{p(x)-2} \nabla v \nabla \varphi \, dx, \quad (2.3)$$

where $I_0(u) = \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} \, dx$.

Now we give the comparison principle as follows.

Theorem 2.2 (1) *Let $u, v \in W^{1,p(x)}(\Omega)$ and (M_0) hold. If $-M(I_0(u))\Delta_{p(x)}(u) \leq -M(I_0(v)) \times \Delta_{p(x)}(v)$ and $(u - v)^+ \in W_0^{1,p(x)}(\Omega)$, then $u \leq v$ in Ω .*

(2) *Under the conditions of (1) above, let in addition $u, v \in C(\overline{\Omega})$ and denote $S = \{x \in \Omega : u(x) = v(x)\}$. If S is a compact subset of Ω , then $S = \emptyset$.*

Proof (1) Taking $\lambda = 0$ in the proof of Theorem 3.2 of [11], we can get the conclusion.

(2) Suppose that S is a compact subset of Ω and $S \neq \emptyset$. Then there is an open subset Ω_3 of Ω such that $S \subset \Omega_3 \subset \overline{\Omega}_3 \subset \Omega$. Thus $u < v$ on $\partial\Omega_3$ and consequently there is an $\varepsilon > 0$ such that $u < v - \varepsilon$ on $\partial\Omega_3$. Noting that $\nabla(v - \varepsilon) = \nabla v$ and applying the conclusion (1) to u and $v - \varepsilon$ on Ω_3 , we obtain $u \leq v - \varepsilon$ in Ω_3 , which contradicts $u = v$ on S . \square

It follows from Theorem 2.2(1) that the solution operator K is increasing under the condition (M_0) , that is, $K(u) \leq K(v)$ if $u \leq v$. We define $T(u) = K(f(x, u))$. It is easy to see that if u is a subsolution (respectively a supersolution) of (1.1), then $u \leq T(u)$ (respectively $u \geq T(u)$), and u is a solution of (1.1) if and only if $u = T(u)$, i.e., u is a fixed point of T .

The basic principle of the sub-supersolution method for (1.1) is the following result.

Theorem 2.3 *Let (M_0) hold and suppose that f satisfies the sub-critical growth condition*

$$|f(x, t)| \leq c_1 + c_2 |t|^{q(x)-1}, \quad \forall x \in \Omega, \forall t \in \mathbb{R},$$

and the function $f(x, t)$ is nondecreasing in $t \in \mathbb{R}$. If there exist a subsolution $u_0 \in W^{1,p(x)}(\Omega)$ and a supersolution $v_0 \in W^{1,p(x)}(\Omega)$ of (1.1) such that $u_0 \leq v^0$, then (1.1) has a minimal solution u^* and a maximal solution v^* in the order interval $[u_0, v^0]$, i.e., $u_0 \leq u^* \leq v^* \leq v^0$ and if u is any solution of (1.1) such that $u_0 \leq u \leq v^0$, then $u^* \leq u \leq v^*$.

Proof Define $T(u) = K(f(x, u))$. Then, under the assumptions of Theorem 2.3, $T : L^{q(x)}(\Omega) \rightarrow L^{q(x)}(\Omega)$ is completely continuous and increasing, $u_0 \leq v^0$, $u_0, v^0 \in L^{q(x)}(\Omega)$, $u_0 \leq T(u_0)$, $v^0 \geq T(v^0)$, and consequently $T : [u_0, v^0] \rightarrow [u_0, v^0]$. It is clear that the cone of all nonnegative functions in $L^{q(x)}(\Omega)$ is normal. Noting the minimal (maximal) fixed point (see [4]) of T is the minimal (maximal) solution of (1.1), so our Theorem 2.3 now follows by applying the well-known fixed point theorem for the increasing operator on the order interval (see, e.g., [4]). \square

In the practical problems, it is often known that the subsolution u_0 and the supersolution v^0 are of class $L^\infty(\Omega)$, so the restriction on the growth condition of f is needless. Hence, the following theorem is more suitable.

Theorem 2.4 *Let (M_0) hold and suppose that $u_0, v^0 \in W^{1,p(x)}(\Omega) \cap L^\infty(\Omega)$, u_0 and v^0 are a subsolution and a supersolution of (1.1) respectively, and $u_0 \leq v^0$. If $f \in C(\overline{\Omega} \times \mathbb{R}, \mathbb{R})$ satisfies the condition*

$$(F_1) \quad f(x, t) \text{ is nondecreasing in } t \in [\inf u_0(x), \sup v^0(x)],$$

then the conclusion of Theorem 2.3 is valid.

The above results show that the general principle of the sub-supersolution method for $p(x)$ -Kirchhoff type equations (1.1) is of the same type as in the case of $p(x)$ -Laplacian type equations. An essential prerequisite for the sub-supersolution method is to find a subsolution u_0 and a supersolution v^0 such that $u_0 \leq v^0$. It is well known that the homogeneity of the p -Laplacian operator and the positivity of the first eigenvalue of p -Laplacian Dirichlet problem play an important role in finding sub- and supersolutions of the p -Laplacian equation [12]. Unlike the p -Laplacian, when $p(x)$ is not identical with a constant, the $p(x)$ -Laplacian operator is inhomogeneous and usually the infimum of its eigenvalues is 0. It is obvious that the eigenvalues of (1.1) are $\mu_j = M(\int_\Omega \frac{1}{p(x)} |\nabla \varphi_j|^{p(x)} dx) \lambda_j$, where λ_j and φ_j are, respectively, the eigenvalues and eigenfunctions of $-\Delta_{p(x)}$ in X . Thus, usually, the infimum of μ_j is also 0. Therefore, it is often difficult to find a subsolution u_0 and a supersolution v^0 of (1.1) with $u_0 \leq v^0$.

At the end of this section, we give a lemma which is useful to find a supersolution of (1.1). We denote by C_0 the best embedding constant of $W_0^{1,1}(\Omega) \subset L^{\frac{N}{N-1}}(\Omega)$.

Lemma 2.1 *Let (M_0) hold, $\mathcal{M} > 0$ and let u be the unique solution of the problem*

$$\begin{cases} -M(t) \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) = \mathcal{M} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} \tag{2.4}$$

Set $h = \frac{m_0 p^-}{2|\Omega|^{1/N} C_0}$. Then, when $\mathcal{M} \geq h$, $|u|_\infty \leq C^* \mathcal{M}^{1/(p^- - 1)}$, and when $\mathcal{M} < h$, $|u|_\infty \leq C^* \mathcal{M}^{1/(p^+ - 1)}$, where C^* and C_* are positive constants depending on p^+ , p^- , N , $|\Omega|$, C_0 and m_0 .

Proof Let u be the solution of (2.4), Theorem 2.2 implies $u \geq 0$. For $k \geq 0$, set $A_k = \{x \in \Omega : u(x) > k\}$. Taking $(u - k)^+$ as a test function in (2.4) and using the Young inequality, we have

$$\begin{aligned} \int_{A_k} |\nabla u|^{p(x)} dx &= \frac{\mathcal{M}}{M(t)} \int_{A_k} (u - k) dx \\ &\leq \frac{\mathcal{M}|\Omega|^{1/N}C_0}{m_0p^-} \int_{A_k} \varepsilon^{p(x)} |\nabla u|^{p(x)} dx + \frac{\mathcal{M}|A_k|^{1/N}C_0}{m_0(p^+)'} \int_{A_k} \varepsilon^{-p'(x)} dx. \end{aligned} \quad (2.5)$$

When $\mathcal{M} \geq h$, taking

$$\varepsilon = \left(\frac{m_0p^-}{2\mathcal{M}|\Omega|^{1/N}C_0} \right)^{1/p^-} = \left(\frac{h}{\mathcal{M}} \right)^{1/p^-},$$

then $\varepsilon \leq 1$ and

$$\frac{\mathcal{M}|\Omega|^{1/N}C_0}{m_0p^-} \int_{A_k} \varepsilon^{p(x)} |\nabla u|^{p(x)} dx \leq \frac{\mathcal{M}|\Omega|^{1/N}C_0}{m_0p^-} \varepsilon^{p^-} \int_{A_k} |\nabla u|^{p(x)} dx = \frac{1}{2} \int_{A_k} |\nabla u|^{p(x)} dx.$$

Consequently, from this and (2.5), it follows that

$$\int_{A_k} |\nabla u|^{p(x)} dx \leq \frac{2\mathcal{M}|A_k|^{1/N}C_0}{m_0(p^+)'} \int_{A_k} \varepsilon^{-p'(x)} dx \leq \frac{2\mathcal{M}C_0\varepsilon^{-(p^-)'}}{m_0(p^+)'} |A_k|^{1+1/N}. \quad (2.6)$$

From (2.5) and (2.6), we have

$$\begin{aligned} \int_{A_k} (u - k) dx &= \frac{M(t)}{\mathcal{M}} \int_{A_k} |\nabla u|^{p(x)} dx \\ &\leq M \left(\frac{2\mathcal{M}C_0\varepsilon^{-(p^-)'}}{p^-m_0(p^+)'} |\Omega|^{1+1/N} \right) \frac{2C_0\varepsilon^{-(p^-)'}}{m_0(p^+)'} |A_k|^{1+1/N}. \end{aligned} \quad (2.7)$$

By Lemma 5.1 in [13, Chapter 2], (2.7) implies that

$$|u|_\infty \leq \gamma(N + 1)|\Omega|^{1/N}, \quad (2.8)$$

where $\gamma = M \left(\frac{2\mathcal{M}C_0\varepsilon^{-(p^-)'}}{p^-m_0(p^+)'} |\Omega|^{1+1/N} \right) \frac{2C_0\varepsilon^{-(p^-)'}}{m_0(p^+)}'$. From (2.7) and (2.8), we obtain

$$|u|_\infty \leq C^* \mathcal{M}^{1/(p^- - 1)},$$

where

$$C^* = \frac{(N + 1)(2C_0)^{(p^-)'}}{(p^+)m_0^{(p^-)'}} |\Omega|^{(p^-)'/N} M \left(\frac{(2\mathcal{M}C_0)^{(p^-)'}}{p^-(p^+)m_0^{(p^-)'}} |\Omega|^{(p^-)'/N} \right).$$

When $\mathcal{M} < h$, taking

$$\varepsilon = \left(\frac{m_0p^-}{2\mathcal{M}|\Omega|^{1/N}C_0} \right)^{1/p^+} = \left(\frac{h}{\mathcal{M}} \right)^{1/p^+}$$

(noting that in this case $\varepsilon > 1$) and using arguments similar to those above, we can obtain

$$|u|_\infty \leq C^* \mathcal{M}^{1/(p^+ - 1)},$$

where

$$C^* = \frac{(N + 1)(2C_0)^{(p^+)'}}{(p^+)m_0^{(p^+)'}(p^-)^{(p^+)'/p^+}} |\Omega|^{(p^+)'/N} M \left(\frac{(2\mathcal{M}C_0)^{(p^+)'}}{p^-(p^+)m_0^{(p^+)'}(p^-)^{(p^+)'/p^+}} |\Omega|^{(p^+)'/N} \right).$$

The proof is complete. □

Remark 2.1 We would like to point out that the fact that a solution of (2.4) is bounded in $L^\infty(\Omega)$ is useful for finding a supersolution of (1.1). Indeed, the fact can be used to estimate the relation of nonlinearity and \mathcal{M} (for details, see the proof of Theorem 4.1).

3 A strong comparison principle for $p(x)$ -Kirchhoff problem

The energy functional associated with the problem (1.1) is

$$J(u) = \widehat{M}(I_0(u)) - \int_\Omega F(x, u) \, dx,$$

where $\widehat{M}(t) = \int_0^t M(\tau) \, d\tau$ and $F(x, u) = \int_0^u f(x, t) \, dt$. In this section, we give a special strong comparison principle for the $p(x)$ -Kirchhoff, which is suitable for finding a positive C^1 local minimizer of the integral functional J in the C^1 topology. In [14], Fan established a Brezis-Nirenberg type theorem (Theorem 1.1 of [14]), which asserts that every local minimizer of J in the $C^1(\Omega)$ topology is also a local minimizer of J in the $W_0^{1,p(x)}(\Omega)$ topology. Applying this theorem, we have the following special form.

Theorem 3.1 *Let (M_0) , (2.1) hold and let $u_0 \in X$ be a local minimizer (resp. a strictly local minimizer) of J in the $C^1(\Omega)$ topology. Then u_0 is a local minimizer (resp. a strictly local minimizer) of J in the X topology.*

Applying Theorem 1.1 of [15], we can easily get the following strong maximum principle.

Theorem 3.2 *Suppose that $p(x) \in C_+(\overline{\Omega}) \cap C^1(\overline{\Omega})$, $u \in X$, $u \geq 0$ and $u \not\equiv 0$ in Ω . If*

$$-M(t) (\operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) - d(x)|u|^{q(x)-2} u) \geq 0,$$

where $t = \int_\Omega (\frac{1}{p(x)} |\nabla u|^{p(x)} + \frac{1}{q(x)} d(x)|u|^{q(x)}) \, dx$, $M(t) \geq m_0 > 0$, $0 \leq d(x) \in L^\infty(\Omega)$, $q(x) \in C(\overline{\Omega})$ with $p(x) \leq q(x) \leq p^*(x)$, then $u > 0$ in Ω .

Now we give a special strong comparison principle for the $p(x)$ -Kirchhoff.

Theorem 3.3 *Let (M_0) hold and suppose that $u, v \in C^1(\overline{\Omega})$, $u \geq v$ in Ω , $g, h \in L^\infty(\Omega)$,*

$$-M(I_0(u)) \Delta_{p(x)} u = g(x) \geq h(x) = -M(I_0(v)) \Delta_{p(x)} v \quad \text{in } \Omega, \tag{3.1}$$

and $g(x) \not\equiv h(x)$ in Ω . If

$$\frac{\partial u}{\partial \mathbf{n}} > 0, \quad \frac{\partial v}{\partial \mathbf{n}} > 0 \quad \text{on } \partial\Omega,$$

where \mathbf{n} is the inward unit normal on $\partial\Omega$, then $u > v$ in Ω and there is a positive constant ε such that

$$\frac{\partial(u-v)}{\partial \mathbf{n}} \geq \varepsilon \quad \text{on } \partial\Omega. \tag{3.2}$$

Proof We denote by \mathbf{n}_y the inward unit normal at $y \in \partial\Omega$. For $\delta > 0$, set $\Omega_\delta = \{x \in \Omega : \text{dist}(x, \partial\Omega) < \delta\}$. Denoting $A(x, \eta) = M(I_0(\eta))|\eta|^{p(x)-2}\eta$, as in the proof of [3], we have

$$u - v \geq 0 \quad \text{in } \Omega_\delta.$$

We claim that $u - v \not\equiv 0$ in Ω_δ . Indeed, if $u \equiv v$ in Ω_δ , then $g \equiv h$ in Ω_δ , and consequently $g(x) \not\equiv h(x)$ in $\Omega \setminus \Omega_\delta$. Take $\varphi \in X$ such that $\varphi > 0$ in Ω , $\varphi = 1$ on $\Omega \setminus \Omega_\delta$. By (3.1) and the property of φ , we have

$$\begin{aligned} \int_{\Omega \setminus \Omega_\delta} g(x)\varphi(x) \, dx &= M\left(\int_{\Omega \setminus \Omega_\delta} \frac{1}{p(x)} |\nabla u|^{p(x)} \, dx\right) \int_{\Omega \setminus \Omega_\delta} |\nabla u|^{p(x)-2} \nabla u \nabla \varphi \, dx = 0 \\ &= M\left(\int_{\Omega \setminus \Omega_\delta} \frac{1}{p(x)} |\nabla v|^{p(x)} \, dx\right) \int_{\Omega \setminus \Omega_\delta} |\nabla v|^{p(x)-2} \nabla v \nabla \varphi \, dx \\ &= \int_{\Omega \setminus \Omega_\delta} h(x)\varphi(x) \, dx, \end{aligned}$$

which contradicts $\int_{\Omega} g(x)\varphi(x) \, dx > \int_{\Omega} h(x)\varphi(x) \, dx$. Hence the claim is true. So, by the well-known strong maximum principle for linear elliptic equations, $u > v$ in Ω_δ and (3.2) holds. Setting $S = \{x \in \Omega : u(x) = v(x)\}$, then S is a compact subset of Ω . By Theorem 2.2(2), $S = \emptyset$, hence $u > v$ in Ω and the proof is complete. \square

The following theorem provides a method to find a positive C^1 local minimizer of the integral functional J in the C^1 topology.

Theorem 3.4 *Let (M_0) hold and suppose that $u_0, v^0 \in X$ are a subsolution and a supersolution of (1.1) respectively, $-M(I(u_0))\Delta_{p(x)}u_0 = g(x)$, $-M(I(v^0))\Delta_{p(x)}v^0 = h(x)$, $g, h \in L^\infty(\Omega)$, $0 \leq g \leq h$, $g(x) \not\equiv h(x)$ and $0 \leq u_0 \leq v^0$ in Ω . Suppose that $p \in C^1(\overline{\Omega})$, $f \in C(\overline{\Omega} \times \mathbb{R}, \mathbb{R})$ satisfies the condition of Theorem 2.3. If neither u_0 nor v^0 is a solution of (1.1), or neither u_0 nor v^0 is a minimizer of J on $[u_0, v^0] \cap X$ in the case of being a solution of (1.1), then there exists $u_* \in [u_0, v^0] \cap C^{1,\alpha}(\overline{\Omega})$ such that $J(u_*) = \inf\{J(u) : u \in [u_0, v^0] \cap X\}$, u_* is a solution of (1.1) and u_* is a local minimizer of J in the C^1 topology.*

Proof The proof is similar to the proof of [3], we omit it here (for details, see the proof of Theorem 3.3 in [3]). \square

4 Applications

As an application of the above abstract theorems, let us consider the following eigenvalue problem:

$$\begin{cases} -M\left(\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx\right) \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) = \lambda f(x, u) + \mu |u|^{q(x)-2} u & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \quad u = 0 & \text{on } \partial\Omega, \end{cases} \quad (4.1)$$

where Ω is a bounded smooth domain in \mathbb{R}^N , $p \in C^1(\overline{\Omega})$, $q \in C(\overline{\Omega})$, $q^- > p^+$, $f \in C(\overline{\Omega} \times \mathbb{R}, \mathbb{R})$, $f(x, t) \geq 0$ for $x \in \Omega$ and $t \geq 0$, $f(x, t)$ is nondecreasing in $t \geq 0$, $\mu \geq 0$ is fixed. The energy functional associated with the problem (4.1) is

$$J_{\lambda}(u) = \widehat{M}\left(\int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx\right) - \lambda \int_{\Omega} F(x, u) dx - \mu \int_{\Omega} \frac{|u|^{q(x)}}{q(x)} dx, \quad \forall u \in X,$$

where $F(x, t) = \int_0^t f(x, s) ds$.

Firstly, we recall the $(PS)_c$ condition and the mountain pass lemma which we shall use later.

Definition 4.1 Let X be a Banach space. We say that I satisfies the $(PS)_c$ condition in X if any sequence $\{u_n\} \subset X$, such that $|I(u_n)| \leq c$ and $I'(u_n) \rightarrow 0$ as $n \rightarrow +\infty$, has a convergent subsequence, where (PS) means Palais-Smale.

Lemma 4.1 (see [16]) *Let X be a Banach space, $\varphi \in C^1(X, \mathbb{R})$, $e \in X$ and $r > 0$ be such that $\|e\| > r$ and*

$$b := \inf_{\|u\|=r} \varphi(u) > \varphi(0) \geq \varphi(e).$$

If φ satisfies the $(PS)_c$ condition with

$$\begin{aligned} c &:= \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \varphi(\gamma(t)), \\ \Gamma &:= \{\gamma \in C([0,1], X) : \gamma(0) = 0, \gamma(1) = e\}, \end{aligned}$$

then c is a critical value of φ .

The main results are the following.

Theorem 4.1 *Suppose that f satisfies the condition either*

- (i) $f(x, 0) \not\equiv 0$ in Ω , or
- (ii) $f(x, 0) \equiv 0$ and there are an open set $U \subset \Omega$, a closed ball $\overline{B}(x_0, \rho) \subset U$, some positive constants $r_0 > 1$ and c_3 such that $f(x, t) \geq c_3 t^{r_0-1}$ for $x \in \overline{B}(x_0, \rho)$ and $t \in [0, 1]$, and $r_0 < p(x)$ for $x \in \partial U$.

Then we have the following assertions:

- (1) *For sufficiently small $\lambda > 0$, (4.1) has a solution u_{λ} which is a local minimizer of J_{λ} in the C^1 topology. Moreover, $\|u_{\lambda}\|_{C^1(\overline{\Omega})} \rightarrow 0$ as $\lambda \rightarrow 0$.*
- (2) *Define $\Lambda_0 = \{\lambda > 0 : (4.1) \text{ has a solution } u_{\lambda} \text{ which is a local minimizer of } J_{\lambda} \text{ in the } C^1 \text{ topology}\}$ and $\Lambda = \{\lambda > 0 : (4.1) \text{ has a solution } u_{\lambda}\}$. Then Λ_0 and Λ are both intervals, $\inf \Lambda_0 = \inf \Lambda = 0$ and $\Lambda_0 \supset \operatorname{int} \Lambda$.*

(3) In addition, suppose that $\mu > 0$, $q(x) < p^*(x)$ for $x \in \overline{\Omega}$ and

$$|f(x, t)| \leq c(1 + |t|^{r(x)}) \quad \text{for } x \in \Omega \text{ and } t \in \mathbb{R},$$

where $r(x) < p^*(x)$ for $x \in \overline{\Omega}$ and $r^+ < q^-$. Then for each $\lambda \in \text{int } \Lambda$, (4.1) has at least two solutions u_λ and v_λ such that $u_\lambda < v_\lambda$ and u_λ is a local minimizer of J_λ in the $W^{1,p(x)}$ topology.

Proof (1) Take $0 < \mathcal{M} < h$, where h is as in Lemma 2.1, and let $v = v_{\mathcal{M}}$ be the unique positive solution of (2.4). Then by Lemma 2.1, $|v|_\infty \leq C_* \mathcal{M}^{1/(p^+-1)}$. Because $q^- > p^+$, we can choose \mathcal{M} small enough such that $\mu(C_* \mathcal{M}^{1/(p^+-1)})^{q^-} < \frac{\mathcal{M}}{2}$, which implies that $\mu |v|^{q(x)-1} < \frac{\mathcal{M}}{2}$. Let $\lambda > 0$ be sufficiently small such that $\lambda f(x, v) < \frac{\mathcal{M}}{2}$. Then for such λ ,

$$-M(I(v)) \Delta_{p(x)} v = \mathcal{M} > \lambda f(x, v) + \mu |v|^{q(x)-2} v,$$

which shows that v is a supersolution of (4.1) and is not a solution of (4.1). By Theorem 3.2, $v > 0$ in Ω and $\frac{\partial v}{\partial \mathbf{n}} > 0$ on $\partial\Omega$.

In the case when f satisfies the condition (i), 0 is a subsolution of (4.1) and 0 does not satisfy the equation in (4.1). Moreover, by Theorem 3.4, (4.1) has a solution $u_\lambda \in [0, v] \cap C^1(\overline{\Omega})$, which is a local minimizer of J_λ in the C^1 topology.

In the case when f satisfies the condition (ii), 0 satisfies the equation in (4.1). We claim that 0 is not a minimizer of J_λ on $[0, v] \cap X$. To see this, noting $J_\lambda(0) = 0$, it is sufficient to show that $\inf_{[0,v] \cap X} J_\lambda(u) < 0$. For $\delta > 0$, denote $U_\delta = \{x \in U : \text{dist}(x, \partial U) < \delta\}$. By the condition (ii), we can find sufficiently small positive constants ρ such that $\overline{B}(x_0, \rho) \subset U \setminus U_\delta$, $r_0 < p^-(U_\delta) := \inf\{p(x) : x \in U_\delta\}$. Define a function $w \in C_0^\infty(U)$ such that $0 \leq w \leq 1$ and $w = 1$ on $U \setminus U_\delta$. Then for sufficiently small $1 > t > 0$, we have that $tw \in [0, v]$ and

$$\begin{aligned} J_\lambda(tw) &\leq \widehat{M} \left(\int_{U_\delta} \frac{t^{p(x)}}{p(x)} |\nabla w|^{p(x)} dx \right) - \lambda \int_{U \setminus U_\delta} F(x, tw) dx \\ &\leq M \left(\int_{U_\delta} \frac{t^{p(x)}}{p(x)} |\nabla w|^{p(x)} dx \right) \int_{U_\delta} \frac{t^{p(x)}}{p(x)} |\nabla w|^{p(x)} dx - \lambda \int_{U \setminus U_\delta} F(x, tw) dx \\ &\leq t^{p^-(U_\delta)} M \left(\int_{U_\delta} \frac{1}{p(x)} |\nabla w|^{p(x)} dx \right) \int_{U_\delta} \frac{1}{p(x)} |\nabla w|^{p(x)} dx - c_1 \lambda t^{r_0} \int_{U \setminus U_\delta} w^{r_0} dx \\ &< 0, \end{aligned}$$

which shows that the claim is true. By Theorem 3.4, there exists $u_\lambda \in [0, v] \cap C^{1,\alpha}(\overline{\Omega})$ such that $J_\lambda(u_\lambda) = \inf_{[0,v] \cap X} J_\lambda(u)$, u_λ is a solution of (4.1) and u_λ is a local minimizer of J_λ in the C^1 topology.

When $\lambda \rightarrow 0$, we can take $\mathcal{M} \rightarrow 0$, consequently $|v_{\mathcal{M}}|_\infty \rightarrow 0$ and $|u_\lambda|_\infty \rightarrow 0$. Furthermore, $\|v_{\mathcal{M}}\|_X \rightarrow 0$ and $\|v_{\mathcal{M}}\|_{C^1(\overline{\Omega})} \rightarrow 0$. Assertion (1) is proved.

(2) The proof is similar to the proof of [3], we omit it here (for details, see the proof of Theorem 4.1 in [3]).

(3) Note that, under additional assumptions, it is easy to verify that $J_\lambda \in C^1(X, \mathbb{R})$ and J_λ satisfies the $(PS)_c$ condition for all λ . Now let $\lambda \in \text{int } \Lambda \subset \Lambda_0$ be given arbitrarily. Take $\lambda_1, \lambda_2 \in \Lambda_0$ with $\lambda_2 < \lambda < \lambda_1$, and let u_{λ_1} , u_λ and u_{λ_2} be the solutions of (4.1) $_{\lambda_1}$, (4.1) $_\lambda$ and

(4.1_{λ₂}) respectively, $u_{λ_2} \leq u_\lambda \leq u_{λ_1}$, and let u_λ be a local minimizer of J_λ in the C^1 topology. Then by Theorem 3.1, u_λ is also a local minimizer of J_λ in the $W^{1,p(x)}$ topology. Define

$$\begin{aligned} \tilde{f}_\lambda(x, t) &= \begin{cases} f(x, t) & \text{if } t > u_\lambda(x), \\ f(x, u_\lambda(x)) & \text{if } t \leq u_\lambda(x), \end{cases} \\ \tilde{g}_\lambda(x, t) &= \begin{cases} t^{q(x)-1} & \text{if } t > u_\lambda(x), \\ (u_\lambda(x))^{q(x)-1} & \text{if } t \leq u_\lambda(x). \end{cases} \end{aligned}$$

Consider the problem

$$\begin{cases} -M \left(\int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx \right) \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) = \lambda \tilde{f}_\lambda(x, u) + \mu \tilde{g}_\lambda(x, u) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega, \end{cases} \quad (4.2)$$

and denote the associated functional to (4.2) by \tilde{J}_λ . It is easy to see that $u_{λ_2}$ and $u_{λ_1}$ are a subsolution and a supersolution of (4.2), respectively. By Theorem 3.4, there exists $u_\lambda^* \in [u_{λ_2}, u_{λ_1}] \cap C^1(\bar{\Omega})$ such that u_λ^* is a solution of (4.2) and is a local minimizer of \tilde{J}_λ in the C^1 topology. By Theorem 2.2(1), we can see that $u_\lambda^* \geq u_\lambda$ and consequently u_λ^* is also a solution of (4.1_λ). If $u_\lambda^* \neq u_\lambda$, then assertion (3) already holds, hence we can assume that $u_\lambda^* = u_\lambda$. Now u_λ is a local minimizer of \tilde{J}_λ in the C^1 topology, and so also in the $W^{1,p(x)}$ topology. We can assume that u_λ is a strictly local minimizer of \tilde{J}_λ in the $W^{1,p(x)}$ topology, otherwise we have obtained assertion (3). It is easy to verify that, under the additional assumptions in the statement (3), $\tilde{J}_\lambda \in C^1(X, \mathbb{R})$ and \tilde{J}_λ satisfies the (PS)_c condition. From $q^- > p^+$, (M_0) and $\mu > 0$, it follows that $\inf\{\tilde{J}_\lambda(u) : u \in X\} = -\infty$. Using Lemma 4.1, we know that (4.2) has a solution v_λ such that $v_\lambda \neq u_\lambda$, as a solution of (4.2), v_λ must satisfy $v_\lambda \geq u_\lambda$, and consequently, by Theorem 3.2 and Theorem 3.3, $v_\lambda > u_\lambda$. Noting that v_λ is also a solution of (4.1_λ) since $v_\lambda \geq u_\lambda$, thus the proof of assertion (3) is complete. \square

Note that in the case of Theorem 4.1(1) and (2), the variational method cannot be used directly because we do not suppose that $q(x) \leq p^*(x)$ and do not restrict the growth rate of f .

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

GD conceived of the study, and participated in its design and coordination and helped to draft the manuscript. XH participated in the design of the study. All authors read and approved the final manuscript.

Acknowledgements

Research was supported by the NSFC (No. 11261052, No. 11101335, No. 11061030).

Received: 4 September 2012 Accepted: 15 November 2012 Published: 5 December 2012

References

1. Diening, L, Harjulehto, P, Hästö, P, Ruzicka, M: Lebesgue and Sobolev Spaces with Variable Exponents. Lecture Notes in Mathematics, vol. 2017. Springer, Berlin (2011)
2. Fan, XL: On nonlocal $p(x)$ -Laplacian Dirichlet problems. *Nonlinear Anal.* **72**, 3314-3323 (2010)
3. Fan, XL: On the sub-supersolution methods for $p(x)$ -Laplacian equations. *J. Math. Anal. Appl.* **330**, 665-682 (2007)
4. Amann, H: Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces. *SIAM Rev.* **18**, 620-709 (1976)
5. Kirchhoff, G: *Mechanik*. Teubner, Leipzig (1883)

6. Dai, G, Ma, R: Solutions for a $p(x)$ -Kirchhoff type equation with Neumann boundary data. *Nonlinear Anal., Real World Appl.* **12**, 2666-2680 (2011)
7. Acerbi, E, Mingione, G: Regularity results for a class of functionals with nonstandard growth. *Arch. Ration. Mech. Anal.* **156**, 121-140 (2001)
8. Fan, XL, Zhao, D: A class of De Giorgi type and Hölder continuity. *Nonlinear Anal.* **36**, 295-318 (1996)
9. Fan, XL: Global $C^{1,\alpha}$ regularity for variable exponent elliptic equations in divergence form. *J. Differ. Equ.* **235**, 397-417 (2007)
10. Dai, G: Three solutions for a nonlocal Dirichlet boundary value problem involving the $p(x)$ -Laplacian. *Appl. Anal.*, 1-20 (2011) (iFirst). doi:10.1080/00036811.2011.602633
11. Ma, R, Dai, G, Gao, C: Existence and multiplicity of positive solutions for a class of $p(x)$ -Kirchhoff type equations. *Bound. Value Probl.* **2012**, 16 (2012)
12. Drábek, P, Hernández, J: Existence and uniqueness of positive solutions for some quasilinear elliptic problems. *Nonlinear Anal.* **44**, 189-204 (2001)
13. Ladyzenskaja, OA, Ural'tzeva, NN: *Linear and Quasilinear Elliptic Equations*. Academic Press, New York (1968)
14. Fan, XL: A Brezis-Nirenberg type theorem on local minimizers for $p(x)$ -Kirchhoff Dirichlet problems and applications. *Differ. Equ. Appl.* **2**(4), 537-551 (2010)
15. Fan, XL, Zhao, YZ, Zhang, QH: A strong maximum principle for $p(x)$ -Laplace equations. *Chin. J. Contemp. Math.* **24**(3), 277-282 (2003)
16. Willem, M: *Minimax Theorems*. Birkhäuser, Boston (1996)

doi:10.1186/1029-242X-2012-283

Cite this article as: Han and Dai: On the sub-supersolution method for $p(x)$ -Kirchhoff type equations. *Journal of Inequalities and Applications* 2012 **2012**:283.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
