

Oral presentation

Open Access

Sparse coding of natural communication signals in midbrain neurons

Katrin Vonderschen* and Maurice J Chacron

Address: Department of Physiology, McGill University, Montreal, Quebec H3G 1Y6, Canada

Email: Katrin Vonderschen* - katrin.vonderschen@mail.mcgill.ca

* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):O3 doi:10.1186/1471-2202-10-S1-O3This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/O3>

© 2009 Vonderschen and Chacron; licensee BioMed Central Ltd.

Sparse neural codes (i.e. codes in which neurons respond only to a few stimuli) have been widely observed across animal taxa [1-4]. Theoretical studies suggest that sparse neural codes critically depend on non-linear mechanisms [5]. However, the network and cellular properties that enable the observed sparse responses remain unclear. We investigated sparse coding in neurons within the mid-brain torus semicircularis (TS) in the weakly electric fish *Apteronotus leptorhynchus*, which is equivalent to the inferior colliculus in the mammal. These fish generate a quasi-sinusoidal electric field via the electric organ discharge (EOD) with a characteristic frequency that varies across individuals. When two individuals come into contact, interference between their EODs will give rise to a beat phenomenon. Male *Apteronotus leptorhynchus* will transiently increase their EOD frequencies in a stereotypical manner during agonist encounters or courtship rituals: these chirps will occur in conjunction with the beat and must be distinguished by either the other male or the female (Figure 1). We performed *in vivo* patch clamp recordings to study TS neural responses to chirps occurring on top of the beat pattern. We found that one neuron type responded almost exclusively to chirps in a most peculiar manner. These neurons had little or no activity during the beat and fired a single action potential in response to the chirp (Figure 2). Chirp detection was negatively correlated with phase locking to the beat suggesting a segregation of information flow in midbrain neurons. Moreover, the chirp detection abilities were highly superior to those found in neurons afferent to TS.

This indicates that separate streams of information arise in the torus and may be used to guide different behaviors by higher brain regions. We investigated the cellular mechanisms underlying this sparse coding. A combination of mathematical modeling and further experiments indicate that shunting inhibition hyperpolarizes the membrane potential below the firing threshold during the beat and increases the membrane conductance which promotes coincidence detection of synaptic input caused by the chirp. We speculate that a potassium conductance then hyperpolarizes the membrane potential after the first action potential: thereby enabling these neurons to




Figure 1
Chirps are transient frequency rises superimposed on top of the beat caused by two fish's EOD.




Figure 2
Response of midbrain neuron to natural communication signal "chirp".

respond to each chirp with a single action potential. Our findings reveal cellular mechanisms for the segregation of information flow and sparsening of neural responses to behaviorally relevant stimuli. These mechanisms may generalize to other sensory systems with which the electrosensory system shares similarities: in particular the auditory system.

Acknowledgements

This research was supported by CIHR, CFI, and CRC (MJC).

References

1. Quiroga RQ, et al.: *Nature* 2005, **435**:1102-1107.
2. Vinje WE, Gallant JL: *Science* 2000, **287**:1273-1276.
3. Perez-Orive J, et al.: *Science* 2002, **297**:359-365.
4. Hahnloser RH, Kozhevnikov AA, Fee MS: *Nature* 2002, **419**:65-70.
5. Kreiman G, et al.: *Neuron* 2006, **49**:433-445.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."
 Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

