

Heterologous expression and transcript analysis of gibberellin biosynthetic genes of grasses reveals novel functionality in the *GA3ox* family

Pearce *et al.*

RESEARCH ARTICLE

Open Access



Heterologous expression and transcript analysis of gibberellin biosynthetic genes of grasses reveals novel functionality in the *GA3ox* family

Stephen Pearce¹, Alison K Huttly², Ian M Prosser², Yi-dan Li^{2,3}, Simon P Vaughan², Barbora Gallova², Archana Patil², Jane A Coghill⁴, Jorge Dubcovsky^{1,5}, Peter Hedden² and Andrew L Phillips^{2*}

Abstract

Background: The gibberellin (GA) pathway plays a central role in the regulation of plant development, with the 2-oxoglutarate-dependent dioxygenases (2-ODDs: *GA20ox*, *GA3ox*, *GA2ox*) that catalyse the later steps in the biosynthetic pathway of particular importance in regulating bioactive GA levels. Although GA has important impacts on crop yield and quality, our understanding of the regulation of GA biosynthesis during wheat and barley development remains limited. In this study we identified or assembled genes encoding the GA 2-ODDs of wheat, barley and *Brachypodium distachyon* and characterised the wheat genes by heterologous expression and transcript analysis.

Results: The wheat, barley and *Brachypodium* genomes each contain orthologous copies of the *GA20ox*, *GA3ox* and *GA2ox* genes identified in rice, with the exception of *OsGA3ox1* and *OsGA2ox5* which are absent in these species. Some additional paralogs of 2-ODD genes were identified: notably, a novel gene in the wheat B genome related to *GA3ox2* was shown to encode a GA 1-oxidase, named as *TaGA1ox-B1*. This enzyme is likely to be responsible for the abundant 1 β -hydroxylated GAs present in developing wheat grains. We also identified a related gene in barley, located in a syntenic position to *TaGA1ox-B1*, that encodes a GA 3,18-dihydroxylase which similarly accounts for the accumulation of unusual GAs in barley grains. Transcript analysis showed that some paralogs of the different classes of 2-ODD were expressed mainly in a single tissue or at specific developmental stages. In particular, *TaGA20ox3*, *TaGA1ox1*, *TaGA3ox3* and *TaGA2ox7* were predominantly expressed in developing grain. More detailed analysis of grain-specific gene expression showed that while the transcripts of biosynthetic genes were most abundant in the endosperm, genes encoding inactivation and signalling components were more highly expressed in the seed coat and pericarp.

Conclusions: The comprehensive expression and functional characterisation of the multigene families encoding the 2-ODD enzymes of the GA pathway in wheat and barley will provide the basis for a better understanding of GA-regulated development in these species. This analysis revealed the existence of a novel, endosperm-specific GA 1-oxidase in wheat and a related GA 3,18-dihydroxylase enzyme in barley that may play important roles during grain expansion and development.

Keywords: Gibberellin, Wheat, Biosynthesis, Signalling, Gene sequences, De novo assembly, Transcriptomics, Heterologous expression, GA 1-oxidase

* Correspondence: andy.phillips@rothamsted.ac.uk

²Department of Plant Biology and Crop Science, Rothamsted Research, Harpenden AL5 2JQ, UK

Full list of author information is available at the end of the article

Background

Gibberellins (GAs) are a group of plant secondary products based on the diterpenoid *ent*-gibberellane skeleton; a small subset of bioactive GAs such as GA₄ and GA₁ act as plant hormones and participate in a wide range of developmental processes. Although classically involved in the promotion of growth processes such as germination and stem elongation, GA signalling has also been shown to be important in root elongation [1], lateral root formation [2], skotomorphogenesis [3], cambial activity [4], leaf expansion [5], trichome development [6], floral induction [7], anther and pollen development (reviewed in Plackett et al.[8]), fruit growth [9] and seed development [10]. Furthermore, GAs mediate environmental effects on growth and development through modulation of both biosynthetic and signalling components [11]. The central components of GA signalling, GRAS-domain proteins containing an N-terminal “DELLA” motif that repress growth, also act as nodes in the interactions with several other plant hormones, including jasmonate [12], brassinosteroids [13] and strigolactones [14]. In short, GAs play a central role in plant development and environmental responses, with impacts on crop yield and quality.


The importance of GA signalling in determining plant stature is clear from evidence in both wild and crop species showing phenotypic effects of genetic variation in GA biosynthetic and signalling genes. In wheat (*Triticum aestivum* L.), semi-dwarfing alleles of the *Rht* DELLA genes were key to increasing yield during the Green Revolution as not only did the shorter stature protect against lodging under high fertiliser application rates, but also enhanced harvest index by reducing straw biomass and, in many genetic backgrounds, increasing grain numbers per ear [15]. In rice (*Oryza sativa* L.), a similar height phenotype was conferred by loss-of-function mutations in a key GA biosynthetic gene, *OsGA20ox2* [16], and there is evidence that semi-dwarfing of barley (*Hordeum vulgare* L.) by the *sdw1/denso* gene is associated with reduced expression of the orthologous *HvGA20ox2* gene [17].

Although *Rht* semi-dwarfing alleles are widespread in modern wheat varieties, the involvement of *Rht* in all GA responses results in pleiotropic effects on many other traits. For example, even mild alleles such as *Rht-B1b* and *Rht-D1b* impart reduced leaf area [18]. These alleles also have a strong effect on coleoptile elongation which prevents deeper sowing under dry conditions [19]. Better targeting of the dwarfing effect to stem tissues might therefore confer significant advantages. In contrast to *Rht*, the enzymes acting during the latter stages of GA biosynthesis are encoded by multiple paralogs with overlapping domains of expression and, in *Arabidopsis*, mutations in individual genes have more localised effects [20–22]. This suggests that the characterisation of the

GA biosynthetic genes of wheat has the potential to identify targets for the development of novel semi-dwarfing alleles with fewer undesirable pleiotropic effects than the current *Rht* alleles.

GAs are also thought to play a critical role in wheat grain development: endogenous GA levels are very high in developing grain and increase during grain expansion [10] and wheat lines containing *Rht* alleles have smaller grains [15]. Despite grain size being an important component of wheat yield and quality, little is known regarding the spatial or temporal regulation of GA biosynthesis and signalling in the grain. A fuller understanding of the role of GA during grain development is required to engineer improvements in this trait in modern wheat varieties.

The GA biosynthetic pathway has been extensively characterized in both rice and *Arabidopsis* (reviewed by Yamaguchi, 2008 [23]), and the early genes in the pathway, from copalyl diphosphate synthase (CPS) to *ent*-kaurenoic acid oxidase (KAO) that produce the GA precursors GA₁₂ and GA₅₃ have also been identified and characterised in wheat [24–26]. The final steps in GA biosynthesis and inactivation are catalysed by soluble 2-oxoglutarate-dependent dioxygenases (2-ODDs) (Fig. 1). GA 20-oxidase catalyses the multi-step oxidation of GA₁₂ and GA₅₃ to form the C₁₉ skeleton, while GA 3-oxidase produces the final bioactive products, GA₄ and GA₁. A third class of 2-ODD, GA 2-oxidase, is involved in inactivation, with two



sub-classes of enzyme that act against either bioactive GAs (GA_4 , GA_1) and their immediate C_{19} precursors (GA_9 , GA_{20}) [27] or against C_{20} -GAs earlier in the pathway (e.g., GA_{12} , GA_{53}) [28]. However, our knowledge of the size, structure and function of the gene families encoding these enzymes in wheat is limited.

Most of the evidence points to dynamic regulation of GA biosynthesis through regulation of the 2-ODD genes which, in contrast to the terpene cyclase and cytochrome P450 genes earlier in the pathway [24], are encoded by small multigene families [23]. Although a small number of the 2-ODDs involved in the GA pathway of wheat have been identified previously [24, 29], this study represents the first comprehensive analysis of the paralogous and homoeologous genes encoding the enzymes in this crop species. In this report we identify and characterise the biochemical function of GA-biosynthetic 2-ODD genes in wheat and demonstrate novel biochemical functions. We also catalogue the orthologous genes, where data is available, in durum wheat, barley and *Brachypodium distachyon*. Finally, we identify tissue-specific patterns of expression in wheat

that suggest specialized roles in plant development for the different paralogs.

Results

Identification of wheat genes involved in GA metabolism

Our strategy to identify the wheat complement of genes of the 2-ODD class from the GA biosynthetic pathway, previously catalogued in rice [30, 31] is outlined in Additional file 1 (Figure S1) and involved first identifying orthologous genes from *Brachypodium* by BLASTP search at www.phytozome.org (Table 1 and Additional file 2) using the rice peptide sequences as queries. Coding sequences from the *Brachypodium* genes were then used to BLAST partially-assembled genomic survey sequence from the International Wheat Genome Sequencing Consortium (IWGSC), generated by Illumina sequencing of DNA from individual wheat chromosome arms [32]. We were thus able to identify high-quality contigs containing wheat orthologs of many of the rice and *Brachypodium* GA 2-ODD genes.

Genes absent from, or incomplete in, the wheat genomic survey data were initially assembled from shotgun

Table 1 Rice, *Brachypodium* and bread wheat genes encoding 2-ODDs from the GA pathway

Target	Rice gene	Brachypodium gene	<i>T. aestivum</i> chromosome arm assembly		
			A	B	D
GA20ox1	<i>Os03g63970</i>	<i>Bradi1g00950</i>	FL (4AL)	FL (5BL)	FL (5DL)
GA20ox2	<i>Os01g66100</i>	<i>Bradi2g57030</i>	Partial ^a (3A)	FL ^a (3B)	FL ^a (3D)
GA20ox3	<i>Os07g07420</i>	<i>Bradi1g56200</i>	Partial ^a (3A)	FL (3B)	FL ^a (3D)
GA20ox4	<i>Os05g34854</i>	<i>Bradi2g24980</i>	FL ^a (1AL)	FL ^{a,b} (1BL)	FL ^{a,b} (1DL)
GA3ox1	<i>Os05g08540</i>	-	-	-	-
GA3ox2	<i>Os01g08220</i>	<i>Bradi2g04840(a) Bradi4g23570(b)</i>	Partial ^{a,c} (3A)	FL (3B)	FL ^a (3D)
GA3ox3	-	-	FL ^a (2AL)	FL ^a (2BL)	FL ^{a,d} (2DL)
GA1ox1	-	-	-	FL (2BL)	-
GA2ox1	<i>Os05g06670</i>	<i>Bradi2g34840</i>	FL ^a (1AS)	FL (1BS)	FL (1DS)
GA2ox2	<i>Os01g22910</i>	<i>Bradi2g12440</i>	-	Partial (7BL)	FL ^a (7D)
GA2ox3	<i>Os01g55240</i>	<i>Bradi2g50280</i>	Partial ^e (3AL)	FL (3B)	FL ^a (3D)
GA2ox4	<i>Os05g43880</i>	<i>Bradi2g19900</i>	FL ^a (1AL)	FL ^a (1B)	FL ^a (5BL)
GA2ox5	<i>Os07g01340</i>	<i>Bradi1g59570</i>	-	-	-
GA2ox6	<i>Os04g44150</i>	<i>Bradi5g16040</i>	FL ^a (2AL)	FL (2BL)	FL ^a (2DL)
GA2ox7	<i>Os01g11150</i>	<i>Bradi2g06670</i>	Partial (3AS)	FL ^a (3B)	FL ^a (3DS)
GA2ox8	<i>Os05g48700</i>	<i>Bradi2g16730(a) Bradi2g16750(b)</i>	FL ^a (1AL)	Partial (1BL)	FL ^{a,b} (5BL)
GA2ox9	<i>Os02g41954</i>	<i>Bradi3g49390</i>	FL (6AL)	Partial (6BL)	FL ^{a,b} (6DL)
GA2ox10	<i>Os05g11810</i>	<i>Bradi2g32580</i>	FL ^a (1AS)	FL (1BS)	FL (1DS)
GA2ox11	-	-	FL (4AS)	FL (4BL)	FL ^a (4DL)
GA2ox12	-	-	-	FL (4BL)	-
GA2ox13	-	-	-	FL (4BL)	-

Notes: FL - full length; ^aGenerated by reassembly of IWGSC chromosome arm reads; ^bMissing data in intron; ^cFL cDNA from cv. Maris Huntsman [29]; ^d7 bp insertion in exon 2; ^eFL cDNA from cv. Avalon (Prosser & Phillips, unpublished). Accession numbers for the wheat genes are in Additional file 2.

genomic reads of wheat cv. Chinese Spring [33] located at <http://www.cerealsdb.uk.net>. The raw 454 reads were identified by BLASTN with the Brachypodium CDS sequences and assembled at high stringency; for most targets this resulted in the identification of genomic contigs covering the bulk of the coding region of the gene; however, the low genome coverage of the raw data, approximately 5x, [33] coupled with the relatively high error rate of 454 sequencing and the hexaploid genome of wheat resulted in contigs that contained ambiguous bases and, in many cases, chimeric sequences derived from more than one homoeolog. Hence, these consensus sequences were used in BLAST searches of the unassembled Illumina data from individual chromosome arms [32], followed by mapping the Illumina reads to the 454 assemblies to generate homoeolog-specific sequences, as annotated in Table 1 and, in greater detail, in Additional file 2. Thus, across the *GA20ox*, *GA3ox* and *GA2ox* gene families, we were able to identify or assemble at least one complete homoeolog, and often all three, for most of the target 2-ODD genes, as described in detail below. In addition, we identified or assembled wheat sequences encoding the GA 13-hydroxylase (*GA13ox*) biosynthetic enzymes and for the *GID1* and *GID2* signalling components.

The genomic survey data from bread wheat cv. Chinese Spring was complemented with assembled RNA-seq data from tetraploid durum wheat (*T. turgidum* L. subsp. *durum* Desf.) cv. Kronos and the diploid progenitor *T. urartu* [34]. In addition, we identified, where possible, likely orthologs of each gene in the genomic survey sequences of *T. urartu* [35] and *Aegilops tauschii* [36], the diploid progenitors of the A and D genomes, respectively, of bread wheat (Additional file 2). Homoeolog-specific sequences from the tetraploid species were assigned to the A or B genomes by BLAST to the bread wheat chromosome arm-specific genomic survey data above (Table 1). Finally, we identified likely orthologs of each of the wheat genes in barley (Additional file 2), within the recent draft genome sequence of this species [37].

Structure and biochemical function of wheat GA 2-ODD genes

We identified likely Brachypodium orthologs for each of the four *GA20ox* genes, two *GA3ox* genes and ten *GA2ox* genes previously described in rice [30, 31]. The only exception was *GA3ox1*, which appears to be absent from Brachypodium, as shown in Table 1 and in the phylogenetic analysis presented in Fig. 2B; however, this species contains two genes related to *GA3ox2*, as discussed below. Brachypodium also contains a tandem duplication of *GA2ox8*. As neither hexaploid bread wheat, nor its tetraploid or diploid progenitors, possesses a fully-sequenced genome, we cannot be unequivocal about the number of GA 2-ODD genes present in wheat.

However, with the exception of *GA2ox5*, we identified in bread wheat at least one homolog of each Brachypodium gene, and usually complete or partial sequence evidence of homoeologs on each of the three chromosomes. Thus we identified in bread wheat four homoeologous sets of *GA20ox* genes, at least two sets of *GA3ox* genes and at least nine sets of *GA2ox* genes, as detailed below.


GA 20-oxidases

Phylogenetic analysis of the 2-ODD genes showed that the grass *GA20ox* genes fall into four paralogous clades (Fig. 2A) each including one of the four rice *GA20ox* genes, *OsGA20ox1* through *OsGA20ox4*. The assignment of the wheat *GA20ox1*, *GA20ox2* and *GA20ox4* genes to the corresponding rice groups is supported by syntenic relationships between rice and wheat chromosomes [38]. In contrast, *GA20ox3* which was expected to be found on wheat chromosome group 2 based on its position in rice and Brachypodium, was found on the three homoeologs of chromosome group 3 [39]. We also identified full-length or partial sequences for all *GA20ox* paralogs in the tetraploid and diploid wheat species and full-length coding sequences from barley (Additional file 2). As previously reported [20], phylogenetic analysis indicates that the four paralogs of *GA20ox* in grass species are not directly orthologous to any of the five paralogs identified in Arabidopsis: four of the five Arabidopsis genes lie in a single clade of the tree (Fig. 2A), suggesting that the expansion in paralogs of *GA20ox* occurred after the separation of the monocot and eudicot lineages.

We have previously reported the biochemical function of all three homoeologs of *TaGA20ox1* by heterologous expression in the pET3d vector [29]. In this study, we present a similar characterization for *TaGA20ox2*, *TaGA20ox3* and *TaGA20ox4*. We expressed the coding regions of one representative homoeolog of each of these three wheat paralogs as fusion proteins in *E. coli* (see Methods) and demonstrated their ability to carry out the series of sequential oxidations of GA_{12} to GA_9 (Additional file 1: Figure S2). These results demonstrate that all four wheat paralogs encode fully active GA 20-oxidases (Fig. 1).

GA 3-oxidases

Phylogenetic relationships between rice, Brachypodium, barley and wheat are less clear in the *GA3ox* gene family. Neither Brachypodium, barley nor either of the wheat polyploid species or their progenitors appear to possess a true ortholog of *OsGA3ox1* (Fig. 2B). However, Brachypodium and wheat possess likely orthologs of *OsGA3ox2*, with three homoeologs in bread wheat as described previously [29]. Similar sequences were also found for the A genome in *T. durum* (Table 1), in *T.*



urartu, *Ae. tauschii* (as a partial genomic sequence), and barley (Additional file 2). Brachypodium contains a second sequence, *Bradi4g23570*, related to *OsGA3ox2*, but the predicted coding regions of the genes from several accessions of this species at www.brachypodium.org contain a frame shift in exon 2 due to the apparent insertion of a G residue at nucleotide 765. However, PCR amplification and sequencing of this region from *B. distachyon* Bd21 genomic DNA clearly showed the inserted base present in the database sequence to be an artefact. Removal of G765 from the database sequence resulted in a complete open reading frame most closely related to *BdGA3ox2a* (*Bradi2g04840*) (Fig. 2B) and *Bradi4g23570* was therefore assigned as *BdGA3ox2b*. Heterologous expression of synthetic coding sequences of *BdGA3ox2a*

and *BdGA3ox2b* in *E. coli* followed by incubation of bacterial lysates of these cultures with radiolabelled substrates showed that both Brachypodium genes encode GA 3-oxidase enzymes, converting GA₉ to GA₄ (Fig. 3G,H).

Syntenic relationships and phylogenetic analysis (Fig. 2B) support the assignment of wheat *GA3ox2* genes on the group 3 chromosomes as orthologous to rice *GA3ox2*. However, in bread wheat we identified four further related sequences encoding potential GA 3-oxidases, all located on the long arms of the group 2 chromosomes, some of which were also found in the tetraploid and diploid species (Additional file 2). Phylogenetic analysis and the location on chromosomes 2AL, 2BL and 2DL suggested that the three most closely-related of these four novel sequences probably form a homoeologous group (Additional file 1:





Figure S12) and therefore were named as *TaGA3ox-A3*, *TaGA3ox-B3* and *TaGA3ox-D3* (Table 1). Near-identical sequences to the A and B homoeologs were identified in the durum wheat transcript assembly [34] and a partial sequence from *T. urartu* was also identified; however, *TaGA3ox3* was not found in the *Ae. tauschii* assembly. The fourth novel gene, located on bread wheat chromosome 2BL and also identified in the durum wheat transcript assembly, was provisionally named *TaGA3ox-B4*; no homoeologs of this sequence were identified in the A or D genomes of wheat or in the A and D diploid progenitor species, although a similar sequence (79.3 % amino acid identity) had been previously identified in barley and annotated as *HvGA3ox1* [39].

We have previously determined the biochemical function of the bread wheat GA3ox2 genes by heterologous expression of the cDNAs in *E. coli* [29]: products from all three homoeologs converted GA₉ to GA₄ and GA₂₀ to GA₁, demonstrating GA 3β-hydroxylase (GA 3-oxidase)

activity. In this study we present the functional characterization of *TaGA3ox3* and *TaGA3ox4* through expression in *E. coli* of synthetic cDNAs. When lysates from induced bacterial cells containing synthetic cDNA constructs were incubated with [1-¹⁴C]GA₉, *TaGA3ox-B3* was shown to encode a functional GA 3-oxidase, converting the substrate to [1-¹⁴C]GA₄ (Fig. 3A), while expression products of *TaGA3ox-A3* did not have any detectable catalytic activity; *TaGA3ox-D3* was not tested as the Chinese Spring sequence contains a 7 bp insertion in exon 2, indicating that this gene is unlikely to be functionally active.

An unexpected result was observed for *TaGA3ox-B4*, which converted [1-¹⁴C]GA₉ to a product with an HPLC retention time different to that of [1-¹⁴C]GA₄ (Fig. 3C). This product was analysed by combined gas chromatography-mass spectroscopy (GC-MS) and had a mass spectrum consistent with [1-¹⁴C]GA₆₁ (1β-hydroxy-GA₉) [40], identifying *TaGA3ox-B4* as a GA 1β-hydroxylase (GA 1-oxidase; Fig. 4), the first time an enzyme with such a



catalytic activity has been described. Based on this result we propose to rename this wheat gene as *TaGA1ox-B1*.

GA_{61} was originally identified in the endosperm of developing grains of bread wheat [41] along with the more abundant $1\beta,3\beta$ -dihydroxylated form, GA_{54} (1-hydroxy- GA_4 ; Fig. 4) [42]. To determine the likely sequence of reactions leading to the production of GA_{54} in grain, expression products of *TaGA3ox-B3* and *TaGA1ox-B1* were separately incubated with ^{14}C -labelled GA_9 , GA_4 and GA_{61} in the presence of co-substrates and cofactors and the products analysed by HPLC. The 3-oxidase, *TaGA3ox-B3*, was active against both GA_9 (producing GA_4) and GA_{61} (producing GA_{54} , Fig. 3A,B) whereas the GA 1-oxidase, *TaGA1ox-B1* acted only upon GA_9 (producing GA_{61}), and not upon GA_4 (Fig. 3C,D). This suggests that the order of reactions in developing wheat grains is $GA_9 \rightarrow GA_{61} \rightarrow GA_{54}$, catalysed by *TaGA1ox-B1* and *TaGA3ox-B3*, respectively (Fig. 4).

As reported above, we identified a published sequence in barley, annotated as *HvGA3ox1* [39] but most closely related to *TaGA1ox1* and *TaGA3ox3* (Fig. 2B) and located on the syntenic barley chromosome arm, 2HL. We investigated the catalytic activity of *HvGA3ox1* and

HvGA3ox2 by heterologous expression of synthetic cDNAs in *E. coli* as above. When incubated with [^{14}C] GA_9 , *HvGA3ox2* yielded GA_4 (Fig. 3E), as expected for a GA 3-oxidase, whereas *HvGA3ox1* generated a product with an HPLC retention time different from both GA_4 and GA_{61} . GC-MS analysis of this novel product revealed that it was GA_{131} ($3\beta,18$ -dihydroxy- GA_9 ; Fig. 3F) [43], thus identifying *HvGA3ox1* as a bifunctional enzyme, a GA $3\beta,18$ -dihydroxylase (GA 3,18-oxidase). We therefore propose to rename *HvGA3ox1* as *HvGA3,18ox1*. It has been previously shown that whereas developing grains of wheat accumulate 1-hydroxy-GAs, grains from barley accumulate 18-hydroxy-GAs including GA_{131} (18-hydroxy- GA_4) and GA_{132} (18-hydroxy- GA_1) [43-45]. It therefore seems highly likely that *HvGA3,18ox1* is the only enzyme required for the production of GA_{131} and GA_{132} from GA_9 and GA_{20} , respectively, in developing barley grains, while biosynthesis of bioactive GA_4 and GA_1 from these substrates in the rest of the plant is catalysed by *HvGA3ox2* (Fig. 4).

As phylogenetic analysis (Fig. 2B) suggested a close relationship between *TaGA3ox3*, *TaGA1ox-B1* and *HvGA3,18ox1*, we investigated their chromosomal locations. All these novel genes are located on the long arms of the group 2 chromosomes of wheat and barley, respectively. To further refine the syntenic relationships, the POPSEQ mapping data of wheat [32] was interrogated and showed the contig containing *TaGA3ox-A3* to be located on chromosome 2AL at 120.3 cM while *TaGA1ox-B1* was on chromosome 2BL at 134.03 cM; the contigs containing *TaGA3ox-B3* and *TaGA3ox-D3* were absent from the POPSEQ data. Predicted genes from the wheat contigs mapped to the same location as *TaGA3ox-A3* and *TaGA1ox-B1* were screened by BLASTN against the pseudomolecule of barley chromosome 2H [37], on which *HvGA3,18ox1* is located at 608.9 Mbp; 95 % of the wheat genes in the same mapping bin as *TaGA3ox-A3* had a top BLAST hit on barley chromosome 2H within 3 cM of *HvGA3,18ox1*, while 90 % of the genes co-locating with *TaGA1ox-B1* also had a top BLAST hit within the same window. This suggests that *TaGA3ox3*, *TaGA1ox-B1* and *HvGA3,18ox1* are in orthologous positions in the wheat and barley genomes and are likely to be derived from a common ancestral gene, as suggested by the phylogenetic analysis (Fig. 2B). Similar BLAST searches of the rice and Brachypodium genomes with the wheat genes flanking *TaGA3ox-A3* and *TaGA1ox-B1* did not identify any linkage to *OsGA3ox1*, *OsGA3ox2*, *BdGA3ox2a* or *BdGA3ox2b*.

GA 2-oxidases

Ten *GA2ox* genes, *OsGA2ox1* through *OsGA2ox10*, have been described in rice (Table 1); although the biochemical function of some of the rice genes has been

demonstrated by heterologous expression in *E. coli* (e.g., *OsGA2ox1*, [46]; *OsGA2ox5* [47]), by transactivation in rice by T-DNA insertion (e.g., *OsGA2ox3*, [46]) or by ectopic expression in transgenic plants (e.g., *OsGA2ox5*, [46]), most have not been fully characterised. Based on phylogenetic analysis of protein sequences from a number of dicot species, Lee and Zeevaart [48] proposed three structural classes of GA2ox enzymes. A phylogenetic analysis of GA2ox sequences from Arabidopsis, rice, and Brachypodium suggested that the grass enzymes can each be assigned to one of these classes (Fig. 2C). Class I, exemplified by AtGA2ox1, -2 and -3, includes the rice and Brachypodium paralogs GA2ox3, -4, -7, -8 and -10; Class II contains AtGA2ox4 and -6, and the grass paralogs GA2ox1 and -2; Class III is represented by AtGA2ox7 and -8 and the grass paralogs GA2ox5, -6 and -9. Previous data suggest that most GA2ox enzymes in Classes I and II almost exclusively use C₁₉-GAs as substrates, while class III enzymes reportedly metabolize only C₂₀-GAs.

In wheat, bioinformatic analysis of assembled and raw chromosome arm data revealed likely orthologs for each of the rice and Brachypodium GA2ox genes (Table 1) with the sole exception of *OsGA2ox5*, which was not detected in any wheat species, or in barley (Additional file 2). In bread wheat, *T. urartu* and *Ae. tauschii* we identified an additional group of GA2ox genes on the homoeologous group 4 chromosomes, that were most similar to *TaGA2ox6*; the bread wheat genes were named as *TaGA2ox-A11*, *-B11* and *-D11* and we identified two further related paralogs in the bread wheat genome assembly, *TaGA2ox-B12* and *TaGA2ox-B13*, both on chromosome 4BL (Table 1) and not detected in the A and D genomes.

In general, the chromosomal locations of the wheat GA2ox genes as indicated by the chromosome arm survey data was as predicted by synteny with rice [38]. However, *GA2ox2* is located on rice chromosome 1 and the orthologous genes in wheat would be expected to be found on the group 3 chromosomes but instead are on the group 7 chromosomes (Table 1). Also, the A and B homoeologs of *TaGA2ox4* and *TaGA2ox8* were found on the long arms of the wheat group 1 chromosomes, as predicted from synteny with rice, but in both cases no homoeolog was found on chromosome 1D, although partial sequences with high nucleotide sequence identity (94–98 %) were identified in the *Ae. tauschii* assembly (Additional file 2). However, genes very closely related to *TaGA2ox4* and *TaGA2ox8* and to the candidate orthologous sequences in *Ae. tauschii* were identified in the chromosome arm assembly for 5BL, and these genes were tentatively named *TaGA2ox-D4(5BL)* and *TaGA2ox-D8(5BL)*.


To confirm the biochemical activities of wheat GA2ox genes we expressed one representative of each paralogous group as a fusion protein in *E. coli* and tested for activity against C₂₀ (GA₁₂) and C₁₉ (GA₉) substrates.

The activity detected in each bacterial lysate was as predicted by the phylogenetic analysis (Fig. 2C): TaGA2ox-D1, -D2, -B3, -D4, -D7, -D8 and -D10 were all active against the C₁₉ substrate, [1-¹⁴C]GA₉, while TaGA2ox-D6 and TaGA2ox-D9 were active against the C₂₀ substrate, [1-¹⁴C]GA₁₂, (Additional file 1: Figure S3); no activity against either substrate was detected for any of the three homoeologs of TaGA2ox11; TaGA2ox-B12 and TaGA2ox-B13 were not tested. TaGA2ox-D2, -B3, -D4 and -D10 also further oxidised the GA₅₁ product of GA₉ to its catabolite, which is almost certainly derived by rearrangement of the ketone, 2-oxo-GA₉, formed by a second round of oxidation at C-2. In contrast to most species, however, we found that some of the wheat enzymes showed markedly reduced substrate specificity towards C₂₀- or C₁₉-GAs. Notably, TaGA2ox-B3, TaGA2ox-D4 and TaGA2ox-D10, all from GA2ox Class I by phylogeny (Fig. 2C), efficiently converted GA₁₂ to GA₁₁₀ (2β-hydroxy-GA₁₂) while TaGA2ox-D6, a Class III enzyme, converted GA₉ to GA₅₁ (2β-hydroxy-GA₉) (Additional file 1: Figure S3). TaGA2ox9, also in Class III, showed partial activity against GA₉, producing an unidentified product with a retention time different to both GA₅₁ and its catabolite.

Transcript levels for GA biosynthetic and signalling gene expression in wheat tissues by RNA-seq

To determine the relative expression levels of the wheat GA genes across the life cycle of wheat, we exploited a dataset of RNA-seq samples derived from five different organs (root, leaf, stem, spike, and grain) each at three developmental stages of bread wheat cv. Chinese Spring, generated as part of the analysis of chromosome 3B [49]. The paired-end RNA-seq reads were mapped to a transcriptome reference consisting of all the wheat coding sequences identified above together with non-redundant cDNA sequences from the IWGSC wheat chromosome arm survey (see Methods). Mean fragments per kb per million mapped reads (FPKM) values (from biological duplicates) for each homoeolog of each gene are presented in Additional file 3 and histograms of expression levels of each gene family, summing the FPKM values from each homoeolog, are shown in Fig. 5.

The genes encoding the early enzymes in GA biosynthesis, catalysing the steps from GGDP to GA₅₃, were found to be expressed in all tissues and stages (Fig. 5A and Additional file 3), although the homoeologous gene sets for *TaCPS*, *TaKS*, *TaKO* and *TaKAO* were more highly expressed in the spike at anthesis than in most other tissues. The *TaKO* genes, particularly *TaKO-D1*, also appear to be very highly expressed late in developing grain at Zadoks stage 85 (Additional file 3), although the physiological basis for this is unclear. GA_{13ox}, which catalyses the 13-hydroxylation of GA₁₂ to form GA₅₃, is encoded by two paralogs as in rice [50]. The homoeologues of



TaGA13ox1 are more highly expressed than those of *TaGA13ox2* in all tissues except mature spikes (Fig. 5A and Additional file 3) and in most organs the *GA13ox* genes have their highest levels of expression in the most mature developmental stage sampled.

The biosynthetic 2-ODD gene families, *GA20ox* and *GA3ox*, showed tissue specific expression profiles (Fig. 5B). *TaGA20ox1* and *TaGA20ox2* were the most highly expressed *GA20ox* genes in vegetative tissues, while *TaGA20ox4* was highest in the spike at anthesis and *TaGA20ox3* expression was very high in and almost completely restricted to expanding grain. *TaGA20ox1* was the most highly expressed *GA20ox* in roots, whereas in leaves and stems *TaGA20ox1* and *TaGA20ox2* showed a contrasting pattern, with the former having higher expression early in development of the organ and the latter being higher towards tissue maturity. The tissue specificity of *GA3ox* paralogs was even more striking: *TaGA3ox2* appeared to be responsible for GA biosynthesis in vegetative and floral organs, while *TaGA1ox-B1* and *TaGA3ox3* were expressed at a very high level and almost exclusively at the mid-way stage of grain development, although *TaGA1ox-B1* also showed some expression in roots (Fig. 5B).

The *GA2ox* gene family, responsible for GA inactivation, showed variation in expression between both genes and tissue/time points (Fig. 5C). Several genes had very low or undetectable expression: no transcripts from *TaGA2ox2* homoeologs were found in any tissue and *TaGA2ox1* was expressed at very low levels. The only transcripts detected from the novel wheat paralogs related to *TaGA2ox6* were found in mature leaves, from *TaGA2ox-B12* (Fig. 5C and Additional file 3). *TaGA2ox3*, *TaGA2ox4* and *TaGA2ox9* were the most highly expressed *GA2ox* genes overall, contributing most to *GA2ox* levels in roots, leaves and stems, while *TaGA2ox9* was the most highly expressed *GA2ox* in the developing spike. In addition, several other *GA2ox* genes, namely *TaGA2ox3*, *TaGA2ox6*, *TaGA2ox7* and *TaGA2ox8*, also contributed to *GA2ox* transcript levels in the spike at anthesis (Fig. 5C and Additional file 3). *TaGA2ox7* was by far the most highly expressed *GA2ox* in developing grain, with some contribution from *TaGA2ox9* at the earliest stage of grain development.

We also investigated expression of the GA signalling components *TaRht*, *TaGID1* and *TaGID2*: these genes were expressed in all tissues studied (Fig. 5D). The genes encoding the GA receptor, *TaGID1*, were most highly expressed in roots, mature leaves and elongating stems while *TaGID2* expression was highest in roots. *TaRht* showed relatively little variation in expression except for a lower level in expanding grain.

There were some noticeable differences between expression levels of homoeologs for each of the studied genes, as shown in Additional file 3. In general, where

one homoeolog was expressed at a significantly higher or lower level than the other two homoeologs, this was reflected across most of the 15 tissue/time samples, as has recently been described for homoeologous gene expression in grain tissues [51]. However, there were clear exceptions to this rule: for example, *TaKO-A1* was most highly expressed in nearly all vegetative, spike and grain samples, but *TaKO-D1* dominated in late grain development. Similarly, *TaCPS-A1* showed the lowest level of expression in most vegetative tissues, but was higher in most reproductive tissues.

qRT-PCR of GA genes in tissues of durum wheat

To support the expression patterns determined by RNA-seq analysis in bread wheat we developed qRT-PCR assays for all the biosynthetic and signalling genes (Additional file 4). As we did not have access to the tissue samples used for the RNA-seq analysis, we assessed transcript levels in existing samples of RNA from shoot, root, grain and spike tissues in the tetraploid (durum) wheat cv. Kronos [34]. Redundant primers were designed to amplify both A and B homoeologous copies of each gene where both sequences were known. Additional file 1 (Figure S4) shows the distribution of expression of each paralog between tissues. The results of this analysis broadly reflect the RNA-seq analysis above: in particular, *TaGA20ox3*, *TaGA1ox1* and *TaGA3ox3* transcripts were detected almost exclusively in grain tissues. *TaGA20ox1* and *TaGA20ox2* were expressed in vegetative tissues, mainly the shoot, although appreciable levels of *TaGA20ox4* were also found in this tissue as well as in the spike. As with the RNA-seq analysis *TaGA3ox2* was expressed in root, shoot and spike tissues but at a much lower level in grain. There was also reasonably good agreement in the expression profiles of the *GA2ox* genes: the qRT-PCR showed *TaGA2ox3* expression to be highest in roots, *TaGA2ox4* highest in shoots, *TaGA2ox7* highest in grain, while *TaGA2ox8* and *TaGA2ox9* expression was highest in the spike.

Expression patterns of Brachypodium GA genes

To gain insight into conservation of expression patterns between members of the grass family, we accessed RNA-seq reads from *Brachypodium distachyon* Bd21 [52] and mapped them to the Brachypodium reference transcriptome from Ensembl Plants (<http://plants.ensembl.org/>). This revealed a high degree of conservation of expression pattern between this species and wheat (Additional file 1: Figure S5): as in wheat, *BdGA20ox1* and *BdGA20ox2* were most highly expressed in vegetative tissues, while *BdGA20ox3* expression was highest in developing seed. Notably, *BdGA20ox4* expression was highly expressed in anthers, suggesting that the high level of *TaGA20ox4* expression observed in wheat spikes close to anthesis (Fig. 5B) may also have been in this organ. Also, while

BdGA3ox2a expression was confined to vegetative tissues, anthers and developing embryos, *BdGA3ox2b* was expressed predominantly in developing seed tissues, specifically the endosperm (Additional file 1: Figure S5B). The expression profiles of the *GA2ox* gene family were somewhat less well conserved between Brachypodium and wheat, although *GA2ox3* and *GA2ox7* were major contributors in both species. The highest level of *GA2ox* expression was in the pistil (*BdGA2ox3* and *BdGA2ox7*), anther (*BdGA2ox3* and *BdGA2ox8*) and in the embryo (*BdGA2ox8* and *BdGA2ox10*) and endosperm (*BdGA2ox7*) of developing seeds (Additional file 1: Figure S5C).

GA gene expression patterns in developing wheat grain

The above results demonstrated that the developing wheat grain exhibits high expression levels of GA biosynthetic and signalling genes, including the novel GA 1-oxidase *TaGA1ox-B1*. To investigate further these patterns of expression within the expanding grain, we generated a set of RNA-seq reads from tissue layers of developing grains of bread wheat cv. Holdfast at 12 days post-anthesis: the endosperm, the inner seed coat/pericarp layer (consisting of the aleurone, nucellar epidermis, integuments, any remaining tube cells, and cross cells) and the outer pericarp layer of immature grain, consisting of the mesocarp parenchyma, hypodermis and epidermis (Additional file 1: Figure S6A). These tissue distributions were confirmed using known cell-specific gene expression patterns (Additional file 1: Figure S6B), although it is likely that there is some cross-contamination

of the samples from adjacent layers. Single-end RNA-seq reads were mapped to the wheat transcriptome reference described above (see Methods), and average RPKM (reads per kilobase per million mapped reads) values for each gene from three biological replicates were calculated. The results in Fig. 6 represent combined counts from A, B and D homoeologs, while expression values for individual genes are presented in Additional file 5.

Taken together, RPKM expression values validated those of the whole grain sample from the previous experiments, showing that the early GA biosynthetic genes *TaKO* and *TaKAO* were highly expressed in the grain and that *TaGA20ox3*, *TaGA20ox4*, *TaGA3ox3*, and *TaGA20ox7* account for the majority of GA 20-oxidase, GA 3-oxidase, and GA 2-oxidase expression, respectively. *TaGA1ox-B1* was also highly expressed in grain. The three GA signalling genes assayed, *TaGID1*, *TaGID2* and *TaRht*, were all expressed in the grain. However, there were very clear distinctions in expression levels of these genes between different grain tissue layers.

While the cyclase genes, *TaCPS* and *TaKS*, were expressed in all three grain layers, *TaKO* and *TaKAO* were more highly expressed in the endosperm and inner pericarp than in the outer layer. In stark contrast, transcripts of both paralogs of *TaGA13ox* were absent from the endosperm and concentrated in the outer pericarp. The later GA biosynthetic genes *TaGA20ox3* and *TaGA3ox3*, and also the novel *TaGA1ox* gene, were found to be more highly expressed in the endosperm than in the seed coat/pericarp layers and, within the seed




Figure 6 Expression of GA biosynthetic and signalling genes in different layers of the developing bread wheat grain at 10 days post-anthesis. Expression levels from each homolog were summed and averaged across replicates, \pm standard error. Tissue layers are as defined in Additional File 1 (Figure S6). *GA2ox11-13* represents the sum of all additional paralogs of *TaGA20ox6*: *GA2ox11*, *-12*, and *-13*. RPKM: Reads per kb per million mapped reads

coat/pericarp, more highly in the inner layer than the outer layer (Fig. 6). *TaGA20ox4* was an exception to this trend, showing highest levels of expression in the outer pericarp layers. A contrasting trend was observed for the GA catabolic gene *TaGA20ox7*, which was expressed at negligible levels in the endosperm, but at higher levels in the seed coat/pericarp tissues and highest in the inner layer (Fig. 6). The GA signalling genes *TaGID1*, *TaGID2* and *TaRht* were expressed almost exclusively in the seed coat/pericarp layers, almost no expression being detected in the endosperm. *TaGID1* and *TaRht* were also predominantly expressed in the outer of the two seed coat/pericarp layers.

Discussion

2-Oxoglutarate-dependent dioxygenases in the GA pathway of wheat

Since GAs contribute to nearly every stage of plant development, modifying their biosynthesis, perception and turnover has great potential to engineer improved varieties of agriculturally important crops. A prerequisite for such an approach is a more complete description and characterization of the components of the GA biosynthesis and signalling pathways. In the current study, we describe the sequence and biochemical activity of the *GA20ox*, *GA3ox* and *GA2ox* gene families in wheat, building upon a previous study which described the genes encoding the enzymes catalysing the earlier stages of GA biosynthesis in wheat [24]. While the absence of a fully-sequenced wheat genome precluded a precise determination of the final number of genes in each family, the transcript and genome assemblies used for this study cover a large proportion of the wheat genes [32, 34]. For missing or incomplete genes, we also successfully re-assembled chromosome arm-specific sequence reads onto consensus sequences derived from assembly of 454 reads, allowing us to further extend the coverage, and also illustrating the potential for improving the wheat genome assembly using existing short read sequence data. Moreover, the number of 2-ODD paralogs found in wheat are very similar to those described for both rice and Brachypodium that have completely sequenced genomes, and in most cases we identified full-length or partial sequences for all three homoeologs, suggesting that the great majority of the genes from the GA pathway were identified in this study.

We identified or assembled full-length or partial sequences for all three homoeologs of four *GA20ox* genes in wheat (Table 1), corresponding to the four characterized members of this family in Brachypodium and rice (Fig. 2A). We confirmed that all four paralogs encode fully functional GA 20-oxidase enzymes, each capable of catalysing the complete series of reactions from GA_{12} to GA_9 . We similarly identified wheat homologs of all but

one of the ten characterized rice *GA2ox* genes and confirmed that representative homoeologs of each of the enzymes encoded by these genes have GA 2-oxidase activity against either C_{19} - or C_{20} -GAs as substrates. However, these assays revealed that the substrate specificities of some of the wheat *GA2ox* enzymes do not conform to the current observations in other species where enzymes in Classes I and II, structurally-related to *Arabidopsis* AtGA2ox1 through AtGA2ox6, are active against C_{19} -GAs whereas enzymes related to AtGA2ox7 and AtGA2ox8 (i.e., from Class III; Fig. 2C) are active only against C_{20} -GAs [48]. Previously, the only identified exception to this pattern was the Class I enzyme CsGA2ox4 from cucumber, which showed weak activity against the C_{20} -GA GA_{12} [53]. However, this study shows that several of the wheat enzymes have broader substrate specificities: TaGA2ox-B3, TaGA2ox-D4 and TaGA2ox-D10, all from *GA2ox* Class I by sequence similarity (Fig. 2C), were also active against C_{20} -GAs, while TaGA2ox-D6, a Class III enzyme, was active against C_{19} -GAs (Additional file 1: Figure S3). The significance of this wider substrate specificity is unclear, particularly since there do not appear to be distinct biological roles for the three structural classes of *GA2ox* enzymes or distinct class-specific expression profiles during development [21].

In addition to the *GA2ox* genes in rice, Brachypodium and barley, we identified five additional *GA2ox* gene sequences in wheat, comprising one homoeologous group on the long arm of the group 4 chromosomes, *TaGA2ox11*, plus two more individual paralogs on chromosome 4B, *TaGA2ox-B12* and *TaGA2ox-B13*. All of these additional genes are closely related to *TaGA2ox6*, suggestive of gene duplication events in the ancestors of wheat. The importance of these GA 2-oxidase genes to wheat development remains unclear, however, since expression of all three homoeologs of *TaGA2ox11* was negligible in all tissues studied and none of the encoded enzymes could be shown to exhibit GA 2-oxidase activity. Expression of the *TaGA2ox-B12* and *TaGA2ox-B13* paralogs was similarly very low in the assayed tissues.

Of the three 2-ODD families, the largest divergence from rice was observed in the GA 3-oxidases. The rice genome contains two *GA3ox* genes [54]; the wheat genes orthologous to *OsGA3ox2* were identified and characterized in a previous study and shown to act principally as GA 3-oxidases, but also with additional minor activities including GA 2-oxidase, 2,3-desaturation and even 13-hydroxylation [29]. We did not identify a clear wheat ortholog of *OsGA3ox1*, either in sequence homology or in syntenic position, and this gene was also absent from the Brachypodium genome. Instead, during the current study, we found novel *GA3ox* gene sequences in bread

wheat, three of which were closely related to one another and most likely represent a homoeologous group (*TaGA3ox3*) while a fourth gene, *TaGA1ox-B1* (initially named *TaGA3ox-B4*) was identified only in the B genome (Fig. 2B); orthologs of these genes were found in durum wheat but not in rice or Brachypodium. Of the *GA3ox3* group, only *TaGA3ox-B3* could be shown to encode an active GA 3-oxidase, converting GA₉ to GA₄; *TaGA3ox-D3* contains a 7 bp insertion in the second exon and while *TaGA3ox-A4* appeared to encode a full-length protein, no enzyme activity could be detected in the *E. coli* expression products. This might be explained by polymorphisms observed in this paralog that result in missense changes in amino acid residues conserved between all grass GA3ox sequences (Additional file 1: Figure S9), but technical problems with heterologous expression cannot be ruled out.

In contrast, expression products of *TaGA3ox-B4* were found to catalyse 1 β -hydroxylation of GA₉ to yield GA₆₁ (Figs. 3B and 4), rather than the 3 β -hydroxylation suggested by its similarity to other GA 3-oxidases. This is the first description of an enzyme with GA 1-oxidase activity, and the gene was therefore renamed as *TaGA1ox-B1*. We found a single ortholog of *TaGA1ox-B1* in durum wheat but not in the A genome or the A and D diploid progenitors, and little sequence data is available for *Ae. speltoides*, the closest living relative of the B genome progenitor. However, a search of the draft barley genome [37] identified two GA3ox-like genes: *HvGA3ox2* was shown to encode a 'normal' GA 3-oxidase (Fig. 3D) and had 95–96 % nucleotide sequence identity with the three wheat *TaGA3ox2* homoeologs. In contrast, *HvGA3ox1* [39] is most closely related to *TaGA1ox-B1* (87.6 % nucleotide sequence identity; Fig. 2B) and is located on the orthologous chromosome arm in barley, 2HL. We showed that *HvGA3ox1* encodes a GA-3 β ,18-dihydroxylase, converting GA₉ to GA₁₃₁ (Figs. 3D and 4) and was therefore renamed as *HvGA3,18ox1*.

Analysis of the POPSEQ mapping data [32] showed that both *TaGA3ox-A3* and *TaGA1ox-B1* were in orthologous positions on the group 2 chromosomes of wheat to *HvGA3,18ox1* on barley chromosome 2H. However, these genes are not orthologous to either *OsGA3ox1* or *OsGA3ox2*, which are located on rice chromosomes 5 and 1, respectively, or to the two *GA3ox2* variants in Brachypodium, where are located on chromosomes 2 and 4 from that species. It seems highly likely, therefore, that the three *Triticeae*-specific genes evolved from an unidentified common ancestor but have subsequently diverged, acquiring different catalytic activities. Amino acid sequence identity ranged from 66 % between *HvGA3,18ox1* and *TaGA3ox-A3* to 79 % between *HvGA3,18ox1* and *TaGA1ox-B1*, whereas amino acid sequence identity between the wheat and barley GA3ox2 enzymes responsible

for bioactive GA production in vegetative tissues is 95–96 %. This suggests the existence of a strong purifying selection of mutations in the *GA3ox2* genes compared with the genes of novel function identified here.

Investigation of the expression patterns of these novel GA3ox-like genes showed that *TaGA1ox-B1* is only expressed in the developing grain of both bread and durum wheat (Fig. 5 and Additional file 1: Figure S4), predominantly in the endosperm (Fig. 6) and that *TaGA3ox-B3* shows a similar expression profile, although with some expression detected in vegetative tissues; the other homoeologues of *TaGA3ox3* are also expressed in developing endosperm, but at a much lower level (Additional file 5). Similarly, inspection of the barley Gene Expression Atlas at <http://www.plexdb.org> revealed that *HvGA3,18ox1* (Affymetrix probeset contig9888_at) is only expressed in the endosperm of developing barley grains. Supporting the expression and functional analyses described above, developing wheat grains accumulate 1 β -hydroxylated GAs, including GA₅₄ (1 β -hydroxy-GA₄) and GA₅₅ (1 β -hydroxy-GA₁) [42]. Given that we showed that *TaGA1ox-B1* is active against GA₉ but not GA₄, whereas *TaGA3ox-B3* is active against both GA₉ and GA₆₁ (1 β -hydroxy-GA₉), the order of reaction *in planta* is likely to be GA₉ \rightarrow GA₆₁ \rightarrow GA₅₄, catalysed by *TaGA1ox-B1* and *TaGA3ox-B3*, respectively (Fig. 4), and it seems likely that these two enzymes also produce the 13-hydroxylated equivalent, GA₅₅, from GA₂₀ in developing grain via the same sequence of reactions. In contrast, 1-hydroxy-GAs have not been identified in developing barley grains, which accumulate a number of 18-hydroxylated GAs, including GA₁₃₁ (18-hydroxy-GA₄) [44], presumably produced through the action of *HvGA3,18ox1* on GA₉ (Fig. 4).

Tissue specificity of GA biosynthesis, signalling and turnover in wheat

Our survey of GA gene expression in five tissues of bread wheat each at three stages of development (Fig. 5) and also different tissues of durum wheat (Additional file 1: Figure S4) and Brachypodium (Additional file 1: Figure S5) indicated that all organs and time points were competent in GA biosynthesis, perception and inactivation, exhibiting measurable transcript levels for at least one paralog of each component assayed. One caveat to consider when drawing conclusions from GA biosynthetic expression data is the existence of feedback and feed-forward mechanisms which act upon the transcription of 2-ODD genes. As part of a homeostatic mechanism, increased concentrations of bioactive GA inhibit *GA20ox* and *GA3ox* expression and promote *GA2ox* expression [23]. Therefore, high expression of GA biosynthetic genes is not necessarily indicative of higher endogenous bioactive GA levels and may instead result from their up-regulation under conditions of low GA. The early biosynthetic genes,

TaCPS, *TaKS*, *TaKO* and *TaKAO*, and the signalling genes *TaRht*, *TaGID1* and *TaGID2*, are expressed at all stages of development, albeit with marked variation in transcript levels between tissues and stages. However, it is clear that the multiple paralogs of the 2-ODD genes encoding *GA20ox*, *GA3ox* and *GA2ox* have more specific patterns of expression. In particular, the expression of *TaGA20ox3*, *TaGA3ox3* and *TaGA1ox1* is very high in the grain, specifically at Zadoks 75 developmental stage, but is much lower in the other tissues assayed. This is similar to the expression patterns observed for *OsGA20ox3* and *OsGA3ox1* in rice [55], although no wheat ortholog of *OsGA3ox1* was identified; *OsGA20ox3* and *TaGA20ox3* are similarly not orthologous by chromosome location, although the phylogenetic relationship appears clear (Fig. 2A).

In the leaf and stem, *TaGA20ox1* is expressed more highly in early, rapidly-elongating stages of development while *TaGA20ox2* appears to be highest in more mature tissue, after the period of maximal elongation growth (Fig. 5B), implying that *TaGA20ox1* might be more important in determining growth rate and therefore final organ size. However, both paralogs of these two genes contribute to plant height in rice [16,56], although only *OsGA20ox2* (*SD-1*) has been exploited in plant breeding [57]. *TaGA20ox1*, -2 and -3 are also expressed to some extent at all stages of spike development, a result consistent with a recent study which showed that these three GA biosynthetic genes are up-regulated by long days in the shoot apical meristem of wheat, where bioactive GA is required for the up-regulation of floral meristem identity genes *SOC1-1* and *LFY* in spike development [7]. Relatively few genes from the *GA2ox* family show high levels of expression in the tissues assayed, and some have narrow expression domains. Thus *TaGA2ox3*, *TaGA2ox4* and *TaGA2ox9* account for most of the vegetative *GA2ox* expression, while *TaGA2ox6* and *TaGA2ox8* predominate in the flowering spike and *TaGA2ox7* in developing grain.

Analysis of GA gene expression in separate tissue layers of the developing grain revealed that the GA biosynthetic genes, notably *TaKO*, *TaGA20ox3*, *TaGA3ox3* and *TaGA1ox1*, were predominantly expressed in the endosperm. In contrast, genes involved in GA perception, signalling and turnover were expressed at only low levels in the endosperm and were instead expressed at their highest levels in the inner and outer seed coat/pericarp layers. Indeed, expression levels of the GA receptor, *TaGID1*, and the central signalling component, *TaRht*, were 26- and 22-fold higher, respectively, in the outer pericarp than in the endosperm (Fig. 6). These results suggest that the main site of GA biosynthesis in the young developing grain is the endosperm tissue while GA signalling occurs predominantly in the outer layers of the wheat grain, possibly implying movement of

bioactive GAs between the tissues. As this period of grain development, Zadoks 73, is one of rapid radial expansion growth, it is tempting to speculate that bioactive GA produced by the endosperm promotes cell expansion in the outer grain layers which otherwise would constrain growth of the endosperm. In support of this, dwarfing mutations in *TaRht* genes that reduce sensitivity to bioactive GAs also reduce grain size [15], implying a role for GAs and GA signalling in grain expansion.

However, no defined role for GA in seed development has been demonstrated in any species. Although developing seeds of plants often accumulate high levels of GAs, in many cases these are inactive or partially active forms. Thus, in rice the *OsGA3ox1* gene that is expressed in developing grain encodes an enzyme with both 3 β -hydroxylase and 2 β -hydroxylase activities [54], although the latter activity is a minor component, at least when assayed *in vitro*. Unlike the other sequenced grass species (maize, sorghum and *Setaria italica*), wheat, barley and Brachypodium lack orthologs of *GA3ox1* but in each case have evolved novel paralogs of the *GA3ox2* gene responsible for GA biosynthesis in vegetative tissues (Fig. 2B and Additional file 1: Figure S7) that have high expression in developing grain. In the case of bread wheat, we show above that two novel paralogs, *TaGA3ox3* and *TaGA1ox1*, combine to produce the unusual gibberellin GA₅₄ (1 β -hydroxy-GA₄) in grain. Similarly, in barley *HvGA3,18ox1* encodes a bifunctional GA 3 β ,18-dihydroxylase; this gene is closely related to *TaGA3ox3* and *TaGA1ox1* and its expression is confined to developing grain. It is presumably, therefore, responsible for the accumulation of 18-hydroxy-GAs, such as GA₁₃₁ (18-hydroxy-GA₄), in seeds of this species. Brachypodium also contains a paralogous gene related to *GA3ox2*, *Bradi4g23570*, that we have named *BdGA3ox2b*. This gene is similarly expressed exclusively in seed tissues (Additional file 1: Figure S5) and encodes an enzyme with only GA 3-oxidase activity. Thus, rice, barley, wheat, and Brachypodium have independently evolved paralogs of *GA3ox* that produce high levels of GAs in developing grain. However, the GAs synthesized in wheat and barley grain exhibit reduced biological activity: GA₁₃₁ is ~4-fold less active than GA₄ [58] and the biological activity of 1-hydroxy GAs is similarly lower than of their non-1-hydroxylated equivalents (personal communication from the late Dr. John Lenton). The biological relevance of these unusual GA-modifying activities is therefore unclear and merits further investigation.

Target genes for novel dwarfing effects in wheat

The introduction of the *Rht* dwarfing alleles during the Green Revolution had a major impact on global wheat productivity because of a reduced incidence of lodging and improved assimilate partitioning [59]. However, because *Rht* is a negative regulator of GA signalling in all

tissues, these alleles are also associated with negative pleiotropic effects, such as the requirement for planting closer to the surface and a slight reduction in grain size [15, 19]. Results from our study confirm that *Rht* was expressed in all tissues assayed, including the outer layers of expanding grain and suggests that alternative dwarfing alleles, more specific and targeted to the stem, may limit these negative pleiotropic effects. Several genetic studies to identify plant height QTLs are being carried out in order to identify alternative dwarfing alleles for wheat, although to date no specific gene has been identified [60].

One solution may lie in the selection of loss-of-function mutations in GA biosynthetic genes using currently available TILLING populations [61, 62]. In rice, the most widely used dwarfing gene is *GA20ox2* [16, 17], where multiple loss-of-function alleles have been identified which reduce height. These mutations have only limited effects on reproductive development, presumably because other *GA 20-oxidase* genes regulate GA biosynthesis in these tissues. In wheat, *TaGA20ox1*, *TaGA20ox2* and *TaGA20ox4* are expressed in the leaf and stem, suggesting that these are promising candidates in which to search for disruptive mutations. However, since all three homoeologs of both *TaGA20ox1* and *TaGA20ox2* are expressed in stem tissues (Additional file 3) multiple mutations may necessary to obtain economically relevant height reductions. Current efforts to sequence the exomes of wheat TILLING lines [63] is accelerating the discovery of mutations for most of the genes presented in this study, which will facilitate the experimental validation of their function and the testing of their effects on plant height and productivity.

Conclusions

In this study we identified the genes encoding 2-ODD enzymes from the GA biosynthetic pathway of bread wheat, through a combination of interrogation of publicly available, partially-assembled genome sequences and *de novo* assembly of shotgun reads from individual chromosome arms. We also catalogued the GA 2-ODD genes from barley and Brachypodium. With a few exceptions, hexaploid bread wheat contains three homoeologs of each paralog of the genes identified in rice. In each case the function of the enzymes encoded by the genes was demonstrated by heterologous expression and their expression was analysed across a range of tissues and developmental stages. Certain paralogs of the *GA20ox*, *GA3ox* and *GA2ox* gene families were shown to be exclusively expressed in developing grain, including a novel, highly-expressed GA 1-oxidase gene in wheat endosperm (and a related GA 3,18-dihydroxylase gene in barley). The presence of these genes likely explains the high levels of 1 β -hydroxylated GAs in wheat grain, and 18-hydroxylated GAs in barley, in both cases the

activities resulting in a reduction in the biological activity of the GAs produced. The pattern of transcript accumulation of the GA genes in early developing grain suggested that, while most GA biosynthesis is carried out in the endosperm, GA inactivation, perception and signalling are confined to the seed coat and pericarp, consistent with a role for GA in grain expansion. This comprehensive identification and characterisation of the GA 2-ODD genes in wheat will provide the basis for a better understanding of GA-regulated development in this species, including the involvement of GA biosynthesis and signalling in grain development.

Methods

Identification and phylogeny of wheat genes for components of the GA pathway

To identify wheat (*Triticum aestivum* L.) genes encoding components of the GA biosynthetic and signalling pathway, sequences from Brachypodium (*Brachypodium distachyon* L.), identified by BLASTP at www.phytozome.org using rice (*Oryza sativa* L.) peptide sequences [31] as queries, were used to search the IWGSC chromosome arm survey assembly at urgi.versailles.inra.fr. In many cases it was not possible to identify all three wheat homoeologs and, therefore, guided assembly of homologous genomic sequence reads from chromosome-arm-specific shotgun sequencing assembled to 454 contigs [32, 33] was carried out using the Geneious software (Biomatters Ltd) as outlined in Additional file 1 (Figure S1). Accession numbers for all full-length 2-ODD sequences identified from wheat and other grasses are shown in Additional file 2. In addition, full-length genomic, coding sequence and polypeptide sequences of all the wheat 2-ODD genes, plus coding sequences of other genes in the pathway that were used for transcript analysis, are provided as FASTA files in Additional file 6.

For phylogenetic analysis, peptide sequences were aligned using MUSCLE [64] and the resulting alignments (Additional file 1: Figures S8-S10) were edited to remove unaligned sequences in the N- and C-terminal regions. For *GA20ox* the N-terminal 82 and C-terminal 31 columns of the alignment were removed; similarly for *GA3ox* the 39 N-terminal and C-terminal 45 columns and for *GA2ox* the N-terminal 69 and C-terminal 49 columns were deleted. Phylogenetic analysis was carried out on the resulting alignments using the PhyML algorithm within TOPALi v2.5 [65], including model optimisation and bootstrapping (100 repeats); an outgroup was used for rooting and phylogenetic trees were drawn using MEGA5 [66]. The trees obtained using Neighbor Joining methods were not significantly different. To simplify presentation, only one homoeolog of each paralogous bread wheat gene was included in the phylogenetic trees presented in Figure 2, but complete versions of

each tree with bootstrap values are shown in Additional file 1 (Figures S11–S13).

Heterologous expression of wheat, barley and Brachypodium GA 2-ODDs

Coding sequences for a single representative homoeolog of each wheat 2-ODD, and of barley (*Hordeum vulgare* L.) and Brachypodium (*Brachypodium distachyon* L.) GA3ox genes, from the GA pathway were synthesised (Genscript, Piscataway, USA) and inserted in-frame as a fusion with thioredoxin in the expression vector pET32b. The constructs were expressed in *E. coli* strain Rosetta2 (DE3) pLysS (Merck Millipore Ltd) by induction of log phase cultures with 0.5 mM IPTG followed by growth at 25 °C for 5–6 h. Cells were harvested by centrifugation, suspended in 100 mM Tris-Cl pH7.5, 5 mM DTT, 10 u.ml⁻¹ DNaseI (Sigma), lysed by freeze-thaw and sonication and assayed using radiolabelled GA substrates in the presence of co-substrates and co-factors as described previously [67]. Control assays were also run to demonstrate that neither *E. coli* proteins nor products encoded by the pET32b vector were active against the GA₁₂ and GA₉ substrates. All products were identified by comparison of HPLC retention times with those of standards, or, in the case of ¹⁴C-labelled GA₅₄, GA₆₁ and GA₁₃₁, by comparison of their mass spectra with published spectra [40] after combined gas chromatography–mass spectrometry of methyl esters trimethylsilyl ethers as described previously [68] using a MAT95XP mass spectrometer coupled to a Trace GC (ThermoElectron).

Mapping RNA-seq reads to the wheat reference transcriptome

Paired-end RNA-seq reads from root, leaf, stem, spike and grain tissues of wheat cv. Chinese Spring [49] were obtained from <http://urgi.versailles.inra.fr/>. Paired-end RNA-seq reads from Brachypodium tissues were downloaded from the Short Read Archive at Genbank (accession number SRA046377). For developing grain samples, bread wheat cv. Holdfast seedlings were vernalized for 8 weeks at 8 °C and transplanted to pots of Rothamsted Prescription Mix including slow-release fertilizer. The plants were grown in a glasshouse with supplemental lighting under a 16 h day (18 °C), 8 h (14 °C) night and tagged at anthesis (anther emergence). Immature grain at 12 days post-anthesis were dissected into endosperm, inner and outer seed coat/pericarp tissues and RNA extracted as described by Wan et al. [69]. Illumina RNA-seq libraries from three biological replicates of each tissue were prepared using the TruSeq mRNA sample kit (Illumina TruSeq RNA sample preparation guide Part 15008136 Rev A, Nov 2010) using 4 µg total RNA. Single end reads, 110 bp, were generated on an Illumina GAIIx sequencer and RNA-seq analysis was carried out within the Galaxy environment

[70]. For all sets of RNA-seq data, sequencing reads were trimmed for quality using Trimmomatic [71] and reads from each library were mapped using BWA-mem using default parameters [72] to a wheat reference transcriptome consisting of the predicted cDNA sequences from the IWGSC assembly (v21) at www.plants.ensembl.org/Triticum_aestivum. Where genes of interest were absent from this reference, full-length coding sequences developed in this work were added to replace either partial or missing sequences as detailed in Additional file 7. Gene sequences from the early part of the GA pathway were obtained from online databases: *TaCPS* [EMBL:GU980886, EMBL:GU980887, EMBL:GU980888]; *TaKS* [GenBank:FR719731, EMBL: GU980889, EMBL: GU980890], *TaKO* [EMBL:GU980893, EMBL: GU980894, EMBL: GU980895]; *TaKAO* [EMBL:GU980891, EMBL:GU980892, EMBL:GU143912]. FPKM (for paired-end reads) and RPKM (for single-end reads) values were generated using the eXpress tool [73].

Availability of supporting data

The RNA-seq data from tissue layers of developing grain described in this work are available in the ArrayExpress database (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-3103. The phylogenetic input data and analysis for the GA20ox, GA3ox and GA2ox trees are available from Dryas (doi:10.5061/dryad.sk2nd).

Additional files

Below is the link to the electronic supplementary material.

Additional file 1: Figure S1. Strategy for identification and assembly of wheat GA 2-ODD sequences. **Figure S2:** HPLC traces of incubations of heterologous expression products of GA20ox genes with [1-¹⁴C]-GA₁₂. **Figure S3:** HPLC traces of incubations of heterologous expression products of GA2ox genes with [1-¹⁴C]-GA₁₂ and [1-¹⁴C]-GA₉. **Figure S4:** Relative expression of GA biosynthetic and signalling genes in durum wheat tissues determined by qRT-PCR. **Figure S5:** Transcript analysis of GA biosynthetic and signalling genes in *Brachypodium distachyon* by RNA-seq. **Figure S6:** Dissection of tissue layers from developing grain of bread wheat. **Figure S7:** Phylogenetic tree of grass GA3ox-like protein sequences. **Figures S8–10:** GA20ox, GA3ox and GA2ox protein alignments, respectively. **Figures S11–13:** Phylogenetic trees of all GA20ox, GA3ox and GA2ox protein sequences, respectively, with bootstrap support values.

Additional file 2: Rice, Brachypodium, barley and wheat genes encoding GA 2-ODDs including accession numbers.

Additional file 3: Homoeolog-specific expression of GA biosynthetic and signalling genes across five tissues of wheat at three developmental stages.

Additional file 4: qRT-PCR primers used in this study and qRT-PCR methods.

Additional file 5: Homoeolog-specific expression of GA biosynthetic and signalling genes in different layers of the wheat grain.

Additional file 6: FASTA files containing genomic, coding sequence and polypeptide sequences of all full-length wheat genes described in this work.

Additional file 7: Sequences in the IWGSC reference used for RNA-seq mapping substituted with full-length CDS sequences from this work.

Abbreviations

2-ODD: 2-oxoglutarate-dependent dioxygenases; FPKM: Fragments per kilobase per million mapped reads; GA: Gibberellin; GA1ox: GA 1-oxidase (GA 1 β -hydroxylase); GA13ox: GA 13-hydroxylase; GA20ox: GA 20-oxidase; GA2ox: GA 2-oxidase (GA 2 β -hydroxylase); GA3ox: GA 3-oxidase (GA 3 β -hydroxylase); GC-MS: Coupled gas chromatography–mass spectrometry; HPLC: High performance liquid chromatography; IWGSC: International Wheat Genome Sequencing Consortium; RPKM: Reads per kilobase per million mapped reads; qRT-PCR: Quantitative reverse transcription-polymerase chain reaction; TILLING: Targeting induced local lesions in genomes.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SP, PH, AH and ALP conceived and designed the experiments; SP, AH, IP, BG, JD and ALP assembled sequences of wheat and barley 2-ODD genes and performed phylogenetic analyses; AH and SV performed dissection and RNA isolation from developing wheat grains; JC carried out RNA-seq library construction and sequencing; SP carried out qRT-PCR on durum wheat samples; YL, AH, AP and ALP performed PCR, DNA sequencing and heterologous expression; PH executed enzyme assays and HPLC and GC-MS analysis of products; SP and ALP analysed RNA-seq data; and SP, PH, AH, JD and ALP drafted the manuscript and generated the figures and tables. All authors read and approved the final manuscript.

Acknowledgments

We thank Rothamsted glasshouse staff for care of plant material, the Rothamsted CSYS Bioinformatics Support Group for advice and provision of Galaxy tools and Stephen Powers for statistical advice. We also thank Prof. Jizeng Jia for prepublication access to assembled genomic sequences from *Triticum urartu* and *Triticum tauschii*. Financial support: work by AKH, IMP, SPV, BG, AP, PH and ALP was carried out under the 20:20 Wheat* Institute Strategic Programme at Rothamsted Research, funded by the Biotechnology and Biological Sciences Research Council of the UK; YL was funded by the OPTICHINA network within the Seventh Framework Programme of the EU. SP and JD acknowledge support from the Howard Hughes Medical Institute, the Gordon and Betty Moore Foundation and from the U.S. Department of Agriculture National Institute of Food and Agriculture grants 2011–67013–30077 and 2011–68002–30029.

Author details

¹Department of Plant Sciences, University of California, Davis, CA 95616, USA. ²Department of Plant Biology and Crop Science, Rothamsted Research, Harpenden AL5 2JQ, UK. ³Biotechnology Research Centre, Jilin Academy of Agricultural Sciences, Changchun 130033, China. ⁴University of Bristol Transcriptomics Facility, School of Biological Sciences, Bristol BS8 1UG, UK. ⁵Howard Hughes Medical Institute, Chevy Chase, MD 20815, USA.

Received: 7 January 2015 Accepted: 1 May 2015

Published online: 05 June 2015

References

- Ubeda-Tomás S, Federici F, Casimiro I, Beemster GTS, Bhalerao R, Swarup R, et al. Gibberellin signaling in the endodermis controls *Arabidopsis* root meristem size. *Curr Biol*. 2009;19:1194–9.
- Gou JQ, Strauss SH, Tsai CJ, Fang K, Chen YR, Jiang XN, et al. Gibberellins regulate lateral root formation in *Populus* through interactions with auxin and other hormones. *Plant Cell*. 2010;22:623–39.
- Alabadí D, Gil J, Blázquez MA, García-Martínez JL. Gibberellins repress photomorphogenesis in darkness. *Plant Physiol*. 2004;134:1050–7.
- Mauriat M, Moritz T. Analyses of GA20ox- and GID1-over-expressing aspen suggest that gibberellins play two distinct roles in wood formation. *Plant J*. 2009;58:989–1003.
- Appleford NEJ, Lenton JR. Gibberellins and leaf expansion in near-sogenic wheat lines containing *Rht1* and *Rht3* dwarfing alleles. *Planta*. 1991;183:229–36.
- Perazza D, Vachon G, Herzog M. Gibberellins promote trichome formation by up-regulating *GLABROUS1* in *Arabidopsis*. *Plant Physiol*. 1998;117:375–83.
- Pearce S, Vanzetti LS, Dubcovsky J. Exogenous gibberellins induce wheat spike development under short days only in the presence of *VERNALIZATION1*. *Plant Physiol*. 2013;163:1433–45.
- Plackett ARG, Thomas SG, Wilson ZA, Hedden P. Gibberellin control of stamen development: a fertile field. *Trends Plant Sci*. 2011;16:568–78.
- Ozga JA, van Huizen R, Reinecke DM. Hormone and seed-specific regulation of pea fruit growth. *Plant Physiol*. 2002;128:1379–89.
- Radley M. The development of wheat grain in relation to endogenous growth substances. *J Exp Bot*. 1976;27:1009–21.
- Hedden P, Thomas SG. Gibberellin biosynthesis and its regulation. *Biochem J*. 2012;444:11–25.
- Yang DL, Yao J, Mei CS, Tong XH, Zeng LJ, Li Q, et al. Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade. *Proc Natl Acad Sci U S A*. 2012;109:E1192–200.
- Li QF, Wang CM, Jiang L, Li S, Sun SSM, He JX. An interaction between BZR1 and DELLAs mediates direct signaling crosstalk between brassinosteroids and gibberellins in *Arabidopsis*. *Sci Signal*. 2012;5:ra72.
- Nakamura H, Xue Y-L, Miyakawa T, Hou F, Qin H-M, Fukui K, et al. Molecular mechanism of strigolactone perception by DWARF14. *Nat Commun*. 2013;4:2613.
- Flintham JE, Börner A, Worland AJ, Gale MD. Optimizing wheat grain yield: effects of *Rht* (gibberellin-insensitive) dwarfing genes. *J Agric Sci*. 1997;128:11–25.
- Spielmeier W, Ellis MH, Chandler PM. Semidwarf (*sd-1*), “green revolution” rice, contains a defective gibberellin 20-oxidase gene. *Proc Natl Acad Sci U S A*. 2002;99:9043–8.
- Jia QJ, Zhang JJ, Westcott S, Zhang XQ, Bellgard M, Lance R, et al. GA-20 oxidase as a candidate for the semidwarf gene *sdw1/denso* in barley. *Funct Integr Genomics*. 2009;9:255–62.
- Calderini DF, Miralles DJ, Sadras VO. Appearance and growth of individual leaves as affected by semidwarfism in isogenic lines of wheat. *Ann Bot*. 1996;77:583–9.
- Ellis MH, Rebetzke GJ, Chandler P, Bonnett D, Spielmeier W, Richards RA. The effect of different height reducing genes on the early growth of wheat. *Funct Plant Biol*. 2004;31:583–9.
- Plackett ARG, Powers SJ, Fernandez-Garcia N, Urbanova T, Takebayashi Y, Seo M, et al. Analysis of the developmental roles of the *Arabidopsis* gibberellin 20-oxidases demonstrates that GA20ox1, -2, and -3 are the dominant paralogs. *Plant Cell*. 2012;24:941–60.
- Rieu I, Eriksson S, Powers SJ, Gong F, Griffiths J, Woolley L, et al. Genetic analysis reveals that C₁₉-GA 2-oxidation is a major gibberellin inactivation pathway in *Arabidopsis*. *Plant Cell*. 2008;20:2420–36.
- Mitchum MG, Yamaguchi S, Hanada A, Kuwahara A, Yoshioka Y, Kato T, et al. Distinct and overlapping roles of two gibberellin 3-oxidases in *Arabidopsis* development. *Plant J*. 2006;45:804–18.
- Yamaguchi S. Gibberellin metabolism and its regulation. *Annu Rev Plant Biol*. 2008;59:225–51.
- Huang Y, Yang W, Pei Z, Guo X, Liu D, Sun J, et al. The genes for gibberellin biosynthesis in wheat. *Funct Integr Genomics*. 2012;12:199–206.
- Wu YS, Zhou K, Toyomasu T, Sugawara C, Oku M, Abe S, et al. Functional characterization of wheat copalyl diphosphate synthases sheds light on the early evolution of labdane-related diterpenoid metabolism in the cereals. *Phytochemistry*. 2012;84:40–6.
- Zhou K, Xu MM, Tiernan M, Xie Q, Toyomasu T, Sugawara C, et al. Functional characterization of wheat *ent-kaurene*-like synthases indicates continuing evolution of labdane-related diterpenoid metabolism in the cereals. *Phytochemistry*. 2012;84:47–55.
- Thomas SG, Phillips AL, Hedden P. Molecular cloning and functional expression of gibberellin 2-oxidases, multifunctional enzymes involved in gibberellin deactivation. *Proc Natl Acad Sci U S A*. 1999;96:4698–703.
- Schomburg FM, Bizzell CM, Lee DJ, Zeevaert JAD, Amasino RM. Overexpression of a novel class of gibberellin 2-oxidases decreases gibberellin levels and creates dwarf plants. *Plant Cell*. 2003;15:151–63.
- Appleford NEJ, Evans DJ, Lenton JR, Gaskin P, Croker SJ, Devos KM, et al. Function and transcript analysis of gibberellin-biosynthetic enzymes in wheat. *Planta*. 2006;223:568–82.
- Lo SF, Yang SY, Chen KT, Hsing YL, Zeevaert JAD, Chen LJ, et al. A novel class of gibberellin 2-oxidases control semidwarfism, tillering, and root development in rice. *Plant Cell*. 2008;20:2603–18.
- Hirano K, Aya K, Hobo T, Sakakibara H, Kojima M, Shim RA, et al. Comprehensive transcriptome analysis of phytohormone biosynthesis and signaling genes in microspore/pollen and tapetum of rice. *Plant Cell Physiol*. 2008;49:1429–50.
- IWGSC. A chromosome-based draft sequence of the hexaploid bread wheat (*Triticum aestivum*) genome. *Science*. 2014;345:1251788.

33. Brenchley R, Spannagl M, Pfeifer M, Barker GLA, D'Amore R, Allen AM, et al. Analysis of the bread wheat genome using whole-genome shotgun sequencing. *Nature*. 2012;491:705–10.
34. Krasileva KV, Buffalo V, Bailey P, Pearce S, Ayling S, Tabbita F, et al. Separating homeologs by phasing in the tetraploid wheat transcriptome. *Genome Biol*. 2013;14:R66.
35. Ling HQ, Zhao SC, Liu DC, Wang JY, Sun H, Zhang C, et al. Draft genome of the wheat A-genome progenitor *Triticum urartu*. *Nature*. 2013;496:87–90.
36. Jia JZ, Zhao SC, Kong XY, Li YR, Zhao GY, He WM, et al. *Aegilops tauschii* draft genome sequence reveals a gene repertoire for wheat adaptation. *Nature*. 2013;496:91–5.
37. Mayer KFX, Waugh R, Langridge P, Close TJ, Wise RP, Graner A, et al. A physical, genetic and functional sequence assembly of the barley genome. *Nature*. 2012;491:711–7.
38. Sorrells ME, La Rota M, Bermudez-Kandianis CE, Greene RA, Kantety R, Munkvold JD, et al. Comparative DNA sequence analysis of wheat and rice genomes. *Genome Res*. 2003;13:1818–27.
39. Spielmeier W, Ellis M, Robertson M, Ali S, Lenton JR, Chandler PM. Isolation of gibberellin and metabolic pathway genes from barley and comparative mapping in barley, wheat and rice. *Theor Appl Genet*. 2004;109:847–55.
40. Gaskin P, MacMillan J. GC-MS of the gibberellins and related compounds: Methodology and a library of spectra. Bristol, U.K: Cantock's Enterprises; 1992.
41. Kirkwood PS, MacMillan J. Gibberellins A₆₀, A₆₁, and A₆₂: partial syntheses and natural occurrence. *J Chem Soc Perkin Trans*. 1982;1:689–97.
42. Gaskin P, Kirkwood PS, Lenton JR, Macmillan J, Radley ME. Identification of gibberellins in developing wheat grain. *Agric Biol Chem*. 1980;44:1589–93.
43. Crow JR, Thomson RJ, Mander LN. Synthesis and confirmation of structure for the gibberellin GA₁₃₁ (18-hydroxy-GA₁). *Org Biomol Chem*. 2006;4:2532–44.
44. Gaskin P, Gilmour SJ, Lenton JR, Macmillan J, Sponsel VM. Endogenous gibberellins and kauranoids identified from developing and germinating barley grain. *J Plant Growth Reg*. 1984;2:229–42.
45. Green LS, Faergestad EM, Poole A, Chandler PM. Grain development mutants of barley - α -amylase production during grain maturation and its relation to endogenous gibberellic acid content. *Plant Physiol*. 1997;114:203–12.
46. Sakamoto T, Kobayashi M, Itoh H, Tagiri A, Kayano T, Tanaka H, et al. Expression of a gibberellin 2-oxidase gene around the shoot apex is related to phase transition in rice. *Plant Physiol*. 2001;125:1508–16.
47. Shan C, Mei ZL, Duan JL, Chen HY, Feng HF, Cai WM. OsGA2ox5, a gibberellin metabolism enzyme, is involved in plant growth, the root gravity response and salt stress. *PLoS One*. 2014;9:e87110.
48. Lee DJ, Zeevaert JAD. Molecular cloning of GA 2-oxidase3 from spinach and its ectopic expression in *Nicotiana glauca*. *Plant Physiol*. 2005;138:243–54.
49. Choulet F, Alberti A, Theil S, Glover N, Barbe V, Daron J, et al. Structural and functional partitioning of bread wheat chromosome 3B. *Science*. 2014;345:1249721.
50. Magome H, Nomura T, Hanada A, Takeda-Kamiya N, Ohnishi T, Shinma Y, et al. *CYP714B1* and *CYP714B2* encode gibberellin 13-oxidases that reduce gibberellin activity in rice. *Proc Natl Acad Sci U S A*. 2013;110:1947–52.
51. Pfeifer M, Kugler KG, Sandve SR, Zhan B, Rudi H, Hvidsten TR, et al. Genome interplay in the grain transcriptome of hexaploid bread wheat. *Science*. 2014;345:1250091.
52. Davidson RM, Gowda M, Moghe G, Lin HN, Vaillancourt B, Shiu SH, et al. Comparative transcriptomics of three Poaceae species reveals patterns of gene expression evolution. *Plant J*. 2012;71:492–502.
53. Lange MJP, Liebrandt A, Arnold L, Chmielewska SM, Felsberger A, Freier E, et al. Functional characterization of gibberellin oxidases from cucumber, *Cucumis sativus* L. *Phytochemistry*. 2013;90:62–9.
54. Itoh H, Ueguchi-Tanaka M, Sentoku N, Kitano H, Matsuoka M, Kobayashi M. Cloning and functional analysis of two gibberellin 3 β -hydroxylase genes that are differently expressed during the growth of rice. *Proc Natl Acad Sci U S A*. 2001;98:8909–14.
55. Sakamoto T, Miura K, Itoh H, Tatsumi T, Ueguchi-Tanaka M, Ishiyama K, et al. An overview of gibberellin metabolism enzyme genes and their related mutants in rice. *Plant Physiol*. 2004;134:1642–53.
56. Oikawa T, Koshioka M, Kojima K, Yoshida H, Kawata M. A role of *OsGA2ox1*, encoding an isoform of gibberellin 20-oxidase, for regulation of plant stature in rice. *Plant Mol Biol*. 2004;55:687–700.
57. Rutger JN. Applications of induced and spontaneous mutation in rice breeding and genetics. *Adv Agron*. 1983;36:383–413.
58. Morrison E, Chandler PM, Thomson RJ, Mander LN. Synthesis and bioactivity of the gibberellin, 18-hydroxy-GA₁ (GA₁₃₂). *Org Biomol Chem*. 2008;6:1416–24.
59. Hedden P. The genes of the Green Revolution. *Trends Genet*. 2003;19:5–9.
60. Griffiths S, Simmonds J, Leverington M, Wang YK, Fish L, Sayers L, et al. Meta-QTL analysis of the genetic control of crop height in elite European winter wheat germplasm. *Mol Breed*. 2012;29:159–71.
61. Botticella E, Sestili F, Hernandez-Lopez A, Phillips A, Lafiandra D. High resolution melting analysis for the detection of EMS induced mutations in wheat *Sbella* genes. *BMC Plant Biol*. 2011;11:156.
62. Jaury C, Paraiso F, Colasuonno P, Tran R, Tsai H, Berardi S, et al. A modified TILLING approach to detect induced mutations in tetraploid and hexaploid wheat. *BMC Plant Biol*. 2009;9:115–28.
63. Henry IM, Nagalakshmi U, Lieberman MC, Ngo KJ, Krasileva KV, Vasquez-Gross H, et al. Efficient genome-wide detection and cataloging of EMS-induced mutations using exome capture and next-generation sequencing. *Plant Cell*. 2014;26:1382–97.
64. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res*. 2004;32:1792–7.
65. Milne I, Lindner D, Bayer M, Husmeier D, McGuire G, Marshall DF, et al. TOPALi v2: a rich graphical interface for evolutionary analyses of multiple alignments on HPC clusters and multi-core desktops. *Bioinformatics*. 2009;25:126–7.
66. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011;28:2731–9.
67. Ward DA, MacMillan J, Gong F, Phillips AL, Hedden P. Gibberellin 3-oxidases in developing embryos of the southern wild cucumber, *Marah macrocarpus*. *Phytochemistry*. 2010;71:2010–8.
68. MacMillan J, Ward DA, Phillips AL, Sánchez-Beltrán MJ, Gaskin P, Lange T, et al. Gibberellin biosynthesis from gibberellin A₁₂-aldehyde in endosperm and embryos of *Marah macrocarpus*. *Plant Physiol*. 1997;113:1369–77.
69. Wan Y, Underwood C, Toole G, Skeggs P, Zhu T, Leverington M, et al. A novel transcriptomic approach to identify candidate genes for grain quality traits in wheat. *Plant Biotechnol J*. 2009;7:401–10.
70. Goecks J, Nekrutenko A, Taylor J, The Galaxy Team: Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. *Genome Biol* 2010;11:R86.
71. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014;30:2114–20.
72. Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics*. 2010;26:589–95.
73. Forster SC, Finkel AM, Gould JA, Hertzog PJ. RNA-eXpress annotates novel transcript features in RNA-seq data. *Bioinformatics*. 2013;29:810–2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

