

POSTER PRESENTATION

Open Access

Effects of reintubation on ventilator associated nosocomial pneumonias incidence in ICU patients with versus without selective digestive decontamination

AM Marrero-Rodríguez^{1*}, ER Argandoña-Primicia¹, ME Gallardo-Santos¹, P Juárez-San Juan¹, CF Lübbe-Vazquez¹, JJ Díaz Díaz¹, P Saavedra², S Ruiz-Santana¹

From ESICM LIVES 2015
Berlin, Germany. 3-7 October 2015

Introduction

Reintubation has been shown, among many others, to be an independent risk factor for development of ventilator-associated pneumonia₁ (VAP) which is a major cause of morbidity and mortality in ICUs. Selective Digestive Decontamination (SDD) is a tool that prevent infections in critically ill patients particularly VAP, that has been used mainly in Holland and in Spain. Despite of the evidence its use remains controversial.

Objectives

To assess the VAP incidence after reintubation in patients admitted to a polyvalent ICU before and after starting a SDD program.

Methods

A retrospective study was conducted in a 30-bed-medical-surgical ICU. We compared all patients admitted to ICU who required reintubation from October 2009 to October 2011 (non-SDD group) to those patients admitted to the ICU from October 2011 to October 2013, after starting a SDD program (SDD group) prospectively collected data. All patients that were expected to require tracheal intubation for longer than 48 hours were given SDD (SDD study group). Nosocomial infections were diagnosed by CDC criteria. In both groups, categorical variables were summarized as frequencies and percentages and the continuous ones as medians and interquartile ranges (IQR). The

percentages were compared using the test of chi-square test or Fisher exact test and medians with the Wilcoxon test for independent samples. For VAP the incidences per 1000 days of mechanical ventilation were obtained. These rates were compared using the Poisson model. Statistical significance was set at $p \leq 0.05$.

Results

Results are shown in Figures 1, and 2. There was a non-significant reduction of VAP per 1000 days of mechanical ventilation (23.2 vs. 35.1) in patients that were reintubated in the SDD period (n=28) compared to those in the non-SDD period (n=28). The risk reduction observed was 34%.

Conclusions

There is a notorius reduction, towards significance in the incidence of VAP per 1000 days of mechanical ventilation in patients reintubated on SDD compared to those reintubated in the non-SDD period.

Authors' details

¹University Hospital of Gran Canaria Dr. Negrín, Intensive Care Unit, Las Palmas de Gran Canaria, Spain. ²Las Palmas of Gran Canaria University, Mathematics and Informatics Department, Las Palmas de Gran Canaria, Spain.

Published: 1 October 2015

Reference

1. Hortal J, Giannella M, Perez MJ, Barrio JM, Desco M, Bouza E, et al: Incidence and risk factors for ventilator-associated pneumonia after major heart surgery. *Intensive care medicine* 2009, **35**(9):1518-25.

¹University Hospital of Gran Canaria Dr. Negrín, Intensive Care Unit, Las Palmas de Gran Canaria, Spain

Full list of author information is available at the end of the article

	Total (n = 56)	SDD		P
		No (n = 28)	Yes (n = 28)	
Age, years	66.0 (49.5 ; 75.4)	64.0 (51.0 ; 76.4)	66.6 (48.0 ; 73.7)	.819
Male / female, %	66.1 / 33.9	64.3 / 35.7	67.9 / 32.1	.949
Apache II score	19.5 (17 ; 22)	19.5 (17 ; 23)	19.5 (18 ; 21)	.915
Cause of admission, n (%)				.029
Cardiac	14 (25.0)	9 (32.1)	5 (17.4)	
Respiratory	5 (8.9)	1 (3.6)	4 (14.3)	
Neurologic	20 (35.7)	6 (21.4)	14 (50.0)	
Digestive	17 (30.4)	12 (42.9)	5 (17.9)	
Other				
Diabetes mellitus, n (%)	18 (32.1)	10 (35.7)	8 (28.6)	.567
COPD, n (%)	7 (12.5)	5 (17.9)	2 (7.1)	.422
Smoker, n (%)	10 (17.9)	7 (25.0)	3 (10.7)	.163
Hypertension, n (%)	30 (53.6)	13 (46.4)	17 (60.7)	.284
Immunosuppression, n (%)	8 (15.1)	5 (18.5)	3 (11.5)	.704
Days in ICU, days	17 (9 ; 30)	17 (9 ; 25)	19 (9 ; 33)	.406
ICU dead patients, n (%)	10 (17.9)	4 (14.3)	6 (21.4)	.485
Glucose at 1 st reintubation	173 (150 ; 199)	168 (149 ; 202)	184 (162 ; 199)	.313
Days of mechanical ventilation (MV)	6 (1 ; 11.25)	6 (1 ; 11)	6 (1 ; 16)	.378
N° of intubations*				.669
2	51 (90.1)	26 (92.9)	25 (89.3)	
3	5 (8.9)	2 (7.1)	3 (10.7)	

Data are shown as medians (IQR) or frequencies (%). SDD: Selective Digestive Decontamination; COPD: Chronic Obstructive Pulmonary Disease.

Figure 1

		SDD		P	RR (95% CI)
		No N = 28	Yes N = 28		
Ventilator-associated pneumonia (VAP)	Days of mechanical ventilation	171	259	.472	0.660*
	Total number of pneumonias *	6	6		(0.213 ; 12.047)
	Pneumonias / 1000 days of MV	35.1	23.2		

Figure 2

doi:10.1186/2197-425X-3-S1-A704

Cite this article as: Marrero-Rodríguez et al.: Effects of reintubation on ventilator associated nosocomial pneumonias incidence in ICU patients with versus without selective digestive decontamination. *Intensive Care Medicine Experimental* 2015 **3**(Suppl 1):A704.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com