

RESEARCH

Open Access



The entropy solution of a hyperbolic-parabolic mixed type equation

Huashui Zhan*

*Correspondence:
Huashuizhan@163.com;
2012111007@xmut.edu.cn
School of Applied
Mathematics, Xiamen
University of Technology,
Xiamen 361024, China

Abstract

The entropy solution of the equation

$$\frac{\partial u}{\partial t} = \Delta A(u) + \operatorname{div}(b(u)), \quad (x, t) \in \Omega \times (0, T),$$

is considered. Besides the usual initial value, only a partial boundary value is imposed. By choosing some special test functions, the stability of the solutions is obtained by Kruzkov's bi-variables method, provided that $\Omega \subset \mathbb{R}^N$ is a unit n -dimensional cube or the half space.

Keywords: Hyperbolic-parabolic mixed type equation, n -dimensional cube, Entropy solution, Kruzkov's bi-variables method

Mathematics Subject Classification: 35L65, 35K85, 35R35

The boundary condition

We consider the equation

$$\frac{\partial u}{\partial t} = \Delta A(u) + \operatorname{div}(b(u)), \quad (x, t) \in \Omega \times (0, T), \quad (1)$$

and assume that

$$A(u) = \int_0^u a(s) ds, \quad a(s) \geq 0, a(0) = 0, \quad (2)$$

where $\Omega \subset \mathbb{R}^N$ is a appropriately smooth open domain. Equation (1) is with hyperbolic-parabolic mixed type, arises from the reaction diffusion problem (Wu et al. 2001), the stationary boundary layer theory (Oleinik and Samokhin 1999).

For Cauchy problem of Eq. (1), the paper (Vol'pert and Hudjaev 1967) was the first one to study its solvability, since then, many papers continued to dedicate to the problem, many excellent results were obtained, one can refer to Wu et al. (2001) and Refs. Bendahamane and Karlsen (2004), Brezis and Crandall (1979), Carrillo (1999), Chen and Dibenedetto (2001), Chen and Perthame (2003), Cockburn and Gripenberg (1999), Evans (1998), Karlsen and Risebro (2003), Kruzkov (1970), Oleinik and Samokhin (1999), Vol'pert (1967), Vol'pert and Hudjaev (1967); Vol'pert and Hudjave (1975), Wu and Yin (1989), Wu et al. (2001), Zhan (2004), Zhao (1985), Zhao and Zhan (2005).

If we want to consider the initial boundary value problem of Eq. (1), the initial value is always imposed

$$u(x, 0) = u_0(x), \quad x \in \Omega. \tag{3}$$

But can we impose Dirichlet homogeneous boundary value

$$u(x, t) = 0, \quad (x, t) \in \partial\Omega \times (0, T), \tag{4}$$

as usual? When the equation is of weakly degenerate, i.e. there is not interior point in the set $\{s : a(s) = 0\}$, we can impose Dirichlet homogeneous boundary condition (4). One can refer to Wu et al. (2001) and the references therein. When the equation is of strongly degenerate, i.e. there is an interior point in the set $\{s : a(s) = 0\}$, there are two ways to deal with the corresponding problem. In one way, the entropy solution u is a BV function, which means that

$$\iint_{Q_T} \left| \frac{\partial u}{\partial t} \right| dx dt \leq c, \quad \iint_{Q_T} \left| \frac{\partial u}{\partial x_i} \right| dx dt \leq c, \quad i = 1, 2, \dots, N.$$

It is well-known that the BV function is the weakest function that one can define the trace on the boundary. In this way, we can directly answer whether (4) is true or not in the sense of the trace, and the general result is that, instead of (4), only a partial boundary value such as

$$u(x, t) = 0, \quad (x, t) \in \Sigma_1 \times (0, T), \tag{5}$$

is imposed, where $\Sigma_1 \subseteq \partial\Omega$ is a relative open subset of $\partial\Omega$. The representative works by Wu and Zhao (1983a, b) had been accomplished in early 1980s, later, one can refer to Yin and Wang (2007). In the other way, the boundary value condition is not directly shown in the sense of the trace as (4), but is elegantly implicitly contained in a family entropy inequalities. Moreover, the entropy solutions defined in this way are only in L^∞ space, the existence of the traditional trace [which was called the strong trace in Kobayasi and Ohwa (2012)] on the boundary is not guaranteed, so the boundary value condition is satisfied in a weaker sense than the sense of the trace, one can refer to Carrillo (1999), Li and Qin (2012), Lions et al. (1994), Kobayasi and Ohwa (2012) for more details.

Recently, by the parabolic regularization method, the author Zhan (2015a) had shown the explicit formula of Σ_1 in (5). Let us give some details.

For small $\eta > 0$, let

$$S_\eta(s) = \int_0^s h_\eta(\tau) d\tau, \quad h_\eta(s) = \frac{2}{\eta} \left(1 - \frac{|s|}{\eta} \right)_+.$$

Obviously $h_\eta(s) \in C(\mathbb{R})$, and

$$h_\eta(s) \geq 0, \quad |sh_\eta(s)| \leq 1, \quad |S_\eta(s)| \leq 1; \quad \lim_{\eta \rightarrow 0} S_\eta(s) = \text{sgns}, \quad \lim_{\eta \rightarrow 0} sS'_\eta(s) = 0. \tag{6}$$

Let

$$\Sigma_{1\eta k} = \{x \in \Sigma, S_\eta(k)[b_i(0) - b_i(k)]n_i(x) > 0\}, \tag{7}$$

$$\Sigma_{2\eta k} = \{x \in \Sigma, S_\eta(k)[b_i(0) - b_i(k)]n_i(x) \leq 0\}. \tag{8}$$

Here and in what follows, $\{n_i\}_{i=1}^N$ is the inner normal vector of Ω . Clearly, $\partial\Omega = \Sigma = \Sigma_{1\eta k} \cup \Sigma_{2\eta k}$. Then

$$\Sigma_1 = \bigcup_{\forall \eta \geq 0, \forall k \in \mathbb{R}} \Sigma_{1\eta k}, \quad \Sigma_2 = \Sigma \setminus \Sigma_1. \tag{9}$$

Basing on (9), if the domain Ω is bounded, the existence of the entropy solution had been proved in Zhan (2015a). Assuming that

$$|\Delta d| \leq c, \quad \frac{1}{\lambda} \int_{\Omega_\lambda} dxdt \leq c, \tag{10}$$

the stability of the solutions also had been proved in Zhan (2015a). Here $d(x) = \text{dist}(x, \partial\Omega)$, and $\Omega_\lambda = \{x \in \Omega, d(x) < \lambda\}$, when λ is small enough. If the domain $\Omega = \mathbb{R}_+^N$ is the half space of \mathbb{R}^N , recently in Zhan (2015b), the author had shown that if $b'_N(0) < 0$, then, $\Sigma_1 = \partial\mathbb{R}_+^N$, we can impose Dirichlet boundary value

$$u(x, t) = 0, \quad (x, t) \in \partial\mathbb{R}_+^N \times (0, T).$$

But if $b'_N(0) \geq 0$, then, $\Sigma_1 = \emptyset$, no any boundary value condition is necessary, the solution of the equation is free from any limitation of the boundary value condition.

Now, inspired by Zhan (2004, 2015a, b) and Zhao and Zhan (2005), we give a new definition of the entropy solution.

Definition 1 A function u is said to be the entropy solution of Eq. (1) with the initial value (3) and with the partial boundary value (5), if

1. u satisfies

$$u \in BV(Q_T) \cap L^\infty(Q_T), \quad \frac{\partial}{\partial x_i} \int_0^u \sqrt{a(s)} ds \in L^2(Q_T).$$

2. For any $\varphi \in C_0^2(Q_T)$, $\varphi \geq 0$, for any $k \in \mathbb{R}$, for any small $\eta > 0$, u satisfies

$$\iint_{Q_T} \left[I_\eta(u - k)\varphi_t - B_\eta^i(u, k)\varphi_{x_i} + A_\eta(u, k)\Delta\varphi - S'_\eta(u - k) \left| \nabla \int_0^u \sqrt{a(s)} ds \right|^2 \varphi \right] dxdt \geq 0, \tag{11}$$

where

$$B_\eta^i(u, k) = \int_k^u b'_i(s)S_\eta(s - k)ds, \quad A_\eta(u, k) = \int_k^u a(s)S_\eta(s - k)ds, \quad I_\eta(u - k) = \int_0^{u-k} S_\eta(s)ds.$$

3. The homogeneous boundary value (5) is satisfied in the sense of that

$$\gamma u |_{\Sigma_{1\eta k}} = 0, \tag{12}$$

for any k, η . Here γu means that the equality is true in the sense of the trace.

4. If the domain Ω is bounded, the initial value is true in the sense that

$$\lim_{t \rightarrow 0} \int_\Omega |u(x, t) - u_0(x)| dx = 0, \quad \text{a.e. } x \in \Omega. \tag{13}$$

If the domain is unbounded, the initial value is true in the sense that

$$\lim_{t \rightarrow 0} \int_{\Omega_R} |u(x, t) - u_0(x)| dx = 0, \text{ a.e. } x \in \Omega, \tag{13}'$$

where $\Omega_R = B(0, R) \cap \Omega$.

The existence of the entropy solution in the sense of Definition 1 can be proved similar as that in Zhan (2015a), we omit the details here. In our paper, we are mainly concern with the stability of the entropy solutions of Eq. (1) without the condition (10). For simplicity, only some special domains, for examples, the unite n -dimensional cube

$$D_1 = \{(x_1, x_2, \dots, x_N) : 0 < x_i < 1, i = 1, 2, \dots, N\}.$$

and the half space \mathbb{R}_+^N , are considered. By choosing special test functions, we will prove the following theorem.

Theorem 1 *Suppose that $A(s)$ and $b_i(s)$ are smooth enough, $\Omega = D_1$ is the unite n -dimensional cube. If Σ_1 is a subset of Σ , let u, v be solutions of Eq. (1) with the different initial values $u_0(x), v_0(x) \in L^\infty(\Omega)$ respectively. Suppose that*

$$\gamma u(x, t) = f(x, t), \gamma v = g(x, t), (x, t) \in \Sigma \times (0, T), \tag{14}$$

and in particular,

$$\gamma u = \gamma v = 0, x \in \Sigma_1. \tag{15}$$

Then

$$\int_{\Omega} |u(x, t) - v(x, t)| dx \leq \int_{\Omega} |u_0 - v_0| dx + \text{ess sup}_{(x,t) \in \Sigma_2 \times (0,T)} |f(x, t) - g(x, t)|, \tag{16}$$

where $(x, t) \in \mathbb{R}^{N+1}$, $\text{ess sup}_{(x,t) \in \Sigma_2 \times (0,T)} |f(x, t) - g(x, t)|$ is in the sense of N -dimensional Hausdorff measure.

Compared Theorem 1 to the results obtained in Zhan (2015a), the essential innovation lies in that, without the condition (10), by skillfully constructing the testing function, we still can obtain the stability. At the last section, we also study the similar problem on half space \mathbb{R}_+^N and get the similar result, this result is just the same as that in Zhan (2015b), but we supply a simpler proof.

Now, let us give some analysis in the boundary value condition (5) or (12) to see the rationality. By the definition of $\Sigma_{1\eta k}$, we know that

$$0 < S_\eta(k)[b_i(0) - b_i(k)]n_i = -kS_\eta(k)b'_i(\zeta)n_i, \quad x \in \Sigma$$

where $\zeta \in (k, 0)$. If we let $\eta \rightarrow 0$. Then

$$b'_i(\zeta)n_i(x) < 0, \quad x \in \Sigma.$$

Let $k \rightarrow 0$. We have

$$b'_i(0)n_i(x) < 0, \quad x \in \Sigma. \tag{17}$$

The last inequality (17) is in according with the classical Fichrea–Oleinik theory, one can refer to the explanation in previous works (Zhan 2015a, b)

Let us come back our definition. On the unite n -dimensilnal cube D_1 , according to the homogeneous boundary value condition (5), and by (17), we have

$$\gamma u = \gamma v = 0, x \in \bigcup_{i=1}^N \left\{ x \in \partial D_1 : \left(x_i - \frac{1}{2} \right) b'_i(0) > 0 \right\}. \tag{18}$$

For example, when we consider the boundary plane $\{x_N = 0\}$, $\mathbf{n} = \{0, 0, \dots, 0, 1\}$, (17) implies $b'_i(0)n_i = b'_N(0) < 0$, (18) is true on $\{x_N = 0\}$, we should give the boundary value on $\{x_N = 0\}$, while, on $\{x_N = 1\}$ we can not give the boundary value. If $b'_N(0) > 0$, then (18) is true on $\{x_N = 1\}$, which implies that we should give the boundary value on $\{x_N = 1\}$, while, on $\{x_N = 0\}$ we can not give the boundary value. Certainly, if $b'_N(0) = 0$, both on $\{x_N = 0\}$ and on $\{x_N = 1\}$, no boundary value condition is imposed.

Kruzkov’s bi-variables method

Let Γ_u be the set of all jump points of $u \in BV(Q_T)$, ν the normal of Γ_u at $X = (x, t)$, $u^+(X)$ and $u^-(X)$ the approximate limits of u at $X \in \Gamma_u$ with respect to $(\nu, Y - X) > 0$ and $(\nu, Y - X) < 0$ respectively. For continuous function $p(u, x, t)$ and $u \in BV(Q_T)$, define

$$\widehat{p}(u, x, t) = \int_0^1 p(\tau u^+ + (1 - \tau)u^-, x, t) d\tau, \tag{19}$$

which is called the composite mean value of p . For a given t , we denote $\Gamma_u^t, H^t, (\nu_1^t, \dots, \nu_N^t)$ and u_\pm^t as all jump points of $u(\cdot, t)$, Housdorff measure of Γ_u^t , the unit normal vector of Γ_u^t , and the asymptotic limit of $u(\cdot, t)$ respectively. Moreover, if $f(s) \in C^1(\mathbb{R})$, $u \in BV(Q_T)$, then $f(u) \in BV(Q_T)$ and

$$\frac{\partial f(u)}{\partial x_i} = \widehat{f}'(u) \frac{\partial u}{\partial x_i}, \quad i = 1, 2, \dots, N, N + 1, \tag{20}$$

where $x_{N+1} = t$ as usual.

Lemma 1 *Let u be the solution of Eq. (1) in the sense of Definition 1. Then*

$$a(s) = 0, \quad s \in I(u^+(x, t), u^-(x, t)) \quad a.e. \text{ on } \Gamma_u, \tag{21}$$

which $I(\alpha, \beta)$ denote the closed interval with endpoints α and β , and (21) is in the sense of Hausdorff measure $H_N(\Gamma_u)$.

This lemma can be proved in a similar way as Zhan (2004); Zhao and Zhan (2005), we omit the details here.

Now, we will show that how Kruzkov’s bi-variables method, which was used to deal with the conservation law equation (Kružkov 1970) originally, can be used to prove the stability of the solutions to Eq. (1). Let u, v be two entropy solutions of Eq. (1) with initial values

$$u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x),$$

and with the boundary values (14)–(15), in particular, $u(x, t) = v(x, t) = 0$, $(x, t) \in \Sigma_1 \times (0, T)$.

By Definition 1, for $\varphi \in C_0^2(Q_T)$, we have

$$\iint_{Q_T} \left[I_\eta(u-k)\varphi_t - B_\eta^i(u,k)\varphi_{x_i} + A_\eta(u,k)\Delta\varphi - S'_\eta(u-k) \left| \nabla \int_0^u \sqrt{a(s)}ds \right|^2 \varphi \right] dxdt \geq 0, \tag{22}$$

$$\iint_{Q_T} \left[I_\eta(v-l)\varphi_\tau - B_\eta^i(v,l)\varphi_{y_i} + A_\eta(v,l)\Delta\varphi - S'_\eta(v-l) \left| \nabla \int_0^v \sqrt{a(s)}ds \right|^2 \varphi \right] dyd\tau \geq 0. \tag{23}$$

Let $\psi(x, t, y, \tau) = \phi(x, t)j_h(x - y, t - \tau)$, where $\phi(x, t) \geq 0$, $\phi(x, t) \in C_0^\infty(Q_T)$, and

$$j_h(x - y, t - \tau) = \omega_h(t - \tau)\prod_{i=1}^N \omega_h(x_i - y_i). \tag{24}$$

Here

$$\omega_h(s) = \frac{1}{h}\omega\left(\frac{s}{h}\right), \omega(s) \in C_0^\infty(R), \omega(s) \geq 0, \omega(s) = 0 \text{ if } |s| > 1, \int_{-\infty}^\infty \omega(s)ds = 1. \tag{25}$$

We choose $k = v(y, \tau)$, $l = u(x, t)$, $\varphi = \psi(x, t, y, \tau)$ in (22) (23), integrate over Q_T , then

$$\begin{aligned} & \iint_{Q_T} \iint_{Q_T} \left[I_\eta(u-v)(\psi_t + \psi_\tau) - (B_\eta^i(u,v)\psi_{x_i} + B_\eta^i(v,u)\psi_{y_i}) + A_\eta(u,v)\Delta_x\psi + A_\eta(v,u)\Delta_y\psi \right] \\ & - S'_\eta(u-v) \left(\left| \nabla \int_0^u \sqrt{a(s)}ds \right|^2 + \left| \nabla \int_0^v \sqrt{a(s)}ds \right|^2 \right) \psi dxdt dyd\tau = 0. \end{aligned} \tag{26}$$

Clearly,

$$\frac{\partial j_h}{\partial t} + \frac{\partial j_h}{\partial \tau} = 0, \quad \frac{\partial j_h}{\partial x_i} + \frac{\partial j_h}{\partial y_i} = 0, \quad i = 1, \dots, N; \quad \frac{\partial \psi}{\partial t} + \frac{\partial \psi}{\partial \tau} = \frac{\partial \phi}{\partial t} j_h, \quad \frac{\partial \psi}{\partial x_i} + \frac{\partial \psi}{\partial y_i} = \frac{\partial \phi}{\partial x_i} j_h.$$

Noticing that

$$\lim_{\eta \rightarrow 0} B_\eta^i(u, v) = \lim_{\eta \rightarrow 0} B_\eta^i(v, u) = \text{sgn}(u - v)(b_i(u) - b_i(v)),$$

as $\eta \rightarrow 0$, we have,

$$\begin{aligned} & \iint_{Q_T} \iint_{Q_T} [B_\eta^i(u, v)\psi_{x_i} + B_\eta^i(v, u)\psi_{y_i}] dxdt dyd\tau \\ & \rightarrow \iint_{Q_T} \iint_{Q_T} \text{sgn}(u - v)[b_i(u) - b_i(v)]\phi_{x_i} j_h dxdt dyd\tau, \end{aligned}$$

as $h \rightarrow 0$, we have

$$\begin{aligned} & \iint_{Q_T} \iint_{Q_T} \text{sgn}(u - v)[b_i(u) - b_i(v)]\phi_{x_i} j_h dxdt dyd\tau \\ & \rightarrow \iint_{Q_T} \text{sgn}(u - v)[b_i(u) - b_i(v)]\phi_{x_i} dxdt. \end{aligned} \tag{27}$$

At the same time, we have

$$\begin{aligned}
 & \iint_{Q_T} \iint_{Q_T} S'_\eta(u - v) \left(\left| \nabla_x \int_0^u \sqrt{a(s)} ds \right|^2 + \left| \nabla_y \int_0^v \sqrt{a(s)} ds \right|^2 \right) \psi dx dt dy d\tau \\
 &= \iint_{Q_T} \iint_{Q_T} S'_\eta(u - v) \left(\left| \nabla_x \int_0^u \sqrt{a(s)} ds \right| - \left| \nabla_y \int_0^v \sqrt{a(s)} ds \right| \right)^2 \psi dx dt dy d\tau \\
 &+ 2 \iint_{Q_T} \iint_{Q_T} S'_\eta(u - v) \nabla_x \int_0^u \sqrt{a(s)} ds \cdot \nabla_y \int_0^v \sqrt{a(s)} ds \psi dx dt dy d\tau, \tag{28}
 \end{aligned}$$

and

$$\begin{aligned}
 & \iint_{Q_T} [A_\eta(u, v) \Delta_x \psi + A_\eta(v, u) \Delta_y \psi] dx dt dy d\tau \\
 &= \iint_{Q_T} \iint_{Q_T} \{A_\eta(u, v) (\Delta_x \phi j_h + 2\phi_{x_i} j_{hx_i} + \phi \Delta j_h) + A_\eta(v, u) \phi \Delta_y j_h\} dx dt dy d\tau \\
 &= \iint_{Q_T} \iint_{Q_T} \{A_\eta(u, v) \Delta_x \phi j_h + A_\eta(u, v) \phi_{x_i} j_{hx_i} + A_\eta(v, u) \phi_{x_i} j_{hy_i}\} dx dt dy d\tau \\
 &- \iint_{Q_T} \iint_{Q_T} \{a(u) \widehat{S_\eta}(u - v) \frac{\partial u}{\partial x_i} - \int_u^v a(s) \widehat{S'_\eta}(s - v) ds \frac{\partial u}{\partial x_i}\} \phi j_{hx_i} dx dt dy d\tau, \tag{29}
 \end{aligned}$$

where

$$\begin{aligned}
 a(u) \widehat{S_\eta}(u - v) &= \int_0^1 a(su^+ + (1 - s)u^-) S_\eta(su^+ + (1 - s)u^- - v) ds, \int_u^v a(s) \widehat{S'_\eta}(s - v) ds \\
 &= \int_0^1 \int_{su^+ + (1-s)u^-}^v a(\sigma) S_\eta(\sigma - su^+ - (1 - s)u^-) d\sigma ds.
 \end{aligned}$$

Now, we will combine the last term on the right hand of (28) with the last term on the right hand side of (29). In details, by Lemma 1, at one hand, we have

$$\begin{aligned}
 & \iint_{Q_T} \iint_{Q_T} \nabla_x \nabla_y \int_v^u \sqrt{a(\delta)} \int_\delta^v \sqrt{a(\sigma)} S'_\eta(\sigma - \delta) d\sigma d\delta \psi dx dt dy d\tau \\
 &= \iint_{Q_T} \iint_{Q_T} \int_0^1 \int_0^1 \sqrt{a(su^+ + (1 - s)u^-)} \sqrt{a(\sigma v^+ + (1 - \sigma)v^-)} \\
 &\quad \times S'_\eta[\sigma v^+ + (1 - \sigma)v^- - su^+ - (1 - s)u^-] ds d\sigma \nabla_x u \nabla_y v dx dt dy d\tau \\
 &= \iint_{Q_T} \iint_{Q_T} \int_0^1 \int_0^1 S'_\eta[\sigma v^+ + (1 - \sigma)v^- - su^+ - (1 - s)u^-] ds d\sigma \\
 &\quad \times \widehat{\sqrt{a(u)}} \nabla_x u \widehat{\sqrt{a(v)}} \nabla_y v dx dt dy d\tau \\
 &= \iint_{Q_T} \iint_{Q_T} \int_0^1 \int_0^1 S'_\eta(v - u) \nabla_x \int_0^u \sqrt{a(s)} ds \nabla_y \int_0^v \sqrt{a(s)} ds dx dt dy d\tau. \tag{30}
 \end{aligned}$$

At the other hand, we have

$$\begin{aligned}
 & \iint_{Q_T} \iint_{Q_T} \nabla_x \nabla_y \int_v^u \sqrt{a(\delta)} \int_\delta^v \sqrt{a(\sigma)} S'_\eta(\sigma - \delta) d\sigma d\delta \psi dx dt dy d\tau \\
 &= \iint_{Q_T} \iint_{Q_T} \int_0^1 \sqrt{a(su^+ + (1 - s)u^-)} \\
 &\quad \times \int_{su^+ + (1-s)u^-}^v \sqrt{a(\sigma)} S'_\eta(\sigma - su^+ - (1 - s)u^-) d\sigma ds \frac{\partial u}{\partial x_i} j_{hx_i} \phi dx dt dy d\tau. \tag{31}
 \end{aligned}$$

By (30) (31), we have

$$\begin{aligned}
 & \iint_{Q_T} \iint_{Q_T} \left(a(u) \widehat{S'_\eta(u-v)} \frac{\partial u}{\partial x_i} - \int_u^v a(s) \widehat{S'_\eta(s-u)} ds \frac{\partial u}{\partial x_i} \right) j_{hx_i} \phi dx dt dy d\tau \\
 & + 2 \iint_{Q_T} \iint_{Q_T} S'_\eta(u-v) \nabla_x \int_0^u \sqrt{a(s)} ds \cdot \nabla_y \int_0^v \sqrt{a(s)} ds \psi dx dt dy d\tau \\
 & = \iint_{Q_T} \iint_{Q_T} \left[\int_0^1 a(su^+ + (1-s)u^-) S_\eta(su^+ + (1-s)u^- - v) ds \right. \\
 & - \int_0^1 \int_{su^+ + (1-s)u^-}^v a(\sigma) S'_\eta(\sigma - su^+ - (1-s)u^-) d\sigma ds \\
 & \left. + 2 \int_0^1 \sqrt{a(su^+ + (1-s)u^-)} \int_{su^+ + (1-s)u^-}^v \sqrt{a(\sigma)} \right. \\
 & \left. \times S'_\eta(\sigma - su^+ - (1-s)u^-) d\sigma ds \right] \frac{\partial u}{\partial x_i} j_{hx_i} \phi dx dt dy d\tau \\
 & = - \iint_{Q_T} \iint_{Q_T} \int_0^1 \int_{su^+ + (1-s)u^-}^v \left[\sqrt{a(\sigma)} - \sqrt{a(su^+ + (1-s)u^-)} \right] \\
 & \times S'_\eta(\sigma - su^+ - (1-s)u^-) d\sigma ds \frac{\partial u}{\partial x_i} j_{hx_i} \phi dx dt dy d\tau \rightarrow 0, \tag{32}
 \end{aligned}$$

as $\eta \rightarrow 0$.

Let us come back (29). Since

$$\lim_{\eta \rightarrow 0} A_\eta(u, v) = \lim_{\eta \rightarrow 0} A_\eta(v, u) = \text{sgn}(u - v)[A(u) - A(v)],$$

we have

$$\lim_{\eta \rightarrow 0} [A_\eta(u, v) \phi_{x_i} j_{hx_i} + A_\eta(u, v) \phi_{y_i} j_{hy_i}] = 0. \tag{33}$$

Combing (26)–(28) with (32)–(33), and letting $\eta \rightarrow 0, h \rightarrow 0$ in (26). We obtain

$$\begin{aligned}
 & \iint_{Q_T} [|u(x, t) - v(x, t)| \phi_t - \text{sgn}(u - v)(b_i(u) - b_i(v)) \phi_{x_i} + |A(u) - A(v)| \Delta \phi] dx dt \\
 & = \iint_{Q_T} \iint_{Q_T} S'_\eta(u - v) \left(\left| \nabla_x \int_0^u \sqrt{a(s)} ds \right| - \left| \nabla_y \int_0^v \sqrt{a(s)} ds \right| \right)^2 \psi dx dt dy d\tau \geq 0. \tag{34}
 \end{aligned}$$

By Kruzkov’s bi-variables method it means that, by a process of limit, we can choose a suitable test function $\phi \in C_0^1(Q_T)$ in (34), to obtain the stability of the solutions.

Proof of Theorem 1

The proof of Theorem 1 Let $x = \{x_1, x_2, \dots, x_i, \dots, N\}$ and define

$$d_i(x_i) = \begin{cases} x_i, & \text{if } 0 \leq x_i \leq \frac{1}{2}, \\ 1 - x_i, & \text{if } \frac{1}{2} < x_i \leq 1. \end{cases} \tag{35}$$

For small enough λ , we set

$$\varphi_{i\lambda}(x_i) = \begin{cases} \sin \frac{1}{\lambda}(d_i(x_i)), & \text{if } 0 \leq d_i(x_i) \leq \frac{\pi\lambda}{2}, \\ 1, & \text{if } d_i(x_i) \geq \frac{\pi\lambda}{2}. \end{cases} \tag{36}$$

Let $0 \leq \eta(t) \in C_0^1(t)$ and choose the test function in (34) as

$$\phi(x, t) = \eta(t) \prod_{j=1}^N \varphi_{j\lambda}(x_j). \tag{37}$$

Then

$$\begin{aligned} \partial_{x_i} \phi(x, t) &= \eta(t) \partial_{x_i} \varphi_{i\lambda}(x_i) \prod_{j=1, j \neq i}^N \varphi_{j\lambda}(x_j) \\ &= \eta(t) \frac{1}{\lambda} \cos \frac{1}{\lambda} (d_i(x_i)) d_{ix_i}(x_i) \prod_{j=1, j \neq i}^N \varphi_{j\lambda}(x_j), \quad 0 \leq d_i(x_i) \leq \frac{\pi\lambda}{2}. \end{aligned} \tag{38}$$

$$\begin{aligned} \Delta \phi(x, t) &= \frac{1}{\lambda} \eta(t) \prod_{j=1, j \neq i}^N \varphi_{j\lambda}(x_j) \left[-\frac{1}{\lambda} \sin \frac{1}{\lambda} (d_i(x_i)) d_{ix_i}^2 + \frac{1}{\lambda} \cos \frac{1}{\lambda} (d_i(x_i)) \Delta d_i(x_i) \right] \\ &= -\frac{1}{\lambda^2} \eta(t) \prod_{j=1, j \neq i}^N \varphi_{j\lambda}(x_j) \sin \frac{1}{\lambda} (d_i(x_i)) d_{ix_i}^2 \leq 0, \quad 0 \leq d_i(x_i) \leq \frac{\pi\lambda}{2}. \end{aligned} \tag{39}$$

For (39) in (34),

$$\iint_{Q_T} [|u(x, t) - v(x, t)| \phi_t - \text{sgn}(u - v) (b_i(u) - b_i(v)) \partial_{x_i} \phi(x, t)] dxdt \geq 0. \tag{40}$$

By that $|\partial_{x_i} \phi(x, t)| \leq \frac{1}{\lambda} |b_i(u) - b_i(v)| \leq c|u - v|$, we have

$$\iint_{Q_T} |u(x, t) - v(x, t)| \phi_t dxdt + c \int_0^T \int_{\Omega_\lambda} \eta(t) \frac{1}{\lambda} |u - v| dxdt \geq 0, \tag{41}$$

where $\Omega_\lambda = \{x \in \Omega : d_i(x_i) < \frac{\lambda\pi}{2}\}$. According to the definition of the trace of BV functions (Enrico 1984), when $x \in \Sigma_1, \gamma u = \gamma v = 0$, let $\lambda \rightarrow 0$ in (41). We have

$$\begin{aligned} \lim_{\lambda \rightarrow 0} \int_0^T \eta(t) \frac{1}{\lambda} \int_{\Omega_\lambda} |u - v| dxdt &= c \int_0^T \eta(t) |u - v|_{\partial\Omega} dt \\ &= c \int_0^T \eta(t) |u - v|_{\partial\Omega} dt \leq c \cdot \text{esssup}_{\Sigma_2 \times (0, T)} |f(x, t) - g(x, t)|. \end{aligned}$$

Let $\lambda \rightarrow 0$ in (41). Then

$$c \cdot \text{esssup}_{\Sigma_2 \times (0, T)} |f(x, t) - g(x, t)| + \iint_{Q_T} |u(x, t) - v(x, t)| \eta'_t dxdt \geq 0. \tag{42}$$

Let $0 < s < \tau < T$, and

$$\eta(t) = \int_{\tau-t}^{s-t} \alpha_\varepsilon(\sigma) d\sigma, \quad \varepsilon < \min\{\tau, T - s\}.$$

Here $\alpha_\varepsilon(t)$ is the kernel of mollifier with $\alpha_\varepsilon(t) = 0$ for $t \notin (-\varepsilon, \varepsilon)$. Then

$$c \cdot \text{esssup}_{\Sigma_2 \times (0, T)} |f(x, t) - g(x, t)| + \int_0^T [\alpha_\varepsilon(t - s) - \alpha_\varepsilon(t - \tau)] |u - v|_{L^1(\Omega)} dt \geq 0,$$

Let $\varepsilon \rightarrow 0$. Then

$$|u(x, \tau) - v(x, \tau)|_{L^1(\Omega)} \leq |u(x, s) - v(x, s)|_{L^1(\Omega)} + c \cdot \text{esssup}_{\Sigma_2 \times (0, T)} |f(x, t) - g(x, t)|$$

and the desired result follows by letting $s \rightarrow 0$.

On the half space

At the last section of the paper, let's consider Eq. (1) on the half space

$$\frac{\partial u}{\partial t} = \Delta A(u) + \text{div}(b(u)), \quad (x, t) \in \mathbb{R}_+^N \times (0, T), \tag{43}$$

with the initial value condition

$$u(x, 0) = u_0(x), \quad x \in \mathbb{R}_+^N, \tag{44}$$

where $\mathbb{R}_+^N = \{x = (x_1, x_2, \dots, x_N) : x_N > 0\}$ is the half space of \mathbb{R}^N . The author Zhan (2015b) had shown that when $b'_N(0) < 0$, we can give Dirichlet homogeneous boundary value

$$u(x, t) = 0, \quad (x, t) \in \partial \mathbb{R}_+^N \times (0, T) = \Sigma \times (0, T). \tag{45}$$

while $b'_N(0) \geq 0$, no any boundary value condition is necessary. Now, we give the definitions of the entropy solutions of the Eq. (43), which are the minor versions of the Definition 2.1 in Zhan (2015b).

Definition 2 Let $b'_N(0) < 0$. A function u is said to be the entropy solution of Eq. (43) with the initial value (44) and the boundary value (45), if

1. u satisfies

$$u \in BV(Q_T) \cap L^\infty(Q_T) \cap L^1(Q_T), \quad \frac{\partial}{\partial x_i} \int_0^u \sqrt{a(s)} ds \in L^2(Q_T).$$

2. For any $\varphi, \varphi \in C_0^2(Q_T), \varphi \geq 0$, for any $k \in \mathbb{R}$, for any small $\eta > 0$, u satisfies

$$\begin{aligned} & \iint_{Q_T} \left[I_\eta(u - k) \varphi_t - B_\eta^i(u, k) \varphi_{x_i} + A_\eta(u, k) \Delta \varphi \right. \\ & \left. - S'_\eta(u - k) \left| \nabla \int_0^u \sqrt{a(s)} ds \right|^2 \varphi \right] dx dt \geq 0. \end{aligned} \tag{46}$$

3. For any positive constant R ,

$$\lim_{t \rightarrow 0} \int_{B_R(x^R)} |u(x, t) - u_0(x)| dx = 0. \tag{47}$$

for any given positive constant R , where $x^R = (0, 0, \dots, 0, R)$, $B_R(x^R) = \{x \in \mathbb{R}_+^N : |x - x^R| < R\}$.

4. The boundary value condition (45) is true in the sense that the traces $\gamma u = \gamma v = 0$ on $\partial \mathbb{R}_+^N$ as usual.

Definition 3 Let $b'_N(0) \geq 0$. A function u is said to be the entropy solution of Eq. (43) with the initial value (44), if u satisfies Definition 2 except the fourth point. In this case, no boundary value condition is required.

Now, we actually are able to prove the existence of the solutions defined as Definition 2–3 in a similar way as Zhan (2015b), we omit the details here. In what follows, we only provide a new and simpler proof of the stability of the solutions.

Theorem 2 *Suppose that $A(s)$ and $b_i(s)$ are smooth enough. Let u, v be solutions of Eq. (1) with the different initial values $u_0(x), v_0(x) \in L^\infty(\mathbb{R}_+^N) \cap L^1(\mathbb{R}_+^N)$ respectively. If $b'_N(0) < 0$, suppose that the traces $\gamma u = \gamma v = 0$ on Σ as (45). Then*

$$\int_{\mathbb{R}_+^N} |u(x, t) - v(x, t)| dx \leq \int_{\mathbb{R}_+^N} |u_0 - v_0| dx. \tag{48}$$

If $b'_N(0) \geq 0$, then

$$\int_{\mathbb{R}_+^N} |u(x, t) - v(x, t)| dx \leq \int_{\mathbb{R}_+^N} |u_0 - v_0| dx + \text{cesssup}_{\Sigma \times (0, T)} |u(x, t) - v(x, t)|. \tag{49}$$

Proof By Kruzkov’s bi-variables method, as we have done in section “Kruzkov’s bi-variables method”, we can get

$$\iint_{Q_T} [|u(x, t) - v(x, t)| \phi_t - \text{sgn}(u - v)(b_i(u) - b_i(v)) \phi_{x_i} + |A(u) - A(v)| \Delta \phi] dx dt \geq 0. \tag{50}$$

Now, we can choose ϕ in (50) by

$$\phi(x, t) = \omega_\lambda(x) \eta(t),$$

where $\eta(t) \in C_0^\infty(0, T)$, and $\omega_\lambda(x) \in C_0^2(\Omega)$ is defined as follows: for any given small enough $0 < \lambda, 0 \leq \omega_\lambda \leq 1, \omega|_{\partial\Omega} = 0$, and when $x_N \geq \lambda$,

$$\omega_\lambda(x) = 1,$$

when $0 \leq x_N \leq \lambda$,

$$\omega_\lambda(x) = \omega_\lambda(x_N) = 1 - \frac{(x_N - \lambda)^2}{\lambda^2}.$$

Clearly,

$$\phi_{x_i} = \eta(t)(\omega_\lambda(x_N))_{x_i} \leq c|\omega'_\lambda(x_N)| \leq \frac{c}{\lambda} \Delta \phi = \eta(t) \Delta(\omega_\lambda(x_N)) = \eta(t) \nabla(\omega'_\lambda(x_N)) \nabla x_N = -\eta(t) \frac{2}{\lambda^2}.$$

Then by (50),

$$\begin{aligned} 0 &\leq \iint_{Q_T} |u(x, t) - v(x, t)| \eta'(t) \omega_\lambda(x) dx dt \\ &\quad - \iint_{Q_T} \left[\text{sgn}(u - v)(b_N(u) - b_N(v)) \eta(t) \frac{\partial \omega_\lambda(x)}{\partial x_N} + |A(u) - A(v)| \eta(t) \Delta \omega_\lambda(x) \right] dx dt \\ &\leq \iint_{Q_T} |u(x, t) - v(x, t)| \eta'(t) \omega_\lambda(x) dx dt + c \int_0^T \eta(t) dt \frac{1}{\lambda} \int_{\{x_N < \lambda\}} |u - v| dx. \end{aligned} \tag{51}$$

Noticing $\lim_{\lambda \rightarrow 0} \omega_\lambda = 1$, by the definition of the trace of a BV function, let $\lambda \rightarrow 0$ in (51).

Then

$$\iint_{Q_T} |u(x, t) - v(x, t)| \eta'(t) dx dt + c \operatorname{esssup} |u - v|_{\Sigma \times (0, T)} \geq 0. \quad (52)$$

Let $0 < s < \tau < T$, and

$$\eta(t) = \int_{\tau-t}^{s-t} \alpha_\epsilon(\sigma) d\sigma, \quad \epsilon < \min\{\tau, T-s\},$$

as before. Let $\epsilon \rightarrow 0$. Then

$$\int_{\Omega} |u(x, s) - v(x, s)| dx \leq \int_{\Omega} |u(x, \tau) - v(x, \tau)| dx + c \operatorname{esssup} |u - v|_{\Sigma \times (0, T)}.$$

Let $s \rightarrow 0$. If $b'_N(0) < 0$, suppose that the traces $\gamma u = \gamma v = 0$ on Σ ,

$$\int_{\mathbb{R}_+^N} |u(x, t) - v(x, t)| dx \leq \int_{\mathbb{R}_+^N} |u_0 - v_0| dx. \quad (53)$$

If $b'_N(0) \geq 0$, no any boundary value condition is necessary, then

$$\int_{\mathbb{R}_+^N} |u(x, t) - v(x, t)| dx \leq \int_{\mathbb{R}_+^N} |u_0 - v_0| dx + c \operatorname{esssup}_{\Sigma \times (0, T)} |u(x, t) - v(x, t)|. \quad (54)$$

we have the conclusion.

Conclusion

The paper shows that there is an essential difference of the boundary conditions between the strongly degenerate parabolic equation and the weakly degenerate parabolic equation. Instead of the whole boundary $\partial\Omega$, only a part of $\partial\Omega$ on which we can impose the boundary value if the well-posedness of the solutions to a strongly parabolic equation is considered. In physics, for example, if we consider a special case of Eq. (1), we consider the nonlinear heat conduction equation

$$u_t = \operatorname{div}(k(u)\nabla u),$$

if $k(0) = 0$, it means there is not heat flux across the boundary. Then the partial boundary $\Sigma_1 = \emptyset$, so there is no any boundary condition is necessary.

Acknowledgments

The paper is supported by NSF of China (No. 11371297), supported by NSF of Fujian Province (No. 2015J01592), China.

Competing interests

The author declares that he has no competing interests.

Received: 9 October 2015 Accepted: 23 September 2016

Published online: 18 October 2016

References

Bendahmane M, Karlsen KH (2004) Reharmonized entropy solutions for quasilinear anisotropic degenerate parabolic equations. *SIAM J Math Anal* 36(2):405–422

- Brezis H, Crandall MG (1979) Uniqueness of solutions of the initial value problem for $u_t - \Delta\varphi(u) = 0$. *J Math Pures et Appl* 58:153–163
- Carrillo J (1999) Entropy solutions for nonlinear degenerate problems. *Arch Ration Mech Anal* 147:269–361
- Chen GQ, Dibenedetto E (2001) Stability of entropy solutions to Cauchy problem for a class of nonlinear hyperbolic-parabolic equations. *SIAM J Math Anal* 33(4):751–762
- Chen GQ, Perthame B (2003) Well-posedness for non-isotropic degenerate parabolic-hyperbolic equations. *Ann I H Poincaré-AN* 20:645–668
- Cockburn B, Gripenberg G (1999) Continuous dependence on the nonlinearities of solutions of degenerate parabolic equations. *J Differ Equ* 151:231–251
- Enrico G (1984) Minimal surfaces and functions of bounded variation. Birkhauser, Bosten
- Evans LC (1998) Weak convergence methods for nonlinear partial differential equations. In: Conference board of the mathematical sciences, regional conferences series in mathematics no. 74
- Karlsen KH, Risebro NH (2003) On the uniqueness of entropy solutions of nonlinear degenerate parabolic equations with rough coefficient. *Discrete Contin Dye Syst* 9(5):1081–1104
- Kobayasi K, Ohwa H (2012) Uniqueness and existence for anisotropic degenerate parabolic equations with boundary conditions on a bounded rectangle. *J Differ Equ* 252:137–167
- Kružkov SN (1970) First order quasilinear equations in several independent variables. *Math USSR-Sb* 10:217–243
- Lions PL, Perthame B, Tadmor E (1994) A kinetic formation of multidimensional conservation laws and related equations. *J Am Math Soc* 7:169–191
- Li Y, Wang Q (2012) Homogeneous Dirichlet problems for quasilinear anisotropic degenerate parabolic- hyperbolic equations. *J Differ Equ* 252:4719–4741
- Oleinik OA, Samokhin VN (1999) Mathematical models in boundary layer theorem. Chapman and Hall/CRC, London
- Vol'pert AI, Hudjave SI (1975) Analysis of classof discontinuous functions and the equations of mathematical physics(Russian), Izda. Nauka Moskwa
- Vol'pert AI (1967) BV space and quasilinear equations. *MatSb* 73:255–302
- Vol'pert AI, Hudjaev SI (1967) On the problem for quasilinear degenerate parabolic equations of second order (Russian). *Mat Sb* 3:374–396
- Wu Z, Zhao J, Yin J, Li H (2001) Nonlinear diffusion equations. Word Scientific Publishing, Singapore
- Wu Z, Yin J (1989) Some properties of functions in BV_x and their applications to the uniqueness of solutions for degenerate quasilinear parabolic equations. *Northeast Math J* 5(4):395–422
- Wu Z, Zhao J (1983) The first boundary value problem for quasilinear degenerate parabolic equations of second order in several variables. *Chin Ann Math* 4B(1):57–76
- Wu Z, Zhao J (1983) Some general results on the first boundary value problem for quasilinear degenerate parabolic equations. *Chin Ann Math* 4B(3):319–328
- Yin J, Wang C (2007) Evolutionary weighted p -Laplacian with boundary degeneracy. *J Differ Equ* 237:421–445
- Zhan H (2004) The study of the Cauchy problem of a second order quasilinear degenerate parabolic equation and the parallelism of a Riemannian manifold, Doctor Thesis, Xiamen University
- Zhan H (2015) Homogeneous Dirichlet condition of anisotropic degenerate parabolic equation. *Bound Value Probl* 22(2015):1–22
- Zhan H (2015) The solutions of a hyperbolic-parabolic mixed type equation on half-space domain. *J Differ Equ* 259:1449–1481
- Zhao J (1985) Uniqueness of solutions of quasilinear degenerate parabolic equations. *Northeastern Math J* 1(2):153–165
- Zhao J, Zhan H (2005) Uniqueness and stability of solution for Cauchy problem of degenerate quasilinear parabolic equations, *Sci China Ser A* 48:583–593

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
