



## RESEARCH

## Open Access

# Succinate dehydrogenase subunit B inhibits the AMPK-HIF-1 $\alpha$ pathway in human ovarian cancer *in vitro*

Lilan Chen<sup>1,2†</sup>, Ting Liu<sup>1,2†</sup>, Shu Zhang<sup>1,2</sup>, Jinhua Zhou<sup>1,2</sup>, Yunfei Wang<sup>1,2</sup> and Wen Di<sup>1,2\*</sup>

## Abstract

**Background:** Ovarian carcinoma is one of the most common gynecological cancers with high mortality rates. Numerous evidences demonstrate that cancer cells undergo metabolic abnormality during tumorigenesis in tumor microenvironment and further facilitate tumor progression. Succinate dehydrogenase (SDH or Complex II) is one of the important enzymes in the tricarboxylic acid (TCA) cycle. Succinate dehydrogenase subunit B (SDHB) gene, which encodes one of the four subunits of SDH, has been recognized as a tumor suppressor. However the role of SDHB in ovarian cancer is still unclear.

**Methods:** Using the *SDHB* specific siRNA and overexpression plasmid, the expression of SDHB was silenced and conversely induced in ovarian cancer cell lines SKOV3 and A2780, respectively. The possible role of SDHB in ovarian cancer was investigated *in vitro*, using proliferation, migration and invasion assays. To explore the mechanism, proliferation and migration related proteins such as Bcl-2, cleaved caspase 3, p-ERK, MMP-2, and p-FAK were examined by western blot. P-P38, p-AMPK $\alpha$ , and HIF-1 $\alpha$  were also examined by western blot. CoCl<sub>2</sub> was used to induce HIF-1 $\alpha$  expression in SKOV3 and A2780 cells.

**Results:** *SDHB* silencing promoted cell proliferation, invasion, and migration, but inhibited apoptosis of SKOV3 and A2780 cells. In contrast, overexpression of SDHB inhibited cell proliferation, invasion, migration, and promoted apoptosis in SKOV3 cells. It was observed that up-regulation of Bcl-2 and MMP-2, activation of p-P38, p-ERK, and p-FAK, inhibition of cleaved caspase 3 in *SDHB*-silenced cells. Meanwhile, decreased Bcl-2 and MMP-2, inhibition of p-P38, p-ERK, and p-FAK, activation of cleaved caspase 3 were shown in *SDHB*-overexpressed SKOV3 cells. HIF-1 $\alpha$ , an essential factor in tumor progression, was up-regulated in *SDHB*-silenced cells with the activation of p-AMPK $\alpha$  and down-regulated in *SDHB*-overexpressed cancer cells with the decreased p-AMPK $\alpha$ . And SDHB was proved to be decreased due to upregulation of HIF-1 $\alpha$  expression in CoCl<sub>2</sub>-treated cancer cells.

**Conclusions:** Our results firstly revealed that SDHB played a key role in cell proliferation, invasion, migration, and apoptosis of human ovarian carcinoma *via* AMPK-HIF-1 $\alpha$  pathway. *SDHB*-overexpression might be a new approach to inhibit tumor progression in human ovarian carcinoma.

**Keywords:** Succinate dehydrogenase subunit B, HIF-1 $\alpha$ , AMPK, Gene silence, Gene overexpression, Ovarian carcinoma

\* Correspondence: diwen163@163.com

†Equal contributors

<sup>1</sup>Department of Obstetrics and Gynecology, Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai 200127, P. R. China<sup>2</sup>Shanghai Key Laboratory of Gynecologic Oncology, Shanghai 200127, P. R. China

## Background

Ovarian carcinoma is one of the leading causes of death among gynecological malignancies [1] that the most of ovarian carcinoma are firstly detected in advanced stages [2] and overall 5-year survival rate is as low as 40% [3]. Current therapeutic strategies against advanced stage ovarian carcinoma include surgical resection along with platinum-based chemotherapy. However, there are serious side effects from platinum-based drugs and surgical intervention [2]. Therefore, it is critical to understand the molecular events leading to initiation and progression of this devastating disease [4].

Succinate dehydrogenase (SDH), known as succinate-ubiquinone oxidoreductase, is a mitochondrial enzyme complex that catalyses the oxidation of succinate to fumarate in the citric acid cycle and participates in the electron transport chain [5]. Mutations of the *SDH* gene determine the genetic basis of paragangliomas/pheochromocytomas (PCCs/PGLs) [6,7] and associated tumour syndromes (Carney-Stratakis syndrome and Carney triad) [8]. Furthermore, mRNA expression of *SDHB* is decreased in recovering Ramos cells in childhood non-Hodgkin lymphoma (NHL) [9]. Reduced *SDHB* expression is associated with growth and de-differentiation of colorectal cancer cells [10]. In addition, loss of *SDHB* expression is associated with an adverse outcome in PCCs/PGLs, indicating *SDHB* might be a predictive marker of adverse outcome both in sporadic and familial PCC/PGL [11]. However, the role of *SDHB* in ovarian carcinoma tumorigenesis, especially its association with cellular proliferation, invasion, and migration are not fully elucidated.

Genetic analysis of hereditary paraganglioma reveals an activation of the hypoxia-response pathway [12]. Immunohistochemical analysis shows strong staining of hypoxia-inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ) and the angiogenic factor, vascular endothelial growth factor in a malignant sporadic pheochromocytoma caused by a germline missense mutation in the *SDHB* gene [13], suggesting that activation of the hypoxia-response pathway may be a common theme underlying SDH (and also FH) mutation [14,15].

Hypoxia-inducible factor-1 (HIF-1), consists of a constitutively expressed  $\beta$ -subunit and an inducible-expressed  $\alpha$ -subunit [16], is a well-established mediator in hypoxia-response pathway. HIF-1 $\alpha$  is accumulated under hypoxic conditions, which activates transcription of target genes involved in angiogenesis, energy metabolism, adaptive survival or apoptosis [17,18]. HIF-1 $\alpha$  is highly expressed in ovarian cancer and is associated with tumour proliferation [19], invasion and metastasis [20,21]. AMP-activated protein kinase (AMPK) is a metabolic sensor that helps maintain cellular energy homeostasis and modulate metabolic stresses such as hypoxia and respiratory impairment. AMPK has been linked to the regulation of tumorigenesis [22] and HIF-1 $\alpha$  mediates the growth

advantage of tumours with reduced AMPK signaling [23]. However, the precise linkage between metabolism dysfunction (HIF-1 $\alpha$ , AMPK) and the propensity for tumorigenesis has not been fully elucidated in ovarian carcinoma. In current study, we aimed to gain a better understanding of the consequences of *SDHB* alteration by gene silencing or overexpression in human ovarian cancer cell lines. Here, we provided the first time that *SDHB* silencing resulted in increased tumour cell proliferation, invasion, migration and decreased apoptosis. Conversely, overexpression of *SDHB* inhibited cell proliferation, invasion, migration, and promoted apoptosis. Further, HIF-1 $\alpha$  and p-AMPK $\alpha$  were found to be upregulated in *SDHB*-silenced, but downregulated in *SDHB*-overexpressed cancer cells. Moreover, *SDHB* was downregulated by hypoxia mimetic CoCl<sub>2</sub> in human ovarian cancer cells. Our data suggested that *SDHB* plays an essential role in ovarian cancer cell proliferation, invasion, migration and apoptosis via AMPK-HIF-1 $\alpha$  pathway. *SDHB*-overexpression might be a potential therapeutic strategy to inhibit tumour progression in human ovarian carcinoma.

## Methods

### Tissue specimens and ethics

A total of 25 tissue specimens were collected from surgical patients enrolled in this study, including 7 normal human ovarian epithelium tissues and 18 ovarian carcinoma tissues between July, 2011 to August, 2012. In addition, 7 metastatic ovarian carcinoma were collected. All specimens, obtained during surgery (Department of Obstetrics and Gynaecology, Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University, China), frozen immediately in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until analysis. All carcinoma patients were newly diagnosed with ovarian tumours, and received no chemical therapy prior to surgery. The study was approved by the Institutional Review Board of Ren Ji Hospital, Shanghai Jiao Tong University School of Medicine. Written informed consents were obtained from all patients. Clinical investigation was conducted according to the principles expressed in the Declaration of Helsinki.

### Cell lines and cell culture

Human epithelial ovarian adenocarcinoma cancer (EOC) cell lines SKOV3 and A2780 obtained from the Cell Bank Chinese Academy of Science (Shanghai, China), cultured in RPMI1640 medium (Hyclone) supplemented with 10% foetal bovine serum (Gibco). These cells were incubated at  $37^{\circ}\text{C}$  in a humidified atmosphere of 95% air and 5% CO<sub>2</sub>. The medium was replaced every 24 h. Hypoxic cells were incubated in the same conditions but in a hypoxic atmosphere with different concentrations of cobalt chloride (CoCl<sub>2</sub>, Sigma) for 24 h.

#### **Transient transfection of SKOV3 and A2780 cells with SDHB siRNA oligonucleotides**

Transient small interfering RNA (siRNA) oligonucleotides were synthesized from Shanghai Integrated Biotech Solutions Co.Ltd. The target sequences were as follows: siSucA 5'-GAT TAA GAA TGA AGT TGA CTC-3' [24], siSucC 5'-GCT CAG AGC TGA ACA TAA TT-3' [24]. As a control for silencing, we constructed a negative control (NC) siRNA (5'-UUUC UCC GAA CGU GUC ACG UTT-3') that did not affect *SDHB* expression. Transient transfection was carried out using Lipofectamine 2000 reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. Cells were collected after 24 h for RNA extraction and 48 h for protein electrophoresis.

#### **Transient transfection of ovarian cancer cell SKOV3 with SDHB overexpression pIRES2-EGFP plasmid**

The full length *SDHB* opening reading frame was subcloned into *pIRES2-EGFP* plasmid. The plasmid was purchased from Shanghai Integrated Biotech Solutions Co., Ltd. SKOV3 cells were transfected with the *pIRES2-EGFP SDHB* vector encoding *SDHB* cDNA using lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. Vector-only transfecting cells were used as controls.

#### **RNA extraction and quantitative real-time reverse transcriptase PCR**

Total RNA, extracted from cancer cell lines and tissue samples using TRIzol Reagent (Invitrogen, Carlsbad, CA), was reverse transcribed using a PrimeScript RT reagent Kit (TaKaRa, Shanghai, China). Resultant complementary DNAs (cDNAs) underwent quantitative real-time reverse transcriptase PCR using a SYBR Green PCR Master Mix Reagent Kit (TaKaRa). Real-time PCR and data collection were performed on an ABI 7500 real-time system (Applied Biosystems). The primers for *SDHB* were 5'-AAA TGT GGC CCC ATG GTA TTG-3' (forward) and 5'-AGA GCC ACA GAT GCC TTC TCT G-3' (reverse). The primers for  $\beta$ -actin, serving as the endogenous controls, were 5'-TGA CGT GGA CAT CCG CAA AG-3' (forward) and 5'-CTG GAA GGT GGA CAG CGA GG-3' (reverse). SDS v.1.4 (Applied Biosystems) software was used to perform comparative delta cycle threshold (Ct) analysis. To minimize random variation in cell experiments, each real-time PCR experiment was performed in triplicate.

#### **Succinate dehydrogenase activity assay**

Succinate dehydrogenase activity was determined, using succinate dehydrogenase kit (Beijing Solarbio Science & Technology Co., Ltd, China). Protein level was quantified by BCA kit (Cwbiotech, China).

#### **ATP assay**

Intracellular ATP concentrations were determined using phosphomolybdic acid colorimetric method (Nanjing Jiancheng Bioengineering Institute, China) according to the manufacturer's protocols. Protein level was quantified by BCA kit (Cwbiotech, China).

#### **Cell proliferation analysis**

Cell proliferation was assessed by Cell Counting kit-8 (CCK-8) (Dojindo). Briefly,  $1 \times 10^3$  SKOV3 and/ or A2780 cells were seeded into 96-well plates treated with *SDHB* siRNA oligonucleotides or *pIRES2-EGFP SDHB* vector. After 1, 2, 3, 4, 5 or 6 day treatment, 10  $\mu$ l CCK8 mixed with 90  $\mu$ l RPMI1640 was added to each well for further 1.5 h. The absorbance (450 nm) from each well was measured at the indicated time points.

#### **Cell invasion and migration assay**

SKOV3 and A2780 cells ( $6 \times 10^4$ ) (48 h post-transfection) were placed into the upper compartment of the chambers (8  $\mu$ m pore size, Millipore) coated with 50  $\mu$ l BD Matrigel (diluted 1:7 in serum-free medium) and placed into 24-well plates. Medium containing 10% foetal bovine serum was added to the lower chamber. After incubation at 37°C for 24 h, cells on the upper side of the membrane were removed using sterile cotton swabs. Cells adhering to the lower surface were fixed in 4% paraformaldehyde and stained with 0.1% crystal violet. Cells were counted using microscope (Nikon TE300, Tokyo, Japan) at 200 $\times$  magnification. Five random fields were selected for examination on each membrane, and results were expressed in terms of the invasion cells per field. Each experiment was conducted in triplicate. Cell migration was performed using a similar approach without Matrigel coating, and the treated cells were incubated for 12 h.

#### **Western blot**

Cells and tissue samples were lysed in RIPA buffer with PMSF as protease inhibitor (Beyotime, Shanghai, China). Proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred on NC or PVDF membrane (Millipore, Bedford, MA). After blocking with 5% dry milk in TBST for 2 h at room temperature, the membranes were incubated with the primary antibodies against *SDHB* (1:1000, Epitomics),  $\beta$ -tubulin (1:4000, Epitomics),  $\beta$ -actin (1:500, Abmart), caspase 3 (1:1000, Cell Signalling Technology), Bcl-2 (1:4000, Epitomics), MMP-2 (1:500, Abcam), FAK (1:1000, CST), p-FAK (1:1000, CST), AMPK $\alpha$  (1:1000, CST), p-AMPK $\alpha$  (1:1000, CST), GAPDH (1:4000, Abmart), P38 (1:1000, CST), p-P38 (1:1000, CST), ERK (1:1000, CST), p-ERK (1:1000, CST), HIF-1 $\alpha$  (1:1000, Epitomics) in dilution buffer overnight at 4°C. Membranes were washed for three times with TBST, then were incubated with IRDye 800CW conjugated goat (polyclonal)

anti-Rabbit IgG or anti-Mouse IgG (1:10000) antibodies for 1 h at room temperature. The expression of specific proteins was detected through the use of Odyssey system following the manufacturer's instructions.

### Statistical analysis

All data were calculated as the means  $\pm$  standard error (SEM). An independent Student *t* test was used to compare the continuous variables between groups using GraphPad Prism 5.0 software (San Diego, CA).  $P < 0.05$  was considered statistical significant.

### Results


#### The effect of *SDHB* silencing on ATP and AMPK/P38 MAPK in human ovarian cancer cells

The role of *SDHB* on ATP and AMPK/P38 MAPK was examined using gene silencing strategy. The efficiency of *SDHB* silencing was confirmed by 24 h *SDHB* siRNA oligonucleotides treatment in SKOV3 or A2780 cells, mRNA level was reduced by 89.80% (siSucA vs NC in SKOV3), 84.89% (siSucC vs NC in SKOV3) ( $P < 0.001$ ), and 83.88% (siSucA vs NC in A2780) ( $P < 0.001$ ), respectively (Figure 1A). Furthermore, *SDHB* protein level and activity were also decreased following 48 h transfection

(Figure 1B,C). Meanwhile, decreased ATP was observed in *SDHB*-silenced cells (Figure 1D). AMPK acts as an energy sensor that modulate metabolic stresses such as hypoxia and respiratory impairment [25,26], exogenous stimuli that increase AMP or decrease ATP could activate AMPK [27]. To examine the effect of *SDHB* silencing on AMPK activation, the level of phosphorylated AMPK $\alpha$  (p-AMPK $\alpha$ ) was analysed. P-AMPK $\alpha$  was increased in the *SDHB*-silenced cells compared to NC group (Figure 1E). In addition, the AMPK/P38 MAPK signalling cascade stimulates glucose uptake during metabolic stress [28,29]. Moreover to confirm the effect of decreased AMPK activity in *SDHB*-silenced cells, phosphorylated p38 MAPK (p-P38 MAPK) level was also examined. P-P38 MAPK was increased in the *SDHB*-silenced cells compared to control cells (Figure 1F).

#### *SDHB* silencing promoted ovarian cancer cell proliferation

CCK8 was used to examine the influence of *SDHB* on cell proliferation. *SDHB* silencing significantly promoted ovarian cancer cell proliferation ( $P < 0.05$ ) (Figure 2A, B). Extracellular signal-regulated kinase (ERK)1/2 mitogen-activated protein (MAP) kinase pathway plays a central role in cell proliferation control [30]. Multiple lines of



**Figure 1** *SDHB* silencing in SKOV3 and A2780 cells. SKOV3 and A2780 cells were treated with siSucA or siSucC oligonucleotides, (A) real-time PCR (24 h) and (B) western blot (48 h) were performed to detect *SDHB* mRNA and protein levels, respectively. 48 h after siRNA transfection, *SDH* activity (C) and ATP level (D) were examined, p-AMPK $\alpha$ , AMPK $\alpha$  (E) and p-P38, P38 (F) were measured in SKOV3 and A2780 cells. Mean  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ .






Figure 2 (See legend on next page.)

(See figure on previous page.)

**Figure 2** *SDHB* silencing contributed to ovarian cancer cell proliferation in SKOV3 and A2780 cells. (A, B) Cell proliferation was assessed following *SDHB* silencing treatment for 6 days by CCK8. Mean  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ . (C) 48 h post transfection, p-ERK, ERK were examined by western blot. (D, E) 48 h post-transfection, SKOV3 and A2780 cells were treated with DDP at indicated concentration for 24 h, cleaved caspase 3, total caspase 3, and Bcl-2 were then analysed by western blot.

evidence have implicated the ERK1/2 MAP kinase pathway in the control of cell proliferation [30,31]. ERK1 and ERK2 are activated in response to virtually all mitogenic factors [30]. ERK signalling is often up regulated in a diverse range of human cancers [32]. It was discovered that the level of p-ERK was increased in the *SDHB*-silenced cells compared to control (Figure 2C), indicating that inhibition of *SDHB* expression promoted tumour cell growth through ERK pathway in SKOV3 and A2780 cells.

#### ***SDHB* silencing prevented apoptosis in human ovarian cancer cells**

Cleaved caspase-3, and Bcl-2 are representative proteins related to cell apoptosis. Cis-platinum (DDP, Sigma) is a chemotherapy for ovarian cancer, by forming a platinum complex inside the cell which binds to DNA and cross-links DNA and causes the cells to undergo apoptosis, or systematic cell death. Cleaved caspase-3 was detected in DDP treated cells for 24 h and its expression was decreased in *SDHB*-silenced cells (Figure 2D). Meanwhile, Bcl-2 was increased in *SDHB*-silenced cells (Figure 2E). The results suggested *SDHB* silencing could prevent ovarian cancer cell apoptosis.

#### ***SDHB* silencing promoted cell invasion and migration of SKOV3 and A2780 cells**

Invasion is among the most significant biological characteristics of malignant tumours. Matrigel cellular invasion assay showed that *SDHB* silencing promoted SKOV3 ( $P < 0.05$ ) and A2780 ( $P < 0.01$ ) invasiveness (Figure 3A). Migration assay showed that *SDHB* silencing upregulated migration ability of SKOV3 ( $P < 0.001$ ) and A2780 cells ( $P < 0.001$ ) (Figure 3B). The active, phosphorylated form of focal adhesion kinase (p-FAK) is a non-receptor protein tyrosine kinase that is involved in cell migration, proliferation, survival and metastasis [33,34]. The extent of p-FAK overexpression correlates with increased metastasis and decreased survival in human ovarian cancer [33,34]. MMP-2 degrades components of the extracellular matrix (ECM) proteins, contributing to tumour invasion and metastasis [35,36]. Up-regulation of p-FAK and MMP-2 expression was observed in *SDHB*-silenced cells (Figure 3C).

#### ***SDHB* overexpression inhibited cell proliferation and promoted apoptosis in SKOV3 cells**

SKOV3 cell line was chosen to overexpress *SDHB* since *SDHB* expression in SKOV3 was lower than that in A2780

(see Additional file 1: Figure S1). After treated with *SDHB*-overexpressed plasmid for 24 h, *SDHB* mRNA level was up regulated 33 fold in SKOV3 ( $P < 0.001$ ) (Figure 4A). Under the same condition, *SDHB* protein expression was increased after 48 h transfection (Figure 4A). CCK8 was used to examine the cell viability. Over-expression of *SDHB* significantly inhibited ovarian cancer cell proliferation (Figure 4B). Likewise, Bcl-2 was decreased in *SDHB*-overexpressed SKOV3 cells (Figure 4C). These results revealed that overexpression of *SDHB* could inhibition cell growth in SKOV3 cancer cells.

#### **Overexpression of *SDHB* inhibited invasion and migration in SKOV3 cells**


*SDHB* overexpression inhibited SKOV3 cell invasion ( $P < 0.05$ ) and migration ( $P < 0.05$ ) ability (Figure 4D). Down-regulation of p-FAK and MMP-2 expression was also found in *SDHB* overexpressed SKOV3 cells (Figure 4E).

#### ***SDHB* affected HIF-1 $\alpha$ level in ovarian cancer cells**

Mitochondrial dysfunction repressed the protein synthesis of HIF-1 $\alpha$  as well as reduced its trans-activation activity through AMPK signalling in human hepatoma HepG2 cells [37]. HIF-1 $\alpha$  was up regulated in *SDHB*-silenced SKOV3 and A2780 cells, but down regulated in *SDHB*-overexpressed SKOV3 cells (Figure 5A,B). Consistent with these results, *SDHB* protein level was down regulated and conversely correlated with HIF-1 $\alpha$  in hypoxic conditions induced by CoCl<sub>2</sub> in ovarian cancer cells in a dose dependent manner (Figure 5C). To verify the effect of *SDHB* on energy metabolism in ovarian cancer cells, p-AMPK $\alpha$  was also analysed in *SDHB* over expressed SKOV3 cells. The level of p-AMPK $\alpha$  was decreased compared to the control (Figure 4F). These results showed *SDHB* might affect ovarian cancer cell phenotype via AMPK-HIF-1 $\alpha$  signalling pathway.

#### ***SDHB* expression in human ovarian carcinoma and normal ovarian epithelium tissues**

To confirm *SDHB* expression in normal and malignant ovarian tissues, mRNA and protein level of *SDHB* were determined. *SDHB* mRNA and protein level was significantly down regulated in human ovarian carcinoma compared with the controls (Figure 6A,B). Interestingly, *SDHB* mRNA expression was 36.80% lower in the corresponding metastasis ( $P < 0.05$ ) than that in the primary ovarian carcinoma tissues (Figure 6C). The level of



*SDHB* in tissue specimens was consistent with our results shown above.

## Discussion

In our study, we aimed to gain a better understanding of the role of *SDHB* in ovarian carcinoma by gene silencing and over expression. *SDHB* is one of the subunits of *SDH* takes parts in TCA [5], ATP is an indicator of energy metabolism. We found that the level ATP was decreased in *SDHB*-silenced cancer cells. On the other hand, p-AMPK $\alpha$  and p-P38 MAPK level were increased in the *SDHB*-silenced cells, suggesting that *SDHB* silencing could activate AMPK pathway in ovarian cancer.

*SDHB* protein is significantly lower compared with other *SDH* subunits in colorectal cancer tissues [10]. Notably, reduced *SDHB* expression in tumour tissues is associated with tumour de-differentiation. Restoration of *SDHB* inhibits the growth of cancer cells [10]. In our current study, *SDHB*-silenced cells increased proliferation in SKOV3 and A2780 cells, accompanied with elevated p-ERK level. Bcl-2 is functioned to oppose the apoptosis pathway of programmed cell death [38,39]. Effector caspases are responsible for initiating the events that lead to the hallmarks of apoptosis. Caspase-9, an essential initiator caspase required for apoptosis through mitochondrial pathway, is activated on the apoptosome




Figure 4 (See legend on next page.)



(See figure on previous page.)


**Figure 4** *SDHB* overexpression inhibited cell proliferation in SKOV3 cells. (A) SKOV3 cells were treated with *pIRES2-SDHB* or empty vector, then *SDHB* mRNA (24 h after treatment) and protein (48 h after treatment) were analysed by real time-PCR or western blot, respectively. (B) 48 h after transfection, cell viability of *SDHB*-overexpressed SKOV3 cells was examined using CCK8 assay for another 6 days. (C) Bcl-2 was downregulated after 48 h transfection in SKOV3 with  $\beta$ -actin as a loading control by western blot. (D) SKOV3 cells were treated with *SDHB* overexpression plasmid for 48 h, cell invasion and migration ability was evaluated as described previously. (E) MMP-2 and p-FAK, FAK were also analysed by western blot. (F) p-AMPK $\alpha$ , AMPK $\alpha$  were measured. Mean  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$ , and \*\*\* $P < 0.001$ .

complex, which directly resulted in cleave and activate effector caspases, such as caspase-3 [40,41]. Anti-apoptotic Bcl-2 level was increased and cleaved caspase 3 was decreased after treated with or without DDP in *SDHB*-silenced SKOV3 and A2780 cells.


FAK and MMPs are associated with metastases [42], secretion and activation of MMP-2 may be responsible for increased motility, invasiveness and metastasis of malignant cells [43]. In addition, increased FAK expression and activity frequently correlate with metastatic disease and poor prognosis [44]. We found that *SDHB* silencing promoted ovarian cancer invasion and migration accompanied with up-regulated expression of MMP-2 and p-FAK. To explore the role of *SDHB* in human ovarian carcinoma, we designed plasmid to induce *SDHB* expression in SKOV3 cells. *SDHB* overexpression inhibited cell proliferation in SKOV3. Cellular invasion and migration were inhibited in *SDHB*-over-expressed cells accompanied with reduced expression of p-FAK and MMP-2. These observations further verified the role of *SDHB* in ovarian carcinoma.

The relationship between HIF-1 $\alpha$  and *SDHB* is controversial in tumorigenesis. *SDHB* is one of the subunits of SDH which takes part in TCA cycle and respiratory

chain [5], AMPK could modulate metabolic stresses such as hypoxia. Study showed mitochondrial dysfunctions could result in reduced HIF-1 $\alpha$  protein synthesis through AMPK-dependent manner in HepG2 cells [37]. Hypoxia is a characteristic of many malignancies arising from various sites [45] and HIF-1 $\alpha$  is a hypoxia responsive factor [16]. Research reported that HIF-1 $\alpha$  was up-regulated in chronically *SDHB*-silenced Hep3B cells [24], HIF-1 $\alpha$  was overexpressed in *SDH*-deficient leiomyomas and renal cell cancer (HLRCC) [15]. While there was also report that increased HIF-1 $\alpha$  was not associated with loss of *SDHB* expression on a series of familial and sporadic tumours [11]. In order to find out the key factor contributed to the phenomenon in our experiment, HIF-1 $\alpha$  was analysed in treated cancer cells. The expression of HIF-1 $\alpha$  was strongly elevated in *SDHB*-silenced ovarian cancer cells. Moreover, *SDHB* was decreased in CoCl<sub>2</sub>-treated cancer cells accompanied by HIF-1 $\alpha$  up-regulation. In accordance with these results, over-expressed *SDHB* could inhibit HIF-1 $\alpha$  expression. The results showed *SDHB* affected ovarian cancer progression by altering HIF-1 $\alpha$  expression. It was suggested that *SDHB* silencing up-regulated HIF-1 $\alpha$  *via* activation of AMPK $\alpha$  in cancers in accordance with the aerobic glycolysis



**Figure 5** *SDHB* modulated HIF-1 $\alpha$  expression in ovarian cancer cells. SKOV3 and/ or A2780 cells were treated with *SDHB* siRNA (A) or over-expression plasmid (B) for 48 h, HIF-1 $\alpha$  expression was analyzed by western blot with  $\beta$ -actin or  $\beta$ -tubulin as a loading control. (C) SKOV3 and A2780 cells were treated with CoCl<sub>2</sub> at indicated concentrations for 24 h, HIF-1 $\alpha$  and *SDHB* were examined by western blot.



**Figure 6** Different expression of SDHB in human ovarian carcinoma tissues. **(A)** The mRNA expression of *SDHB* in ovarian carcinoma tissues (n = 18) and normal ovarian epithelium tissues (n = 7) by real-time PCR (N = normal ovarian epithelium, M = ovarian carcinoma). **(B)** SDHB protein level in ovarian carcinoma and normal ovarian epithelium by western blot, M1 and M2 represent different patients. **(C)** *SDHB* mRNA expression in primary ovarian carcinoma (n = 7) and the corresponding metastasis (n = 7) by real-time PCR. Mean  $\pm$  SEM. \* $P < 0.05$  and \*\* $P < 0.01$ .

(Warburg effect) [46]. To verify the effect of *SDHB* on energy metabolism in ovarian cancer, p-AMPK $\alpha$  was also analysed in *SDHB*-overexpressed SKOV3 cells. The level of p-AMPK $\alpha$  was decreased compared to control. In addition, HIF-1 $\alpha$  is decreased by inhibiting ERK pathway in cervical carcinoma CaSki cells [47], but apoptosis is enhanced by HIF-1 $\alpha$  knockdown in pancreatic cancerous BxPC-3 cells [48]. HIF-1 $\alpha$  also promotes cell migration by regulating MMP-2 [49] and FAK [50], which is in line with our current data. These results showed *SDHB* might affect cancer cell proliferation, invasion, migration, and apoptosis *via* AMPK-HIF-1 $\alpha$  in ovarian carcinoma.

Finally, we compared the expression of *SDHB* between ovarian carcinoma and normal ovarian epithelium tissues. *SDHB* mRNA and protein was decreased in human ovarian carcinoma compared with normal ovarian epithelium. In addition, *SDHB* mRNA expression was decreased in the corresponding metastasis compared with the primary ovarian carcinoma, suggesting *SDHB* contributed to tumour metastasis. Moreover, it was well known that *SDHB*-associated PCCs/PGLs often metastasise in a familial setting [11]. Our current data suggested that *SDHB* play an important role in cancer progression. Enhanced *SDHB* expression inhibited cell proliferation,

invasion, migration and promoted apoptosis in human ovarian carcinoma. The relationship between *SDHB* expression and clinical characteristics in human ovarian carcinoma will be determined in our future studies.

## Conclusions

In current study, we demonstrated that *SDHB* played an important role in cellular proliferation, invasion, migration and apoptosis *via* AMPK-HIF-1 $\alpha$  pathway in human ovarian carcinoma. Overexpression of *SDHB* might be an effective therapeutic strategy for treatment of ovarian carcinoma.

## Additional file

**Additional file 1: Figure S1.** *SDHB* expression in SKOV3 and A2780. (A, B) Real-time PCR and western blot were used to analyse *SDHB* mRNA and protein expression in SKOV3 and A2780 cells.

## Abbreviations

MMP-2: Matrix metalloproteinase-2; FAK: Focal adhesion kinase; Bcl-2: B-cell lymphoma-2; CoCl<sub>2</sub>: Cobalt chloride; DDP: Cis-platinum; RIPA: Radio immuno-precipitation assay; PMSF: Phenylmethanesulfonyl fluoride; TBST: Tris-buffered saline with Tween-20; NC: Nitrocellulose membrane; PVDF: Polyvinylidene fluoride.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

LLC, WD and SZ conceived and designed the experiments, LLC and TL performed the experiments, LLC and TL analysed the data, JHZ and YFW contributed reagents/materials/analysis tools, LLC wrote the paper. All authors have approved the final manuscript.

### Acknowledgements

This work was supported by grants from *Shanghai Committee of Science and Technology* (Grant No. 10dz2212100, Grant No.12411950200), *National Natural Science Foundation* (Grant No.81072137, Grant No. 81272882), *Shanghai Health Bureau Key Disciplines and Specialties Foundation*. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Received: 20 September 2014 Accepted: 27 November 2014

Published online: 10 December 2014

### References

- Jelovac D, Armstrong DK: Recent progress in the diagnosis and treatment of ovarian cancer. *CA Cancer J Clin* 2011, **61**(3):183–203.
- Cannistra SA: Cancer of the ovary. *N Engl J Med* 2004, **351**(24):2519–2529.
- Baldwin LA, Huang B, Miller RW, Tucker T, Goodrich ST, Podzielinski I, DeSimone CP, Ueland FR, van Nagell JR, Seamon LG: Ten-year relative survival for epithelial ovarian cancer. *Obstet Gynecol* 2012, **120**(3):612–618.
- Zhu J, Zhang S, Gu L, Di W: Epigenetic silencing of DKK2 and Wnt signal pathway components in human ovarian carcinoma. *Carcinogenesis* 2012, **33**(12):2334–2343.
- Gaude E, Frezza C: Defects in mitochondrial metabolism and cancer. *Cancer Metab* 2014, **2**:10.
- Favier J, Briere JJ, Strompf L, Amar L, Filali M, Jeunemaitre X, Rustin P, Gimenez-Roqueplo AP: Hereditary paraganglioma/pheochromocytoma and inherited succinate dehydrogenase deficiency. *Horm Res* 2005, **63**(4):171–179.
- Baysal BE: On the association of succinate dehydrogenase mutations with hereditary paraganglioma. *Trends Endocrinol Metab* 2003, **14**(10):453–459.
- Gaal J, Stratakis CA, Carney JA, Ball ER, Korpershoek E, Lodish MB, Levy I, Xekouki P, van Nederveen FH, den Bakker MA: SDHB immunohistochemistry: a useful tool in the diagnosis of Carney-Stratakis and Carney triad gastrointestinal stromal tumors. *Mod Pathol* 2011, **24**(1):147–151.
- Kusao I, Troelstrup D, Shiramizu B: Possible Mitochondria-Associated Enzymatic Role in Non-Hodgkin Lymphoma Residual Disease. *Cancer Growth Metastasis* 2008, **1**:3–8.
- Zhang D, Wang W, Xiang B, Li N, Huang S, Zhou W, Sun Y, Wang X, Ma J, Li G: Reduced succinate dehydrogenase B expression is associated with growth and de-differentiation of colorectal cancer cells. *Tumour Biol* 2013, **34**(4):2337–2347.
- Blank A, Schmitt AM, Korpershoek E, van Nederveen F, Rudolph T, Weber N, Strelbe RT, de Krijger R, Komminoth P, Perren A: SDHB loss predicts malignancy in pheochromocytomas/sympathetic paragangliomas, but not through hypoxia signalling. *Endocr Relat Cancer* 2010, **17**(4):919–928.
- Gimenez-Roqueplo AP, Favier J, Rustin P, Mourad JJ, Plouin PF, Corvol P, Rotig A, Jeunemaitre X: The R22X mutation of the SDHD gene in hereditary paraganglioma abolishes the enzymatic activity of complex II in the mitochondrial respiratory chain and activates the hypoxia pathway. *Am J Hum Genet* 2001, **69**(6):1186–1197.
- Gimenez-Roqueplo AP, Favier J, Rustin P, Rieubland C, Kerlan V, Plouin PF, Rotig A, Jeunemaitre X: Functional consequences of a SDHB gene mutation in an apparently sporadic pheochromocytoma. *J Clin Endocrinol Metab* 2002, **87**(10):4771–4774.
- Dahia PL, Ross KN, Wright ME, Hayashida CY, Santagata S, Barontini M, Kung AL, Sanso G, Powers JF, Tischler AS: A HIF1alpha regulatory loop links hypoxia and mitochondrial signals in pheochromocytomas. *PLoS Genet* 2005, **1**(1):72–80.
- Pollard PJ, Briere JJ, Alam NA, Barwell J, Barclay E, Wortham NC, Hunt T, Mitchell M, Olpin S, Moat SJ: Accumulation of Krebs cycle intermediates and over-expression of HIF1alpha in tumours which result from germline FH and SDH mutations. *Hum Mol Genet* 2005, **14**(15):2231–2239.
- Allen M, Louise Jones J: Jekyll and Hyde: the role of the microenvironment on the progression of cancer. *J Pathol* 2011, **223**(2):162–176.
- Semenza GL: Defining the role of hypoxia-inducible factor 1 in cancer biology and therapeutics. *Oncogene* 2010, **29**(5):625–634.
- Chen CL, Chu JS, Su WC, Huang SC, Lee WY: Hypoxia and metabolic phenotypes during breast carcinogenesis: expression of HIF-1alpha, GLUT1, and CAIX. *Virchows Arch* 2010, **457**(1):53–61.
- Zhu P, Ning Y, Yao L, Chen M, Xu C: The proliferation, apoptosis, invasion of endothelial-like epithelial ovarian cancer cells induced by hypoxia. *J Exp Clin Cancer Res* 2010, **29**:124.
- Ji F, Wang Y, Qiu L, Li S, Zhu J, Liang Z, Wan Y, Di W: Hypoxia inducible factor 1alpha-mediated LOX expression correlates with migration and invasion in epithelial ovarian cancer. *Int J Oncol* 2013, **42**(5):1578–1588.
- Yeh YM, Chuang CM, Chao KC, Wang LH: MicroRNA-138 suppresses ovarian cancer cell invasion and metastasis by targeting SOX4 and HIF-1alpha. *Int J Cancer* 2013, **133**(4):867–878.
- Shackelford DB, Shaw RJ: The LKB1-AMPK pathway: metabolism and growth control in tumour suppression. *Nat Rev Cancer* 2009, **9**(8):563–575.
- Faubert B, Boily G, Izreig S, Griss T, Samborska B, Dong Z, Dupuy F, Chambers C, Fuerth BJ, Viollet B: AMPK is a negative regulator of the Warburg effect and suppresses tumor growth in vivo. *Cell Metab* 2013, **17**(1):113–124.
- Cervera AM, Apostolova N, Crespo FL, Mata M, McCreath KJ: Cells silenced for SDHB expression display characteristic features of the tumor phenotype. *Cancer Res* 2008, **68**(11):4058–4067.
- Fujii N, Jessen N, Goodyear LJ: AMP-activated protein kinase and the regulation of glucose transport. *Am J Physiol Endocrinol Metab* 2006, **291**(5):E867–877.
- Hardie DG: The AMP-activated protein kinase pathway—new players upstream and downstream. *J Cell Sci* 2004, **117**(Pt 23):5479–5487.
- Shaw RJ: Glucose metabolism and cancer. *Curr Opin Cell Biol* 2006, **18**(6):598–608.
- Li J, Miller EJ, Ninomiya-Tsuji J, Russell RR 3rd, Young LH: AMP-activated protein kinase activates p38 mitogen-activated protein kinase by increasing recruitment of p38 MAPK to TAB1 in the ischemic heart. *Circ Res* 2005, **97**(9):872–879.
- Pelletier A, Joly E, Prentki M, Coderre L: Adenosine 5'-monophosphate-activated protein kinase and p38 mitogen-activated protein kinase participate in the stimulation of glucose uptake by dinitrophenol in adult cardiomyocytes. *Endocrinology* 2005, **146**(5):2285–2294.
- Fremin C, Meloche S: From basic research to clinical development of MEK1/2 inhibitors for cancer therapy. *J Hematol Oncol* 2010, **3**:8.
- Meloche S, Pouyssegur J: The ERK1/2 mitogen-activated protein kinase pathway as a master regulator of the G1- to S-phase transition. *Oncogene* 2007, **26**(22):3227–3239.
- Poulikakos PI, Solit DB: Resistance to MEK inhibitors: should we co-target upstream? *Sci Signal* 2011, **4**(166):e16.
- Schlaepfer DD, Mitra SK, Ilic D: Control of motile and invasive cell phenotypes by focal adhesion kinase. *Biochim Biophys Acta* 2004, **1692**(2–3):77–102.
- McLean GW, Carragher NO, Avizienyte E, Evans J, Brunton VG, Frame MC: The role of focal-adhesion kinase in cancer - a new therapeutic opportunity. *Nat Rev Cancer* 2005, **5**(7):505–515.
- Hoekstra R, Eskens FA, Verweij J: Matrix metalloproteinase inhibitors: current developments and future perspectives. *Oncologist* 2001, **6**(5):415–427.
- Lengyel E, Schmalfeldt B, Konik E, Spathe K, Harting K, Fenn A, Berger U, Fridman R, Schmitt M, Prechtel D: Expression of latent matrix metalloproteinase 9 (MMP-9) predicts survival in advanced ovarian cancer. *Gynecol Oncol* 2001, **82**(2):291–298.
- Hsu CC, Wang CH, Wu LC, Hsia CY, Chi CW, Yin PH, Chang CJ, Sung MT, Wei YH, Lu SH: Mitochondrial dysfunction represses HIF-1alpha protein synthesis through AMPK activation in human hepatoma HepG2 cells. *Biochim Biophys Acta* 2013, **1830**(10):4743–4751.
- McDonnell TJ, Deane N, Platt FM, Nunez G, Jaeger U, McKearn JP, Korsmeyer SJ: bcl-2-immunoglobulin transgenic mice demonstrate extended B cell survival and follicular lymphoproliferation. *Cell* 1989, **57**(1):79–88.
- Vaux DL, Cory S, Adams JM: Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature* 1988, **335**(6189):440–442.

40. Green DR: Apoptotic pathways: paper wraps stone blunts scissors. *Cell* 2000, **102**(1):1–4.
41. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH: Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biol* 2013, **14**:32.
42. Fidler IJ, Radinsky R: Genetic control of cancer metastasis. *J Natl Cancer Inst* 1990, **82**(3):166–168.
43. Tester AM, Ruangpanit N, Anderson RL, Thompson EW: MMP-9 secretion and MMP-2 activation distinguish invasive and metastatic sublines of a mouse mammary carcinoma system showing epithelial-mesenchymal transition traits. *Clin Exp Metastasis* 2000, **18**(7):553–560.
44. Luo M, Guan JL: Focal adhesion kinase: a prominent determinant in breast cancer initiation, progression and metastasis. *Cancer Lett* 2010, **289**(2):127–139.
45. Cassavaugh J, Lounsbury KM: Hypoxia-mediated biological control. *J Cell Biochem* 2011, **112**(3):735–744.
46. Vander Heiden MG, Cantley LC, Thompson CB: Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 2009, **324**(5930):1029–1033.
47. Shin JM, Jeong YJ, Cho HJ, Park KK, Chung IK, Lee IK, Kwak JY, Chang HW, Kim CH, Moon SK: Melittin suppresses HIF-1alpha/VEGF expression through inhibition of ERK and mTOR/p70S6K pathway in human cervical carcinoma cells. *PLoS One* 2013, **8**(7):e69380.
48. He G, Jiang Y, Zhang B, Wu G: The effect of HIF-1alpha on glucose metabolism, growth and apoptosis of pancreatic cancerous cells. *Asia Pac J Clin Nutr* 2014, **23**(1):174–180.
49. Hanna SC, Krishnan B, Bailey ST, Moschos SJ, Kuan PF, Shimamura T, Osborne LD, Siegel MB, Duncan LM, O'Brien ET 3rd: HIF1alpha and HIF2alpha independently activate SRC to promote melanoma metastases. *J Clin Invest* 2013, **123**(5):2078–2093.
50. Karna E, Szoka L, Palka J: Thrombin-dependent modulation of beta1-integrin-mediated signaling up-regulates prolidase and HIF-1alpha through p-FAK in colorectal cancer cells. *Mol Cell Biochem* 2012, **361**(1–2):235–241.

doi:10.1186/s13048-014-0115-1

Cite this article as: Chen *et al.*: Succinate dehydrogenase subunit B inhibits the AMPK-HIF-1 $\alpha$  pathway in human ovarian cancer *in vitro*. *Journal of Ovarian Research* 2014 **7**:115.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

