

RESEARCH ARTICLE

Open Access

NAT1 and NAT2 genetic polymorphisms and environmental exposure as risk factors for oesophageal squamous cell carcinoma: a case-control study

Marco Matejic^{1,2}, Matjaz Vogelsang^{1,2}, Yabing Wang^{1,2} and Iqbal M Parker^{1,2*}

Abstract

Background: Tobacco smoking and red meat consumption are some of the known risk factors associated with the development of oesophageal cancer. N-acetyltransferases (*NAT1* and *NAT2*) play a key role in metabolism of carcinogenic arylamines present in tobacco smoke and overcooked red meat. We hypothesized that *NAT1* and *NAT2* genetic polymorphisms may influence the risk of oesophageal cancer upon exposure to environmental carcinogens.

Methods: Single nucleotide polymorphisms (SNPs) in the *NAT1* and *NAT2* genes were investigated by genotyping 732 cases and 768 healthy individuals from two South African populations to deduce the acetylator phenotype (slow, intermediate or rapid) from the combination of the genotyped SNPs.

Results: The 341 CC genotype (rs1801280) was significantly associated with a reduced risk for oesophageal cancer in the Mixed Ancestry population (OR = 0.31; 95% CI 0.11-0.87). The *NAT2* slow/intermediate acetylator status significantly increased the risk among cigarette smokers in the Black population (OR = 2.76; 95% CI 1.69-4.52), as well as among alcohol drinkers in the Mixed Ancestry population (OR = 2.77; 95% CI 1.38-5.58). Similarly, the *NAT1* slow/intermediate acetylator status was a risk factor for tobacco smokers in the Black population (OR = 3.41; 95% CI 1.95-5.96) and for alcohol drinkers in the Mixed Ancestry population (OR = 3.41; 95% CI 1.70-6.81). In a case-only analysis, frequent red meat consumption was associated with a significantly increased cancer risk for *NAT2* slow/intermediate acetylators in the Mixed Ancestry population (OR = 3.55; 95% CI 1.29-9.82; $P = 0.019$), whereas daily white meat intake was associated with an increased risk among *NAT1* slow/intermediate acetylators in the Black population (OR = 1.82; 95% CI 1.09-3.04; $P = 0.023$).

Conclusions: Our findings indicate that N-acetylation polymorphisms may modify the association between environmental risk factors and oesophageal cancer risk and that N-acetyltransferases may play a key role in detoxification of carcinogens. Prevention strategies in lifestyle and dietary habits may reduce the incidence of oesophageal cancer in high-risk populations.

Keywords: N-acetyltransferases, Single nucleotide polymorphism, Acetylator phenotype, Environmental carcinogens, Oesophageal cancer

* Correspondence: Iqbal.Parker@icgeb.org

¹International Centre for Genetic Engineering and Biotechnology, Cape Town Component, Observatory, UCT Medical Campus, Anzio Road, Observatory 7925, Cape Town, South Africa

²Division of Medical Biochemistry and IDM, UCT Faculty of Health Sciences, Cape Town, South Africa

Background

Oesophageal squamous cell carcinoma (OSCC) is the most common histological subtype of oesophageal cancer in developing countries and the main cause of cancer death in Southern Africa, with incidence and mortality rates approximately 7-fold higher among men and 4-fold higher among women than those in Northern Africa [1]. The prevalence of oesophageal cancer varies widely between different regions of South Africa, with exceptionally high-incidence rates recorded in the Black African populations of KwaZulu-Natal and the Eastern Cape [2]. The wide geographical variation in incidence of OSCC is most likely due to both genetic and environmental factors [3]. A number of epidemiological studies have pointed to tobacco smoking and alcohol consumption as major environmental risk factors for OSCC [4-6], and recent studies have also reported a significant association between red meat intake and an increased risk of oesophageal cancer [7,8].

Heterocyclic amines (HCAs) and nitrosamines are among the major arylamine carcinogens present in tobacco smoke and overcooked meat [9,10]. In humans these compounds are metabolized by the Phase I and Phase II xenobiotic metabolizing enzymes. The N-acetyltransferases (NATs) are the main phase II xenobiotic metabolizing enzymes involved in either detoxification or activation of arylamines, depending on the nature of the substrate and the organ involved [11]. The human NATs are encoded by two genes, *NAT1* [HGNC:7645; Ensembl:ENSG00000171428] and *NAT2* [HGNC:7646; Ensembl:ENSG00000156006], separated by 170 kb on chromosome 8p21.3 – 23.1 [12].

Single nucleotide polymorphisms (SNPs) in the *NAT1* and *NAT2* genes alter the enzyme activity and are classified into rapid, intermediate or slow acetylators [13]. The relationship between *NAT2* polymorphisms and *NAT2* acetylation activity has been widely investigated and a strong genotype-phenotype correlation exists [14]. The common SNPs in the coding region of the *NAT2* gene, 191G>A, 341T>C, 590G>A, and 857G>A (referred to as *NAT2*14*, *NAT2*5*, *NAT2*6*, and *NAT2*7* alleles respectively) lead to a significant decrease in acetylation activity compared to the wild-type allele (*NAT2*4*), and are thus designated as slow acetylator alleles [15]. A recent study reported that genotyping of these SNPs is sufficient to predict the *NAT2* acetylator status with high accuracy [16]. The *NAT2*5*, *NAT2*6*, and *NAT2*7*, and *NAT2*14* alleles were observed at frequencies of 20%, 15%, 1.7%, and 2.5% in the South African Black population, respectively [17]. The relationship between *NAT1* polymorphisms and *NAT1* acetylation activity, on the other hand, has been poorly investigated with inconsistent genotype-phenotype correlation for SNPs that lie outside the coding region [18]. The *NAT1* polymorphisms in Black South Africans are solely represented by *NAT1*4* (wild-type), *NAT1*10*, and *NAT1*3* alleles, with frequencies of 49%, 50%, and 1%,

respectively [19]. The *NAT1*10* allele, inferred by the simultaneous presence of two SNPs (1088T>A and 1095C>A) within the 3'-UTR of the *NAT1* gene, leads to an alteration of the consensus polyadenylation signal and is associated with a significantly increased acetylation activity compared to the wild-type allele [20]. The *NAT1*3* allele is inferred by the sole presence of 1095C>A, and does not result in a significantly elevated acetylation activity compared to the wild-type allele [18].

The *NAT1* and *NAT2* polymorphic variants may influence the metabolism of arylamine carcinogens and modulate the individual susceptibility to cancer upon exposure to environmental risk factors [21]. Genetic polymorphisms in the *NAT1* and *NAT2* loci have been associated with susceptibility to bladder, colon and rectal, breast, head and neck, lung and prostate cancers (reviewed in [22]). However, more consistent results were observed when the polymorphisms were investigated in association with environmental exposure [23]. Previous studies have reported conflicting results with regard to the role of *NAT1* and *NAT2* polymorphisms in oesophageal carcinogenesis [24-29].

We believe that genetic polymorphisms associated with low N-acetylation activity may lead to an impaired deactivation of arylamine carcinogens making subjects with slow or intermediate acetylator status more susceptible to environmental or dietary carcinogens than those with the rapid acetylator status. This study investigated the role of *NAT1* and *NAT2* functional polymorphisms as independent risk factors for OSCC and as modifiers of the association between environmental risk factors and OSCC in two indigenous populations of South Africa.

Methods

Study group

The study group included 1500 South African subjects subdivided into 463 OSCC patients and 480 control individuals from the Black population, and 269 OSCC patients and 288 control individuals from the Mixed Ancestry population. The Black individuals were mostly Xhosa speakers residing in the Western Cape Province of South Africa. The Mixed Ancestry individuals in the Western Cape is an admixed population that originated about 300 years ago from the union of different ethnic groups, receiving ancestral contribution from the indigenous Khoisan, sub-Saharan Africans, Europeans, Indonesians and Malaysians [30].

All patients selected for the study were histologically diagnosed with squamous cell carcinoma of the oesophagus, recruited between 2000 and 2012 from Groote Schuur and Tygerberg Hospitals in Cape Town. The control group included healthy volunteers with no history of any cancer and no familial history of oesophageal cancer. They were frequency matched to cases for geographical location, ethnicity, sex, and age. Because oesophageal cancer is

a disease of middle-age or the elderly, the minimum age of control individuals set for the study was 50. However, the control group had a higher proportion of individuals below the age of 60, while the patients were mostly over the age of 60. As a result, the mean age of the patients was approximately three years higher than that in control individuals in both the Black and Mixed Ancestry population groups (Table 1).

Ethics approval was obtained from the joint University of Cape Town/Groote Schuur Hospital Research Ethics Committee and the University of Stellenbosch/Tygerberg Hospital Ethics Committee. Written informed consent was obtained from all study participants.

Definition of variables

The study participants completed a standard questionnaire to collect demographic (i.e. ethnicity, origin, language, age, gender), lifestyle (i.e. smoking history, alcohol consumption), and dietary (i.e. frequency and type of meat intake) information.

The smoking habits were reported as smoking of cigarettes, hand-rolled cigarettes, and pipes (all of which referred to as 'cigarettes' in the text). Smoking is a qualitative variable and subjects were classified as current smokers (currently smoking at recruitment or stopped smoking not earlier than one year prior to recruitment), former smokers (smoked regularly at some point in their lives for at least six months or quit smoking at least one year prior to recruitment), or never smokers (never smoked). Current and former smokers were also asked to report the number of cigarettes smoked per day and subjects were grouped into those who smoked <10 cigarettes/day and those who smoked ≥10 cigarettes/day.

Alcohol consumption status was defined according to the frequency of alcohol (wine or beer) consumed per week and subjects were subdivided into those who consumed more than one litre of wine/beer per week and those who consumed less than one litre of wine/beer per week.

The study participants were also asked to report their average intake of red and white meat, fish, vegetables, and fried food and grouped into those who consumed each food category less than three times per week and those who consumed them (almost) daily. Red meat included beef, veal, pork, lamb, and offal. White meat included chicken, turkey, and duck. Data on food consumption was not available for control individuals, and only 299 Black patients and 158 Mixed Ancestry patients (out of 463 and 269 total patients, respectively) provided information about meat, fish, vegetables, and fried food intake.

Isolation and purification of DNA

Peripheral blood samples were collected from each subject enrolled in the study and stored at -20°C prior to DNA extraction. Genomic DNA was extracted using a

standard protocol [31]. All DNA samples were diluted to a final concentration of 20 ng/μl in 96-well plates and stored at -20°C until genotyping.

SNP selection and genotyping

Genotyping was determined blinded as to case-control status and environmental exposure of study participants.

Four *NAT2* SNPs; 191G>A (rs1801279), 341T>C (rs1801280), 590G>A (rs1799930), and 857G>A (rs1799931), were genotyped by TaqMan allele discrimination assay. Each reaction was carried out in a 2.5 μl volume containing 1X Universal PCR Master Mix, 1X Drug Metabolism Genotyping Assay Mix (Applied Biosystems), and 20 ng of DNA. The thermal cycling conditions consisted of an initial denaturation step at 95°C for 10 min, followed by 50 cycles of denaturation at 92°C for 15 sec and annealing/extension at 60°C for 90 sec. The amplification reaction and fluorescence measurement was performed in a Roche LightCycler 480 II, and genotypes were assigned using SP4 1.5.0 software (Roche). Five *NAT2* alleles were identified: *NAT2*4* (wild-type), *NAT2*5*, *NAT2*6*, *NAT2*7* and *NAT2*14* (slow acetylator alleles). The acetylator phenotype was determined as follows: (i) subjects with two *NAT2*4* alleles were classified as "rapid" acetylators; (ii) subjects with any combination of two slow acetylator alleles were "slow" acetylators; (iii) subjects with one wild-type allele and one slow acetylator allele were "intermediate" acetylators. Individuals carrying alleles deducing either slow or intermediate acetylator status were merged into one group, referred to as *NAT2* "slow/intermediate" acetylators, and compared to rapid acetylators (*NAT2*4/*4*).

Two *NAT1* SNPs; 1095C>A (rs15561) and 1088T>A (rs1057126), were determined by TaqMan assay as previously described [32]. We used a three-probe system to genotype the two variants simultaneously in a single reaction. This was necessary because the 1088/1095 SNPs are too close to each other to use the conventional two-probe TaqMan assay. Primers and hybridization probes were designed according to the recommendations by Roche Diagnostics Inc. [see Additional file 1]. 1X Universal PCR Master Mix (Applied Biosystems) was used as PCR solution and primers, probes, and DNA were added to final concentrations of 300 nM, 100 nM, and 8 ng/μl, respectively. The amplification reaction was carried out in a Roche LightCycler 480 II instrument with an initial hold step at 95°C for 10 min followed by 40 cycles of a two-step reaction (92°C for 15 s, 60°C for 1 min). Three *NAT1* alleles were identified: *NAT1*4* (wild-type), *NAT1*3* and *NAT1*10* (rapid acetylator allele). Individuals with both 1095C>A and 1088T>A were referred to as *NAT1*10* carriers, and those with only 1095C>A were *NAT1*3* carriers. Subjects were classified as slow, intermediate, or rapid acetylators if they had zero, one, or two copies

Table 1 Characteristics of the population study and effect of tobacco smoking, alcohol consumption, and red meat intake on OSCC risk

Demographic/risk factors	Variables	Black				Mixed Ancestry			
		Controls (%)	Cases (%)	OR (95% CI) ^a	P-value	Controls (%)	Cases (%)	OR (95% CI) ^a	P-value
Gender	Males	235 (49)	229 (49)			178 (62)	177 (66)		
	Females	245 (51)	234 (51)			110 (38)	92 (34)		
Age	Mean (SD) ^b	56.7 (15.0)	59.6 (10.7)			57.7 (14.3)	60.7 (10.3)		
Tobacco smoking	Never smokers	258 (54)	181 (39)	1 (Ref)		62 (22)	15 (6)	1 (Ref)	
	Smokers ^c	222 (46)	280 (60)	1.79 (1.39-2.33)	<0.0001	226 (78)	250 (93)	4.57 (2.53-8.27)	<0.0001
	Unknown	0 (0)	2 (0.4)			0 (0)	4 (1)		
Total cigarettes/day	0	258 (54)	181 (39)	1 (Ref)		62 (21)	15 (6)	1 (Ref)	
	<10	137 (28)	179 (39)	1.86 (1.39-2.49)	<0.0001	118 (41)	85 (31)	2.98 (1.59-5.59)	0.001
	≥10	73 (15)	95 (20)	1.85 (1.29-2.66)	0.001	90 (31)	160 (59)	7.35 (3.95-13.6)	<0.0001
	Unknown	12 (2)	8 (2)			18 (6)	9 (3)		
Alcohol drinking	Non-drinkers	201 (42)	175 (38)	1 (Ref)		115 (40)	51 (19)	1 (Ref)	
	Drinkers ^d	278 (58)	286 (62)	1.18 (0.91-1.53)	0.211	172 (60)	215 (80)	2.82 (1.92-4.15)	<0.0001
	Unknown	1 (0.2)	2 (0.4)			1 (0.3)	3 (1)		
Tobacco smoking + alcohol drinking	Never smokers ^e	257 (53)	180 (39)	1 (Ref)		62 (21)	15 (6)	1 (Ref)	
	Smokers/non-drinkers	41 (8)	33 (7)	1.15 (0.70-1.89)	0.583	75 (26)	37 (14)	2.04 (1.02-4.06)	0.042
	Smokers/drinkers	181 (38)	247 (53)	1.95 (1.49-2.55)	<0.0001	150 (52)	213 (79)	5.87 (3.22-10.7)	<0.0001
	Unknown	1 (0.2)	3 (0.6)			1 (0.3)	4 (1)		
Red meat intake	3 times/week or less	–	261 (56)			–	108 (40)		
	Daily or almost daily	–	37 (8)			–	50 (19)		
	Unknown	–	165 (36)			–	111 (41)		
White meat intake	3 times/week or less	–	151 (33)			–	75 (28)		
	Daily or almost daily	–	148 (32)			–	82 (30)		
	Unknown	–	164 (35)			–	112 (42)		
Fish	3 times/week or less	–	289 (62)			–	146 (54)		
	Daily or almost daily	–	9 (2)			–	11 (4)		
	Unknown	–	165 (36)			–	112 (42)		
Vegetables	3 times/week or less	–	243 (52)			–	54 (20)		
	Daily or almost daily	–	53 (11)			–	101 (37)		
	Unknown	–	165 (36)			–	114 (42)		

Table 1 Characteristics of the population study and effect of tobacco smoking, alcohol consumption, and red meat intake on OSCC risk (Continued)

Fried food	3 times/week or less	–	291 (63)	–	135 (50)
	Daily or almost daily	–	5 (1)	–	20 (7)
	Unknown	–	167 (36)	–	114 (42)

^aCrude odds ratio was calculated.

^bSD = standard deviation of the mean.

^cSmokers = current and former smokers.

^dDrinkers = light to heavy alcohol drinkers.

^eNever smokers (including drinkers and non-drinkers).

Ref = reference allele.

of the *NAT1*10* allele, respectively. Individuals with either zero or one copy of *NAT1*10* allele were merged into one group, referred to as *NAT1* “slow/intermediate” acetylators, and compared to rapid acetylators (*NAT1*10/*10*).

Statistical analysis

Genotype counts among controls were tested for deviation from Hardy-Weinberg Equilibrium using the Chi-Square test with one degree of freedom [33].

The genetic association analyses were performed assuming a multiplicative genetic model for any risk allele or risk category. Where sample sizes were sufficiently large, we tested for differences in frequency distribution between cases and control individuals using Pearson's Chi-Square (χ^2) test of association. Crude odds ratio (OR), adjusted odds ratio, 95% confidence interval (95% CI), and *P*-value for association were estimated for the tested allele or category. The adjustment of the OR for potential confounders such as age, gender, tobacco smoking, and alcohol consumption was computed by using multivariate conditional logistic analysis [the disease status (case/control) was checked against each polymorphism (GG/GA/AA) for all the cases and controls used in the study by considering also other variables such as smoking, alcohol consumption, age and sex]. All reported *P*-values are two-sided and a *P*-value <0.05 was considered as significant. A conservative *P*-value threshold of 0.0042 (=0.05/[2x6]) was used to assess the significance of association applying Bonferroni correction for multiple testing of the six genotyped SNPs in the two populations.

Case-only analyses were performed to evaluate the interaction between the acetylation polymorphism and meat consumption by testing for frequency of red and white meat intake (<3 times per week/almost daily).

All statistical analyses were performed using SPSS 19 software package (SPSS, Chicago, IL). Linkage disequilibrium and haplotype analyses were performed on the control groups using Haploview [34]. The pairwise linkage disequilibrium was measured by variables D^I and r^2 . Pearson's χ^2 was used to test for differences in haplotype distribution between patients and control individuals using UNPHASED [35].

Results

Demographic details and environmental exposure

The distribution of demographic details and lifestyle factors among cases and control individuals enrolled in the study is shown in Table 1. The percentages of tobacco smokers, alcohol drinkers, and of subjects that consumed red meat, fried food, and vegetables daily were higher in the Mixed Ancestry group than in the Black group, while

no significant difference was observed for daily consumption of white meat and fish.

Tobacco smoking was associated with a significantly increased risk of OSCC among both the Black (OR = 1.79; 95% CI 1.39-2.33; *P* < 0.0001) and Mixed Ancestry (OR = 4.57; 95% CI 2.53-8.27; *P* < 0.0001) individuals. In the Mixed Ancestry group, the risk associated with tobacco smoking increased with the number of cigarettes smoked per day, with a 2.98-fold higher risk (95% CI 1.59-5.59; *P* = 0.001) among those smoking <10 cigarettes/day, and a 7.35-fold higher risk (95% CI 3.95-13.66; *P* < 0.0001) among those smoking ≥10 cigarettes/day, compared to never smokers. No cumulative effect of tobacco smoking on OSCC risk was observed in the Black group.

A higher proportion of alcohol drinkers was observed among patients compared to control individuals in both ethnic groups, although the difference was significant only in the Mixed Ancestry group (OR = 2.82; 95% CI 1.92-4.15; *P* < 0.0001). The analysis of the combined effect of tobacco smoking and alcohol consumption showed a significantly higher OSCC risk for smokers who drank alcohol (‘smokers/drinkers’) compared to ‘never smokers’ (including drinkers and non-drinkers) among both Black (OR = 1.95; 95% CI 1.49-2.55; *P* < 0.0001) and Mixed Ancestry (OR = 5.87; 95% CI 3.22-10.71; *P* < 0.0001) individuals. No increased risk was observed for ‘smokers/non-drinkers’ in the Black group (*P* = 0.583), while only a weak association was detected in the Mixed Ancestry group (OR = 2.04; 95% CI 1.02-4.06; *P* = 0.042).

Linkage disequilibrium analysis

The results of the linkage disequilibrium analysis (LD) and Hardy-Weinberg equilibrium tests in the two South African populations showed that the distribution of genotypes for each SNP in the control groups were in Hardy-Weinberg equilibrium (*P* > 0.05; see Additional file 2). The *NAT1* 1088T>A and 1095C>A SNPs occurred at high frequency (MAF ≥ 0.401) and were in complete LD in both ethnic groups (D^I = 1.0). For the *NAT2* SNPs, only 341T>C (*NAT2*5*) and 590G>A (*NAT2*6*) were present at relatively high frequencies (MAF ≥ 0.214) and strong LD (D^I = 1.0) existed between these two polymorphisms in both population groups. Linkage disequilibrium of 857G>A (*NAT2*7*) and 191G>A (*NAT2*14*) with other SNPs was not considered because of the low frequency of these two polymorphic variants in both population groups (MAF ≤ 0.063). No significant LD was observed between SNPs in the *NAT1* locus and the two most common SNPs in the *NAT2* locus (341T>C and 590G>A) in either ethnic group, thereby forming two main LD blocks, Block 1 (including *NAT1* SNPs) and Block 2 (including *NAT2* SNPs), as shown in Figure 1.




Figure 1 Linkage disequilibrium between *NAT1* and *NAT2* SNPs in Black and Mixed Ancestry South Africans. Plots show r^2 and D' values for pairwise LD between *NAT1* and *NAT2* variants in the South African Black and Mixed Ancestry population groups. The left triangle in each plot represents *NAT1* SNPs, while the right triangle represents *NAT2* SNPs. Colour intensity of squares (black to white) indicates the strength of LD (high to low) by D' ; while numbers within squares refer to r^2 values for pairwise SNPs. D' and r^2 refer to different statistical methods to measure linkage disequilibrium between alleles; r^2 is preferred to predict one allele given the other, whereas D' is mainly used to assess recombination patterns such as haplotype blocks. Linkage disequilibrium analysis was carried out in control groups by using Haploview [34].

***NAT1* and *NAT2* polymorphisms, environmental exposure, and OSCC risk**

A significant association was detected among carriers of the 341 CC variant genotype (*NAT2**5/*5) compared to carriers of the wild-type (341 TT) genotype (OR = 0.31; 95% CI 0.11-0.87; $P = 0.026$) in the Mixed Ancestry population, with a borderline association in the Black population (OR = 0.55; 95% CI 0.30-0.99; $P = 0.05$) (Table 2). The *NAT2* 341 C allele (*NAT2**5) was associated with a reduced risk for OSCC in the Mixed Ancestry group (OR = 0.57; 95% CI 0.38-0.87; $P = 0.010$), while no significant association was found in the Black group (OR = 0.82; 95% CI 0.64-1.06; $P = 0.132$; see Additional file 3). However, none of the associations observed in the Mixed Ancestry population had significance below the Bonferroni corrected P -value of 0.0042.

The distribution of *NAT2* acetylator phenotypes (rapid, intermediate, and slow acetylators) between cases and control individuals was significantly different in the Mixed Ancestry group, while no significant difference was observed in the Black group, even when intermediate and slow acetylators were merged into one group and compared with the rapid acetylators (Table 3). For the Mixed Ancestry individuals, the *NAT2* slow/intermediate acetylators were at reduced risk of OSCC compared with rapid acetylators (OR = 0.44; 95% CI 0.27-0.70; $P = 0.001$), while the *NAT1* slow/intermediate acetylator status was not a risk factor for OSCC (OR = 0.85; 95% CI 0.43-1.68; $P = 0.638$). For the Black individuals, the ORs were 0.78 (95% CI 0.55-1.11; $P = 0.166$) for *NAT1* slow/intermediate acetylators and 0.82 (95% CI 0.54-1.23; $P = 0.333$) for

NAT2 slow/intermediate acetylators compared to their respective rapid acetylator groups.

The effect of *NAT1* and *NAT2* acetylator phenotypes on the association of tobacco smoking and alcohol consumption with OSCC risk was investigated by case-control analysis (Tables 4 and 5).

In the Black group, smoking was associated with a significantly increased risk of OSCC for the *NAT2* slow/intermediate acetylators, with an OR of 2.76 (95% CI 1.69-4.52; $P < 0.0001$) among smokers compared to never smokers, while tobacco smoking was not a risk factor for the *NAT2* rapid acetylators (OR = 2.58; 95% CI 0.80-8.25; $P = 0.111$; Table 4). Similarly, the smoking-associated risk for OSCC was significantly higher for *NAT1* slow/intermediate acetylators (OR = 3.41; 95% CI 1.95-5.96; $P < 0.0001$), whereas no significant association was observed for *NAT1* rapid acetylators (OR = 1.71; 95% CI 0.73-4.04; $P = 0.218$; Table 5). However, heterogeneity testing for *NAT2* slow/intermediate acetylators vs rapid acetylators ($I^2 = 0\%$; $P = 0.917$) and for *NAT1* slow/intermediate acetylators vs rapid acetylators ($I^2 = 42.9\%$; $P = 0.185$) indicated that smoking might increase the risk for OSCC in Black subjects regardless of their acetylator status. The effect of *NAT1* and *NAT2* polymorphisms on the smoking-associated risk of OSCC was not evaluated in the Mixed Ancestry group because of the extremely low number of non-smokers in this group.

The alcohol-related risk of OSCC in the Mixed Ancestry population was significantly increased for *NAT1* slow/intermediate acetylators (OR = 3.41; 95% CI 1.70-6.81; $P = 0.001$) and *NAT2* slow/intermediate acetylators

Table 2 Genotype frequency of NAT2 and NAT1 polymorphisms and association with OSCC in Black and Mixed Ancestry South Africans

Gene	SNP	Polymorphism	Black				Mixed Ancestry			
			Controls (%)	Cases (%)	OR (95% CI) ^a	P-value	Controls (%)	Cases (%)	OR (95% CI) ^a	P-value
NAT2	590G>A	G / G	296 (62.3)	257 (57.2)	1 (Ref)	-	172 (59.7)	159 (60.0)	1 (Ref)	-
	(NAT2*6)	G / A	155 (32.6)	163 (36.3)	1.20 (0.85 - 1.69)	0.299	105 (36.4)	93 (35.1)	0.69 (0.39 - 1.21)	0.192
	rs1799930	A / A	24 (5.0)	29 (6.4)	1.46 (0.72 - 2.94)	0.291	11 (3.8)	13 (4.9)	1.30 (0.34 - 4.91)	0.697
	341T>C	T / T	247 (52.1)	237 (53.0)	1 (Ref)	-	130 (45.1)	147 (55.7)	1 (Ref)	-
	(NAT2*5)	T / C	179 (37.8)	178 (39.8)	0.97 (0.69 - 1.36)	0.856	125 (43.4)	101 (38.2)	0.59 (0.34 - 1.04)	0.069
	rs1801280	C / C	48 (10.1)	32 (7.1)	0.55 (0.30 - 0.99)	0.05	33 (11.5)	16 (6.1)	0.31 (0.11 - 0.87)	0.026
	857G>A	G / G	471 (98.9)	440 (98.2)	1 (Ref)	-	265 (92.0)	237 (90.5)	1 (Ref)	-
	(NAT2*7)	G / A	5 (1.0)	7 (1.6)	0.95 (0.19 - 4.77)	0.948	23 (8.0)	24 (9.2)	1.32 (0.54 - 3.23)	0.548
	rs1799931	A / A	0 (0)	1 (0.2)	-	-	0 (0)	1 (0.4)	-	-
	191G>A	G / G	417 (87.6)	402 (90.1)	1 (Ref)	-	275 (95.5)	244 (92.8)	1 (Ref)	-
	(NAT2*14)	G / A	58 (12.2)	41 (9.2)	0.78 (0.46 - 1.33)	0.364	13 (4.5)	18 (6.8)	1.46 (0.48 - 4.49)	0.508
	rs1801279	A / A	1 (0.2)	3 (0.7)	-	-	0 (0)	1 (0.4)	-	-
NAT1	1088T>A	T / T	89 (19.2)	100 (22.8)	1 (Ref)	-	106 (37.3)	81 (31.4)	1 (Ref)	-
	(NAT1*10)	T / A	219 (47.3)	196 (44.7)	0.77 (0.50 - 1.18)	0.226	128 (45.1)	130 (50.4)	1.64 (0.90 - 2.98)	0.103
	rs1057126	A / A	155 (33.5)	142 (32.4)	1.07 (0.67 - 1.70)	0.781	50 (17.6)	47 (18.2)	1.57 (0.73 - 3.40)	0.249
	1095C>A	C / C	86 (18.6)	95 (21.7)	1 (Ref)	-	97 (34.1)	69 (26.7)	1 (Ref)	-
	(NAT1*10, NAT1*3)	C / A	220 (47.5)	198 (45.2)	0.79 (0.51 - 1.23)	0.299	130 (45.8)	138 (53.5)	1.83 (0.99 - 3.39)	0.054
	rs15561	A / A	157 (33.9)	145 (33.1)	1.10 (0.69 - 1.75)	0.681	57 (20.1)	51 (19.8)	1.78 (0.82 - 3.84)	0.143

Samples with unknown genotype are not shown.

^aOdds ratio was adjusted for age, gender, tobacco smoking, and alcohol consumption status; odds ratio was not calculated for subgroups with low sample size.

*Indicates the allele for each of the genes.

Table 3 Distribution of NAT1 and NAT2 acetylation polymorphisms and association with OSCC in the Black and Mixed Ancestry populations

Gene	Acetylator status	Black				Mixed Ancestry			
		Controls (%)	Cases (%)	OR (95% CI) ^a	P-value	Controls (%)	Cases (%)	OR (95% CI) ^a	P-value
NAT2	Rapid	87 (18)	90 (19)	1 (Ref)		49 (17)	57 (21)	1 (Ref)	
	Intermediate	231 (48)	192 (41)	0.77 (0.50-1.20)	0.248	124 (43)	113 (42)	0.47 (0.28-0.80)	0.004
	Slow	155 (32)	161 (35)	0.88 (0.56-1.39)	0.588	115 (40)	91 (34)	0.40 (0.24-0.68)	0.001
	Unknown	7 (1)	20 (4)			0 (0)	8 (3)		
	Slow + intermediate	386 (82)	353 (79)	0.82 (0.54-1.23)	0.333	239 (83)	204 (78)	0.44 (0.27-0.70)	0.001
NAT1	Rapid	155 (32)	142 (31)	1 (Ref)		50 (17)	47 (17)	1 (Ref)	
	Intermediate	219 (46)	196 (42)	0.72 (0.49-1.05)	0.084	128 (44)	130 (48)	1.04 (0.50-2.17)	0.908
	Slow	89 (18)	100 (22)	0.94 (0.59-1.48)	0.781	106 (37)	81 (30)	0.64 (0.29-1.37)	0.249
	Unknown	17 (3)	25 (5)			4 (1)	11 (4)		
	Slow + intermediate	308 (66)	296 (68)	0.78 (0.55-1.11)	0.166	234 (82)	211 (81)	0.85 (0.43-1.68)	0.638

^aOdds ratio was adjusted for age, gender, tobacco smoking, and alcohol consumption status.

Ref = reference allele.

NAT2 rapid = NAT2*4/*4.

NAT2 intermediate = NAT2*4/*5, NAT2*4/*6, NAT2*4/*7, or NAT2*4/*14.

NAT2 slow = any combination of NAT2*5, NAT2*6, NAT2*7, and NAT2*14.

NAT1 rapid = NAT1*10/*10.

NAT1 intermediate = NAT1*10/*4 or NAT1*10/*3.

NAT1 slow = NAT1*4/*4, NAT1*4/*3, or NAT1*3/*3.

*Indicates the allele for each of the genes.

(OR = 2.77; 95% CI 1.38-5.58; $P = 0.004$), whereas no increased risk was observed for alcohol drinkers who were either NAT1 rapid acetylators (OR = 0.80; 95% CI 0.19-3.44; $P = 0.764$) or NAT2 rapid acetylators (OR = 2.43; 95% CI 0.65-9.15; $P = 0.188$). However, heterogeneity testing for NAT2 slow/intermediate acetylators vs rapid acetylators ($I^2 = 0\%$; $P = 0.864$) suggested that alcohol may be a significant risk factor for OSCC in both NAT2 acetylator groups, while substantial heterogeneity between the NAT1 acetylator phenotypes ($I^2 = 68.1\%$; $P = 0.077$) suggested that NAT1 acetylator status may modify the alcohol-associated risk of OSCC among Mixed Ancestry individuals. In the Black population, the acetylation polymorphism had no effect on the association between alcohol intake and OSCC risk (Tables 4 and 5).

A case-only analysis was performed to investigate the interaction of NAT1 and NAT2 acetylator phenotypes with meat consumption on susceptibility to OSCC. In the Mixed Ancestry patients, daily intake of red meat was significantly more frequent among NAT2 slow/intermediate acetylators than among NAT2 rapid acetylators (OR = 3.55; 95% CI 1.29-9.82; $P = 0.019$), whereas no significant difference was observed among Black patients (OR = 1.67; 95% CI 0.61-4.58; $P = 0.316$; Table 4). No interaction between NAT1 acetylator phenotypes and red meat consumption was observed among either Black patients (OR = 0.93; 95% CI 0.43-2.0; $P = 0.851$) or Mixed Ancestry patients (OR = 1.32; 95% CI 0.49-3.55; $P = 0.586$). We also observed a statistically significant interaction between daily consumption of white meat and the NAT1 slow acetylator status in Black

patients (OR = 1.82; 95% CI 1.09-3.04; $P = 0.023$), while such interaction was not significant in Mixed Ancestry patients (OR = 1.58; 95% CI 0.68-3.67; $P = 0.288$; Table 5).

Discussion

We determined the distribution of the NAT1*3, NAT1*4, NAT1*10, NAT2*4, NAT2*5, NAT2*6, NAT2*7, and NAT2*14 alleles in the Black and Mixed Ancestry populations of South Africa, accounting for the majority of the NAT1 and NAT2 alleles observed in Black South Africans and Caucasians [19].

The overall frequency of the NAT2 slow acetylator alleles in Black Xhosa and Mixed Ancestry healthy individuals (57.2% and 61.5%, respectively) was similar to that previously reported in the Black Tswana and Mixed Ancestry populations from South Africa (59% and 54%, respectively) [19,36]. The allele frequency of NAT1*10 observed in Black Xhosa (57.1%) was similar to that previously reported in Black Tswana (50.5%), whereas the frequency in Mixed Ancestry individuals (40.1%) was closer to that observed in Black South Africans than that of UK Caucasians (18.7%) [19].

The high LD observed between 341T>C (NAT2*5) and 590G>A (NAT2*6) in both South African populations has also been reported in a Northern Indian population [26]. Although studies reported a strong LD between NAT1*10 and NAT2*4 in Caucasians [37,38], we found no significant LD between NAT1 and NAT2 alleles in South Africans.

Table 4 Association of NAT2 acetylator phenotype with OSCC risk upon exposure to environmental risk factors in Black and Mixed Ancestry South Africans

Population	Risk factors	Variables	NAT2 rapid acetylators				NAT2 slow/intermediate acetylators			
			Controls	Cases	OR (95% CI) ^a	P-value	Controls	Cases	OR (95% CI) ^a	P-value
Black										
	Tobacco smoking	Never smokers	49	39	1 (Ref)	0.111	206	132	1 (Ref)	<0.0001
		Smokers ^b	38	51	2.58 (0.80-8.25)		180	220	2.76 (1.69-4.52)	
	Tot cigarettes/day	0	49	39	1 (Ref)	0.189	206	132	1 (Ref)	<0.0001
		≤9	23	31	2.23 (0.67-7.36)		112	143	3.27 (1.93-5.54)	
		≥10	13	19	3.79 (0.92-15.7)		0.066	58	72	
	Alcohol drinking	Non-drinkers	34	39	1 (Ref)	0.903	164	127	1 (Ref)	0.675
		Drinkers ^c	53	51	0.94 (0.34-2.62)		221	224	0.91 (0.59-1.41)	
	Red meat intake ^d	3 times/week or less	–	55	–	–	–	198	1 (Ref)	–
		Daily or almost daily	–	5	–	–	–	31	1.67 (0.61-4.58)	0.316
	White meat intake ^d	3 times/week or less	–	33	–	–	–	113	1 (Ref)	–
		Daily or almost daily	–	27	–	–	–	117	1.28 (0.72-2.29)	0.397
Mixed Ancestry										
	Tobacco smoking	Never smokers	8	2	1 (Ref)	–	54	12	1 (Ref)	–
		Smokers ^b	41	53	–		185	191	–	
	Tot cigarettes/day	0	8	2	1 (Ref)	–	54	12	1 (Ref)	–
		≤9	22	25	–		96	58	–	
		≥10	16	27	–		74	129	–	
	Alcohol drinking	Non-drinkers	15	9	1 (Ref)	0.188	100	41	1 (Ref)	0.004
		Drinkers ^c	34	47	2.43 (0.65-9.15)		138	162	2.77 (1.38-5.58)	
	Red meat intake ^d	3 times/week or less	–	31	–	–	–	75	1 (Ref)	–
		Daily or almost daily	–	5	–	–	–	43	3.55 (1.29-9.82)	0.019
	White meat intake ^d	3 times/week or less	–	20	–	–	–	51	1 (Ref)	–
		Daily or almost daily	–	16	–	–	–	66	1.80 (0.81-4.02)	0.149

Samples with unknown acetylator status are not shown.

Ref = reference allele.

^aOR was adjusted for age, gender, smoking and alcohol consumption status; OR was not calculated for subgroups with low sample size.

^bSmokers = current and former smokers.

^cDrinkers = light to heavy alcohol drinkers.

^dCase-only analysis has been performed for red meat intake due to lack of data from control individuals.

Table 5 Association of NAT1 acetylator phenotype with OSCC risk upon exposure to environmental risk factors in Black and Mixed Ancestry South Africans

Population	Risk factors	Variables	NAT1 rapid acetylators				NAT1 slow/intermediate acetylators			
			Controls	Cases	OR (95% CI) ^a	P-value	Controls	Cases	OR (95% CI) ^a	P-value
Black										
	Tobacco smoking	Never smokers	84	61	1 (Ref)	0.218	166	109	1 (Ref)	<0.0001
		Smokers ^b	71	81	1.71 (0.73-4.04)		142	186	3.41 (1.95-5.96)	
	Tot cigarettes/day	0	84	61	1 (Ref)	0.208	166	109	1 (Ref)	<0.0001
		≤9	45	50	1.80 (0.72-4.53)		87	123	3.96 (2.19-7.13)	
		≥10	24	31	2.06 (0.73-5.84)		45	57	2.97 (1.47-5.99)	
	Alcohol drinking	Non-drinkers	66	53	1 (Ref)	0.707	130	112	1 (Ref)	0.570
		Drinkers ^c	89	89	1.17 (0.52-2.59)		177	182	0.87 (0.54-1.40)	
Red meat intake ^d	3 times/week or less	–	81		–	168	1 (Ref)			
		Daily or almost daily	–	12		–	23	0.93 (0.43-2.0)	0.851	
	White meat intake ^d	3 times/week or less	–	56		–	87	1 (Ref)		
		Daily or almost daily	–	38		–	104	1.82 (1.09-3.04)	0.023	
Mixed Ancestry										
	Tobacco smoking	Never smokers	17	1	1 (Ref)	–	43	14	1 (Ref)	–
		Smokers ^b	33	46	–		191	194	–	
	Tot cigarettes/day	0	17	1	1 (Ref)	–	43	14	1 (Ref)	–
		≤9	17	24	–		101	58	–	
		≥10	11	20	–		77	133	–	
	Alcohol drinking	Non-drinkers	21	12	1 (Ref)	0.764	93	39	1 (Ref)	0.001
		Drinkers ^c	29	35	0.80 (0.19-3.44)		140	170	3.41 (1.70-6.81)	
Red meat intake ^d	3 times/week or less	–	–	22		–	81	1 (Ref)		
		Daily or almost daily	–	7		–	42	1.32 (0.49-3.55)	0.586	
	White meat intake ^d	3 times/week or less	–	16		–	56	1 (Ref)		
		Daily or almost daily	–	13		–	66	1.58 (0.68-3.67)	0.288	

Samples with unknown acetylator status are not shown.

Ref = reference allele.

^aOR was adjusted for age, gender, smoking and alcohol consumption status; OR was not calculated for subgroups with low sample size.

^bSmokers = current and former smokers.

^cDrinkers = light to heavy alcohol drinkers.

^dCase-only analysis has been performed for red meat intake due to lack of data from control individuals.

This study confirmed previous findings on the association between tobacco smoking and a high risk of OSCC in the Black and Mixed Ancestry populations from South Africa, as well as a cumulative risk associated with the number of cigarettes smoked per day in the Mixed Ancestry population [4]. Alcohol consumption was an independent risk factor only in the Mixed Ancestry population, but no association was observed in the Black Xhosa population, which is in agreement with the results from a previous study conducted in the same ethnic groups [4]. We also confirmed the previously reported combined effect of alcohol consumption and tobacco smoking on the enhanced oesophageal cancer risk in Black South Africans [39]. Furthermore, tobacco smoking, alcohol consumption, and frequent red meat intake were more prevalent in the Mixed Ancestry population than in the Black population, indicating that Mixed Ancestry South Africans may be more exposed to environmental carcinogens than Black South Africans.

The *NAT2* 341T>C (*NAT2**5) polymorphism and *NAT2* slow/intermediate acetylator status were associated with a significantly reduced risk of OSCC in the Mixed Ancestry population. Because *NAT2**5 is a slow acetylator allele and NATs are able to activate arylamine carcinogens to reactive intermediates [11,15], the *NAT2* slow acetylator status may result in an enzyme with impaired N-acetylation capacity, possibly leading to reduced production of reactive intermediates and ultimately preventing DNA damage and carcinogenesis. *NAT2**5 is the most frequent *NAT2* allele in the South African population, and has the greatest reduction in acetylation activity among *NAT2* slow acetylator alleles [40]. These findings may explain why the association with OSCC was observed among the *NAT2**5 allele carriers and not among subjects carrying other *NAT2* slow acetylator alleles. A recent study in a Chinese population found the CC genotype of the *NAT2* rs1565684 T > C polymorphism to be associated with a borderline significantly increased risk for OSCC (OR = 1.77; 95% CI 0.97–3.21; $P = 0.063$). Although this variant has been reported to be functional and in LD with another important *NAT2* polymorphism (rs4345600; $r^2 = 0.845$) in the Han Chinese, the etiology of *NAT2* rs1565684 on oesophageal carcinogenesis is still unknown [41]. In accordance with a previous study on oesophageal adenocarcinoma [29], we found no statistically significant association between *NAT1* polymorphisms (*NAT1**10 and *3) and OSCC risk in both South African populations.

When we evaluated the association between the *NAT1* and *NAT2* acetylator phenotypes and OSCC risk in conjunction with environmental exposure, the risk associated with tobacco smoking and alcohol intake appeared to be higher among slow/intermediate acetylators than among rapid acetylators, although this difference was not significant. The lack of a significant association between smoking

and alcohol intake with OSCC among rapid acetylators is probably due to the small sample size of these subgroups and the resulting low statistical power to detect significant associations. The interaction of the *NAT1* and *NAT2* polymorphisms with tobacco smoking or red meat intake is likely to be due to the role of NATs in the metabolism of HCAs and nitrosamines, which are the most relevant carcinogen in tobacco smoke and overcooked meat [9,10]. Our finding of an interaction between *NAT1* acetylator status and white meat intake in the Black population is in agreement with a study showing that heat-treated poultry products have significant amounts of HCAs based on the cooking methods and conditions [42]. Furthermore, a previous cohort study reported a significant positive association between poultry intake and oesophageal adenocarcinoma risk [43]. Slow and intermediate acetylators may be unable to detoxify carcinogenic metabolites efficiently; therefore, an impaired detoxification of HCA- and nitrosamine-derived metabolites may explain the greater risk of OSCC associated with low N-acetylation activity upon exposure to environmental carcinogens. However, no study to date has biologically demonstrated that low N-acetylation activity leads to an impaired deactivation of arylamine carcinogens in humans.

A recent cohort study identified statistically significant positive associations of both heme iron (HR = 1.67; 95% CI 1.05–2.68; $P = 0.022$) and processed red meat intake (HR = 2.27; 95% CI 1.33–3.89; $P = 0.004$) with oesophageal adenocarcinoma [44]. Heme iron is an organic form of iron and is mainly provided by red meat. It has been demonstrated that heme iron specifically contributes to carcinogenesis by increasing oxidative stress or by catalysing endogenous formation of nitroso-compounds [45,46]. Thus, the N-nitrosation induced by heme iron may increase the level of nitrosamines in humans and ultimately contribute to oesophageal carcinogenesis in slow/intermediate acetylators [44].

We observed that *NAT1* and *NAT2* acetylator phenotypes modify the alcohol-associated risk of OSCC in the Mixed Ancestry population. Ethanol from alcoholic beverages can cause local irritation of the upper gastrointestinal tract and can act as solvent for tobacco-related carcinogens, facilitating their uptake through the oesophageal mucosa [47]. Furthermore, studies *in vitro* and *in vivo* reported that acetaldehyde, a metabolite of ethanol oxidation, has direct mutagenic and carcinogenic properties on human tissues [48]. A previous study did not find a modulating effect of *NAT2* polymorphisms on the association between alcohol consumption and OSCC risk [26]; however, another study suggested a direct role of *NAT2* in the metabolism of alcohol-derived carcinogens [49]. Impairment of aldehyde dehydrogenase 2 (*ALDH2*), a major enzyme involved in alcohol metabolism, has been associated with acetaldehyde-derived DNA damage in the oesophagus

following alcohol ingestion [50]. It is therefore possible that the alcohol-associated risk of OSCC observed among slow/intermediate acetylators in the Mixed Ancestry population may occur by a similar mechanism.

This study assessed the role of *NAT1* polymorphisms on susceptibility to OSCC, the interaction of NATs with alcohol and meat consumption on OSCC risk, and the role of NATs on cancer susceptibility in African populations. A limitation of the study was the lack of data for control individuals with regard to red and white meat intake and the cooking and preparation methods of the meat. Therefore, some caution must be taken when drawing interpretations of the results on the association between meat consumption and oesophageal cancer risk.

Conclusions

In conclusion, we provide evidence of tobacco smoking, alcohol consumption, and meat intake as independent risk factors for OSCC in South Africa. The main finding of this study is that *NAT1* and *NAT2* polymorphisms may modulate the risk of OSCC upon exposure to environmental risk factors in two indigenous populations of South Africa, indicating that NATs play a key role in detoxification of carcinogens responsible for the initiation of oesophageal cancer. Our findings may offer opportunities to adopt prevention strategies in lifestyle and dietary habits aimed at reducing the exposure to environmental risk factors and at preventing the occurrence of oesophageal cancer in high-risk populations of South Africa.

Additional files

Additional file 1: Primers and fluorogenic probes for *NAT110 and *NAT1**3 allele determination by TaqMan assay.**

Additional file 2: Linkage disequilibrium analysis for *NAT1* and *NAT2* SNPs in Black and Mixed Ancestry South Africans.

Additional file 3: Allele frequency of *NAT2* and *NAT1* polymorphisms and association with OSCC in Black and Mixed Ancestry South Africans.

Abbreviations

NATs: N-acetyltransferases; OSCC: Oesophageal squamous cell carcinoma; HCAs: Heterocyclic amines; SNPs: Single nucleotide polymorphisms; LD: Linkage disequilibrium.

Competing interests

The authors declare they have no competing interests.

Authors' contributions

MM collected samples and data, conceived and designed the study, carried out the molecular genetic experiments, and wrote the manuscript. MV performed the statistical analysis and helped to draft the manuscript. YW analyzed the data and helped to perform the genetic association analysis. MIP participated in the design of the study and coordinated the work. All authors read and approved the final manuscript.

Acknowledgments

This work was supported by the International Centre for Genetic Engineering and Biotechnology (ICGEB), the South African Medical Research Council and

the University of Cape Town. MM is a recipient of an ICGEB postdoctoral fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Received: 2 September 2014 Accepted: 20 February 2015
Published online: 18 March 2015

References

- Jemal A, Bray F, Forman D, O'Brien M, Ferlay J, Center M, et al. Cancer burden in Africa and opportunities for prevention. *Cancer*. 2012;118:4372–84.
- Sammon AM. Carcinogens and endemic squamous cancer of the oesophagus in Transkei, South Africa. Environmental initiation is the dominant factor; tobacco or other carcinogens of low potency or concentration are sufficient for carcinogenesis in the predisposed mucosa. *Med Hypotheses*. 2007;69:125–31.
- Hendricks D, Parker MI. Oesophageal Cancer in Africa. *IUBMB Life*. 2002;53:263–8.
- Dandara C, Li DP, Walther G, Parker MI. Gene-environment interaction: the role of *SULT1A1* and *CYP3A5* polymorphisms as risk modifiers for squamous cell carcinoma of the oesophagus. *Carcinogenesis*. 2006;27:791–7.
- Islami F, Fedirko V, Tramacere I, Bagnardi V, Jenab M, Scotti L, et al. Alcohol drinking and esophageal squamous cell carcinoma with focus on light-drinkers and never-smokers: a systematic review and meta-analysis. *Int J Cancer*. 2011;129:2473–84.
- Steevens J, Schouten LJ, Goldbohm RA, van den Brandt PA. Alcohol consumption, cigarette smoking and risk of subtypes of oesophageal and gastric cancer: a prospective cohort study. *Gut*. 2010;59:39–48.
- Keszei AP, Schouten LJ, Goldbohm RA, van den Brandt PA. Red and processed meat consumption and the risk of esophageal and gastric cancer subtypes in The Netherlands Cohort Study. *Ann Oncol*. 2012;23:2319–26.
- O'Doherty MG, Cantwell MM, Murray LJ, Anderson LA, Abnet CC, FINBAR Study Group. Dietary fat and meat intakes and risk of reflux esophagitis, Barrett's esophagus and esophageal adenocarcinoma. *Int J Cancer*. 2011;129:1493–502.
- Turesky RJ. Formation and biochemistry of carcinogenic heterocyclic aromatic amines in cooked meats. *Toxicol Lett*. 2007;168:219–27.
- Jakszyn P, Agudo A, Ibáñez R, García-Closas R, Pera G, Amiano P, et al. Development of a food database of nitrosamines, heterocyclic amines, and polycyclic aromatic hydrocarbons. *J Nutr*. 2004;134:2011–4.
- Hein DW. Molecular genetics and function of *NAT1* and *NAT2*: role in aromatic amine metabolism and carcinogenesis. *Mutat Res*. 2002;506–507:65–77.
- Hickman D, Risch A, Buckle V, Spurr NK, Jeremiah SJ, McCarthy A, et al. Chromosomal localization of human genes for arylamine N-acetyltransferase. *Biochem J*. 1994;297:441–5.
- Hein DW, Doll MA, Fretland AJ, Leff MA, Webb SJ, Xiao GH, et al. Molecular genetics and epidemiology of the *NAT1* and *NAT2* acetylation polymorphisms. *Cancer Epidemiol Biomarkers Prev*. 2000;9:29–42.
- Taja-Chayeb L, González-Fierro A, Miguez-Muñoz C, Trejo-Becerril C, Cruz-Hernandez Ede L, Cantu D, et al. Acetylator status and N-acetyltransferase 2 gene polymorphisms; phenotype-genotype correlation with the sulfamethazine test. *Pharmacogenet Genomics*. 2011;21:894–901.
- Fretland AJ, Leff MA, Doll MA, Hein DW. Functional characterization of human N-acetyltransferase 2 (*NAT2*) single nucleotide polymorphisms. *Pharmacogenetics*. 2001;11:207–15.
- Hein DW, Doll MA. Accuracy of various human *NAT2* SNP genotyping panels to infer rapid, intermediate and slow acetylator phenotypes. *Pharmacogenomics*. 2012;13:31–41.
- Adams CH, Werely CJ, Victor TC, Hoal EG, Rossouw G, van Helden PD. Allele frequencies for glutathione S-transferase and N-acetyltransferase 2 differ in African population groups and may be associated with oesophageal cancer or tuberculosis incidence. *Clin Chem Lab Med*. 2003;41:600–5.
- Zhu Y, States JC, Wang Y, Hein DW. Functional effects of genetic polymorphisms in the N-acetyltransferase 1 coding and 3' untranslated regions. *Birth Defects Res A Clin Mol Teratol*. 2011;91:77–84.
- Loktionov A, Moore W, Spencer SP, Vorster H, Nell T, O'Neill IK, et al. Differences in N-acetylation genotypes between Caucasians and Black South Africans: implications for cancer prevention. *Cancer Detect Prev*. 2002;26:15–22.
- Wang D, Para MF, Koletar SL, Sadee W. Human N-acetyltransferase 1 *10 and *11 alleles increase protein expression through distinct mechanisms and associate with sulfamethoxazole-induced hypersensitivity. *Pharmacogenet Genomics*. 2011;21:652–64.

21. Zang Y, Doll MA, Zhao S, States JC, Hein DW. Functional characterization of single-nucleotide polymorphisms and haplotypes of human N-acetyltransferase 2. *Carcinogenesis*. 2007;28:1665–71.
22. Agúndez JA. Polymorphisms of human N-acetyltransferases and cancer risk. *Curr Drug Metab*. 2008;9:520–31.
23. Nöthlings U, Yamamoto JF, Wilkens LR, Murphy SP, Park SY, Henderson BE, et al. Meat and heterocyclic amine intake, smoking, *NAT1* and *NAT2* polymorphisms, and colorectal cancer risk in the multiethnic cohort study. *Cancer Epidemiol Biomarkers Prev*. 2009;18:2098–106.
24. Morita S, Yano M, Tsujinaka T, Ogawa A, Taniguchi M, Kaneko K, et al. Association between genetic polymorphisms of glutathione S-transferase P1 and N-acetyltransferase 2 and susceptibility to squamous-cell carcinoma of the esophagus. *Int J Cancer*. 1998;79:517–20.
25. Shibuta J, Eto T, Kataoka A, Inoue H, Ueo H, Suzuki T, et al. Genetic polymorphism of N-acetyltransferase 2 in patients with esophageal cancer. *Am J Gastroenterol*. 2001;96:3419–24.
26. Jain M, Kumar S, Lal P, Tiwari A, Ghoshal UC, Mittal B. Association of genetic polymorphisms of N-acetyltransferase 2 and susceptibility to esophageal cancer in north Indian population. *Cancer Invest*. 2007;25:340–6.
27. Lee JM, Lee YC, Yang SY, Shi WL, Lee CJ, Luh SP, et al. Genetic polymorphisms of p53 and GSTP1, but not NAT2, are associated with susceptibility to squamous-cell carcinoma of the esophagus. *Int J Cancer*. 2000;89:458–64.
28. Malik MA, Upadhyay R, Modi DR, Zargar SA, Mittal B. Association of NAT2 gene polymorphisms with susceptibility to esophageal and gastric cancers in the Kashmir Valley. *Arch Med Res*. 2009;40:416–23.
29. Wideroff L, Vaughan TL, Farin FM, Gammon MD, Risch H, Stanford JL, et al. GST, NAT1, CYP1A1 polymorphisms and risk of esophageal and gastric adenocarcinomas. *Cancer Detect Prev*. 2007;31:233–6.
30. de Wit E, Delport W, Rugamika CE, Meintjes A, Möller M, van Helden PD, et al. Genome-wide analysis of the structure of the South African Coloured Population in the Western Cape. *Hum Genet*. 2010;128:145–53.
31. Gustafson S, Proper JA, Bowie EJ, Sommer SS. Parameters affecting the yield of DNA from human blood. *Anal Biochem*. 1987;165:294–9.
32. Doll MA, Hein DW. Rapid genotype method to distinguish frequent and/or functional polymorphisms in human N-acetyltransferase-1. *Anal Biochem*. 2002;301:328–32.
33. Rodríguez S, Gaunt TR, Day IN. Hardy-Weinberg equilibrium testing of biological ascertainment for Mendelian randomization studies. *Am J Epidemiol*. 2009;169:505–14.
34. Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics*. 2005;21:263–5.
35. Dudbridge F. Likelihood-based association analysis for nuclear families and unrelated subjects with missing genotype data. *Hum Hered*. 2008;66:87–98.
36. Meyer D, Parkin DP, Seifart HI, Maritz JS, Engelbrecht AH, Werely CJ, et al. *NAT2* slow acetylator function as a risk indicator for age-related cataract formation. *Pharmacogenetics*. 2003;13:285–9.
37. Smelt VA, Mardon HJ, Sim E. Placental expression of arylamine N-acetyltransferases: evidence for linkage disequilibrium between *NAT1*10* and *NAT2*4* alleles of the two human arylamine N-acetyltransferase loci *NAT1* and *NAT2*. *Pharmacol Toxicol*. 1998;83:149–57.
38. Cascorbi I, Roots I, Brockmöller J. Association of *NAT1* and *NAT2* polymorphisms to urinary bladder cancer: Significantly reduced risk in subjects with *NAT1*10*. *Cancer Res*. 2001;61:5051–6.
39. Pacella-Norman R, Urban MI, Sitas F, Carrara H, Sur R, Hale M, et al. Risk factors for oesophageal, lung, oral and laryngeal cancers in black South Africans. *Br J Cancer*. 2002;86:1751–6.
40. Hein DW, Doll MA, Rustan TD, Ferguson RJ. Metabolic activation of N-hydroxyarylamines and N-hydroxyarylamides by 16 recombinant human *NAT2* allozymes: effects of 7 specific *NAT2* nucleic acid substitutions. *Cancer Res*. 1995;55:3531–6.
41. Wang L, Tang W, Chen S, Sun Y, Fan Y, Shi Y, et al. N-acetyltransferase 2 polymorphisms and risk of esophageal cancer in a Chinese population. *PLoS One*. 2014;9:e87783.
42. Skog K, Solyakov A. Heterocyclic amines in poultry products: a literature review. *Food Chem Toxicol*. 2002;40:1213–21.
43. González CA, Jakszyn P, Pera G, Agudo A, Bingham S, Palli D, et al. Meat intake and risk of stomach and esophageal adenocarcinoma within the European Prospective Investigation into Cancer and Nutrition (EPIC). *J Natl Cancer Inst*. 2006;98:345–54.
44. Jakszyn P, Luján-Barroso L, Agudo A, Bueno-de-Mesquita HB, Molina E, Sánchez MJ, et al. Meat and heme iron intake and esophageal adenocarcinoma in the European Prospective Investigation into Cancer and Nutrition study. *Int J Cancer*. 2013;133:2744–50.
45. Tappel A. Heme of consumed red meat can act as a catalyst of oxidative damage and could initiate colon, breast and prostate cancers, heart disease and other diseases. *Med Hypotheses*. 2007;68:562–4.
46. Cross AJ, Pollock JR, Bingham SA. Haem, not protein or inorganic iron, is responsible for endogenous intestinal N-nitrosation arising from red meat. *Cancer Res*. 2003;63:2358–60.
47. Seitz HK, Stickel F, Homann N. Pathogenetic mechanisms of upper aerodigestive tract cancer in alcoholics. *Int J Cancer*. 2004;108:483–7.
48. Salaspuro M. Acetaldehyde as a common denominator and cumulative carcinogen in digestive tract cancers. *Scand J Gastroenterol*. 2009;44:912–25.
49. Chen C, Ricks S, Doody DR, Fitzgibbons ED, Porter PL, Schwartz SM. N-Acetyltransferase 2 polymorphisms, cigarette smoking and alcohol consumption, and oral squamous cell cancer risk. *Carcinogenesis*. 2001;22:1993–9.
50. Yukawa Y, Ohashi S, Amanuma Y, Nakai Y, Tsurumaki M, Kikuchi O, et al. Impairment of aldehyde dehydrogenase 2 increases accumulation of acetaldehyde-derived DNA damage in the esophagus after ethanol ingestion. *Am J Cancer Res*. 2014;4:279–84.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

