

## RESEARCH

## Open Access

# Stochastic delay evolution equations driven by sub-fractional Brownian motion

Zhi Li<sup>1\*</sup>, Guoli Zhou<sup>2</sup> and Jiaowan Luo<sup>3</sup>\*Correspondence:  
lizhi\_csu@126.com<sup>1</sup>School of Information and Mathematics, Yangtze University, Jingzhou, 434023, China  
Full list of author information is available at the end of the article**Abstract**

In this paper, we investigate the existence, uniqueness and exponential asymptotic behavior of mild solutions to stochastic delay evolution equations perturbed by a sub-fractional Brownian motion  $S_Q^H(t): dX(t) = (AX(t) + f(t, X_t)) dt + g(t) dS_Q^H(t)$  with index  $H \in (1/2, 1)$ .

**MSC:** 60H15; 60G15; 60H05**Keywords:** existence and uniqueness; stochastic delay evolution equations; sub-fractional Brownian motion; exponential decay in mean square

## 1 Introduction

The fractional Brownian motion (fBm for short) has become an object of intense study, due to its interesting properties and its applications in various scientific areas including telecommunications, turbulence and finance. The fBm with Hurst index  $H \in (0, 1)$  is a continuous centered Gaussian process  $B^H = \{B^H(t), t \geq 0\}$ , starting from zero, with covariance

$$R_H(t, s) = \mathbb{E}[B^H(t)B^H(s)] = \frac{1}{2} [t^{2H} + s^{2H} + |t - s|^{2H}] \quad (1.1)$$

for all  $s, t \geq 0$ . For  $H = \frac{1}{2}$ ,  $B^H$  coincides with the standard Brownian motion  $B$ .  $B^H$  is neither a semimartingale nor a Markov process when  $H \neq \frac{1}{2}$ . The fBm is a suitable generalization of the standard Brownian motion, but exhibits long-range dependence, self-similarity and which has stationary increments. On the other hand, based on the sufficient study of fBm, many authors have proposed to use more general self-similar Gaussian processes and random fields as stochastic models. Such applications have raised many interesting theoretical questions about self-similar Gaussian processes and fields in general. Therefore, some other generalizations of the fBm were introduced. However, contrast to the extensive studies on fBm, there has been little systematic investigation on other self-similar Gaussian processes. The main reason for this is the complexity of dependence structures for self-similar Gaussian processes which do not have stationary increments.

As an extension of Brownian motion, recently, Bojdecki *et al.* [1] introduced and studied a rather special class of self-similar Gaussian process. This process arises from occupation time fluctuations of branching particle systems with Poisson initial condition. This process is called the sub-fractional Brownian motion (sub-fBm in short). The sub-fBm with index

$H \in (0, 1)$  is mean zero Gaussian  $S^H = \{S^H(t), t \geq 0\}$  starting from zero, with covariance

$$C_H(t, s) = \mathbb{E}[S^H(t)S^H(s)] = s^{2H} + t^{2H} - \frac{1}{2}[(s + t)^{2H} + |t - s|^{2H}] \tag{1.2}$$

for all  $s, t \geq 0$ . For  $H = \frac{1}{2}$ ,  $S^H$  coincides with the standard Brownian motion  $B$ .  $S^H$  is neither a semimartingale nor a Markov process when  $H \neq \frac{1}{2}$ , so many of the power techniques from stochastic analysis are not available when dealing with  $S^H$ . The sub-fBm has properties analogous to those of fBm (self-similarity, long-range dependence, Hölder paths), but it does not have stationary increments. More works for sub-fractional Brownian motion can be found in Bojdecki *et al.* [2, 3], Tudor [4–9] and Yan and Shen [10].

The sub-fractional Brownian motion satisfies the following estimates:

$$[(2 - 2^{2H-1}) \wedge 1] |t - s|^{2H} \mathbb{E}|S^H(t) - S^H(s)|^2 \leq [(2 - 2^{2H-1}) \wedge 1] |t - s|^{2H}. \tag{1.3}$$

Thus, Kolmogorov’s continuity criterion implies that sub-fBm is Hölder continuous of order  $\gamma$  for any  $\gamma < H$  on any finite interval. Therefore, if  $u$  is a stochastic process with Hölder continuous trajectories of order  $\beta > 1 - H$ , then the path-wise Riemann-Stieltjes integral  $\int_0^T u_t(\omega) dS^H(t)(\omega)$  exists for all  $T \geq 0$  (see Young [11]). Yan *et al.* [12] have used the divergence operator to define the stochastic integrals with respect to sub-fBm with  $H > 1/2$  and showed that if  $f \in C^2(\mathbb{R})$  and  $1/2 < H < 1$ , then

$$f(S^H(t)) = f(0) + \int_0^t f'(S^H(s)) dS^H(s) + H(2 - 2^{2H-1}) \int_0^t f''(S^H(s)) s^{2H-1} ds.$$

In 2012, Shen and Chen [13] defined a stochastic integral with respect to sub-fractional Brownian motion  $S^H$  with index  $H \in (0, 1/2)$  that extends the divergence integral from Malliavin calculus, and established versions of the formulas of Itô and Tanaka that hold for all  $H \in (0, 1/2)$ . Tudor [6, 8] characterized the domain of the Wiener integral with respect to a sub-fractional Brownian motion  $S^H(s)$  with index  $H \in (0, 1)$  and stochastic differential equations driven by sub-fractional Brownian motion has also been consider by Mendy [14].

In this paper, we desire to define a stochastic integral with respect to  $Q$ -sub-fractional Brownian motion  $S_Q^H$  in infinite dimensional space and consider the following stochastic semilinear delay evolution equation:

$$\begin{cases} dX(t) = (AX(t) + f(t, X_t)) dt + g(t) dS_Q^H(t), \\ X(s) = \varphi(s), \quad -r \leq s \leq 0, r \geq 0, \end{cases} \tag{1.4}$$

under suitable conditions on the operator  $A$ , the coefficient functions  $f, g$ , and the initial value  $\varphi$ . Here  $S_Q^H(t)$  denotes an  $Q$ -sub-fBm with  $H \in (1/2, 1)$ . The purpose of this paper is to investigate the existence and uniqueness of mild solutions to the stochastic delay differential equation (1.4) and to study its longtime behavior as well.

The contents of the paper are as follows. In Section 2, some necessary preliminaries on the stochastic integration with respect to sub-fBm are established. Also a technical lemma which is crucial in our stability analysis is proved. In Section 3, the existence and uniqueness of mild solutions are proved. In Section 4, we prove that a mild solution, when it

exists, is also a weak solution. In Section 5, we establish some sufficient conditions ensuring the exponential decay to zero in mean square of the mild solution of our delay model. In Section 6, an example is given to illustrate the effectiveness of our results.

## 2 Preliminaries

In this section we introduce the sub-fractional Brownian motion as well as the Wiener integral with respect to it. We also establish some important results which will be needed throughout this paper.

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a complete probability space. Now we aim at introducing the Wiener integral with respect to one-dimensional sub-fBm  $S^H$ . Fix a time interval  $[0, T]$ . We denote by  $\mathcal{E}$  the linear space of  $\mathbb{R}$ -valued step functions on  $[0, T]$ , that is,  $\varphi \in \mathcal{E}$  if

$$\varphi(t) = \sum_{i=1}^{n-1} x_i \mathbf{I}_{[t_i, t_{i+1})}(t),$$

where  $t \in [0, T]$ ,  $x_i \in \mathbb{R}$ , and  $0 = t_1 < t_2 < \dots < t_n = T$ . For  $\varphi \in \mathcal{E}$  we define its Wiener integral with respect to  $S^H$  as

$$\int_0^T \varphi(s) dS^H(s) = \sum_{i=1}^n x_i (S_{t_{i+1}}^H - S_{t_i}^H).$$

Let  $\mathcal{H}_{S^H}$  be the canonical Hilbert space associated to the sub-fBm  $S^H$ . That is,  $\mathcal{H}_{S^H}$  is the closure of the linear span  $\mathcal{E}$  with respect to the scalar product

$$\langle \mathbf{I}_{[0,t]}, \mathbf{I}_{[0,s]} \rangle_{\mathcal{H}_{S^H}} = C_H(t, s).$$

We know that the covariance of sub-fractional Brownian motion can be written as

$$\mathbb{E}[S^H(t)S^H(s)] = \int_0^t \int_0^s \phi_H(u, v) du dv = C_H(s, t), \tag{2.1}$$

where  $\phi_H(u, v) = H(2H - 1)[|u - v|^{2H-2} - (u + v)^{2H-2}]$ .

Equation (2.1) implies that

$$\langle \varphi, \psi \rangle_{\mathcal{H}_{S^H}} = \int_0^t \int_0^t \varphi_u \psi_v \phi_H(u, v) du dv \tag{2.2}$$

for any pair step functions  $\varphi$  and  $\psi$  on  $[0, T]$ . Consider the kernel

$$n_H(t, s) = \frac{2^{1-H} \sqrt{\pi}}{\Gamma(H - \frac{1}{2})} s^{3/2-H} \left( \int_s^t (x^2 - s^2)^{H-3/2} dx \right) \mathbf{I}_{[0,t]}(s). \tag{2.3}$$

By Dzharparidze and Van Zanten [15], we have

$$C_H(t, s) = c_H^2 \int_0^{s \wedge t} n_H(t, u) n_H(s, u) du, \tag{2.4}$$

where

$$c_H^2 = \frac{\Gamma(1 + 2H) \sin(\pi H)}{\pi}.$$

Property (2.4) implies that  $C_H(s, t)$  is non-negative definite. Consider the linear operator  $n_H^*$  from  $\mathcal{E}$  to  $L^2([0, T])$  defined by

$$n_H^*(\varphi)(s) := c_H \int_s^r \varphi_r \frac{\partial n_H}{\partial r}(r, s) dr.$$

Using (2.2) and (2.4) we have

$$\begin{aligned} \langle n_H^* \varphi, n_H^* \psi \rangle_{L^2([0, T])} &= c_H^2 \int_0^T \left( \int_s^T \varphi_r \frac{\partial n_H}{\partial r}(r, s) dr \right) \left( \int_s^T \psi_u \frac{\partial n_H}{\partial u}(u, s) du \right) ds \\ &= c_H^2 \int_0^T \int_0^T \left( \int_0^{r \wedge u} \frac{\partial n_H}{\partial r}(r, s) \frac{\partial n_H}{\partial u}(u, s) ds \right) \varphi_r \psi_u dr du \\ &= c_H^2 \int_0^T \int_0^T \frac{\partial^2 n_H}{\partial r \partial u}(r, u) \varphi_r \psi_u dr du \\ &= H(2H - 1) \int_0^T \int_0^T [|u - r|^{2H-2} - (u + r)^{2H-2}] \varphi_r \psi_u dr du \\ &= \langle \varphi, \psi \rangle_{\mathcal{H}_{S^H}}. \end{aligned} \tag{2.5}$$

As a consequence, the operator  $n_H^*$  provides an isometry between the Hilbert space  $\mathcal{H}_{S^H}$  and  $L^2([0, T])$ . Hence, the process  $W$  defined by

$$W(t) := S^H \left( (n_H^*)^{-1}(\mathbf{I}_{[0, t]}) \right)$$

is a Wiener process, and  $S^H$  has the following Wiener integral representation:

$$S^H(t) = c_H \int_0^t n_H(t, s) dW(s)$$

because  $(n_H^*)(\mathbf{I}_{[0, t]})(s) = c_H n_H(t, s)$ . By Dzharaparidze and Van Zanten [15], we have

$$W(t) = \int_0^t \psi_H(t, s) dS^H(s),$$

where

$$\begin{aligned} \psi_H(t, s) &= \frac{s^{H-1/2}}{\Gamma(3/2 - H)} \left[ t^{H-3/2} (t^2 - s^2)^{1/2-H} - (H - 3/2) \int_s^t (x^2 - s^2)^{1/2-H} x^{H-3/2} dx \right] \\ &\quad \times \mathbf{I}_{[0, t]}(s). \end{aligned}$$

In addition, for any  $\varphi \in \mathcal{H}_{S^H}$ ,

$$\int_0^T \varphi(s) dS^H(s) = \int_0^T (n_H^* \varphi)(t) dW(t)$$

if and only if  $n_H^* \varphi \in L^2([0, T])$ .

Also denoting  $L_{\mathcal{H}_{S^H}}^2([0, T]) = \{\varphi \in \mathcal{H}_{S^H}, n_H^* \varphi \in L^2([0, T])\}$ . Since  $H > 1/2$ , we have by

(2.5) and Lemma 2.1 of [16],

$$L^2([0, T]) \subset L^{\frac{1}{H}}([0, T]) \subset L^2_{\mathcal{H}_s, H}([0, T]). \tag{2.6}$$

Moreover, the following useful result holds:

**Lemma 2.1** (Nualart [17]) *For  $\varphi \in L^{1/H}([0, T])$ ,*

$$H(2H - 1) \int_0^T \int_0^T |\varphi_r| |\varphi_u| |u - r|^{2H-2} dr du \leq C_H \|\varphi\|_{L^{\frac{1}{H}}([0, T])},$$

where  $C_H = (\frac{H(2H-1)}{B(2-2H, H-\frac{1}{2})})^{1/2}$ , with  $B$  denoting the beta function.

Next we are interested in considering a sub-fBm with values in Hilbert space and giving the definition of the corresponding stochastic integral.

Let  $(U, \|\cdot\|_U, \langle \cdot, \cdot \rangle_U)$  and  $(K, \|\cdot\|_K, \langle \cdot, \cdot \rangle_K)$  be two separable Hilbert spaces. Let  $L(K, U)$  denote the space of all bounded linear operators from  $K$  to  $U$ . Let  $Q \in L(K, K)$  be a non-negative self-adjoint operator. Denote by  $L^0_Q(K, U)$  the space of all  $\xi \in L(K, U)$  such that  $\xi Q^{\frac{1}{2}}$  is a Hilbert-Schmidt operator. The norm is given by

$$\|\xi\|_{L^0_Q(K, U)}^2 = \|\xi Q^{\frac{1}{2}}\|_{HS}^2 = \text{tr}(\xi Q \xi^*).$$

Then  $\xi$  is called a  $Q$ -Hilbert-Schmidt operator from  $K$  to  $U$ .

Let  $\{S_n^H(t)\}_{n \in \mathbb{N}}$  be a sequence of one-dimensional standard sub-fractional Brownian motions mutually independent on  $(\Omega, \mathcal{F}, \mathbb{P})$ . When one considers the following series:

$$\sum_{n=1}^{\infty} S_n^H(t) e_n, \quad t \geq 0,$$

where  $\{e_n\}_{n \in \mathbb{N}}$  is a complete orthonormal basis in  $K$ , this series does not necessarily converge in the space  $K$ . Thus we consider a  $K$ -valued stochastic process  $S_Q^H(t)$  given formally by the following series:

$$S_Q^H(t) = \sum_{n=1}^{\infty} S_n^H(t) Q^{\frac{1}{2}} e_n, \quad t \geq 0.$$

If  $Q$  is a non-negative self-adjoint trace class operator, then this series converges in the space  $K$ , that is, we have  $S_Q^H(t) \in L^2(\Omega, K)$ . Then we say that the above  $S_Q^H(t)$  is a  $K$ -valued  $Q$ -cylindrical sub-fractional Brownian motion with covariance operator  $Q$ . For example, if  $\{\sigma_n\}_{n \in \mathbb{N}}$  is a bounded sequence of non-negative real numbers such that  $Qe_n = \sigma_n e_n$ , assuming that  $Q$  is a nuclear operator in  $K$  (that is,  $\sum_{n=1}^{\infty} \sigma_n < \infty$ ), then the stochastic process

$$S_Q^H(t) = \sum_{n=1}^{\infty} S_n^H(t) Q^{\frac{1}{2}} e_n = \sum_{n=1}^{\infty} \sqrt{\sigma_n} S_n^H(t) e_n, \quad t \geq 0,$$

is well defined as a  $K$ -valued  $Q$ -cylindrical sub-fractional Brownian motion.

Let  $\varphi : [0, T] \rightarrow L^0_Q(K, U)$  such that

$$\sum_{n=1}^{\infty} \|n_H^*(\varphi Q^{1/2} e_n)\|_{L^2([0,T];U)} < \infty. \tag{2.7}$$

**Definition 2.1** Let  $\varphi : [0, T] \rightarrow L^0_Q(K, U)$  satisfy (2.7). Then its stochastic integral with respect to the sub-fBm  $S_Q^H$  is defined, for  $t \geq 0$ , as follows:

$$\begin{aligned} \int_0^t \varphi(s) dS_Q^H(s) &:= \sum_{n=1}^{\infty} \int_0^t \varphi(s) Q^{1/2} e_n dS_n^H(s) \\ &= \sum_{n=1}^{\infty} \int_0^t (n_H^*(\varphi Q^{1/2} e_n))(s) dW(s). \end{aligned}$$

Notice that if

$$\sum_{n=1}^{\infty} \|\varphi Q^{1/2} e_n\|_{L^{1/H}([0,T];U)} < \infty, \tag{2.8}$$

then in particular (2.7) holds, which follows immediately from (2.6).

The following lemma is obtained as a simple application of Lemma 2.1.

**Lemma 2.2** For any  $\varphi : [0, T] \rightarrow L^0_Q(K, U)$  such that (2.8) holds, and for any  $\alpha, \beta \in [0, T]$  with  $\alpha > \beta$ ,

$$\mathbb{E} \left\| \int_{\alpha}^{\beta} \varphi(s) dS_Q^H(s) \right\|_U^2 \leq C_H(\alpha - \beta)^{2H-1} \sum_{n=1}^{\infty} \int_{\alpha}^{\beta} \|\varphi(s) Q^{1/2} e_n\|_U^2 ds.$$

If, in addition,

$$\sum_{n=1}^{\infty} \|\varphi(s) Q^{1/2} e_n\|_U^2 \text{ is uniformly convergent for } t \in [0, T], \tag{2.9}$$

then

$$\mathbb{E} \left\| \int_{\alpha}^{\beta} \varphi(s) dS_Q^H(s) \right\|_U^2 \leq C_H(\alpha - \beta)^{2H-1} \int_{\alpha}^{\beta} \|\varphi(s)\|_{L^0_Q(K,U)}^2 ds. \tag{2.10}$$

*Proof* Let  $\{e_n\}_{n \in \mathbb{N}}$  be the complete orthonormal basis of  $K$  introduced above. Applying Lemma 2.1 we can obtain

$$\begin{aligned} \mathbb{E} \left\| \int_{\alpha}^{\beta} \varphi(s) dS_Q^H(s) \right\|_U^2 &= \mathbb{E} \left\| \sum_{n=1}^{\infty} \int_{\alpha}^{\beta} \varphi(s) Q^{1/2} e_n dS_n^H(s) \right\|_U^2 \\ &= \sum_{n=1}^{\infty} \mathbb{E} \left\| \int_{\alpha}^{\beta} \varphi(s) Q^{1/2} e_n dS_n^H(s) \right\|_U^2 \\ &= \sum_{n=1}^{\infty} H(2H - 1) \int_{\alpha}^{\beta} \int_{\alpha}^{\beta} \|\varphi(t) Q^{1/2} e_n\|_U \|\varphi(s) Q^{1/2} e_n\|_U \end{aligned}$$

$$\begin{aligned}
 & \times [|t-s|^{2H-2} - (t+s)^{2H-2}] dt ds \\
 & \leq \sum_{n=1}^{\infty} H(2H-1) \int_{\alpha}^{\beta} \int_{\alpha}^{\beta} \|\varphi(t)Q^{1/2}e_n\|_U \|\varphi(s)Q^{1/2}e_n\|_U \\
 & \quad \times |t-s|^{2H-2} dt ds \\
 & \leq C_H \sum_{n=1}^{\infty} \left( \int_{\alpha}^{\beta} \|\varphi(s)Q^{1/2}e_n\|_U^{1/H} ds \right)^{2H} \\
 & \leq C_H(\alpha-\beta)^{2H-1} \sum_{n=1}^{\infty} \int_{\alpha}^{\beta} \|\varphi(s)Q^{1/2}e_n\|_U^2 ds.
 \end{aligned}$$

The second assertion is an immediate consequence of the Weierstrass  $M$ -test. □

**Remark 2.1** If  $\{\sigma_n\}_{n \in \mathbb{N}}$  is a bounded sequence of non-negative real numbers such that the nuclear operator  $Q$  satisfies  $Qe_n = \sigma_n e_n$ , assuming that there exists a positive constant  $k_{\varphi}$  such that

$$\|\varphi(t)\|_{L^2_Q(K,U)} \leq k_{\varphi}, \quad \text{uniformly in } t \in [0, T],$$

then (2.9) holds automatically.

### 3 Existence and uniqueness of mild solution

We denote by  $C(a, b; L^2(\Omega; U)) = C(a, b; L^2(\Omega, \mathcal{F}, \mathbb{P}; U))$  the Banach space of all continuous functions from  $[a, b]$  into  $L^2(\Omega; U)$  equipped with sup norm. Let us consider two fixed real numbers  $r \geq 0$  and  $T > 0$ . If  $x \in C(-r, T; L^2(\Omega; U))$  for each  $t \in [0, T]$  we denote  $x_t \in C(-r, 0; L^2(\Omega; U))$  the function defined by  $x_t(\theta) = x(t + \theta)$ , for  $\theta \in [-r, 0]$ .

In this section we consider the existence and uniqueness of mild solutions to the following stochastic evolution equation with delays:

$$\begin{cases} dX(t) = (AX(t) + f(t, X_t)) dt + g(t) dS_Q^H(t), & t \in [0, T], \\ X(t) = \varphi(t), & t \in [-r, 0], \end{cases} \tag{3.1}$$

where  $S_Q^H(t)$  is the sub-fractional Brownian motion which was introduced in the previous section, the initial data  $\varphi \in C(-r, 0; L^2(\Omega; U))$ , and  $A : \text{Dom}(A) \subset U \rightarrow U$  is the infinitesimal generator of a strongly continuous semigroup  $S(\cdot)$  on  $U$ , that is, for  $t \geq 0$ , we have

$$\|S(t)\|_U \leq Me^{\rho t}, \quad M \geq 1, \rho \in \mathbb{R}.$$

$f : [0, T] \times C(-r, 0; U) \rightarrow U$  is a family of nonlinear operators defined for almost every  $t$  which satisfy:

- (f.1) The mapping  $t \in [0, T] \rightarrow f(t, \xi) \in U$  is Lebesgue measurable for all  $\xi \in C(-r, 0; L^2(\Omega; U))$ .
- (f.2) There exists a constant  $C > 0$  such that for any  $x, y \in C(-r, T; U)$  and  $t \in [0, T]$ ,

$$\int_0^t \|f(s, x_s) - f(s, y_s)\|_U^2 ds \leq C \int_{-r}^t \|x(s) - y(s)\|_U^2 ds.$$

$$(f.3) \int_0^T \|f(s, 0)\|_U^2 ds < \infty.$$

Moreover, for  $g : [0, T] \rightarrow L^0_Q(K, U)$  we assume the following conditions: for the complete orthonormal basis  $\{e_n\}_{n \in \mathbb{N}}$  in  $K$ , we have:

$$(g.1) \sum_{n=1}^\infty \|g Q^{1/2} e_n\|_{L^2([0, T]; U)} < \infty.$$

$$(g.2) \sum_{n=1}^\infty \|g(t) Q^{1/2} e_n\|_U \text{ is uniformly convergent for } t \in [0, T].$$

**Definition 3.1** A  $U$ -valued process  $X(t)$  is called a mild solution of (3.1) if  $X \in C(-r, T; L^2(\Omega; U))$ ,  $X(t) = \varphi(t)$ , for  $t \in [-r, 0]$  and for  $t \in [0, T]$ , satisfies

$$\begin{aligned} X(t) &= S(t)\varphi(0) + \int_0^t S(t-s)f(s, X_s) ds \\ &\quad + \int_0^t S(t-s)g(s) dS_Q^H(t) \quad \mathbb{P}\text{-a.s.} \end{aligned} \tag{3.2}$$

Notice that, thanks to (g.1) and the fact that  $H \in (1/2, 1)$ , (2.8) holds, which implies that the stochastic integral in (3.2) is well defined since  $S(\cdot)$  is a strongly continuous semigroup. Moreover, (g.1) together with (g.2) immediately imply that, for every  $t \in [0, T]$ ,

$$\int_0^t \|g(s)\|_{L^0_Q(K, U)}^2 ds < \infty.$$

**Theorem 3.1** Under the assumptions on  $A$  and conditions (f.1)-(f.3) and (g.1)-(g.2), for every  $\varphi \in C(-r, 0; L^2(\Omega, U))$  there exists a unique mild solution  $X$  to (3.1).

*Proof* We can assume that  $\rho > 0$ , otherwise we can take  $\rho_0 > 0$  such that, for  $t \geq 0$ ,  $\|S(t)\|_U \leq Me^{\rho_0 t}$ .

We start the proof by checking the uniqueness of solutions. Assume that  $X, Y$  are two mild solutions of (3.1). Then

$$\begin{aligned} \mathbb{E} \|X(t) - Y(t)\|_U^2 &\leq t \mathbb{E} \int_0^t \|S(t-s)(f(s, X_s) - f(s, Y_s))\|_U^2 ds \\ &\leq tM^2 e^{2\rho t} \mathbb{E} \int_0^t \|f(s, X_s) - f(s, Y_s)\|_U^2 ds \\ &\leq tM^2 e^{2\rho t} C \int_0^t \mathbb{E} \|X(s) - Y(s)\|_U^2 ds \\ &\leq tM^2 e^{2\rho t} C \int_0^t \sup_{0 \leq \tau \leq s} \mathbb{E} \|X(\tau) - Y(\tau)\|_U^2 ds, \end{aligned}$$

and therefore, since  $X = Y$  over the interval  $[-r, 0]$ , by taking the supremum in the above inequality,

$$\sup_{0 \leq \theta \leq t} \mathbb{E} \|X(\theta) - Y(\theta)\|_U^2 \leq TM^2 e^{2\rho T} C \int_0^t \sup_{0 \leq \tau \leq s} \mathbb{E} \|X(\tau) - Y(\tau)\|_U^2 ds.$$

The Gronwall's lemma implies now the uniqueness result.

Now we prove the existence of solutions to problem (3.1).



First of all, we check that the well-defined stochastic integral possesses the repaired regularity. To this end, let us consider  $\sigma > 0$  small enough. We have

$$\begin{aligned} & \mathbb{E} \left\| \int_0^{t+\sigma} S(t+\sigma-s)g(s) dS_Q^H(s) - \int_0^t S(t-s)g(s) dS_Q^H(s) \right\|_U^2 \\ & \leq 2\mathbb{E} \left\| \int_0^{t+\sigma} (S(t+\sigma-s) - S(t-s))g(s) dS_Q^H(s) \right\|_U^2 \\ & \quad + 2\mathbb{E} \left\| \int_t^{t+\sigma} S(t-s)g(s) dS_Q^H(s) \right\|_U^2 \\ & := J_1 + J_2. \end{aligned}$$

Applying inequality (2.10) to  $J_1$ , we obtain

$$\begin{aligned} J_1 & \leq 2C_H t^{2H-1} \int_0^t \|S(t-s)(S(\sigma) - Id)g(s)\|_{L^0_Q(K,U)}^2 ds \\ & \leq C_H t^{2H-1} M^2 e^{2\rho t} \int_0^t \|(S(\sigma) - Id)g(s)\|_{L^0_Q(K,U)}^2 ds \rightarrow 0 \end{aligned}$$

when  $\sigma \rightarrow 0$  thanks to the Lebesgue majorant theorem, since, for every  $s$  fixed,

$$S(\sigma)g(s) \rightarrow g(s), \quad \|S(\sigma)g(s)\|_{L^0_Q(K,U)} \leq Me^{\rho\sigma} \|g(s)\|_{L^0_Q(K,U)}.$$

Applying now (2.10) to  $J_2$ , we have

$$J_2 \leq 2C_H \sigma^{2H-1} M^2 e^{2\rho\sigma} \int_t^{t+\sigma} \|g(s)\|_{L^0_Q(K,U)}^2 ds \rightarrow 0$$

when  $\sigma \rightarrow 0$ . Therefore the stochastic integral belongs to the space  $C(-r, T; L^2(\Omega; U))$ .

We denote  $X_0 = 0$  and define by recurrence a sequence  $\{X^n\}_{n \in \mathbb{N}}$  of processes as

$$\begin{cases} X^n(t) = S(t)\varphi(0) + \int_0^t S(t-s)f(s, X_s^{n-1}) ds + \int_0^t S(t-s)g(s) dS_Q^H(s), & t \in [0, T], \\ X^n(t) = \varphi(t), & t \in [-r, 0]. \end{cases} \quad (3.3)$$

The sequence (3.3) is well defined, since  $X^0 = 0 \in C(-r, T; L^2(\Omega; U))$  and given  $X^{n-1} \in C(-r, T; L^2(\Omega; U))$ , let us check that  $X^n \in C(-r, T; L^2(\Omega; U))$  as well. To this end, let us consider  $\sigma > 0$  sufficiently small. Then

$$\begin{aligned} \|X^n(t+\sigma) - X^n(t)\|_U^2 & \leq 2 \left\| \int_0^t (S(t+\sigma-s) - S(t-s))f(s, X_s^{n-1}) ds \right\|_U^2 \\ & \quad + 2 \left\| \int_t^{t+\sigma} S(t+\sigma-s)f(s, X_s^{n-1}) ds \right\|_U^2 \\ & := I_1 + I_2. \end{aligned}$$

On the one hand,

$$\mathbb{E}I_1 \leq 2M^2 t e^{2\rho t} \mathbb{E} \int_0^t \|(S(\sigma) - Id)f(s, X_s^{n-1})\|_U^2 ds \rightarrow 0$$

when  $\sigma \rightarrow 0$  thanks to the Lebesgue majorant theorem, since, for each  $s$  fixed,

$$S(\sigma)f(s, X_s^{n-1}) \rightarrow f(s, X_s^{n-1}), \quad \|S(\sigma)f(s, X_s^{n-1})\|_U \leq Me^{\rho\sigma} \|f(s, X_s^{n-1})\|_U$$

and

$$\mathbb{E} \int_0^t \|f(s, X_s^{n-1})\|_U^2 ds \leq C \mathbb{E} \int_{-r}^t \|X^{n-1}(s)\|_U^2 ds + \mathbb{E} \int_0^t \|f(s, 0)\|_U^2 ds$$

due to conditions (f.2) and (f.3) and the fact that  $X^{n-1} \in C(-r, T; L^2(\Omega; U))$ .

On the other hand,

$$\begin{aligned} I_2 &\leq 2\sigma M^2 e^{2\rho\sigma} \int_t^{t+\sigma} \|f(s, X_s^{n-1}) - f(s, 0)\|_U^2 ds + 2\sigma M^2 e^{2\rho\sigma} \int_t^{t+\sigma} \|f(s, 0)\|_U^2 ds \\ &\leq 2\sigma M^2 e^{2\rho\sigma} C \int_{-r}^{t+\sigma} \|X^{n-1}(s)\|_U^2 ds + 2\sigma M^2 e^{2\rho\sigma} \int_t^{t+\sigma} \|f(s, 0)\|_U^2 ds, \end{aligned}$$

so that

$$\mathbb{E} I_2 \leq 2\sigma M^2 e^{2\rho\sigma} C \int_{-r}^{t+\sigma} \mathbb{E} \|X^{n-1}(s)\|_U^2 ds + 2\sigma M^2 e^{2\rho\sigma} \int_t^{t+\sigma} \|f(s, 0)\|_U^2 ds \rightarrow 0$$

when  $\sigma \rightarrow 0$ .

Next, we want to show that  $\{X^n\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $C(-r, T; L^2(\Omega; U))$ .

Firstly, for  $t \in [0, T]$  and  $n \in \mathbb{N}$ , since  $X^n = X^{n-1}$  on  $[-r, 0]$ , we have

$$\|X^{n+1}(t) - X^n(t)\|_U^2 \leq tM^2 e^{2\rho t} C \int_0^t \|X^n(s) - X^{n-1}(s)\|_U^2 ds$$

and this implies

$$\mathbb{E} \|X^{n+1}(t) - X^n(t)\|_U^2 \leq tM^2 e^{2\rho t} C \int_0^t \sup_{0 \leq \tau \leq s} \|X^n(\tau) - X^{n-1}(\tau)\|_U^2 ds.$$

Defining

$$G^n(t) = \sup_{0 \leq \theta \leq t} \mathbb{E} \|X^{n+1}(\theta) - X^n(\theta)\|_U^2,$$

we obtain

$$G^n(t) \leq k \int_0^t G^{n-1}(s) ds, \quad \forall n \geq 2$$

for  $k = TM^2 e^{2\rho T} C$ . Consequently, by iteration we can obtain for  $\forall t \in [0, T]$ ,

$$G^n(t) \leq \frac{k^{n-1} T^{n-1}}{(n-1)!} G^1(T), \quad \forall n \geq 2.$$

Since  $X^{n+1}(t) = X^n(t)$ ,  $\forall t \in [-r, 0]$ , the last estimate implies that  $\{X^n\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $C(-r, T; L^2(\Omega; U))$ .

Finally, we check that the limit  $X$  of the sequence  $\{X^n\}_{n \in \mathbb{N}}$  is a solution of (3.1). But this is straightforward, taking into account that  $X^n$  is defined by (3.3) and that  $f$  satisfies (f.2), so that, in particular, when  $n \rightarrow \infty$ ,

$$\mathbb{E} \left\| \int_0^t S(t-s) (f(s, X_s^{n-1}) - f(s, X_s)) ds \right\|_U^2 \leq tM^2 e^{2\rho t} C \int_0^t \mathbb{E} \|X^{n-1}(s) - X(s)\|_U^2 ds \rightarrow 0,$$

and therefore  $X$  is the unique (mild) solution of (3.1). □

#### 4 Existence of weak solutions

In this section we prove that the mild solution to system (3.1) is also a weak solution. First of all we recall the definition of weak solution according to Da Prato and Zabczyk [18].

**Definition 4.1** An  $U$ -valued process  $X(t)$ ,  $t \in [-r, T]$  is called a weak solution of (3.1) if  $X(t) = \varphi(t)$ , for  $t \in [-r, 0]$ , and for all  $\xi \in D(A^*)$  and all  $t \in [0, T]$ ,

$$\begin{aligned} \langle X(t), \xi \rangle_U &= \langle \varphi(0), \xi \rangle_U + \int_0^t (\langle X(s), A^* \xi \rangle_U + \langle f(s, X_s), \xi \rangle_U) ds \\ &\quad + \int_0^t \langle g(s), \xi \rangle_U dS_Q^H(s) \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

**Theorem 4.1** Under the assumptions of Theorem 3.1, the mild solution of (3.1) is also a weak solution.

*Proof* For each  $\xi \in D(A^*)$  it follows that

$$\begin{aligned} &\mathbb{E} \left[ \int_0^t \langle X(s), A^* \xi \rangle_U ds - \int_0^t \langle S(s)\varphi(0), A^* \xi \rangle_U ds - \int_0^t \int_0^s \langle S(s-\tau)f(\tau, X_\tau), A^* \xi \rangle_U d\tau ds \right. \\ &\quad \left. - \int_0^t \int_0^s \langle S(s-\tau)g(\tau), A^* \xi \rangle_U dS_Q^H(\tau) ds \right] \\ &\leq \int_0^t \mathbb{E} \left[ \langle X(s), A^* \xi \rangle_U - \langle S(s)\varphi(0), A^* \xi \rangle_U - \int_0^s \langle S(s-\tau)f(\tau, X_\tau), A^* \xi \rangle_U d\tau \right. \\ &\quad \left. - \int_0^s \langle S(s-\tau)g(\tau), A^* \xi \rangle_U dS_Q^H(\tau) \right] ds \\ &= \int_0^t \mathbb{E} \left[ \left\langle X(s) - S(s)\varphi(0) - \int_0^s S(s-\tau)f(\tau, X_\tau) d\tau \right. \right. \\ &\quad \left. \left. - \int_0^s S(s-\tau)g(\tau) dS_Q^H(\tau), A^* \xi \right\rangle_U \right] ds \\ &= 0. \end{aligned}$$

Thus, for a.e.  $\omega \in \Omega$ , we have

$$\begin{aligned} \int_0^t \langle X(s), A^* \xi \rangle_U ds &= \int_0^t \langle S(s)\varphi(0), A^* \xi \rangle_U ds + \int_0^t \int_0^s \langle S(s-\tau)f(\tau, X_\tau), A^* \xi \rangle_U d\tau ds \\ &\quad + \int_0^t \int_0^s \langle S(s-\tau)g(\tau), A^* \xi \rangle_U dS_Q^H(\tau) ds. \end{aligned} \tag{4.1}$$

Now we use the fact that, for  $\xi \in D(A^*)$ ,  $\frac{d}{dt}S^*(t)\xi = S^*(t)A^*\xi$ . We can obtain

$$\int_0^t \langle S(s)\varphi(0), A^*\xi \rangle_U ds = \int_0^t \langle \varphi(0), S^*(s)A^*\xi \rangle_U ds = \langle S(t)\varphi(0) - \varphi(0), \xi \rangle_U.$$

On the other hand, using Fubini's theorem we have

$$\begin{aligned} & \int_0^t \int_0^s \langle S(s-\tau)f(\tau, X_\tau), A^*\xi \rangle_U d\tau ds \\ &= \int_0^t \int_\tau^t \langle \mathbf{I}_{(0,s]}(\tau)f(\tau, X_\tau), S^*(s-\tau)A^*\xi \rangle_U ds d\tau \\ &= \int_0^t \langle S(t-\tau)f(\tau, X_\tau) - f(\tau, X_\tau), \xi \rangle_U d\tau. \end{aligned}$$

Finally,

$$\begin{aligned} & \int_0^t \int_0^s \langle S(s-\tau)g(\tau), A^*\xi \rangle_U dS_Q^H(\tau) ds \\ &= \int_0^t \int_\tau^t \langle \mathbf{I}_{(0,s]}(\tau)g(\tau), S^*(s-\tau)A^*\xi \rangle_U ds dS_Q^H(\tau) \\ &= \int_0^t \langle g(\tau), S^*(t-\tau)\xi - \xi \rangle_U dS_Q^H(\tau) \\ &= \int_0^t \langle S(t-\tau)g(\tau), \xi \rangle_U dS_Q^H(\tau) - \int_0^t \langle g(\tau), \xi \rangle_U dS_Q^H(\tau). \end{aligned}$$

Therefore by (4.1) for a.e.  $\omega \in \Omega$ , it follows that

$$\begin{aligned} \int_0^t \langle AX(s), \xi \rangle_U ds &= \int_0^t \langle X(s), A^*\xi \rangle_U ds \\ &= \langle S(t)\varphi(0) - \varphi(0), \xi \rangle_U + \int_0^t \langle S(t-\tau)f(\tau, X_\tau) - f(\tau, X_\tau), \xi \rangle_U d\tau \\ &\quad + \int_0^t \langle S(t-\tau)g(\tau), \xi \rangle_U dS_Q^H(\tau) - \int_0^t \langle g(\tau), \xi \rangle_U dS_Q^H(\tau) \\ &= \langle X(t) - \varphi(0), \xi \rangle_U + \int_0^t (\langle X(\tau), A^*\xi \rangle_U + \langle f(\tau, X_\tau), \xi \rangle_U) d\tau \\ &\quad + \int_0^t \langle g(\tau), \xi \rangle_U dS_Q^H(\tau). \end{aligned}$$

Consequently, it follows that almost surely

$$\langle X(t), \xi \rangle_U = \langle \varphi(0), \xi \rangle_U + \int_0^t (\langle X(s), A^*\xi \rangle_U + \langle f(s, X_s), \xi \rangle_U) ds + \int_0^t \langle g(s), \xi \rangle_U dS_Q^H(s),$$

which means that  $X(t)$  is the weak solution to (3.1). □

### 5 Exponential decay of solutions in mean square

In this section we investigate the exponential stability of the mild solutions to (3.1). We impose the following conditions:

**Condition 1** The operator  $A$  is a closed linear operator generating a strongly continuous semigroup  $S(t)$ ,  $t \geq 0$ , on the separable Hilbert space  $U$  and satisfies

$$\|S(t)\|_U \leq Me^{-\lambda t}, \quad \forall t \geq 0, \text{ where } M \geq 1, \lambda > 0.$$

**Condition 2** There exists a constant  $C \geq 0$  such that for any  $x, y \in C(-r, T; U)$  and for all  $t \geq 0$ ,

$$\int_0^t e^{ms} \|f(s, x_s) - f(s, y_s)\|_U^2 \leq C \int_{-r}^t e^{ms} \|x(s) - y(s)\|_U^2 ds \quad \text{for all } 0 \leq m \leq \lambda$$

and

$$\int_0^\infty e^{\lambda s} \|f(s, 0)\|_U^2 ds < \infty.$$

**Condition 3** In addition to assumptions (g.1) and (g.2), assume

$$\int_0^\infty e^{\lambda s} \|g(s)\|_{L^0_Q(K,U)}^2 ds < \infty.$$

The following theorem shows the exponential decay to zero in mean square, with an explicit exponential decay rate  $\gamma$ .

**Theorem 5.1** *In addition to Conditions 1-3, assume that the mild solution  $X(t)$  of system (3.1) corresponding to initial function  $\varphi \in C(-r, 0; L^2(\Omega; U))$ , exists for all  $t \geq -r$ , and that*

$$\lambda^2 > 6CM^2. \tag{5.1}$$

*Then there exists a constant  $\gamma > 0$  such that*

$$\limsup_{t \rightarrow \infty} \left(\frac{1}{t}\right) \log \mathbb{E} \|X(t)\|_U^2 \leq -\gamma.$$

*In other words, every mild solution exponentially decays to zero in mean square.*

*Proof* By (5.1) we can choose an  $\theta > 0$  such that  $\gamma = \lambda - \theta - 6CM^2\lambda^{-1} > 0$ . Then for this  $\gamma$  we have

$$\begin{aligned} \mathbb{E} \|X(t)\|_U^2 &\leq 3\mathbb{E} \|S(t)\varphi(0)\|_U^2 + 3\mathbb{E} \left\| \int_0^t S(t-s)f(s, X_s) ds \right\|_U^2 \\ &\quad + 3\mathbb{E} \left\| \int_0^t S(t-s)g(s) dS_Q^H(s) \right\|_U^2. \end{aligned}$$

Therefore, by Condition 1 and Lemma 2.2 we have

$$\begin{aligned} \mathbb{E} \|X(t)\|_U^2 &\leq 3\mathbb{E} \|S(t)\varphi(0)\|_U^2 + 3M^2 \int_0^t e^{-\lambda(t-s)} ds \int_0^t e^{-\lambda(t-s)} \mathbb{E} \|f(s, X_s)\|_U^2 ds \\ &\quad + 3C_H M^2 t^{2H-1} \int_0^t e^{-2\lambda(t-s)} \|g(s)\|_{L^0_Q(K,U)}^2 ds \end{aligned}$$

$$\begin{aligned} &\leq 3M^2 e^{-2\lambda t} \mathbb{E} \|\varphi(0)\|_U^2 + \frac{3}{\lambda} M^2 \int_0^t e^{-\lambda(t-s)} \mathbb{E} \|f(s, X_s)\|_U^2 ds \\ &\quad + 3C_H M^2 t^{2H-1} \int_0^t e^{-2\lambda(t-s)} \|g(s)\|_{L^0_Q(K,U)}^2 ds \end{aligned}$$

and

$$\begin{aligned} e^{\lambda t} \mathbb{E} \|X(t)\|_U^2 &\leq 3M^2 \mathbb{E} \|\varphi(0)\|_U^2 + \frac{3}{\lambda} M^2 \int_0^t e^{\lambda s} \mathbb{E} \|f(s, X_s)\|_U^2 ds \\ &\quad + 3C_H M^2 t^{2H-1} \int_0^t e^{\lambda s} \|g(s)\|_{L^0_Q(K,U)}^2 ds. \end{aligned}$$

Then, for the chosen parameter  $\theta$ , we obtain

$$\begin{aligned} e^{(\lambda-\theta)t} \mathbb{E} \|X(t)\|_U^2 &\leq 3M^2 e^{-\theta t} \mathbb{E} \|\varphi(0)\|_U^2 + \frac{3}{\lambda} M^2 \int_0^t e^{(\lambda-\theta)s} \mathbb{E} \|f(s, X_s)\|_U^2 ds \\ &\quad + 3C_H M^2 t^{2H-1} e^{-\theta t} \int_0^t e^{\lambda s} \|g(s)\|_{L^0_Q(K,U)}^2 ds. \end{aligned}$$

On the one hand, Condition 3 ensures the existence of a positive constant  $C_1$  such that

$$3C_H M^2 t^{2H-1} e^{-\theta t} \int_0^t e^{\lambda s} \|g(s)\|_{L^0_Q(K,U)}^2 ds \leq C_1 \quad \text{for all } t \geq 0,$$

hence,

$$e^{(\lambda-\theta)t} \mathbb{E} \|X(t)\|_U^2 \leq C_1 + 3M^2 \mathbb{E} \|\varphi(0)\|_U^2 + \frac{3}{\lambda} M^2 \int_0^t e^{(\lambda-\theta)s} \mathbb{E} \|f(s, X_s)\|_U^2 ds.$$

On the other hand, in view of Condition 2, there exists another positive constant  $C_2$  such that

$$\begin{aligned} \int_0^t e^{(\lambda-\theta)s} \mathbb{E} \|f(s, X_s)\|_U^2 ds &\leq 2C \int_{-r}^0 e^{(\lambda-\theta)s} \mathbb{E} \|\varphi(s)\|_U^2 ds + 2C \int_0^t e^{(\lambda-\theta)s} \mathbb{E} \|X(s)\|_U^2 ds \\ &\quad + 2 \int_0^t e^{(\lambda-\theta)s} \|f(s, 0)\|_U^2 ds \\ &\leq C_2 + 2C \int_{-r}^0 \mathbb{E} \|\varphi(s)\|_U^2 ds + 2C \int_0^t e^{(\lambda-\theta)s} \mathbb{E} \|X(s)\|_U^2 ds. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} e^{(\lambda-\theta)t} \mathbb{E} \|X(t)\|_U^2 &\leq C_1 + 3M^2 \lambda^{-1} C_2 + 6CM^2 \lambda^{-1} \int_{-r}^0 \mathbb{E} \|\varphi(s)\|_U^2 ds \\ &\quad + 6CM^2 \lambda^{-1} \int_0^t e^{(\lambda-\theta)s} \mathbb{E} \|X(s)\|_U^2 ds \\ &= C_3 + 6CM^2 \lambda^{-1} \int_0^t e^{(\lambda-\theta)s} \mathbb{E} \|X(s)\|_U^2 ds, \end{aligned}$$

where  $C_3$  is a suitable positive constant. Therefore, Gronwall's lemma implies that

$$e^{(\lambda-\theta)t} \mathbb{E} \|X(t)\|_U^2 \leq C_3 e^{6CM^2 \lambda^{-1} t}.$$

Consequently,

$$\mathbb{E} \|X(t)\|_U^2 \leq C_3 e^{(6CM^2\lambda^{-1}-\lambda+\theta)t} = C_3 e^{-\gamma t}.$$

The proof is complete. □

**Remark 5.1** A remarkable fact is that the decay rate  $\gamma$  is independent of  $H$ . Indeed, in the case of considering a  $Q$ -Brownian motion, *i.e.*, the case  $H = 1/2$ , instead of our sub-fBm  $S_Q^H$ , the condition on  $\lambda$  would be exactly (5.1) (to check this assertion, it is enough to take into account the isometry for classical Wiener integrals). In other words, whenever the stochastic integral is well defined, and under Conditions 1-3, the rate of the exponential decay to zero in mean square is insensitive to the parameter  $H$ .

**Remark 5.2** Theorem 5.1 remains true if we replace the first part of Condition 2 by Condition 4 below.

**Condition 4** For any  $x \in C(-r, T; U)$ ,

$$\int_0^t e^{ms} \|f(s, X_s)\|_U^2 \leq C_2 + 2C \int_{-r}^t e^{ms} \|X(s)\|_U^2 ds \quad \text{for all } 0 \leq m \leq \lambda.$$

### 6 An example

In this section, an example is provided to illustrate the obtained theory.

Let  $K = L^2(0, \pi)$  and  $e_n = \sqrt{\frac{2}{\pi}} \sin(nx)$ ,  $n \in \mathbb{N}$ . Then  $\{e_n\}_{n \in \mathbb{N}}$  is a complete orthonormal basis in  $K$ . Let  $U = L^2(0, \pi)$  and  $A = \frac{\partial^2}{\partial x^2}$  with domain  $D(A) = L_0^1(0, \pi) \cap L^2(0, \pi)$ . Then it is well known that  $Au = -\sum_{n=1}^\infty n^2 \langle u, e_n \rangle_U e_n$  for any  $u \in U$ , and  $A$  is the infinitesimal generator of a strongly continuous semigroup of bounded linear operators  $S(t) : U \rightarrow U$ , where  $S(t)u = \sum_{n=1}^\infty e^{-n^2 t} \langle u, e_n \rangle_U e_n$  and for  $\forall t \geq 0$ ,  $\|S(t)\|_U \leq e^{-t}$ . In order to define the operator  $Q : K \rightarrow K$ , we choose a sequence  $\{\sigma_n\}_{n \geq 1} \subset \mathbb{R}^+$  and set  $Qe_n = \sigma_n e_n$ , and assume that  $\text{tr}(Q) = \sum_{n=1}^\infty \sqrt{\sigma_n} < \infty$ . Define the process  $S_Q^H(s)$  by

$$S_Q^H(t) = \sum_{n=1}^\infty \sqrt{\sigma_n} S_n^H(t) e_n,$$

where  $H \in (1/2, 1)$  and  $\{S_n^H\}_{n \in \mathbb{N}}$  is a sequence of two-sided one-dimensional sub-fractional Brownian motions mutually independent.

Then we consider the following stochastic evolution equation:

$$\begin{cases} du(t, x) = [\frac{\partial^2}{\partial x^2} u(t, x) + b(t) \cdot u(t, x(t-r))] dt \\ \quad + g(t) dS_Q^H(t), \quad t \in [0, T], x \in [0, \pi], \\ u(t, 0) = u(t, \pi) = 0, \quad t \in [0, T], \\ u(t, x) = \varphi(t, x), \quad t \in [-\tau, 0], x \in [0, \pi], \end{cases} \tag{6.1}$$

where  $r > 0$  and  $b, g : \mathbb{R}^+ \rightarrow \mathbb{R}$  are continuous functions such that  $g$  satisfies Condition 3 above and  $b$  satisfies

$$\int_0^\infty e^{\lambda s} |b(s)|^2 ds < \infty.$$

Observe that the fact  $\int_0^\infty e^{\lambda s} |b(s)|^2 ds < \infty$  implies that  $b(t)$  is bounded for all  $t \geq 0$ . Denote by  $b_0$  the smallest upper bound of the  $b$ . Taking

$$f(t, \varphi_t)(\eta) = \sin(t) \cdot \varphi(\eta_t).$$

Thus, for any  $x, y \in C(-r, T; U)$ , and for all  $t \geq 0$ , one has

$$\int_0^t \|f(s, x_s) - f(s, y_s)\|_U^2 ds \leq b_0^2 \int_{-r}^t \|x(s) - y(s)\|^2 ds$$

and

$$\int_0^t e^{ms} \|f(s, x_s) - f(s, y_s)\|_U^2 ds \leq b_0^2 \int_{-r}^t e^{ms} \|x(s) - y(s)\|_U^2 ds.$$

Then we can check that there exists a unique mild solution to (6.1) according to Theorem 3.1.

If we assume, in addition, that

$$b_0^2 < \frac{1}{6},$$

then any mild solution to (6.1) decays exponentially to zero in mean square.

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

**Author details**

<sup>1</sup>School of Information and Mathematics, Yangtze University, Jingzhou, 434023, China. <sup>2</sup>School of Mathematics and Statistics, Chongqing University, Chongqing, 400044, China. <sup>3</sup>School of Mathematics and Information Sciences, Guangzhou University, Guangzhou, 510006, China.

**Acknowledgements**

The authors would like to thank the referees for their detailed comments and valuable suggestions which considerably improved the presentation of the paper. This research is partially supported by the NNSF of China (No. 11271093), the Science Research Project of Hubei Provincial Department of Education (No. Q20141306) and Open Research Fund Program of Institute of Applied Mathematics, Yangtze University (KF1509).

Received: 3 August 2014 Accepted: 7 January 2015 Published online: 20 February 2015

**References**

1. Bojdecki, TLG, Gorostiza, LG, Talarczyk, A: Sub-fractional Brownian motion and its relation to occupation times. *Stat. Probab. Lett.* **69**, 405-419 (2004)
2. Bojdecki, TLG, Gorostiza, LG, Talarczyk, A: Limit theorems for occupation time fluctuations of branching systems I: long-range dependence. *Stoch. Process. Appl.* **116**, 1-18 (2006)
3. Bojdecki, TLG, Gorostiza, LG, Talarczyk, A: Some extensions of fractional Brownian motion and sub-fractional Brownian motion related to particle systems. *Electron. Commun. Probab.* **12**, 161-172 (2007)
4. Tudor, C: Some properties of the sub-fractional Brownian motion. *Stochastics* **79**, 431-448 (2007)
5. Tudor, C: Some aspects of stochastic calculus for the sub-fractional Brownian motion. *An. Univ. Bucur., Mat.* **57**, 199-230 (2008)
6. Tudor, C: Inner product spaces of integrands associated to sub-fractional Brownian motion. *Stat. Probab. Lett.* **78**, 2201-2209 (2008)
7. Tudor, C: Sub-fractional Brownian Motion as a Model in Finance. University of Bucharest (2008)
8. Tudor, C: On the Wiener integral with respect to a sub-fractional Brownian motion on an interval. *J. Math. Anal. Appl.* **351**, 456-468 (2009)
9. Tudor, C: Berry-Esseen bounds and almost sure CLT for the quadratic variation of the sub-fractional Brownian motion. *J. Math. Anal. Appl.* **375**, 667-676 (2011)
10. Yan, L, Shen, G: On the collision local time of sub-fractional Brownian motions. *Stat. Probab. Lett.* **80**, 296-308 (2010)



11. Young, LC: An inequality of the Hölder type connected with Stieltjes integration. *Acta Math.* **67**, 251-282 (1936)
12. Yan, L, Shen, G, He, K: Itô's formula for a sub-fractional Brownian motion. *Commun. Stoch. Anal.* **5**, 135-159 (2011)
13. Shen, G, Chen, C: Stochastic integration with respect to the sub-fractional Brownian motion with  $H \in (0, 1/2)$ . *Stat. Probab. Lett.* **82**(2), 240-251 (2012)
14. Mendy, I: A Stochastic differential equation driven by a sub-fractional Brownian motion. Preprint (2010)
15. Dzharidze, K, Van Zanten, H: A series expansion of fractional Brownian motion. *Probab. Theory Relat. Fields* **103**, 39-55 (2004)
16. Mendy, I: Parametric estimation for sub-fractional Ornstein-Uhlenbeck process. *J. Stat. Plan. Inference* **143**, 663-674 (2013)
17. Nualart, D: *The Malliavin Calculus and Related Topics*, 2nd edn. Springer, Berlin (2006)
18. Da Prato, G, Zabczyk, J: *Stochastic Equations in Infinite Dimensions*. Cambridge University Press, Cambridge (1992)

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

Submit your next manuscript at ▶ [springeropen.com](http://springeropen.com)

---