

EEB-S-13-00485 2nd revision

Singlet oxygen scavenging by leaf flavonoids contributes to sunlight acclimation in *Tilia platyphyllos*

Petra Majer¹, Susanne Neugart², Angelika Krumbein², Monika Schreiner² and Éva Hideg^{3,1,*}

¹ Institute of Plant Biology, Biological Research Centre, Hungarian Academy of Sciences, PO Box 521, H-6701 Szeged, Hungary; email: pmajer@brc.hu

² Department Quality, Leibniz-Institute of Vegetable and Ornamental Crops, Theodor-Echtermeyer-Weg 1, 14979 Großbeeren, Germany; emails: neugart@igzev.de (S.N.), krumbein@igzev.de (A.K.), Schreiner@igzev.de (M.S.)

³ Institute of Biology, University of Pécs, Ifjúság u. 6, H-7624 Pécs, Hungary; email: ehideg@gamma.ttk.pte.hu

* corresponding author: Dr. Éva Hideg

email: ehideg@gamma.ttk.pte.hu

tel.: +3672 503600 ext. 24229

1 **Abstract**

2 Both high photosynthetically active radiation (PAR) and ultraviolet radiation (UV) are
3 capable of causing photooxidative stress, but leaves are equipped with an array of protective
4 mechanisms making life under full sunlight possible. Comparing acclimation strategies of
5 *Tilia platyphyllos* leaves we found that sun leaves were better protected against stress than
6 shade leaves by having (i) more efficient regulated non-photochemical quenching (ii) a
7 higher capacity to neutralize singlet oxygen, a reactive oxygen species known to be capable
8 of promoting oxidative damage by excess PAR and (iii) containing more UV absorbing
9 pigments. HPLC-MSⁿ analysis showed both quantitative and qualitative differences in higher
10 flavonoid contents: Sun leaves contained 4.2-times more flavonoids than shade leaves and the
11 quercetin:kaempferol ratio was also higher in the former. In addition, sun leaves also
12 contained significant amounts of myricetin, which was detectable only in traces in shade
13 leaves. Flavonols were mainly present as rhamnosides and *in vitro* tests of these compounds
14 showed that quercetin and myricetin glycosides were much better singlet oxygen antioxidants
15 than kaempferol glycosides. Thus a shift from monohydroxylated flavonols (kaempferol
16 derivatives) towards dihydroxylated quercetin or trihydroxylated myricetin improves the
17 singlet oxygen targeted antioxidant potential of leaves and potentially improves protection
18 against photoinhibition by high PAR. On the other hand, experiments with pure test
19 compounds also showed that multihydroxylated flavonol rhamnosides do not have better UV-
20 B absorption than monohydroxylated ones. Thus the observed difference in flavonoid
21 composition is not expected to contribute to the observed better UV-B absorbing capacity of
22 sun leaves. Our data suggest that responses to high intensity PAR and to solar UV-B are
23 closely onnected and that UV-inducible flavonols play a key role in the successful
24 acclimation of sun leaves to high PAR as efficient singlet oxygen antioxidants.

25

26

27 **Key words**

28 singlet oxygen; sunlight acclimation; UV; flavonoid; quercetin; kaempferol; myricetin

29

30 **1. Introduction**

31 Plants grown in a natural environment are exposed to variable light conditions both in
32 terms of photosynthetically active radiation (PAR, 400-700 nm) and solar ultraviolet (UV,
33 290-400 nm) radiation. Light conditions which individual plant leaves experience vary

1 significantly depending on weather conditions, position, canopy shading, etc. Reflecting
2 long-term acclimative adaptation mechanisms, leaves grown in shaded areas of a plant differ
3 from the sun grown leaves in morphology, physiology, chemical composition and anatomy
4 (Boardman, 1977; Lichtenthaler et al. 1981; Hallik et al., 2012). Photoprotective mechanism
5 are important constituents of light acclimation and high solar radiation tolerance includes
6 both physical and biochemical defenses (Niyogi, 1999; Takahashi et al. 2011). The former
7 include leaf and chloroplast movements, epidermal screening (Solovchenko and Merzlyak,
8 2008) and dissipation of excess energy (Müller et al. 2001). Chemical defense pathways
9 equip the leaf for situations when physical mechanisms fail to protect the photosynthetic
10 electron transport system from over-excitation and, potentially from photoinhibition (Long et
11 al. 1994; Powles, 1984) via the formation of damaging reactive oxygen species (ROS).
12 During photoinhibition by excess PAR singlet oxygen ($^1\text{O}_2$) is photoproduced by the reaction
13 centre chlorophyll of Photosystem (PS) II (Macpherson et al. 1993; Hideg et al. 1994).
14 Shorter wavelength (higher energy) solar UV-B radiation (290-315 nm) is also a potential
15 inducer of oxidative stress (Jansen et al. 1998). Laboratory experiments with high UV doses
16 showed the presence of both $^1\text{O}_2$ and superoxide anion radicals ($\text{O}_2^{\cdot-}$) in irradiated spinach
17 leaves (Barta et al. 2004), although the damage initiated by $^1\text{O}_2$ and derived reactive
18 compounds was more characteristic to visible light induced photoinhibition (Hideg et al.,
19 2000, Barta et al., 2004). While biochemical defence against $\text{O}_2^{\cdot-}$ is mainly achieved by a
20 specific enzyme, superoxide dismutase, which is included in the supportive water-water cycle
21 (Asada, 1999a), damage by $^1\text{O}_2$ can only be controlled by non-enzymatic antioxidants.
22 Singlet oxygen quenching plant antioxidants include lipid-soluble compounds, e.g. β -
23 carotene, α -tocopherol, plastoquinones, and water-soluble molecules, such as ascorbate,
24 vitamin B6 and flavonoids (Triantaphylides and Havaux, 2009).

25 Flavonoids are secondary plant metabolites within the group of phenolics and are
26 present in plants mainly as glycosides (see Hernandez et al. 2009 for a recent review).
27 Transcriptome analyses of UV-B photoreceptor (UV RESISTANCE LOCUS8 (UVR8);
28 Oravecz et al. 2006) mutants exposed to low fluence rates of UV-B indoors have shown that
29 UVR8 is required for the induction of genes with important functions in UV protection
30 including flavonoid synthesis pathways (Brown et al. 2005; Favory et al., 2009). Field studies
31 of silver birch leaves have shown that besides UV-B, solar UV-A also regulates gene
32 expression and the accumulation of flavonoids (Morales et al. 2010) and a recent comparative
33 study of genes encoding flavonoid biosynthesis proteins in sun exposed wild type and uvr8

1 mutant *Arabidopsis* leaves suggested an interaction of UVR8 with other photoreceptors to
2 modulate UV-A responses in the presence of UV-B (Morales et al. 2013).

3 Antioxidant properties of flavonoids have been extensively studied *in vitro* due to
4 their nutritional and medical relevance (Rice-Evans et al. 1995, 1996, Pietta 2000; Scalbert et
5 al. 2005). Flavonoids in leaves are typically regarded as UV-B regulated compounds, because
6 higher UV-B radiation leads to an enhanced biosynthesis and accumulation through the
7 induction of the phenylpropanoid pathway (Jansen et al. 2008; Kim et al. 2008; Zhang and
8 Björn, 2009). Increased leaf flavonoid production in response to high solar irradiances or
9 UV-B radiation has primarily been considered as physical defence mechanism, due to the UV
10 screening function of these compounds, but flavonoids are also known as antioxidants
11 (Harborne 1986).

12 Whether the primary role of flavonoids in UV defence is direct screening or radical
13 scavenging is a controversial issue and the general view of flavonoids acting predominantly
14 as radiation shielding screen has been challenged by experimental evidence of their strong
15 antioxidant activities and presence in mesophyll layers and chloroplasts (for reviews see
16 Hernandez et al. 2009; Winkel-Shirley 2002, Agati and Tattini 2010; Agati et al. 2012,
17 Brunetti et al. 2013). Flavonoids are synthesized by a multi-enzyme complex located on the
18 cytoplasmic surface of the endoplasmic reticulum and are transported to the vacuolar
19 compartment or to the cell wall (Winkel-Shirley 2002). However, flavonoids were also found
20 in the chloroplasts of many vascular plants (Saunders and McClure 1976) and recently their
21 *in situ* synthesis was also proved in isolated kidney bean chloroplasts (Zaprometov and
22 Nicolaeva, 2003).

23 In addition to good total antioxidant capacities which have been the focus of
24 nutritional studies for decades (Rice-Evans et al. 1996, Sekher Pannala et al. 2001, Brunetti et
25 al. 2013, Chen and Chen 2013), flavonoids were shown to be good $^1\text{O}_2$ antioxidants *in vitro*
26 (Tournaire, 1993) as well as in leaves of a Mediterranean shrub, *Phillyrea latifolia* (Agati et
27 al. 2007). In plants $^1\text{O}_2$ is primarily associated with damage by excess PAR and the well-
28 documented inducibility of flavonoids by UV suggests a link between responses to these two
29 spectrally distinct environmental factors. The aim of the present study was to test the
30 hypothesis whether ambient solar UV radiation would promote defence against
31 photoinhibition *in vivo*, via enhancing flavonoid synthesis. We studied large leaved linden
32 (*Tilia platyphyllos*) leaves from sun exposed and shaded areas of the same tree. Sun and
33 shade leaves have already been studied extensively for decades but the novel aspect of our
34 work is a comparative study of their $^1\text{O}_2$ antioxidant capacities and UV absorbing

1 characteristics including both leaf extracts and pure test compounds representing flavonoids
2 in which the two leaf types differ mostly. Members of the *Tilia* genus have high flavonoid
3 content not only in flowers but in leaves and bracts as well (Toker et al. 2001) making this
4 species a good model for the present study.

6 **2. Materials and Methods**

7 **2.1. Plant material**

8 Large-leaved linden (*Tilia platyphyllos* Scop.) leaves were collected in Szeged, Hungary
9 (46.2460 N, 20.1673 E) in July from sun exposed and predominantly shaded parts of the
10 same tree (referred to as sun leaves and shade leaves). The leaves of the two groups were
11 carefully selected to avoid variance originating from size, age or other factors. On cloudless
12 days, typical photon flux densities reaching sunny and shaded areas at noon were 1700-1800
13 and 80-100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PAR, respectively, as measured with a Cole Parmer radiometer
14 (Cole-Parmer Instrument Co. Ltd., London, UK). Typical ambient daily doses of biologically
15 effective UV-B radiation on the Northern hemisphere (at N 46°) in summer are around 7.5 kJ
16 $\text{m}^{-2} \text{d}^{-1}$ (Bassman et al. 2001).

17 Chlorophyll fluorescence measurements were carried out on detached leaves following a 30
18 min dark adaptation period (see below). These measurements were followed by freeze-drying
19 of the leaves (Christ Alpha 2-4 LD Freeze dryer, SciQuip Ltd, UK) at -85°C and 0.060 mbar
20 and dry weight (DW) determination. Freeze dried samples were kept at -80°C until use.

22 **2.2. Photochemical yield and non-photochemical quenching measurements**

23 Photochemical yield and non-photochemical quenching were determined by chlorophyll *a*
24 fluorescence measurements of sun and shade leaves using the MAXI-version of the Imaging-
25 PAM (Heinz Walz GmbH, Effeltrich, Germany). Before starting the measurements, detached
26 *Tilia platyphyllos* leaves were kept in darkness for 30 min. Of these dark adapted samples,
27 minimum and maximum fluorescence yields (F_0 and F_m , respectively) were determined
28 before and after a saturating pulse, respectively. Following this, F and F_m' yields were
29 obtained in a similar way at the end of short (30 sec) blue actinic light irradiations of stepwise
30 increased intensity (Schreiber 2004). The quantum yields of PS II photochemistry,
31 $Y(\text{II}) = (F_m' - F) / F_m$, non-regulated dissimilative processes, $Y(\text{NO}) = F / F_m$ and regulated
32 energy dissipation, $Y(\text{NPQ}) = F / F_m' - Y(\text{NO})$, characteristic to each illumination step, were
33 calculated according to Klughammer and Schreiber (2008). As these quantum yields

1 represent the partitioning of absorbed energy, $Y(\text{II}) + Y(\text{NO}) + Y(\text{NPQ}) = 1$ at each radiation
2 wavelength.

3

4 **2.3. Chemicals**

5 Myricetin, dihydroquercetin, kaempferol quercetin-3-glucoside, kaempferol-3-glucoside,
6 quercetin-3-rhamnoside and myricetin-3-rhamnoside standards were purchased from Carl
7 Roth GmbH, (Karlsruhe, Germany). Kaempferol-3,7-dirhamnoside was purchased from
8 PhytoLab GmbH & Co. KG (Vestenbergsgreuth, Germany).

9

10 **2.4. Flavonol aglycon and flavonol glycoside content determination**

11 For the flavonol aglycones 0.5 g of lyophilised leaf sample was hydrolysed with 50 %
12 aqueous methanol and 1.6 M HCl in double determination (Krumbein et al. 2007). After
13 refluxing at 90 °C for 2 hours, the extract was cooled down to room temperature, adjusted to
14 100 mL and sonicated for 5 min. The extract was then filtered through a 0.45 µm PTFE filter
15 for HPLC analysis.

16 The flavonol glycosides were prepared using a modified method following Schütz et al.
17 (2004). For this a lyophilised leaf sample (0.5 g) was extracted with 15 mL of 60 % aqueous
18 methanol on a magnetic stirrer plate for 1.5 hours in double determination. The extract was
19 filtered through a fluted filter and subsequently evaporated to dryness. The residue was
20 dissolved in 5 mL of distilled water and then filtered through a CME filter (cellulose mixed
21 ether-membrane filter) for the HPLC analysis.

22 An HPLC series 1100 from Agilent (Waldbronn, Germany) consisting of a degaser, binary
23 pump, autosampler, column oven and photodiode array detector was used to quantify the
24 flavonol aglycones and flavonoid glycosides. For identification purposes, an ion trap mass
25 spectrometer (Agilent series 1100 MSD) was used with ESI as an ion source in negative
26 ionisation mode. Nitrogen was the dry gas (12 l/min, 350 °C) and nebulizer gas (40 psi).
27 Helium was the inert collision gas in the ion trap. Details of the analysis of flavonol
28 aglycones and flavonol glycosides are given below.

29 The flavonol aglycones were quantified after acid hydrolysis using a modified HPLC-DAD-
30 ESI-MSⁿ method (Krumbein et al. 2007). The extracts were separated on a Prodigy (ODS 3,
31 150 x 3.0 mm, 5 µm, 100 Å) column (Phenomenex, Aschaffenburg, Germany) with a security
32 guard C18 (ODS 3, 4 x 3.0 mm, 5 µm, 100 Å) at a temperature of 25 °C using a
33 water/acetonitrile gradient. Solvent A consisted of 99.5 % water and 0.5 % acetic acid;
34 solvent B contained 100 % acetonitrile. The following gradient was used for eluent B: 30-

1 35 % (0-5 min), 35-39 % (5-17 min), 39-90 % (17-21 min), 90 % isocratic (21-26 min), 90-
2 30 % (26-29 min), 30 % isocratic (29-34 min). Flow was performed using 0.3 mL min^{-1} , and
3 the measured detector wavelength was 370 nm. The standards dihydroquercetin, kaempferol
4 and myricetin were used to obtain an external calibration curve in the range of 0.1 to 10 mg
5 100 mL^{-1} . The total flavonol concentration was calculated as the sum of the concentration of
6 the individual flavonol aglycones quercetin, kaempferol and isorhamnetin. Quercetin,
7 kaempferol and myricetin were identified as deprotonated molecular ions and characteristic
8 mass fragment ions by HPLC-DAD-ESI-MS². The mass optimisation was performed for
9 quercetin $[\text{M-H}]^- m/z$ 301.

10 The flavonol glycosides were analysed using the same column and eluent composition as that
11 for aglycone determination. The following gradient was used for eluent B (100 %
12 acetonitrile) at a temperature of 30 °C: 5-7 % (0-12 min), 7-9 % (12-25 min), 9-12 % (25-
13 45 min), 12-15 % (45-100 min), 15 % isocratic (100-150 min), 15-50 % (150-155 min), 50 %
14 isocratic (155-165 min), 50-5 % (165-170 min), 5 % isocratic (170-175 min) (Schmidt et al.
15 2010). The flow was performed using 0.4 mL min^{-1} , and the measured detector wavelength
16 for the quantification was set at 370 nm for non-acylated flavonol glycosides and 330 nm for
17 acylated flavonol glycosides. The standards quercetin-3-glucoside, kaempferol-3-glucoside,
18 quercetin-3-rhamnoside, myricetin-3-rhamnoside and kaempferol-3,7-dirhamnoside were
19 used to obtain an external calibration curve in the range of 0.1 to 10 mg 100 mL^{-1} . Mass
20 optimisation for the ion optics of the mass spectrometer was performed for quercetin m/z 301.
21 In addition, arbitrary m/z 1000 was used as the target mass in auto mode to include higher
22 mass fragments. The MSⁿ experiments were performed in auto or manual mode up to MS⁴ in
23 a scan from m/z 200 to 2000. Mass fragments used for identification are described in Schmidt
24 et al. (2010).

25

26 **2.5. Singlet oxygen scavenging capacity measurements**

27 Freeze dried samples were extracted using MeOH : H₂O (in 1 : 1 ratio) then centrifuged
28 ($6,000 \times g$ for 10 min at 4 °C) and supernatants were used for measurements. Methanol
29 extractions were used because these provided higher extraction yields than water only (Hideg
30 and Majer, 2010) and made comparisons between ROS scavenging capacity measurements
31 and flavonoid analysis possible.

32 Measurements were based on the leaf extract's ability to scavenge ¹O₂ and therefore prevent
33 the oxidation of 1,3-diphenylisobenzofuran (DPBF) dye (Young et al. 1973). The reaction
34 mixture contained 20 μM methylene blue dye as ¹O₂ source and 100 μM DPBF in 30%

1 methanol. DPBF oxidation was followed by a decrease of absorption at 410 nm using a
2 Shimadzu (UV-1601) spectrophotometer. The $^1\text{O}_2$ scavenging capacity of leaf extracts and
3 test compounds were compared to that of Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-
4 carboxylic acid) and expressed as μM Trolox equivalents mg^{-1} DW and μM Trolox
5 equivalents μM^{-1} flavonoid, respectively.

6 7 **2.6. Absorbance spectra measurements**

8 Ground freeze dried samples of sun and shade *T. platyphyllos* leaves were extracted into
9 acidified methanol (MeOH : H₂O : HCl in 79 : 20 : 1 ratio), vortexed and centrifuged (10,000
10 x g, 10 min, 4 °C) and clear supernatants (1 mg DW mL⁻¹) were used for measuring
11 absorbance spectra (Mirecki and Teramura, 1984) with a Shimadzu UV-1601
12 spectrophotometer. Pure flavonoids were extracted into methanol and the solutions were
13 diluted to 20 $\mu\text{g mL}^{-1}$

14 15 **2.7. Statistical analysis**

16 For photosynthesis and singlet oxygen antioxidant measurements five leaves were collected
17 from the sun exposed and five from shaded parts of the same tree. Antioxidant capacity
18 measurements of pure flavonol compounds were repeated three times. Results are given in
19 tables or in figures as means \pm standard deviations. The significance of differences between
20 data sets was assessed using unpaired Students *t*-tests and significantly different data sets
21 ($p < 0.05$) are labelled with asterisks. Data were analysed with GraphPad QuickCalcs
22 (GraphPad Software Inc., La Jolla, CA, USA). Graphs were created with SigmaPlot 12
23 (Systat Software Inc., San Jose, CA, USA).

24 25 **3. Results and Discussion**

26 **3.1. Photosynthetic response to solar radiation: sun leaves have more efficient energy** 27 **dissipation than shade leaves**

28 The partitioning of absorbed radiation energy in leaves provides information on
29 physical photoprotective capacities. Abilities of sun and shade *Tilia platyphyllos* leaves to
30 tolerate high solar irradiances were characterized by measuring ratios of energy used in
31 photochemistry and dissipated in non-photochemical quenching according to Klughammer
32 and Schreiber (2008). Ratios of photochemical yield $Y(\text{II})$ (circles in Fig.1A) and non-
33 photochemical yields $Y(\text{NPQ})+Y(\text{NO})$ (squares in Fig.1A) show that at irradiances higher
34 than 80 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PAR sun leaves had higher photochemical yields and put relatively less

1 quanta into alternative dissipation pathways than shade leaves. At relatively low PAR (80
2 $\mu\text{mol m}^{-2} \text{s}^{-1}$) sun and shade leaves dissipated similar ratios of absorbed energy in non-
3 regulated processes (Fig.1B). Sun leaves acclimated to higher PAR, for example to 510 or
4 $1050 \mu\text{mol m}^{-2} \text{s}^{-1}$, by maintaining the same Y(NO) as under low PAR. Under high PAR,
5 which is the natural environment of these leaves Y(II) decreased to nearly zero at the expense
6 of Y(NPQ) and all absorbed energy was dissipated (see upper row in Fig.1B). On the other
7 hand, when shade leaves were subjected to high PAR, Y(NPQ) pathways were not capable of
8 balancing the decrease in Y(II) and non regulated quenching increased (lower row in Fig.1B).
9 The same tendency was observed by Hallik et al. (2012) in *Tilia cordata* leaves grown at
10 different PAR, where shade leaves had higher values of Y(NO) but lower values of Y(II)
11 compared to leaves grown at higher irradiance levels.

12 Biochemically, Y(NPQ) is realized through regulated reactions such as dissipating
13 absorbed energy in the light harvesting complex (Demmig-Adams and Adams, 1996a,
14 1996b) and energy transfer from the reaction centre chlorophyll to zeaxanthin (Müller et al.,
15 2001). High Y(NO) levels may indicate photodamage (Klughammer and Schreiber, 2008). In
16 their natural growth site, shade leaves are sometimes exposed to high irradiances (e.g. sudden
17 sun flecks), so in this case these fluorescence parameters measured after exposure to high
18 PAR reflect their capacity to tolerate rare fluctuations in radiation. On the other hand, sun
19 leaves are typically grown in high irradiance radiation conditions and therefore have
20 acclimated to these conditions. In our study, *T. platyphyllos* sun leaves acclimated to high
21 light with efficient physical defence pathways to avoid oxidative damage due to over-
22 excitation. Hallik et al. (2011) proposed that as high light radiation grown *T. cordata* leaves
23 have a higher capacity for photochemistry, this could provide them with a protection against
24 photodamage, while shade leaves having high values of Y(NO) are in a greater danger from
25 radiation caused stress effects. Szöllősi et al. (2010) concluded the same for the sun leaves of
26 another tree species *Quercus petraea*, showing that these leaves were able to prevent
27 photodamage by having higher Y(II) and Y(NPQ). We studied the physiological need of sun
28 and shade *T. platyphyllos* leaves for additional protection by analyzing their flavonoid
29 content. This was followed by a comparison of UV absorbing and specific $^1\text{O}_2$ antioxidant
30 capacities of sun and shade leaf extracts and pure flavonoid test compounds identified as leaf
31 components.

32

33 ***3.2. Flavonoid response to solar radiation: quantitative and qualitative differences between***
34 ***sun and shade leaves***

1 After aglyconation of flavonoids, three flavonols were identified by HPLC-DAD-ESI-
2 MS/MS: myricetin, quercetin and kaempferol (Table 1). In sun leaves the total flavonoid
3 content (21.5 mg g⁻¹ DW) was 4.2-times higher than in shade leaves (5.1 mg g⁻¹ DW), and
4 significantly higher concentrations were detected of quercetin and myricetin (Table 1). In sun
5 leaves, quercetin was the most abundant flavonol, followed by kaempferol and myricetin. In
6 shade leaves the quantity of kaempferol exceeded that of quercetin, and myricetin could be
7 detected only in trace amounts. The difference in quercetin concentrations was the most
8 notable distinction between the two leaf types: sun leaves contained 22-times more of this
9 flavonol. Toker et al. (2001) reported nine different flavonoid compounds in leaf extracts of
10 large-leaved linden (three of them unidentified), which were various kaempferol and
11 quercetin glycosides. Among these, quercetin compounds were more abundant, representing
12 ca. 62% of the flavonoid content, which is very similar to the proportion of this flavonol
13 found among the aglycones in our sun leaf samples (70%) (Table 1). To our best knowledge,
14 ours is the first observation of myricetin in *Tilia platyphyllos* leaves.

15 A high quercetin to kaempferol ratio in response to either UV-B radiation or sunlight
16 has already been reported in a large variety of plants including petunia, Arabidopsis, white
17 mustard, white clover and bilberry (Ryan et al. 1998, 2001; Reifenrath and Müller, 2007;
18 Hofmann et al. 2000; Jaakola et al. 2004). A common response to UV light is a more
19 pronounced increase in quercetin, while the amount of kaempferol increases to a smaller
20 extent or remains unaffected (Winter and Rostás 2008; Reifenrath and Müller 2007; Hofmann
21 et al. 2000). In agreement with these published data we detected similar amounts of
22 kaempferol in *Tilia platyphyllos* sun and shade leaves and 22-times more quercetin in sun
23 leaves than in shade leaves (Table 1).

24 In addition to overall ratios of flavonols detected in acidified leaf samples, their
25 naturally occurring glycosides were identified using HPLC-DAD-ESI-MSⁿ (Table 2). We
26 found that the most prominent flavonoid glycosides in *Tilia platyphyllos* leaves were present
27 as rhamnosides (Tables S1 and S2). Shade leaves were dominated by two kaempferol
28 glycosides, kaempferol-3-*O*-rutinoside-7-*O*-rhamnoside and kaempferol-3-*O*-rhamnoside-7-
29 *O*-rhamnoside, which made-up approximately 70% of total flavonoid content (Table S1).
30 This result is in agreement with the identification of kaempferol as the main aglycone in
31 acidified extracts from shade leaves (Table 1). Also matching the aglycone pattern,
32 kaempferol was less dominant in sun leaves which contained mainly myricetin and quercetin
33 glycosides (Table 2). Eleven compounds were identified in extracts, but most were present in
34 trace amounts only. The quercetin-3-*O*-rhamnoside-7-*O*-rhamnoside made-up approximately

1 18% of glycosylated flavonols in sun exposed leaves (Table S2). This flavonol glycoside has
2 already been reported to be the main flavonoid in *T. platyphyllos* leaves (Toker et al. 2001)
3 although the dependence on light conditions during growth was not analysed. In our leaf
4 samples kaempferol glycosides were partly acylated with coumaric acid. Some quercetin
5 glycosides were acylated with caffeic acid (Tables S1 and S2). Tiliroside, a coumaric acid
6 acylated kaempferol glycoside was also identified by Aguirre-Hernández et al. (2010) in
7 flowers and bracts of different cultivars of *Tilia americana* var. *mexicana*.

8 9 **3.3. Sun leaves have more and better singlet oxygen neutralizing antioxidant flavonols** 10 **than shade leaves**

11 Singlet oxygen scavenging capacities of sun and shade *Tilia platyphyllos* leaves
12 were measured *in vitro*, using a photometric assay. As shown in Fig.2A, methanol-water
13 extracts from sun leaves had a 5-fold higher capacity to quench $^1\text{O}_2$ than extracts from shade
14 leaves. To estimate contributions of flavonols to $^1\text{O}_2$ antioxidant capacities of sun and shade
15 leaves, we tested pure flavonols using the same methodology. Figure 2B shows that
16 flavonoid aglycones were stronger scavengers of $^1\text{O}_2$ than Trolox (6-hydroxy-2,5,7,8-
17 tetramethylchroman-2-carboxylic acid), which is a vitamin E derivative frequently used for
18 quantifying antioxidant capacities due to its strong reactivity to $^1\text{O}_2$ (Jung and Min 2009).
19 Among the three studied flavonol aglycones, myricetin was the strongest $^1\text{O}_2$ antioxidant
20 while quercetin had less scavenging capacity and kaempferol was the weakest.

21 The general structure of major flavonols detected in *Tilia platyphyllos* leaves is shown
22 in Fig.3. The top row demonstrates that basic structures differ only in the number of hydroxyl
23 groups on the B-ring: kaempferol having one hydroxyl group at position 4', quercetin having
24 two groups at positions 3' and 4' and myricetin having three groups at the 3', 4' and 5'-
25 positions. According to earlier studies on structure-dependent antioxidant activity relations of
26 flavonoids, hydroxyl groups located on the B-ring enhance reactivity to superoxide radicals
27 (Hu et al. 1995) or to a synthetic free radical (2,2'-azinobis-(3-ethylbenzothiazoline-6-
28 sulfonic acid, ABTS) used in assessing total antioxidant capacities (Heim et al. 2002).
29 Tournaire et al. (1993) studied flavonoid aglycone reactivities to $^1\text{O}_2$ *in vitro* and found a
30 similar importance of B-ring hydroxyl groups for physical quenching. In accordance, we
31 found that myricetin which contains three of these hydroxyl groups but was not included in
32 the above study had the highest reactivity to $^1\text{O}_2$. In the study of Tournaire et al. (1993)
33 quercetin, with a catechol structure on ring B had almost 4-times higher reactivity to $^1\text{O}_2$ was
34 than kaempferol lacking this structure. Our measurements gave a smaller ratio, 1.3 (Fig.2B)

1 which could be due to differences in the applied solvents (methanol and deuterated methanol,
2 respectively). In addition, our photometric method measures a decrease in $^1\text{O}_2$ concentration
3 and thus it is more likely to detect only the chemical quenching of this ROS via oxidation,
4 while Tournaire et al. determined total reactivities including both chemical and physical
5 quenching.

6 The importance of a hydroxyl group activating the double bond of the C-ring in
7 flavonols in chemical $^1\text{O}_2$ quenching was established by Tournaire et al. (1993) and it is
8 supported by a strong decrease in $^1\text{O}_2$ neutralizing capacity when the hydroxyl group is
9 glycosylated with a sugar moiety in flavonol glycosides. In general, glycosylated forms are
10 weaker antioxidants than their respective aglycons (Ratty and Das 1988; Montoro et al.
11 2005). Differences in $^1\text{O}_2$ antioxidant capacities are shown in Fig.2B where flavonol
12 aglycons are compared to their corresponding glycosides. Flavonol rhamnosides were
13 chosen due to the abundance of this sugar moiety in large leaved linden flavonoids. In
14 addition to confirming the lower $^1\text{O}_2$ quenching of glycosides compared to corresponding
15 aglycons, as already reported for quercetin and quercetin-rutinoside (Tournaire et al. 1993),
16 the aim of these measurements was to test whether the observed advantage of quercetin and
17 myricetin over kaempferol is maintained when these flavonols react with $^1\text{O}_2$ as glycosylated
18 forms. This was certainly the case for rhamnosides, although the myricetin > quercetin >
19 kaempferol order of $^1\text{O}_2$ antioxidant capacities changed to quercetin-rhamnoside > myricetin-
20 rhamnoside > kaempferol-dirhamnoside (Fig.2B). Because kaempferol-3-rhamnoside as
21 matching pair for quercetin-3-rhamnoside was not available as pure test compound, two other
22 corresponding glycosides, quercetin- and kaempferol-3-glucoside were also compared. These
23 data showed that regardless of the nature of glycosylation compounds containing the
24 dihydroxy-flavonol are better $^1\text{O}_2$ quenchers than mono-hydroxy-flavonols. This is a new
25 finding in respect of specific $^1\text{O}_2$ neutralizing and it is in agreement with Zietz et al. (2010)
26 reporting a higher total antioxidant capacity of quercetin glycosides than their corresponding
27 kaempferol glycosides. We observed several differences between flavonoid compositions of
28 sun and shade leaves which contribute to the higher $^1\text{O}_2$ neutralizing capacity of the former
29 (Fig. 2A). These include (1) the presence of myricetin-glycosides (9%) in sun leaves as
30 opposed to shade leaves (2) the increase in the ratio of quercetin to kaempferol glycosides
31 from 0.01 in shaded leaves to 0.9 in sun leaves and (3) the larger variety of myricetin and
32 quercetin glycosides detected in the sun leaves than in shade leaves (Table 2).

33

1 **3.4. Higher amounts of flavonols in sun leaves do not provide better UV-B screening**

2 In addition to the need of higher antioxidant capacities, higher amounts of flavonoids
3 in sun leaves than in shade leaves can also be interpreted as physical defence against the UV
4 region of the spectrum, i.e. screening harmful radiation by absorption (Caldwell et al., 1983;
5 Burchard et al. 2000). To see whether the observed higher concentrations of flavonols serve
6 this purpose in sun *T. platyphyllos* leaves, we compared UV absorbing properties of leaf
7 extracts and of pure compounds. Figure 4A shows that extracts from sun leaves and shade
8 leaves were different in their UV absorbing properties. These were total leaf extracts,
9 containing both the epidermis which is best known to be responsible for UV screening
10 (Robberecht and Caldwell, 1983) and mesophyll layers which were also reported to increase
11 their flavonoid content in response to high solar radiation (Tattini et al. 2005).

12 Figure 4B illustrates that kaempferol, quercetin and myricetin aglycones absorb UV
13 light mainly in the 350-400 nm spectral region, with slightly different wavelength maxima.
14 Maximum absorption of quercetin-3-glucoside and kaempferol-3-glucoside was shifted by
15 approximately 10 nm towards shorter wavelengths compared to their corresponding
16 aglycones (Fig 4C). Absorption maxima of the studied flavonol rhamnosides were between
17 345-355 nm with slight differences in peaks (Fig.4D). Differential spectra in Figs.4E and 4F
18 show that a selective increase in quercetin over kaempferol improves UV screening in the
19 UV-A (315-400 nm) but not in the UV-B (290-315 nm) region of the solar spectrum.
20 Therefore, higher amounts and larger variety of myricetin and quercetin glycosides than
21 kaempferol glycosides cannot be accounted for the observed better UV-B absorption of sun
22 leaves as compared to shade leaves. This result suggests that the increase in quercetin to
23 kaempferol glycoside ratios observed in response to UV radiation (Ryan et al. 1998, 2001;
24 Reifenrath and Müller 2007; Hofmann et al. 2000) or sunlight (Jaakola et al. 2004) is unlikely
25 to serve solely the purpose of UV-B screening.

26 27 **3.5. Hypothesis: solar UV facilitates the acclimation of sun leaves to high PAR via** 28 **promoting ¹O₂ antioxidant flavonols**

29 In summary, comparing flavonol contents of sun and shade *T. platyphyllos* leaves we
30 found a more pronounced difference in quercetin and myricetin glycosides than in
31 kaempferol glycosides. A general increase in flavonoids is likely to improve UV screening
32 but the above change in flavonols cannot be interpreted as the main constituent of
33 photoprotection against potentially damaging UV-B. Other phenolic compounds such as
34 hydroxycinnamic acid derivatives would serve specific UV-B screening functions much

1 better (Harborne and Williams 2000) while multihydroxylated flavonols promote acclimation
2 to sunlight by other means. To support a role different from direct UV absorption, Götz et al.
3 (2010) showed that the action spectrum of quercetin biosynthesis induction in *Arabidopsis*
4 leaves did not match the absorption spectrum characteristic to flavonols, but quasi-linearly
5 increased with higher energy radiation.

6 We found that the observed changes in flavonol glycoside composition gave better
7 antioxidant defence against $^1\text{O}_2$ to *T. platyphyllos* sun leaves as compared to shade leaves.
8 Improved protection against reactive oxygen species has been established as defence against
9 potential oxidative damage by high PAR or UV-B radiation (for reviews see Mittler 2002;
10 Apel and Hirt 2004; Asada, 1999b). However, $^1\text{O}_2$ is unique among pro-oxidants associated
11 with light stress and thus accumulation of effective $^1\text{O}_2$ antioxidants may signify more
12 specific plant responses. In earlier laboratory experiments we detected $^1\text{O}_2$ when tobacco and
13 spinach leaves were subject to excess PAR or high doses of UV-A but not when stress
14 brought about by UV-B only (Hideg et al. 2000, Barta et al. 2004).

15 Although ambient solar UV-B alone is not damaging to plants at middle latitudes of
16 the Northern hemisphere, it may enhance the effect of other environmental factors and
17 contribute to a synergistic stress (Hideg et al. 2013). In this picture, flavonols in sun leaves
18 are important for the successful avoidance of photoinhibition as effective $^1\text{O}_2$ antioxidants
19 rather than physical UV-B screeners. Morales et al. (2013) recently found that solar UV-A
20 elicited some of the same metabolic responses as solar UV-B in *Arabidopsis* leaves and
21 predicted an interaction of UVR8 with other photoreceptors to modulate UV-A responses in
22 the presence of UV-B. Our data support this concept and also suggest another link between
23 responses to various spectral components of sunlight. We hypothesise that in addition to the
24 above link between responses to UV-A and to UV-B, responses to high intensity PAR and to
25 UV are also connected and that solar UV-induced changes in leaf flavonol profile promote
26 acclimation to high PAR by helping to prevent $^1\text{O}_2$ mediated oxidative stress. Molecular
27 mechanisms behind this connection are subject to further studies.

28 29 **4. Supplementary data**

30 Table S1: Fragmentation pattern of flavonol glycosides and other phenolic compounds
31 present in shaded *Tilia platyphyllos* leaves resulting from HPLC-DAD-ESI-MSⁿ
32 measurements.

1 Table S2: Fragmentation pattern of flavonol glycosides and other phenolic compounds
2 present in sun exposed *Tilia platyphyllos* leaves resulting from HPLC-DAD-ESI-MSn
3 measurements.

4

5 **5. Acknowledgements**

6 Cooperation between participating laboratories was aided by COST Action FA0906
7 UV4growth. Work in Hungary was also supported by the Hungarian Scientific Grant Agency
8 (grant number OTKA NN85349). We thank the valuable assistance of Andrea Jankowsky
9 (IGZ, Großbeeren, Germany) in the preparation of leaf samples for the HPLC and Dr. Csaba
10 Tömböly (BRC Szeged, Hungary) for making a freeze dryer available. P.M. acknowledges
11 the research grant of the ‘Scientia Amabilis Foundation for Hungarian Plant Physiology’ and
12 the Short Term Science Mission grant of COST.

13

14

15

References

- Agati, G., Matteini, P., Goti, A., Tattini, M., 2007. Chloroplast-located flavonoids can scavenge singlet oxygen. *New Phytol.* 174, 77–89.
- Agati, G., Tattini, M., 2010. Multiple functional roles of flavonoids in photoprotection. *New Phytol.* 186, 786-793.
- Agati, G., Azzarello, E., Pollastri, S., Tattini, M., 2012. Flavonoids as antioxidants in plants: location and functional significance. *Plant Sci.* 196, 67-76.
- Aguirre- Hernández, E., González-Trujano, M.E., Martínez, A.L., Moreno J, Kite G, Terrazas T, Soto-Hernández M. 2010. HPLC/MS analysis and anxiolytic effect of quercetin and kaempferol flavonoids from *Tilia ammericana* var. *Mexicana*. *J. Ethnopharmacol.* 127, 91-97.
- Apel ,K., Hirt, H., 2004. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373-399.
- Asada, K., 1999a. The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons. *Ann. Rev. Plant Physiol.* 50, 601-639.
- Asada, K., 1999b. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* 141, 391-396.
- Barta, Cs., Kálai, T., Hideg, K., Vass, I., Hideg, É., 2004. Differences in the ROS generating efficacy of various ultraviolet wavelengths in detached spinach leaves. *Funct. Plant Biol.* 31, 23-28.
- Bassman, J.H, Robberecht, R., Edwards, G.E., 2001. Effects of enhanced UV-B radiation on growth and gas exchange in *Populus deltoides* Bartr. ex Marsh. *Int. J. Plant Sci.* 162, 103-110.
- Boardman, N.K., 1977. Comparative photosynthesis of sun and shade plants. *Annu. Rev. Plant Physiol.* 28, 355-377.
- Brown, B.A., Cloix, C., Jiang, G.H., Kaiserli, E., Herzyk, P., Kliebenstein, D.J., Jenkins, G.I., 2005. A UV-B-specific signaling component orchestrates plant UV protection. *Proc. Natl. Acad. Sci. USA* 102, 18225-18230.
- Brunetti, C., Di Ferdinando, M., Fini, A., Pollastri, S., Tattini, M., 2013. Flavonoids as antioxidants and developmental regulators: rel. significance in plants and humans. *Int. J. Mol. Sci.* 14, 3540-3555.
- Burchard, P., Bilger, W., Weissenböck, G., 2000. Contribution of hydroxycinnamates and flavonoids to epidermal shielding of UV-A and UV-B radiation in developing rye primary

- leaves as assessed by ultraviolet-induced chlorophyll fluorescence measurements. *Plant Cell Environ.* 23, 1373-1380.
- Caldwell, M.M., Robberecht, R., Flint, S.D., 1983. Internal filters: Prospects for UV-acclimation in higher plants. *Physiol. Plant.* 58, 445-450.
- Chen, A., Chen, Y., 2013. A review of the dietary flavonoid, kaempferol on human health and cancer chemoprotection. *Food Chem.* 138, 2099-2107.
- Demmig-Adams, B., Adams, W.W., 1996a. Xanthophyll cycle and light stress in nature: uniform response to excess direct sunlight among higher plant species. *Planta* 198, 460-470.
- Demmig-Adams, B., Adams, W.W., 1996b. The role of xanthophyll cycle carotenoids in the protection of photosynthesis. *TIPS* 1, 21-26.
- Favory, J.J., Stec, A., Gruber, H., Rizzini, L., Oravecz, A., Funk, M., Albert, A., Cloix, C., Jenkins, G.I., Oakeley, E.J., Seidlitz, H.K., Nagy, F., Ulm, R., 2009. Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress acclimation in *Arabidopsis*. *EMBO J.* 28, 591-601.
- Götz, M., Albert, A., Stich, S., Heller, W., Scherb, H., Krins, A., Langebartels, C., Seidlitz, H.K., Ernst, D., 2010. PAR modulation of the UV-dependent levels of flavonoid metabolites in *Arabidopsis thaliana* (L.) Heynh. leaf rosettes: cumulative effects after a whole vegetative growth period. *Protoplasma* 243, 95-103.
- Hallik, L., Niinemets, U., Kull, O., 2012. Photosynthetic acclimation to light in woody and herbaceous species: a comparison of leaf structure, pigment content and chlorophyll fluorescence characteristics measured in the field. *Plant Biol.* 14, 88-99.
- Harborne, J.B., Williams, C.A., 2000. Advances in flavonoid research since 1992. *Phytochemistry* 55, 481-504.
- Harborne, J.B., 1986. Nature, distribution and function of plant flavonoids. *Prog. Clin. Biol. Res.* 213, 15-24.
- Heim, K.E., Tagliaferro, A.R., Bobilya, D.J., 2002. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *J. Nutr. Biochem.* 13, 572-584.
- Hernández, I., Alegre, L., Van Breusegem, F., Munné-Bosch, S., 2009. How relevant are flavonoids as antioxidants in plants? *TIPS* 14, 125-132.
- Hideg, É., Jansen, M.A.K., Strid, Å., 2013. UV-B, ROS and stress; inseparable companions or loosely linked associates? *TIPS* 18, 107-115.

- Hideg, É., Majer, P., 2010. Factors contributing to the high light tolerance of leaves in vivo - Involvement of photo-protective energy dissipation and singlet oxygen scavenging. *Acta Biol. Hung.* 61, 69-80.
- Hideg, É., Spetea, C., Vass, I., 1994. Singlet oxygen production in thylakoid membranes during photoinhibition as detected by EPR spectroscopy. *Photosynth. Res.* 39, 191-199.
- Hideg, É., Kálai, T., Hideg, K., Vass, I., 2000. Do oxidative stress conditions impairing photosynthesis in the light manifest as photoinhibition? *Philos. T. Roy. Soc. B.* 355, 1511-1516.
- Hofmann, R.W., Swinny, E.E., Bloor, S.J., Markham, K.R., Ryan, K.G., Campbell, B.D., Jordan, B.R., Fountain, D.W., 2000. Responses of nine *Trifolium repens* L. populations to ultraviolet-B radiation: differential flavonol glycoside accumulation and biomass production. *Annal. Bot.* 86, 527-537.
- Hu, J.P., Calomme, M., Lasure, A., De Bruyne, T., Peters, L., Vlietinck, A., Van den Berghe, D.A., 1995. Structure-activity relationship of flavonoids with superoxide scavenging activity. *Biol. Trace Elem. Res.* 47, 327-331.
- Jaakola, L., Määttä-Riihinen, K., Kärenlampi, S., Hohtola, A., 2004. Activation of flavonoid biosynthesis by solar radiation in bilberry (*Vaccinium myrtillus* L.) leaves. *Planta* 218, 721-728.
- Jansen, M.A.K., Gaba, V., Greenberg, B.M., 1998. Higher plants and UV-B radiation: balancing damage, repair and acclimation. *TIPS* 3, 131-135.
- Jansen, M.A.K., Hectors, K., O'Brien, N.M., Guisez, Y., Potters, G., 2008. Plant stress and human health: Do human consumers benefit from UV-B acclimated crops? *Plant Sci.* 175, 449-458.
- Jung, M.Y., Min, D.B., 2009. ESR study of the singlet oxygen quenching and protective activity of Trolox on the photodecomposition of riboflavin and lumiflavin in aqueous buffer solutions. *J. Food Sci.* 74, 449-455.
- Kim, B.G., Kim, J.H., Kim, J., Lee, C., Ahn, J.H., 2008. Accumulation of flavonols in response to ultraviolet-B irradiation in soybean is related to induction of flavanone 3-beta-hydroxylase and flavonol synthase. *Mol. Cells* 25, 247-252.
- Klughammer, Ch., Schreiber, U., 2008. Saturation pulse method for assessment of energy conversion in PSI. *PAM Appl. Notes* 1, 11-14.
- Krumbein, A.; Saeger-Fink, H.; Schonhof, I., 2007. Changes in quercetin and kaempferol concentrations during broccoli head ontogeny in three broccoli cultivars. *J. Appl. Bot. Food Qual.* 81, 136-139.

- Lichtenthaler, H.K., Buschmann, C., Döll, M., Fietz, H.-J., Bach, T., Kozel, U., Meier, D., Rahmsdorf, U., 1981. Photosynthetic activity, chloroplast ultrastructure, and leaf characteristics of high-light and low-light plants and of sun and shade leaves. *Photosynth. Res.* 2, 115-141.
- Long, S.P., Humphries, S., Falkowski, P.G., 1994. Photoinhibition of photosynthesis in nature. *Ann. Rev. Plant Physiol.* 45, 633-662.
- Macpherson, A.N., Telfer, A., Barber, J., Truscott, T.G., 1993. Direct detection of singlet oxygen from isolated photosystem II reaction centers. *BBA* 1143, 301-309.
- Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *TIPS* 7, 405-410.
- Mirecki, R.M., Teramura, A.H., 1984. Effects of Ultraviolet-B Irradiance on Soybean: V. The Dependence of Plant Sensitivity on the Photosynthetic Photon Flux Density during and after Leaf Expansion. *Plant Physiol.* 74, 475-480.
- Montoro, P., Braca, A., Pizza, C., De Tommasi, N., 2005. Structure-antioxidant activity relationships of flavonoids isolated from different plant species. *Food Chem.* 92, 349-355.
- Morales, L.O., Tegelberg, R., Brosché, M., Keinänen, M., Lindfors, A., Aphalo, P.J., 2010. Effects of solar UV-A and UV-B radiation on gene expression and phenolic accumulation in *Betula pendula* leaves. *Tree Physiol.* 30, 923-934.
- Morales, L.O., Brosché, M., Vainonen, J., Jenkins, G.I., Wargent, J.J., Sipari, N., Strid, Å., Lindfors, A.V., Tegelberg, R., Aphalo, P.J., 2013. Multiple Roles for UV RESISTANCE LOCUS 8 in Regulating Gene Expression and Metabolite Accumulation in *Arabidopsis* under Solar UV Radiation. *Plant Physiol.* 161, 744-759.
- Müller, P., Li, X.-P., Niyogi, K., 2001. Non-photochemical quenching. A response to excess light energy. *Plant Physiol.* 125, 1558-1556.
- Niyogi, K.K., 1999. Photoprotection revisited: genetic and molecular approaches. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50, 333-359.
- Oravec, A., Baumann, A., Mate, Z., Brzezinska, A., Molinier, J., Oakeley, E.J., Ádám, É., Schäfer, E., Nagy, F., Ulm, R., 2006. CONSTITUTIVELY PHOTOMORPHOGENIC1 is required for the UV-B response in *Arabidopsis*. *Plant Cell* 18, 1975-1990.
- Pietta, P.-G., 2000. Flavonoids as antioxidants. *J. Nat. Prod.* 63, 1035-1042.
- Powles, S.B., 1984. Photoinhibition of photosynthesis induced by visible light. *Annu. Rev. Plant Physiol.* 35, 15-44.
- Ratty, A.K., Das, N.P., 1988. Effects of flavonoids on nonenzymatic lipid peroxidation: structure-activity relationship. *Biochem. Med. Metabol. Biol.* 39, 69-79.

- Reifenrath, K., Müller, C., 2007. Species-specific and leaf-age dependent effects of ultraviolet radiation on two Brassicaceae. *Phytochem.* 68, 875-885.
- Rice-Evans, C.A., Miller, N.J., Bolwell, P.G., Bramley, P.M., Pridham, J.B., 1995. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Rad. Res.* 22, 375-383.
- Rice-Evans, C.A., Miller, N.J., Paganga, G., 1996. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Rad. Biol. Med.* 20, 933-956.
- Robberecht, R., Caldwell, M.M., 1983. Protective mechanisms and acclimation to solar ultraviolet radiation in *Oenothera stricta*. *Plant Cell Environ.* 6, 477-485.
- Ryan, K.G., Markham, K.R., Bloor, S.J., Bradley, J.M., Mitchell, K.A., Jordan, B.R., 1998. UV-B radiation induced increase in quercetin : kaempferol ratio in wild-type and transgenic lines of petunia. *Photochem. Photobiol.* 68, 323-330.
- Ryan, K.G., Swimmy, E.E., Winefield, C., Markham, K.E., 2001. Flavonoids and UV photoprotection in *Arabidopsis* mutants. *Z. Naturforsch.* 56, 745-754.
- Saunders, J.A., McClure, J.W., 1976. The distribution of flavonoids in chloroplasts of twenty five species of vascular plants. *Phytochem.* 15, 809-810.
- Scalbert, A., Manach, C., Morand, C., Rémésy, C., Jiménez, L., 2005. Dietary polyphenols and the prevention of diseases. *Crit. Rev. Food Sci. Nutr.* 45, 287-306.
- Schmidt, S., Zietz, M., Schreiner, M., Rohn, S., Kroh, L.W., Krumbein, A., 2010. Genotypic and climatic influences on the concentration and composition of flavonoids in kale (*Brassica oleracea* var. *sabellica*). *Food Chem.* 119, 1293-1299.
- Schreiber, U., 2004. Pulse-Amplitude (PAM) fluorometry and saturation pulse method. In: Papageorgiou, G., Govindjee (Eds.), *Chlorophyll fluorescence: a signature of photosynthesis*, Kluwer Academic Publishers, Dordrecht, The Netherlands, pp, 279-319.
- Schütz, K., Kammerer, D., Carle, R., Schieber, A., 2004. Identification and quantification of caffeoylquinic acids and flavonoids from artichoke (*Cynara scolymus* L.) heads, juice, and pomace by HPLC-DAD-ESI/MS(n). *J. Agric. Food Chem.* 52, 4090-4096.
- Sekher Pannala, A., Chan, T.S., O'Brien, P.J., Rice-Evans, C.A., 2001. Flavonoid B-ring chemistry and antioxidant activity: fast reaction kinetics. *Bioch. Biophys. Res. Commun.* 282, 1161-1168.
- Solovchenko, A., Merzlyak, M., 2008. Screening of visible and UV radiation as a photoprotective mechanism in plants. *Russ. J. Plant Physiol.* 55, 719-737.
- Szőllősi, E., Oláh, V., Kanalas, P., Kis, J., Fenyvesi, A., Mészáros. I., 2010. Seasonal

- variation of leaf ecophysiological traits within the canopy of *Quercus petraea* trees. *Acta Biol. Hung.* 61, 192-208.
- Takahashi, S., Milward, S.E., Yamori, W., Evans, J.R., Hillier, W., Badger, M.R., 2010. The solar action spectrum of photosystem II damage. *Plant Physiol.* 153, 988-993.
- Tattini, M., Guidi, L., Morassi-Bonzi, L., Pinelli, P., Remorini, D., Degl'Innocenti, E., Giordano, C., Massai, R., Agati, G., 2005. On the role of flavonoids in the integrated mechanisms of response of *Ligustrum vulgare* and *Phillyrea latifolia* to high solar radiation. *New Phytol.* 167, 457-470.
- Toker, G., Aslan, M., Yeşilada, E., Memişoğlu, M., Ito, S., 2001. Comparative evaluation of the flavonoid content in officinal *Tiliae flos* and Turkish lime species for quality assessment. *J. Pharm. Biomed. Anal.* 26, 111-121.
- Tournaire, C., Croux, S., Maurette, M.T., Beck, I., Hocquaux, M., Braun, A.M., Oliveros, E., 1993. Antioxidant activity of flavonoids: efficiency of singlet oxygen (Δg) quenching. *J. Photochem. Photobiol. B.* 19, 205-215.
- Triantaphylidès, C., Havaux, M., 2009. Singlet oxygen in plants: production, detoxification and signaling. *TIPS* 14, 219-28.
- Winkel-Shirley, B., 2002. Biosynthesis of flavonoids and effects of stress. *Curr. Opin. Plant Biol.* 5, 218-223.
- Winter, T.R., Rostás, M., 2008. Ambient ultraviolet radiation induces protective responses in soybean but does not attenuate indirect defense. *Environ. Pollut.* 155, 290-297.
- Young, R.H., Brewer, D., Keller, R.A., 1973. Determination of rate constants of reaction and lifetimes of singlet oxygen in solution by a flash photolysis technique. *JACS* 95, 375-379.
- Zaprometov, M.N., Nikolaeva, T.N., 2003. Chloroplasts isolated from kidney bean leaves are capable of phenolic compound biosynthesis. *Russ. J. Plant Physiol.* 50, 623-626.
- Zhang, W.J., Björn, L.O. 2009. The effect of ultraviolet radiation on the accumulation of medicinal compounds in plants. *Fitoterapia* 80, 207-218.
- Zietz, M., Weckmüller, A., Schmidt, S., Rohn, S., Schreiner, M., Krumbein, A., Kroh, L.W. 2010. Genotypic and climatic influence on the antioxidant activity of flavonoids in kale (*Brassica oleracea* var. *sabellica*). *J. Agric. Food Chem.* 58, 2123-2130.

Figure legends

Figure 1

Photosystem II quantum yield components of sun (empty symbols) and shade (full symbols) *Tilia platyphyllos* leaves. (A): light response curves of photochemistry (circles) and a sum of non-photochemical quenching components (squares). (B): Energy partitioning between photochemistry (black segments, Y(II)), regulated (dark gray, Y(NPQ)) and non-regulated (light gray, Y(NO)) non-photochemical quenching at three different PAR (photosynthetically active radiation) level in sun (upper row) and shade (lower row) leaves.

Figure 2

Singlet oxygen ($^1\text{O}_2$) neutralizing antioxidant capacities of (A) sun and shade *Tilia platyphyllos* leaf extracts and (B) various flavonol glycosides (standards). Error bars represent standard deviations and significantly different values ($p < 0.05$) are marked with asterisks or with different letters in figures A and B, respectively. $n=5$ (A) and $n=3$ (B).

Figure 3

Structure of major flavonol aglycones and flavonol glucosides and flavonol rhamnosides detected in *Tilia platyphyllos* leaves.

Figure 4

Absorbance spectra (A) of *Tilia* leaf extracts normalized to the maximum absorption of the sun leaf extract and (B-D) of pure flavonol compounds normalized to the absorption maximum of quercetin. (E-F) Differential spectra calculated from data in figures (A-D).

Figure 1
[Click here to download high resolution image](#)





Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)










Flavonol	 <p>KAEMPFEROL</p>	 <p>QUERCETIN</p>	 <p>MYRICETIN</p>
Flavonol glucoside	 <p>KAEMPFEROL-3-O-glucoside (astragalin)</p>	 <p>QUERCETIN-3-O-glucoside (isoquercitrin)</p>	
Flavonol rhamnoside	 <p>KAEMPFEROL-3,7-di-rhamnoside (kaempferitrin)</p>	 <p>QUERCETIN-3-O-rhamnoside (quercitrin)</p>	 <p>MYRICETIN-3-O-rhamnoside (myricitrin)</p>

Figure 4
[Click here to download high resolution image](#)



Tables

Table 1

Flavonoid aglycones of *Tilia platyphyllos* sun and shade leaves

flavonoid (mg g⁻¹ dry weight)	sun leaf	shade leaf
total content	21.5 ± 5.9	5.1 ± 1.3 *
quercetin	15.2 ± 4.6	0.7 ± 0.3 *
kaempferol	4.7 ± 0.7	4.3 ± 1.0
myricetin	1.6 ± 0.6	0.1 ± 0.03 *

Table 2

Percentage composition of flavonoid glycosides and other phenolic compounds detected in *Tilia platyphyllos* sun and shade leaves

compounds	sun leaves	shade leaves
quercetin glycosides	39%	0.5%
kaempferol glycosides	42%	77.5%
myricetin glycosides	9%	traces
chlorogenic acid	3%	6%
unidentified phenol	1%	5%
unidentified compounds	6%	11%

See Table S1 (shaded leaves of *Tilia*) and S2 (sun exposed leaves of *Tilia*) in Supplementary material for a more detailed list of compounds including HPLC-MS data for their identification.

Supplemental file

Singlet oxygen scavenging by leaf flavonoids contributes to sunlight acclimation in *Tilia platyphyllos*

Petra Majer, Susanne Neugart, Angelika Krumbein, Monika Schreiner and Éva Hideg

Table S1: Fragmentation pattern of flavonol glycosides and other phenolic compounds present in shaded linden leaves resulting from HPLC-DAD-ESI-MSⁿ measurements.

Shaded leaves MW	MS	MS ²	MS ³	RT (min)	λ_{\max} (nm)	Compound name	Peak area in %
354	707 , 191 (33), 353 (13)	353	191 , 161 (2), 135 (1)	20.1	240, 303sh, 333	Chlorogenic acid	5
354	707 ,353 (77)	353	173 ,179 (57)	23.7	228, 304sh, 330	Chlorogenic acid	1
?	591	295 , 179 (24), 255 (9)	179 (81), 133 , 115 (14)	34.1	241, 300sh, 331	Unknown phenol	5
594	593 , 675 (95)	447(56), 431 (65), 285		48.9	240, 274sh, 313	Kaempferol-3- <i>O</i> -rutinoside	0.5
610	609 , 575 (62)	463 (52), 447 , 301 (47)	301	50.8	270, 296, 353	Quercetin-3- <i>O</i> -caffeoyl-7- <i>O</i> - rhamnoside	0.5
756	755	593 , 447 (58), 285 (27)	285	63.9	230, 270, 348	Kaempferol-3- <i>O</i> -rutinoside-7- <i>O</i> -glucoside	2
594	593	447 , 431 (68), 285 (36)	284	64.6	230, 270, 350	Kaempferol-3- <i>O</i> -glucoside-7- <i>O</i> - rhamnoside	1
564	563	430 (47), 417 , 285 (18)	284	66.4	237, 265, 351	Kaempferol-3- <i>O</i> -coumaroyl- pentoside	3
740	739	593 , 431 (22), 285 (23)	285	84.3	268, 320sh, 346	Kaempferol-3- <i>O</i> -rutinoside-7- <i>O</i> -rhamnoside	39
578	577	431 , 285 (29)	285	87.1	270, 322sh, 349	Kaempferol-3- <i>O</i> -rhamnoside-7- <i>O</i> -rhamnoside	32

MW = Molecular weight, RT = Retention time, λ_{\max} = Absorption maximum in nm, sh = shoulder

The numbers in brackets indicate the relative intensity of the fragment ions. Bold fragment ions conform 100 percent.

Further quercetin and kaempferol glycosides in traces (<0.5%).

Table S2: Fragmentation pattern of flavonol glycosides and other phenolic compounds present in sun exposed linden leaves resulting from HPLC-DAD-ESI-MSⁿ measurements.

Sun leaves MW	MS	MS ²	MS ³	RT (min)	λ_{\max} (nm)	Compound name	Peak area in %
354	707	353	191 , 179 (3), 135 (2)	19.7	247, 303sh, 334	Chlorogenic acid	2
354	707 , 353 (91)	353		20.7	244, 301sh, 328	Chlorogenic acid	1
?	591	477 , 359 (14), 295 (27)	359 , 271 (55), 315 (5)	33.8	243, 331	Unknown phenol	1
596	595	462	315 , 179 (2)	46.8	228, 347	Myricetin-3- <i>O</i> -rhamnoside-7- <i>O</i> -pentoside	1
610	609	463 (76), 447 , 301 (29)	301	49.0	271, 357	Quercetin-3- <i>O</i> -caffeoyl-7- <i>O</i> -rhamnoside (isomer)	5
610	609	463 (8), 447 , 301 (33)	301	51.9	260, 355	Quercetin-3- <i>O</i> -caffeoyl-7- <i>O</i> -rhamnoside (isomer)	2
580	579	447 , 433 (58), 301 (15)	301	54.7	259, 349	Quercetin-3- <i>O</i> -rhamnoside-7- <i>O</i> -pentoside	5
594	593	447 , 285 (5)	285 , 327 (45)	57.6	258, 358	Kaempferol-3- <i>O</i> -glucoside-7- <i>O</i> -rhamnoside	1
450	449	316		58.1	273, 352	Myricetin-3- <i>O</i> -pentoside	2
464	463	316		60.0	268, 360	Myricetin-3- <i>O</i> -rhamnoside	6
564	563	431 (55), 417 , 285 (13)	284	63.3	268, 355	Kaempferol-3- <i>O</i> -coumaroyl-pentoside	7
594	593	447 , 431 (57), 285 (25)	284	64.6	268, 350	Kaempferol-3- <i>O</i> -coumaroyl-glucoside (tiliroside)	3
594	593	447 , 301 (49)	301	65.4	269, 352	Quercetin-3- <i>O</i> -rhamnoside-7- <i>O</i> -rhamnoside	18
726	725	579 (75), 447 , 301 (52)	301	68.3	267, 353	Quercetin-3- <i>O</i> -rhamnoside-7- <i>O</i> -rhamnoside-pentoside	3
434	433	301		76.5	260, 353	Quercetin-3- <i>O</i> -pentoside	6
740	739	593 , 431 (23), 285 (28)	285	83.5	270, 321sh, 349	Kaempferol-3- <i>O</i> -rutinoside-7- <i>O</i> -rhamnoside	13
710	709 , 577 (91)	563 , 431 (37), 285 (25)	285	86.5	260, 349	Kaempferol-3- <i>O</i> -rhamnoside-pentoside-7- <i>O</i> -rhamnoside	4

Sun leaves MW continued	MS	MS ²	MS ³	RT (min)	λ_{\max} (nm)	Compound name	Peak area in %
578	577	431 , 285 (28)	285	87.4	269, 353	Kaempferol-3- <i>O</i> -rhamnoside-7- O-rhamnoside	13
432	431	285		118.8	270, 322sh, 353	Kaempferol-3- <i>O</i> -rhamnoside	1

MW = Molecular weight, RT = Retention time, λ_{\max} = Absorption maximum in nm, sh = shoulder

The numbers in brackets indicate the relative intensity of the fragment ions. Bold fragment ions conform 100 percent.

Further myricetin, quercetin and kaempferol glycosides in traces (< 1%).