

Hindawi Publishing Corporation
 Abstract and Applied Analysis
 Volume 2007, Article ID 61296, 11 pages
 doi:10.1155/2007/61296

Research Article

Continuous-Time Multiobjective Optimization Problems via Invexity

Valeriano A. De Oliveira and Marko A. Rojas-Medar

Received 29 March 2006; Accepted 1 December 2006

Recommended by Nikolaos S. Papageorgiou

We introduce some concepts of generalized invexity for the continuous-time multiobjective programming problems, namely, the concepts of Karush-Kuhn-Tucker invexity and Karush-Kuhn-Tucker pseudoinvexity. Using the concept of Karush-Kuhn-Tucker invexity, we study the relationship of the multiobjective problems with some related scalar problems. Further, we show that Karush-Kuhn-Tucker pseudoinvexity is a necessary and sufficient condition for a vector Karush-Kuhn-Tucker solution to be a weakly efficient solution.

Copyright © 2007 V. A. De Oliveira and M. A. Rojas-Medar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this work, we regard the continuous-time multiobjective optimization problem,

$$\begin{aligned} \text{Minimize } \phi(x) &= \left(\int_0^T f_1(x(t), t) dt, \dots, \int_0^T f_p(x(t), t) dt \right) \\ &\text{subject to } g_i(x(t), t) \leq 0 \quad \text{a.e. in } [0, T], \quad i = 1, \dots, m, \quad x \in X. \end{aligned} \quad (\text{CMP})$$

Here X is a nonempty open convex subset of the Banach space $L_\infty^n[0, T]$, $\phi : X \rightarrow \mathbb{R}^p$, $f_j(x(t), t) = \xi_j(x)(t)$, $j \in J := \{1, \dots, p\}$, $g_i(x(t), t) = \gamma_i(x)(t)$, $i \in I := \{1, \dots, m\}$, where $\xi_j : X \rightarrow \Lambda_1^1[0, T]$, $j \in J$, and $\gamma_i : X \rightarrow \Lambda_1^1[0, T]$, $i \in I$. $L_\infty^n[0, T]$ denotes the space of all n -dimensional vector-valued Lebesgue measurable functions, which are essentially bounded,

2 Abstract and Applied Analysis

defined on the compact interval $[0, T] \subset \mathbb{R}$, with norm $\|\cdot\|_\infty$ defined by

$$\|x\|_\infty = \max_{1 \leq j \leq n} \operatorname{ess\,sup} \{ |x_j(t)|, 0 \leq t \leq T \}, \quad (1.1)$$

where for each $t \in [0, T]$, $x_j(t)$ is the j th component of $x(t) \in \mathbb{R}^n$ and $\Lambda_1^m[0, T]$ denotes the space of all m -dimensional vector-valued functions which are essentially bounded and Lebesgue measurable, defined on $[0, T]$, with the norm $\|\cdot\|_1$ defined by

$$\|y\|_1 = \max_{1 \leq j \leq m} \int_0^T |y_j(t)| dt. \quad (1.2)$$

The mono-objective version of this class of problems was introduced by Bellman [1] in connection with production-inventory “bottleneck processes.” He considered a type of optimization problems, which is now known as continuous-time linear programming, formulated its dual and provided duality relations. He also suggested some computational procedures.

Since then, various authors have extended his theory to wider classes of continuous-time problems (e.g., [2–12]). In these articles, the authors study the mono-objective case, but in many applications it is necessary to minimize not only one objective. So, the multiobjective problem is more general and more suitable for many applications. The development of the necessary and sufficient optimality conditions for (CMP) was done in [13].

Our aim in this paper is to provide necessary and sufficient conditions for global optimality of a vector Karush-Kuhn-Tucker solution as well for a vector Karush-Kuhn-Tucker solution to solve a related scalar problem. Our results extend the finite dimensional case studied in [14, 15] and the continuous-time mono-objective case studied in [16].

For more literature about these issues, we refer the reader to [14–16] and the bibliography cited therein.

We organized this work into four sections. In Section 2, we give some preliminaries. We introduce KKT pseudoinvexity for (CMP) and state our first mean result in Section 3. The notion of KKT invexity and our second mean result are given in Section 4.

2. Preliminaries

Let V be an open subset of \mathbb{R}^n containing the set $\{x(t) \in \mathbb{R}^n : x \in X, t \in [0, T]\}$. We assume that $f_j, j \in J$, and $g_i, i \in I$, are real functions defined on $V \times [0, T]$. The functions $t \mapsto f_j(x(t), t), j \in J$, and $t \mapsto g_i(x(t), t), i \in I$, are assumed to be Lebesgue measurable and integrable for all $x \in X$. We assume also that the functions $f_j, j \in J$, and $g_i, i \in I$, are continuously differentiable with respect to their first arguments.

Let \mathbb{F} be the set of all feasible solutions of problem (CMP) (which we suppose non empty), that is, let

$$\mathbb{F} = \{x \in X : g_i(x(t), t) \leq 0 \text{ a.e. in } [0, T], i \in I\}. \quad (2.1)$$

Given $x \in \mathbb{F}$, for each $i \in I$ we denote by $A_i(x)$ the subset of $[0, T]$ where the i th constraint is active, that is,

$$A_i(x) = \{t \in [0, T] : g_i(x(t), t) = 0\}. \quad (2.2)$$

In this paper, all vectors are column vectors. We use a prime to denote transposition. Besides, given $w \in \mathbb{R}^p$, $w \leq 0$ means that $w_i \leq 0$ for $i = 1, 2, \dots, p$, and $w < 0$ means that $w_i < 0$ for $i = 1, 2, \dots, p$.

In what follows, we state a result which will be useful for the proof of our results. This result can be viewed as a generalized Motzkin theorem of the alternative. It is the continuous-time analogue of the theorem given by Mangasarian [17, page 66] and its proof is almost identical to the one given in the Mangasarian's book.

THEOREM 2.1. *Let $Z \subseteq L_\infty^n[0, T]$ be a nonempty convex subset. Let $p : W \times [0, T] \rightarrow \mathbb{R}^m$ and $q : W \times [0, T] \rightarrow \mathbb{R}^k$ be mappings given by $p(z(t), t) = \pi(z)(t)$ and $q(z(t), t) = B(t)z(t) - b(t)$, respectively, where $W \subseteq \mathbb{R}^n$ is an open subset, π is a mapping from Z into $\Lambda_1^m[0, T]$, $B(t)$ is a $k \times n$ matrix, and $b(t) \in \mathbb{R}^k$. Assume that p is convex with respect to its first argument in W throughout $[0, T]$ and that there does not exist $v \in L_\infty^k[0, T] \setminus \{0\}$, $v(t) \geq 0$ a.e. in $[0, T]$ such that*

$$B'(t)v(t) = 0 \quad \text{a.e. in } [0, T]. \quad (2.3)$$

Then exactly one of the following systems is consistent:

- (I) $p(z(t), t) < 0$, $B(t)z(t) \leq b(t)$ a.e. in $[0, T]$ has a solution $z \in Z$;
- (II) $\int_0^T \{u'(t)p(z(t), t) + v'(t)[B(t)z(t) - b(t)]\} dt \geq 0$ for all $z \in Z$, for some $u \in L_\infty^m[0, T]$, $u(t) \geq 0$, $u(t) \neq 0$ a.e. in $[0, T]$ and for some $v \in L_\infty^k[0, T]$, $v(t) \geq 0$ a.e. in $[0, T]$.

Proof. Quite similar to the proof of Theorem 3.4 in [10, page 137]. □

In the two following definitions, we define as in [13] a weakly efficient solution and a vector Karush-Kuhn-Tucker solution.

Definition 2.2. A feasible solution y is said to be a *weakly efficient solution* of (CMP) if and only if there does not exist another feasible solution x such that $\phi(x) < \phi(y)$.

Definition 2.3. A feasible solution y is said to be a *vector Karush-Kuhn-Tucker solution* (or *vector KKT solution*) for problem (CMP) if there exist $\mu \in \mathbb{R}^p$ and $\lambda \in L_\infty^m[0, T]$ such that

$$\int_0^T \left[\sum_{j \in J} \mu_j \nabla f'_j(y(t), t) + \sum_{i \in I} \lambda_i(t) \nabla g'_i(y(t), t) \right] h(t) dt = 0 \quad \forall h \in L_\infty^n[0, T], \quad (2.4)$$

$$\lambda_i(t) g_i(y(t), t) = 0 \quad \text{a.e. in } [0, T], \quad i \in I, \quad (2.5)$$

$$\lambda_i(t) \geq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \quad (2.6)$$

$$\mu_j \geq 0, \quad j \in J, \quad \mu \neq 0. \quad (2.7)$$

At last we give a constraint qualification in the continuous-time setting.

Definition 2.4. The constraints g_i , $i \in I$, satisfy (CQ) at $y \in \mathbb{F}$ if there do not exist $v_i \in L_\infty[0, T]$, $v_i(t) \geq 0$ a.e. in $[0, T]$, $i \in I$, not all zero, such that

$$\sum_{i \in I} \int_{A_i(y)} v_i(t) \nabla g_i(y(t), t) h(t) dt \geq 0 \quad \forall h \in L_\infty^n[0, T]. \quad (2.8)$$

3. KKT-pseudoinvexity and optimality conditions

In this section, we introduce the notion of Karush-Kuhn-Tucker pseudoinvexity for (CMP). Further, we state and prove a result which provides necessary and sufficient conditions for global optimality of a vector Karush-Kuhn-Tucker solution.

Definition 3.1. The problem (CMP) is said to be *Karush-Kuhn-Tucker pseudoinvex* (or *KKT-pseudoinvex*) if there exists a function $\eta : V \times V \times [0, T] \rightarrow \mathbb{R}^n$ such that $t \mapsto \eta(x(t), y(t), t) \in L_\infty^n[0, T]$, and

$$\phi(x) < \phi(y) \implies \int_0^T \nabla f'_j(y(t), t) \eta(x(t), y(t), t) dt < 0, \quad j \in J, \quad (3.1)$$

$$-\nabla g'_i(y(t), t) \eta(x(t), y(t), t) \geq 0 \quad \text{a.e. in } A_i(y), \quad i \in I, \quad (3.2)$$

for all $x, y \in \mathbb{F}$.

THEOREM 3.2. *Assume that the constraints g_i , $i \in I$, satisfy (CQ) at each $y \in \mathbb{F}$. Then every vector KKT-solution is a weak efficient solution of (CMP) if and only if (CMP) is KKT-pseudoinvex.*

Proof. Let y be a vector KKT-solution and suppose that (CMP) is KKT-pseudoinvex. Suppose that there exists a feasible solution x such that $\phi(x) < \phi(y)$. As (CMP) is KKT-pseudoinvex, using (3.1), we obtain

$$\int_0^T \nabla f'_j(y(t), t) \eta(x(t), y(t), t) dt < 0, \quad j \in J. \quad (3.3)$$

Since y is a vector KKT-solution, there exist $\mu \in \mathbb{R}^p$ and $\lambda \in L_\infty^m[0, T]$ satisfying (2.4)–(2.7). By (2.7) and (3.3), we have

$$\int_0^T \sum_{j \in J} \mu_j \nabla f'_j(y(t), t) \eta(x(t), y(t), t) dt < 0. \quad (3.4)$$

Using (2.4) with $h(t) = \eta(x(t), y(t), t)$, $t \in [0, T]$, we obtain

$$\int_0^T \sum_{i \in I} \lambda_i(t) \nabla g'_i(y(t), t) \eta(x(t), y(t), t) dt > 0. \quad (3.5)$$

By the other hand, from (2.6) and (3.2), since by (2.5) $\lambda_i(t) = 0$, $t \notin A_i(y)$, $i \in I$, it follows that

$$\int_0^T \sum_{i \in I} \lambda_i(t) \nabla g'_i(y(t), t) \eta(x(t), y(t), t) dt \leq 0, \quad (3.6)$$

which is a contradiction to (3.5). Therefore y is a weakly efficient solution.

Conversely, suppose that every vector KKT-solution is a weakly efficient solution. Let $x, y \in \mathbb{F}$ be such that $\phi(x) < \phi(y)$. Then y is not a weakly efficient solution, so that, by hypothesis, y is not a vector Karush-Kuhn-Tucker solution. So the system

$$\begin{aligned} \int_0^T \left[\sum_{j \in J} \mu_j \nabla f'_j(y(t), t) + \sum_{i \in I} \lambda_i(t) \nabla g'_i(y(t), t) \right] h(t) dt &= 0 \quad \forall h \in L_\infty^n[0, T], \\ \lambda_i(t) g_i(y(t), t) &= 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ \lambda_i(t) &\geq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ \mu_j &\geq 0, \quad j \in J, \quad \mu \neq 0, \end{aligned} \quad (3.7)$$

has no solution $(\mu, \lambda) \in \mathbb{R}^p \times L_\infty^m[0, T]$. Equivalently, the system

$$\begin{aligned} \int_0^T \left[\sum_{j \in J} \mu_j \nabla f'_j(y(t), t) + \sum_{i \in I} \lambda_i(t) \chi_i(t) \nabla g'_i(y(t), t) \right] h(t) dt &= 0 \quad \forall h \in L_\infty^n[0, T], \\ \lambda_i(t) &\geq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ \mu_j &\geq 0, \quad j \in J, \quad \mu \neq 0, \end{aligned} \quad (3.8)$$

has no solution $(\mu, \lambda) \in \mathbb{R}^p \times L_\infty^m[0, T]$, where $\chi_i : [0, T] \rightarrow \mathbb{R}$ is defined, for each $i \in I$, by

$$\chi_i(t) = \begin{cases} 1 & \text{if } t \in A_i(y), \\ 0 & \text{if } t \notin A_i(y). \end{cases} \quad (3.9)$$

As the constraint qualification holds by hypothesis, the condition (2.3) in Theorem 2.1 is verified. Applying that theorem, it follows that there exists $h \in L_\infty^n[0, T]$ such that

$$\begin{aligned} \int_0^T \nabla f'_j(y(t), t) h(t) dt &< 0, \quad j \in J, \\ \chi_i(t) \nabla g'_i(y(t), t) h(t) &\leq 0 \quad \text{a.e. in } [0, T], \quad i \in I. \end{aligned} \quad (3.10)$$

Define $\eta(x(t), y(t), t) = h(t)$ a.e. in $[0, T]$. Therefore

$$\begin{aligned} \int_0^T \nabla f'_j(y(t), t) \eta(x(t), y(t), t) dt &< 0, \quad j \in J, \\ -\nabla g'_i(y(t), t) \eta(x(t), y(t), t) &\geq 0 \quad \text{a.e. in } A_i(y), \quad i \in I. \end{aligned} \quad (3.11)$$

Thus there exists a function $\eta: V \times V \times [0, T] \rightarrow \mathbb{R}^n$ such that $t \mapsto \eta(x(t), y(t), t) \in L_\infty^n[0, T]$ and

$$\begin{aligned} \phi(x) < \phi(y) &\implies \int_0^T \nabla f'_j(y(t), t) \eta(x(t), y(t), t) dt < 0, \quad j \in J, \\ -\nabla g'_i(y(t), t) \eta(x(t), y(t), t) &\geq 0 \quad \text{a.e. in } A_i(y), \quad i \in I, \end{aligned} \quad (3.12)$$

for all $x, y \in \mathbb{F}$, so that (CMP) is KKT-pseudoinvex. \square

4. Karush-Kuhn-Tucker invexity and scalar problems

In this section, we generalize to the continuous-time context the notion of Karush-Kuhn-Tucker invexity introduced in [15] for finite-dimensional multiobjective problems. In addition, we give necessary and sufficient conditions for a vector KKT-solution to solve a related weighting scalar problem.

We will regard the following weighting scalar problem related with (CMP):

$$\begin{aligned} \text{Minimize } \Phi(x) &= \int_0^T \sum_{j \in J} \mu_j f_j(x(t), t) dt \\ \text{subject to } g_i(x(t), t) &\leq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \quad x \in X, \end{aligned} \quad (4.1)$$

where $\mu_j \in \mathbb{R}$, $j \in J$. This is one of the most known scalar problems associated with multiobjective optimization problems.

THEOREM 4.1. *Every optimal solution of a weighting scalar problem with $\mu_j \geq 0$, $j \in J$, not all zero, is a weakly efficient solution of (CMP).*

Proof. Let y be an optimal solution of a scalar problem with $\mu_j \geq 0$, $j \in J$, not all zero, and let us suppose that there exists $x \in \mathbb{F}$ such that $\phi(x) < \phi(y)$. Then $\phi_j(x) < \phi_j(y)$, $j \in J$, so that

$$\mu_j \phi_j(x) \leq \mu_j \phi_j(y), \quad j \in J, \quad (4.2)$$

since $\mu_j \geq 0$, $j \in J$. Provided μ_j , $j \in J$, are not all zero, there exists at least one $j \in J$ such that $\mu_j > 0$. Therefore, in the inequalities above, at least one holds strictly. So summing over J ,

$$\sum_{j \in J} \mu_j \phi_j(x) < \sum_{j \in J} \mu_j \phi_j(y), \quad (4.3)$$

which contradicts the optimality of y . \square

In order to establish the reciprocal of Theorem 4.1, we need some qualifications on the constraints and, furthermore, we need some generalized convexity hypothesis.

Definition 4.2. The problem (CMP) is said to be *Karush-Kuhn-Tucker invex* (or *KKT-invex*) if there exists a function $\eta : V \times V \times [0, T] \rightarrow \mathbb{R}^n$ such that $t \mapsto \eta(x(t), y(t), t) \in L_\infty^n[0, T]$ and

$$\begin{aligned} \phi_j(x) - \phi_j(y) &\geq \int_0^T \nabla f_j'(y(t), t) \eta(x(t), y(t), t) dt, \quad j \in J, \\ -\nabla g_i'(y(t), t) \eta(x(t), y(t), t) &\geq 0 \quad \text{a.e. in } A_i(y), \quad i \in I, \end{aligned} \quad (4.4)$$

for all $x, y \in \mathbb{F}$.

THEOREM 4.3. *Assume that the constraints g_i , $i \in I$, satisfy (CQ) at each $y \in \mathbb{F}$. If (CMP) is KKT-invex, then every weakly efficient solution solves a weighting scalar problem with $\mu_j \geq 0$, $j \in J$, not all zero.*

Proof. Let y be a weakly efficient solution of (CMP). Then, by [13, Theorem 3.3, page 9], there exist $\mu \in \mathbb{R}^p$ and $\lambda \in L_\infty^m[0, T]$ satisfying

$$\begin{aligned} \int_0^T \left[\sum_{j \in J} \mu_j \nabla f_j'(y(t), t) + \sum_{i \in I} \lambda_i(t) \nabla g_i'(y(t), t) \right] h(t) dt &= 0 \quad \forall h \in L_\infty^n[0, T], \\ \lambda_i(t) g_i(y(t), t) &= 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ \mu_j \geq 0, \quad j \in J, \quad \lambda_i(t) \geq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ (\mu, \lambda(t)) &\neq 0 \quad \text{a.e. in } [0, T]. \end{aligned} \quad (4.5)$$

If $\mu = 0$, then $\lambda(t) \neq 0$ a.e in $[0, T]$, and

$$\int_0^T \left[\sum_{i \in I} \lambda_i(t) \nabla g_i'(y(t), t) \right] h(t) dt = 0 \quad \forall h \in L_\infty^n[0, T]. \quad (4.6)$$

From $\lambda_i(t) g_i(y(t), t) = 0$, a.e. in $[0, T]$, $i \in I$, it follows that $\lambda_i(t) = 0$, $t \notin A_i(y)$. So we have

$$\sum_{i \in I} \int_{A_i(y)} \lambda_i(t) \nabla g_i'(y(t), t) h(t) dt = 0 \quad \forall h \in L_\infty^n[0, T], \quad (4.7)$$

with $\lambda_i(t) \geq 0$ a.e. in $[0, T]$, $i \in I$, not all zero, which contradicts the hypothesis that (CQ) holds at y . Thus y is a vector KKT-solution, that is, there exist $\mu \in \mathbb{R}^p$ and $\lambda \in L_\infty^m[0, T]$ satisfying (2.4)–(2.7).

As by (2.7) $\mu_j \geq 0$, $j \in J$, and by (2.6) $\lambda_i(t) \geq 0$ a.e. in $[0, T]$, $i \in I$, using (4.4) we obtain

$$\int_0^T \sum_{j \in J} \mu_j [f_j(x(t), t) - f_j(y(t), t)] dt \geq \int_0^T \sum_{j \in J} \mu_j \nabla f_j'(y(t), t) \eta(x(t), y(t), t) dt, \quad (4.8)$$

$$-\lambda_i(t) \nabla g_i'(y(t), t) \eta(x(t), y(t), t) \geq 0 \quad \text{a.e. in } A_i(y), \quad i \in I, \quad (4.9)$$

for all $x \in \mathbb{F}$. Remembering that $\lambda_i(t) = 0$, $t \notin A_i(y)$, $i \in I$, integrating the inequalities in (4.9) over $[0, T]$ and summing over I , we obtain

$$-\int_0^T \sum_{i \in I} \lambda_i(t) \nabla g'_i(y(t), t) \eta(x(t), y(t), t) dt \geq 0. \quad (4.10)$$

From (4.8) and (4.10), it follows that

$$\begin{aligned} & \int_0^T \sum_{j \in J} \mu_j [f_j(x(t), t) - f_j(y(t), t)] dt \\ & \geq \int_0^T \left[\sum_{j \in J} \mu_j \nabla f'_j(y(t), t) + \sum_{i \in I} \lambda_i(t) \nabla g'_i(y(t), t) \right] \eta(x(t), y(t), t) dt, \end{aligned} \quad (4.11)$$

for all $x \in \mathbb{F}$. Setting $h(t) = \eta(x(t), y(t), t)$ a.e. in $[0, T]$, it follows from (2.4) that the integral in the second line above is null. Thus y solves a weighting scalar problem with $\mu_j \geq 0$, $j \in J$, not all zero. \square

THEOREM 4.4. *If (CMP) is KKT-invex, then every vector KKT-solution solves a weighting scalar problem with $\mu_j \geq 0$, $j \in J$, not all zero.*

Proof. Similar to the proof of Theorem 4.3. \square

The theorems above show us that under the assumptions that (CMP) is KKT-invex and the constraints satisfy (CQ) at each $y \in \mathbb{F}$, the sets of vector KKT-solutions, weakly efficient solutions, and optimal solutions of weighting scalar problems are equal.

Under the hypothesis that the constraints satisfy (CQ), we can establish a reciprocal of Theorem 4.4.

THEOREM 4.5. *Assume that the constraints g_i , $i \in I$, satisfy (CQ) at each $y \in \mathbb{F}$. Then every vector KKT-solution solves a weighting scalar problem if and only if (CMP) is KKT-invex.*

Proof. The sufficiency part was proved in Theorem 4.4. Let us proceed to the necessity part.

Assume that every vector KKT-solution solves a weighting scalar problem. Then the system

$$\begin{aligned} & \int_0^T \left[\sum_{j \in J} \mu_j \nabla f'_j(y(t), t) + \sum_{i \in I} \lambda_i(t) \nabla g'_i(y(t), t) \right] h(t) dt = 0 \quad \forall h \in L_\infty^n[0, T], \\ & \lambda_i(t) g_i(y(t), t) = 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ & \lambda_i(t) \geq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ & \mu_j \geq 0, \quad j \in J, \quad \mu \neq 0, \\ & \int_0^T \sum_{j \in J} \mu_j [f_j(x(t), t) - f_j(y(t), t)] dt < 0, \end{aligned} \quad (4.12)$$

has no solution $(\mu, \lambda) \in \mathbb{R}^p \times L_\infty^m[0, T]$ for any $x, y \in \mathbb{F}$. Equivalently, the system

$$\begin{aligned} \int_0^T \left[\sum_{j \in J} \mu_j \nabla f'_j(y(t), t) + \sum_{i \in I} \lambda_i(t) \chi_i(t) \nabla g_i(y(t), t) \right] h(t) dt &= 0 \quad \forall h \in L_\infty^n[0, T], \\ \lambda_i(t) &\geq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \\ \mu_j &\geq 0, \quad j \in J, \quad \mu \neq 0, \\ \int_0^T \sum_{j \in J} \mu_j [f_j(x(t), t) - f_j(y(t), t)] dt &< 0, \end{aligned} \quad (4.13)$$

has no solution $(\mu, \lambda) \in \mathbb{R}^p \times L_\infty^m[0, T]$ for any $x, y \in \mathbb{F}$, where χ_i is defined as in the proof of Theorem 3.2. It is easy to see that this implies that the system

$$\begin{aligned} \int_0^T \begin{bmatrix} \nu & \mu_1 & \cdots & \mu_p \end{bmatrix} \begin{bmatrix} 0 & 1 \\ \nabla f'_1(y(t), t) & f_1(x(t), t) - f_1(y(t), t) \\ \vdots & \vdots \\ \nabla f'_p(y(t), t) & f_p(x(t), t) - f_p(y(t), t) \end{bmatrix} \begin{bmatrix} h(t) \\ \alpha \end{bmatrix} dt \\ + \int_0^T \begin{bmatrix} \lambda_1(t) & \cdots & \lambda_m(t) \end{bmatrix} \begin{bmatrix} \chi_1(t) \nabla g'_1(y(t), t) & 0 \\ \vdots & \vdots \\ \chi_m(t) \nabla g'_m(y(t), t) & 0 \end{bmatrix} \begin{bmatrix} h(t) \\ \alpha \end{bmatrix} dt = 0 \\ \forall h \in L_\infty^n[0, T], \quad \forall \alpha \in \mathbb{R}, \\ \nu > 0, \quad \mu_j \geq 0, \quad j \in J, \quad \mu \neq 0, \quad \lambda_i(t) \geq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \end{aligned} \quad (4.14)$$

has no solution $(\nu, \mu, \lambda) \in \mathbb{R}^{p+1} \times L_\infty^m[0, T]$ for any $x, y \in \mathbb{F}$. Since (CQ) holds by hypothesis, the condition (2.3) in Theorem 2.1 is satisfied. Therefore, applying that theorem, it follows that there exist $\alpha \in \mathbb{R}$ and $h \in L_\infty^n[0, T]$ such that

$$\begin{aligned} \int_0^T \begin{bmatrix} 0 & 1 \\ \nabla f'_1(y(t), t) & f_1(x(t), t) - f_1(y(t), t) \\ \vdots & \vdots \\ \nabla f'_p(y(t), t) & f_p(x(t), t) - f_p(y(t), t) \end{bmatrix} \begin{bmatrix} h(t) \\ \alpha \end{bmatrix} dt < 0, \\ \chi_i(t) \nabla g'_i(y(t), t) h(t) \leq 0 \quad \text{a.e. in } [0, T], \quad i \in I, \end{aligned} \quad (4.15)$$

for any $x, y \in \mathbb{F}$. Then $\alpha < 0$ and we can set $\alpha = -1$. Defining

$$\eta(x(t), y(t), t) = h(t) \quad \text{a.e. in } [0, T], \quad (4.16)$$

we have

$$\int_0^T \nabla f'_j(y(t), t) \eta(x(t), y(t), t) dt < \int_0^T [f_j(x(t), t) - f_j(y(t), t)] dt, \quad j \in J, \quad (4.17)$$

$$\nabla g'_i(y(t), t) \eta(x(t), y(t), t) \leq 0 \quad \text{a.e. in } A_i(y), \quad i \in I.$$

Therefore (CMP) is KKT-invex. \square

Acknowledgments

The first author was supported by BIG/UNICAMP and CNPq-Brazil, Grant no. 141168/2003-0. The second author was partially supported by Ministerio De Ciencia y Tecnología, Spain, Grant BFM 2003-06579 and by CNPq-Brazil, Grant no. 301354/03-0.

References

- [1] R. Bellman, "Bottleneck problems and dynamic programming," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 39, pp. 947–951, 1953.
- [2] E. J. Anderson and A. B. Philpott, "On the solutions of a class of continuous linear programs," *SIAM Journal on Control and Optimization*, vol. 32, no. 5, pp. 1289–1296, 1994.
- [3] A. J. V. Brandão, M. A. Rojas-Medar, and G. N. Silva, "Nonsmooth continuous-time optimization problems: necessary conditions," *Computers & Mathematics with Applications*, vol. 41, no. 12, pp. 1477–1486, 2001.
- [4] R. C. Grinold, "Continuous programming part one: linear objectives," *Journal of Mathematical Analysis and Applications*, vol. 28, pp. 32–51, 1969.
- [5] N. Levinson, "A class of continuous linear programming problems," *Journal of Mathematical Analysis and Applications*, vol. 16, no. 1, pp. 73–83, 1966.
- [6] M. C. Pullan, "An algorithm for a class of continuous linear programs," *SIAM Journal on Control and Optimization*, vol. 31, no. 6, pp. 1558–1577, 1993.
- [7] T. W. Reiland and M. A. Hanson, "Generalized Kuhn-Tucker conditions and duality for continuous nonlinear programming problems," *Journal of Mathematical Analysis and Applications*, vol. 74, no. 2, pp. 578–598, 1980.
- [8] M. A. Rojas-Medar, A. J. V. Brandão, and G. N. Silva, "Nonsmooth continuous-time optimization problems: sufficient conditions," *Journal of Mathematical Analysis and Applications*, vol. 227, no. 2, pp. 305–318, 1998.
- [9] G. J. Zalmi, "Optimality conditions and Lagrangian duality in continuous-time nonlinear programming," *Journal of Mathematical Analysis and Applications*, vol. 109, no. 2, pp. 426–452, 1985.
- [10] G. J. Zalmi, "A continuous-time generalization of Gordan's transposition theorem," *Journal of Mathematical Analysis and Applications*, vol. 110, no. 1, pp. 130–140, 1985.
- [11] G. J. Zalmi, "The Fritz John and Kuhn-Tucker optimality conditions in continuous-time nonlinear programming," *Journal of Mathematical Analysis and Applications*, vol. 110, no. 2, pp. 503–518, 1985.
- [12] G. J. Zalmi, "Sufficient optimality conditions in continuous-time nonlinear programming," *Journal of Mathematical Analysis and Applications*, vol. 111, no. 1, pp. 130–147, 1985.
- [13] L. B. dos Santos, A. J. V. Brandão, R. Osuna-Gómez, and M. A. Rojas-Medar, "Nonsmooth continuous-time multiobjective optimization problems," Tech. Rep., State University of Campinas, São Paulo, Brazil, 2005. http://www.ime.unicamp.br/rel_pesq/2005/rp35-05.html.

- [14] R. Osuna-Gómez, A. Rufián-Lizana, and P. Ruíz-Canales, “Invex functions and generalized convexity in multiobjective programming,” *Journal of Optimization Theory and Applications*, vol. 98, no. 3, pp. 651–661, 1998.
- [15] R. Osuna-Gómez, A. Beato-Moreno, and A. Rufián-Lizana, “Generalized convexity in multi-objective programming,” *Journal of Mathematical Analysis and Applications*, vol. 233, no. 1, pp. 205–220, 1999.
- [16] V. A. de Oliveira and M. A. Rojas-Medar, “Continuous-time optimization problems involving invex functions,” *Journal of Mathematical Analysis and Applications*, vol. 327, no. 2, pp. 1320–1334, 2007.
- [17] O. L. Mangasarian, *Nonlinear Programming*, vol. 10 of *Classics in Applied Mathematics*, SIAM, Philadelphia, Pa, USA, 1994.

Valeriano A. De Oliveira: Departamento de Matemática Aplicada, Instituto de Matemática, Estatística e Computação Científica, UNICAMP, CP 6065, 13083-859 Campinas, SP, Brazil
Email address: vantunes@ime.unicamp.br

Marko A. Rojas-Medar: Departamento de Matemática Aplicada, Instituto de Matemática, Estatística e Computação Científica, UNICAMP, CP 6065, 13083-859 Campinas, SP, Brazil
Email address: marko@ime.unicamp.br



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

