

Hindawi Publishing Corporation
 Discrete Dynamics in Nature and Society
 Volume 2009, Article ID 205481, 9 pages
 doi:10.1155/2009/205481

Research Article

Permanence of a Discrete Periodic Volterra Model with Mutual Interference

Lijuan Chen¹ and Liujuan Chen²

¹ College of Mathematics and Computer Science, Fuzhou University, Fuzhou, Fujian 350002, China

² Department of Mathematics and Physics, Fujian Institute of Education, Fuzhou, Fujian 350001, China

Correspondence should be addressed to Lijuan Chen, chenlijuan@fzu.edu.cn

Received 5 December 2008; Accepted 12 February 2009

Recommended by Antonia Vecchio

This paper discusses a discrete periodic Volterra model with mutual interference and Holling II type functional response. Firstly, sufficient conditions are obtained for the permanence of the system. After that, we give an example to show the feasibility of our main results.

Copyright © 2009 L. Chen and L. Chen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In 1971, Hassell introduced the concept of mutual interference between the predators and preys. Hassell [1] established a Volterra model with mutual interference as follows:

$$\begin{aligned}\dot{x}(t) &= xg(x) - \varphi(x)y^m, \\ \dot{y}(t) &= y(-d + k\varphi(x)y^{m-1} - q(y)),\end{aligned}\tag{1.1}$$

where m denote mutual interference constant and $0 < m \leq 1$.

Motivated by the works of Hassell [1], Wang and Zhu [2] considered the following Volterra model with mutual interference and Holling II type functional response:

$$\begin{aligned}\dot{x}(t) &= x(t)(r_1(t) - b_1(t)x(t)) - \frac{c_1(t)x(t)}{k + x(t)}y^m(t), \\ \dot{y}(t) &= y(t)(-r_2(t) - b_2(t)y(t)) + \frac{c_2(t)x(t)}{k + x(t)}y^m(t).\end{aligned}\tag{1.2}$$

Sufficient conditions which guarantee the existence, uniqueness, and global attractivity of positive periodic solution are obtained by employing Mawhin's continuation theorem and constructing suitable Lyapunov function.

On the other hand, it has been found that the discrete time models governed by difference equations are more appropriate than the continuous ones when the populations have nonoverlapping generations. Discrete time models can also provide efficient computational models of continuous models for numerical simulations (see [3–15]). However, to the best of the author's knowledge, until today, there are still no scholars propose and study a discrete-time analogue of system (1.2). Therefore, the main purpose of this paper is to study the following discrete periodic Volterra model with mutual interference and Holling II type functional response:

$$\begin{aligned} x(n+1) &= x(n) \exp \left\{ r_1(n) - b_1(n)x(n) - \frac{c_1(n)}{k+x(n)}y^m(n) \right\}, \\ y(n+1) &= y(n) \exp \left\{ -r_2(n) - b_2(n)y(n) + \frac{c_2(n)x(n)}{k+x(n)}y^{m-1}(n) \right\}, \end{aligned} \quad (1.3)$$

where $x(n)$ is the density of prey species at n th generation and $y(n)$ is the density of predator species at n th generation. Also, $r_1(n)$, $b_1(n)$ denote the intrinsic growth rate and density-dependent coefficient of the prey, respectively, $r_2(n)$, $b_2(n)$ denote the death rate and density-dependent coefficient of the predator, respectively, $c_1(n)$ denote the capturing rate of the predator and $c_2(n)$ represent the transformation from preys to predators. Further, m is mutual interference constant and k is a positive constant. In this paper, we always assume that $\{r_i(n)\}$, $\{b_i(n)\}$, $\{c_i(n)\}$, $i = 1, 2$, are positive T -periodic sequences and $0 < m < 1$. Here, for convenience, we denote $\bar{f} = (1/T)\sum_{n=0}^{T-1} f(n)$, $f^M = \sup_{n \in I_T} \{f(n)\}$, and $f^L = \inf_{n \in I_T} \{f(n)\}$, where $I_T = \{0, 1, 2, \dots, T-1\}$.

This paper is organized as follows. In Section 2, we will introduce a definition and establish several useful lemmas. The permanence of system (1.3) is then studied in Section 3. In Section 4, we give an example to show the feasibility of our main results.

From the view point of biology, we only need to focus our discussion on the positive solution of system (1.3). So it is assumed that the initial conditions of (1.3) are of the form

$$x(0) > 0, \quad y(0) > 0. \quad (1.4)$$

One can easily show that the solution of (1.3) with the initial condition (1.4) are defined and remain positive for all $n \in N$ where $N = \{0, 1, 2, \dots\}$.

2. Preliminaries

In this section, we will introduce the definition of permanence and several useful lemmas.

Definition 2.1. System (1.3) is said to be permanent if there exist positive constants x^* , y^* , x_* , y_* , which are independent of the solution of the system, such that for any positive

solution $(x(n), y(n))$ of system (1.3) satisfies

$$\begin{aligned} x_* &\leq \liminf_{n \rightarrow \infty} x(n) \leq \limsup_{n \rightarrow \infty} x(n) \leq x^*, \\ y_* &\leq \liminf_{n \rightarrow \infty} y(n) \leq \limsup_{n \rightarrow \infty} y(n) \leq y^*. \end{aligned} \quad (2.1)$$

Lemma 2.2. *Assume that $x(n)$ satisfies*

$$x(n+1) \leq x(n) \exp\{a(n) - b(n)x(n)\} \quad \forall n \geq n_0, \quad (2.2)$$

where $\{a(n)\}$ and $\{b(n)\}$ are positive sequences, $x(n_0) > 0$ and $n_0 \in \mathbb{N}$. Then, one has

$$\limsup_{n \rightarrow \infty} x(n) \leq B, \quad (2.3)$$

where $B = \exp(a^M - 1)/b^L$.

Lemma 2.3. *Assume that $x(n)$ satisfies*

$$x(n+1) \geq x(n) \exp\{a(n) - b(n)x(n)\} \quad \forall n \geq n_0, \quad (2.4)$$

where $\{a(n)\}$ and $\{b(n)\}$ are positive sequences, $x(n_0) > 0$ and $n_0 \in \mathbb{N}$. Also, $\limsup_{n \rightarrow \infty} x(n) \leq B$ and $b^M B / a^L > 1$. Then, one has

$$\liminf_{n \rightarrow \infty} x(n) \geq A, \quad (2.5)$$

where $A = (a^L / b^M) \exp(a^L - b^M B)$.

Proof. The proofs of Lemmas 2.2 and 2.3 are very similar to that of [8, Lemmas 1 and 2], respectively. So, we omit the detail here. \square

The following Lemma 2.4 is Lemma 2.2 of Fan and Li [12].

Lemma 2.4. *The problem*

$$x(n+1) = x(n) \exp\{a(n) - b(n)x(n)\}, \quad (2.6)$$

with $x(0) = x_0 > 0$ has at least one periodic positive solution $x^*(n)$ if both $b : \mathbb{Z} \rightarrow \mathbb{R}^+$ and $a : \mathbb{Z} \rightarrow \mathbb{R}$ are T -periodic sequences with $\bar{a} > 0$. Moreover, if $b(n) = b$ is a constant and $a^M < 1$, then $bx(n) \leq 1$ for n sufficiently large, where $x(n)$ is any solution of (2.6).

The following comparison theorem for the difference equation is Theorem 2.1 of L. Wang and M. Q. Wang [15, page 241].

Lemma 2.5. *Suppose that $f : Z^+ \times [0, +\infty)$ and $g : Z^+ \times [0, +\infty)$ with $f(n, x) \leq g(n, x)$ ($f(n, x) \geq g(n, x)$) for $n \in Z^+$ and $x \in [0, +\infty)$. Assume that $g(n, x)$ is nondecreasing with respect to the argument x . If $x(n)$ and $u(n)$ are solutions of*

$$x(n+1) = f(n, x(n)), \quad u(n+1) = g(n, u(n)), \quad (2.7)$$

respectively, and $x(0) \leq u(0)$ ($x(0) \geq u(0)$), then

$$x(n) \leq u(n), \quad (x(n) \geq u(n)) \quad (2.8)$$

for all $n \geq 0$.

3. Permanence

In this section, we establish a permanent result for system (1.3).

Proposition 3.1. *If $(H_1) : (1-m)r_2^M < 1$ holds, then for any positive solution $(x(n), y(n))$ of system (1.3), there exist positive constants x^* and y^* , which are independent of the solution of the system, such that*

$$\limsup_{n \rightarrow \infty} x(n) \leq x^*, \quad \limsup_{n \rightarrow \infty} y(n) \leq y^*. \quad (3.1)$$

Proof. Let $(x(n), y(n))$ be any positive solution of system (1.3), from the first equation of (1.3), it follows that

$$x(n+1) \leq x(n) \exp \{r_1(n) - b_1(n)x(n)\}. \quad (3.2)$$

By applying Lemma 2.2, we obtain

$$\limsup_{n \rightarrow \infty} x(n) \leq x^*, \quad (3.3)$$

where

$$x^* = \frac{1}{b_1^L} \exp(r_1^M - 1). \quad (3.4)$$

Denote $P(n) = (1/y(n))^{1-m}$. Then, from the second equation of (1.3), it follows that

$$P(n+1) = P(n) \exp \left\{ (1-m)r_2(n) + \frac{(1-m)b_2(n)}{1-\sqrt[m]{P(n)}} - \frac{(1-m)c_2(n)x(n)}{k+x(n)} P(n) \right\}, \quad (3.5)$$

which leads to

$$P(n+1) \geq P(n) \exp \{ (1-m)r_2(n) - (1-m)c_2^M P(n) \}. \quad (3.6)$$

Consider the following auxiliary equation:

$$Z(n+1) = Z(n) \exp \{ (1-m)r_2(n) - (1-m)c_2^M Z(n) \}. \quad (3.7)$$

By Lemma 2.4, (3.7) has at least one positive T -periodic solution and we denote one of them as $Z^*(n)$. Now (H_1) and Lemma 2.4 imply $(1-m)c_2^M Z(n) \leq 1$ for n sufficiently large, where $Z(n)$ is any solution of (3.7). Consider the following function:

$$g(n, Z) = Z \exp \{ (1-m)r_2(n) - (1-m)c_2^M Z \}. \quad (3.8)$$

It is not difficult to see that $g(n, Z)$ is nondecreasing with respect to the argument Z . Then, applying Lemma 2.5 to (3.6) and (3.7), we easily obtain that $P(n) \geq Z^*(n)$. So $\liminf_{n \rightarrow \infty} P(n) \geq (Z^*(n))^L$, which together with that transformation $P(n) = (1/y(n))^{1-m}$, produces

$$\limsup_{n \rightarrow \infty} y(n) \leq \frac{1}{{}^{1-m}\sqrt{(Z^*(n))^L}} \triangleq y^*. \quad (3.9)$$

Thus, we complete the proof of Proposition 3.1. \square

Proposition 3.2. *Assume that*

$$(H_2) : \left(r_1(n) - \frac{c_1(n)}{k} (y^*)^m \right)^L > 0 \quad (3.10)$$

holds, then for any positive solution $(x(n), y(n))$ of system (1.3), there exist positive constants x_ and y_* , which are independent of the solution of the system, such that*

$$\liminf_{n \rightarrow \infty} x(n) \geq x_*, \quad \liminf_{n \rightarrow \infty} y(n) \geq y_*, \quad (3.11)$$

where y^ can be seen in Proposition 3.1.*

Proof. Let $(x(n), y(n))$ be any positive solution of system (1.3). From (H_2) , there exists a small enough positive constant ε such that

$$\left(r_1(n) - \frac{c_1(n)}{k} (y^* + \varepsilon)^m \right)^L > 0. \quad (3.12)$$

Also, according to Proposition 3.1, for above ε , there exists $N_1 > 0$ such that for $n \geq N_1$,

$$y(n) \leq y^* + \varepsilon. \quad (3.13)$$

Then, from the first equation of (1.3), for $n \geq N_1$, we have

$$x(n+1) \geq x(n) \exp \left\{ r_1(n) - \frac{c_1(n)}{k} (y^* + \varepsilon)^m - b_1(n)x(n) \right\}. \quad (3.14)$$

Let $a_1(n, \varepsilon) = r_1(n) - (c_1(n)/k)(y^* + \varepsilon)^m$, so the above inequality follows that

$$x(n+1) \geq x(n) \exp \{ a_1(n, \varepsilon) - b_1(n)x(n) \}. \quad (3.15)$$

Because $(a_1(n, \varepsilon))^L < r_1^L$ and $b_1^L < b_1^M$, we have

$$\frac{b_1^M}{(a_1(n, \varepsilon))^L} x^* > \frac{b_1^M \exp(r_1^M - 1)}{r_1^L b_1^L} > 1. \quad (3.16)$$

Here, we use the fact $\exp(r_1^M - 1) > r_1^M$. From (3.12) and (3.15), by Lemma 2.3, we have

$$\liminf_{n \rightarrow \infty} x(n) \geq \frac{(a_1(n, \varepsilon))^L}{b_1^M} \exp \{ (a_1(n, \varepsilon))^L - b_1^M x^* \}. \quad (3.17)$$

Setting $\varepsilon \rightarrow 0$ in the above inequality leads to

$$\liminf_{n \rightarrow \infty} x(n) \geq \frac{a_1^L}{b_1^M} \exp \{ a_1^L - b_1^M x^* \} \triangleq x_*, \quad (3.18)$$

where

$$a_1(n) = r_1(n) - \frac{c_1(n)}{k} (y^*)^m. \quad (3.19)$$

For above ε , there exists $N_2 > N_1$ such that for $n \geq N_2$, $x(n) \geq x_* - \varepsilon$. So from (3.5), we obtain that

$$P(n+1) \leq P(n) \exp \left\{ (1-m)(r_2(n) + b_2(n)(y^* + \varepsilon)) - \frac{(1-m)c_2(n)(x_* - \varepsilon)}{k + x_* - \varepsilon} P(n) \right\}. \quad (3.20)$$

Consider the following auxiliary equation:

$$W(n+1) = W(n) \exp \left\{ (1-m)(r_2(n) + b_2(n)(y^* + \varepsilon)) - \frac{(1-m)c_2(n)(x_* - \varepsilon)}{k + x_* - \varepsilon} W(n) \right\}. \quad (3.21)$$

By Lemma 2.4, (3.21) has at least one positive T -periodic solution and we denote one of them as $W^*(n)$.

Let

$$R(n) = \ln(P(n)), \quad Y(n) = \ln(W^*(n)). \quad (3.22)$$

Then,

$$\begin{aligned} R(n+1) - R(n) &\leq (1-m)(r_2(n) + b_2(n)(y^* + \varepsilon)) - \frac{(1-m)c_2(n)(x_* - \varepsilon)}{k + x_* - \varepsilon} \exp\{R(n)\}, \\ Y(n+1) - Y(n) &= (1-m)(r_2(n) + b_2(n)(y^* + \varepsilon)) - \frac{(1-m)c_2(n)(x_* - \varepsilon)}{k + x_* - \varepsilon} \exp\{Y(n)\}. \end{aligned} \quad (3.23)$$

Set

$$U(n) = R(n) - Y(n). \quad (3.24)$$

Then,

$$U(n+1) - U(n) \leq -\frac{(1-m)c_2(n)(x_* - \varepsilon)}{k + x_* - \varepsilon} \exp\{Y(n)\} [\exp\{U(n)\} - 1]. \quad (3.25)$$

In the following we distinguish three cases.

Case 1. $\{U(n)\}$ is eventually positive. Then, from (3.25), we see that $U(n+1) < U(n)$ for any sufficiently large n . Hence, $\lim_{n \rightarrow \infty} U(n) = 0$, which implies that

$$\limsup_{n \rightarrow \infty} P(n) \leq (W^*(n))^M. \quad (3.26)$$

Case 2. $\{U(n)\}$ is eventually negative. Then, from (3.24), we can also obtain (3.26).

Case 3. $\{U(n)\}$ oscillates about zero. In this case, we let $\{U(n_{st})\}$ ($s, t \in N$) be the positive semicycle of $\{U(n)\}$, where $U(n_{s1})$ denotes the first element of the s th positive semicycle of $\{U(n)\}$. From (3.25), we know that $U(n+1) < U(n)$ if $U(n) > 0$. Hence, $\limsup_{n \rightarrow \infty} U(n) = \limsup_{s \rightarrow \infty} U(n_{s1})$. From (3.25), and $U(n_{s1} - 1) < 0$, we can obtain

$$\begin{aligned} U(n_{s1}) &\leq \frac{(1-m)c_2(n_{s1})(x_* - \varepsilon)}{k + x_* - \varepsilon} \exp\{Y(n_{s1})\} [1 - \exp\{U(n_{s1})\}], \\ &\leq \frac{(1-m)c_2^M(x_* - \varepsilon)}{k + x_* - \varepsilon} (W^*(n))^M. \end{aligned} \quad (3.27)$$

From (3.22) and (3.24), we easily obtain

$$\limsup_{n \rightarrow \infty} P(n) \leq (W^*(n))^M \exp\left\{\frac{(1-m)c_2^M(x_* - \varepsilon)}{k + x_* - \varepsilon} (W^*(n))^M\right\}. \quad (3.28)$$




Figure 1: Dynamics behavior of system (1.3) with initial condition $(x(0), y(0)) = (0.6, 0.03)$.

Setting $\varepsilon \rightarrow 0$ in the above inequality leads to

$$\limsup_{n \rightarrow \infty} P(n) \leq (W^*(n))^M \exp \left\{ \frac{(1-m)c_2^M x_*}{k+x_*} (W^*(n))^M \right\} \triangleq P^*, \quad (3.29)$$

which together with that transformation $P(n) = (1/y(n))^{1-m}$, we have

$$\liminf_{n \rightarrow \infty} y(n) \geq \frac{1}{\sqrt[1-m]{P^*}} \triangleq y_*. \quad (3.30)$$

Thus, we complete the proof of Proposition 3.2. \square

Theorem 3.3. *Assume that (H_1) and (H_2) hold, then system (1.3) is permanent.*

It should be noticed that, from the proofs of Propositions 3.1 and 3.2, one knows that under the conditions of Theorem 3.3, the set $\Omega = \{(x, y) \mid x_* \leq x \leq x^*, y_* \leq y \leq y^*\}$ is an invariant set of system (1.3).

4. Example

In this section, we give an example to show the feasibility of our main result.

Example 4.1. Consider the following system

$$\begin{aligned} x(n+1) &= x(n) \exp \left\{ 0.7 + 0.1 \sin(n) - 0.7x(n) - \frac{0.4y^{0.6}(n)}{2.4+x(n)} \right\}, \\ y(n+1) &= y(n) \exp \left\{ -(0.8 + 0.1 \cos(n)) - (1.1 + 0.1 \sin(n))x(n) - \frac{0.6y^{-0.4}(n)}{2.4+x(n)} \right\}, \end{aligned} \quad (4.1)$$

where $r_1(n) = 0.7 + 0.1 \sin(n)$, $b_1(n) = 0.7$, $c_1(n) = 0.4$, $r_2(n) = 0.8 + 0.1 \cos(n)$, $b_2(n) = 1.1 + 0.1 \sin(n)$, $c_2(n) = 0.6$, $m = 0.6$, $k = 2.4$.

By simple computation, we have $y^* \approx 1.1405$. Thus, one could easily see that

$$\left(r_1(n) - \frac{c_1(n)}{k} (y^*)^m \right)^L \approx 0.4197 > 0, \quad (1 - m)(r_2(n))^M = 0.36 < 1. \quad (4.2)$$

Clearly, conditions (H_1) and (H_2) are satisfied, then system (1.3) is permanent.


Figure 1 shows the dynamics behavior of system (1.3).

Acknowledgments

The authors would like to thank the referees for their helpful comments and suggestions which greatly improve the presentation of the paper. This work was supported by the Program for New Century Excellent Talents in Fujian Province University (NCETFJ) and the Foundation of Fujian Education Bureau (JB08028).

References

- [1] M. P. Hassell, "Density-dependence in single-species populations," *The Journal of Animal Ecology*, vol. 44, no. 1, pp. 283–295, 1975.
- [2] K. Wang and Y. Zhu, "Global attractivity of positive periodic solution for a Volterra model," *Applied Mathematics and Computation*, vol. 203, no. 2, pp. 493–501, 2008.
- [3] F. Chen, "Permanence of a discrete N -species cooperation system with time delays and feedback controls," *Applied Mathematics and Computation*, vol. 186, no. 1, pp. 23–29, 2007.
- [4] F. Chen, "Permanence of a discrete n -species food-chain system with time delays," *Applied Mathematics and Computation*, vol. 185, no. 1, pp. 719–726, 2007.
- [5] F. Chen, L. Wu, and Z. Li, "Permanence and global attractivity of the discrete Gilpin-Ayala type population model," *Computers & Mathematics with Applications*, vol. 53, no. 8, pp. 1214–1227, 2007.
- [6] F. Chen, "Permanence and global attractivity of a discrete multispecies Lotka-Volterra competition predator-prey systems," *Applied Mathematics and Computation*, vol. 182, no. 1, pp. 3–12, 2006.
- [7] M. Fan and K. Wang, "Periodic solutions of a discrete time nonautonomous ratio-dependent predator-prey system," *Mathematical and Computer Modelling*, vol. 35, no. 9–10, pp. 951–961, 2002.
- [8] X. Yang, "Uniform persistence and periodic solutions for a discrete predator-prey system with delays," *Journal of Mathematical Analysis and Applications*, vol. 316, no. 1, pp. 161–177, 2006.
- [9] Y. Li, "Positive periodic solutions of discrete Lotka-Volterra competition systems with state dependent and distributed delays," *Applied Mathematics and Computation*, vol. 190, no. 1, pp. 526–531, 2007.
- [10] Y. Chen and Z. Zhou, "Stable periodic solution of a discrete periodic Lotka-Volterra competition system," *Journal of Mathematical Analysis and Applications*, vol. 277, no. 1, pp. 358–366, 2003.
- [11] Z. Zhou and X. Zou, "Stable periodic solutions in a discrete periodic logistic equation," *Applied Mathematics Letters*, vol. 16, no. 2, pp. 165–171, 2003.
- [12] Y.-H. Fan and W.-T. Li, "Permanence for a delayed discrete ratio-dependent predator-prey system with Holling type functional response," *Journal of Mathematical Analysis and Applications*, vol. 299, no. 2, pp. 357–374, 2004.
- [13] H.-F. Huo and W.-T. Li, "Existence and global stability of periodic solutions of a discrete ratio-dependent food chain model with delay," *Applied Mathematics and Computation*, vol. 162, no. 3, pp. 1333–1349, 2005.
- [14] Y. Li and L. Zhu, "Existence of positive periodic solutions for difference equations with feedback control," *Applied Mathematics Letters*, vol. 18, no. 1, pp. 61–67, 2005.
- [15] L. Wang and M. Q. Wang, *Ordinary Difference Equation*, Xinjiang University Press, Xinjiang, China, 1991.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

