

Research Article

Epidemiologic Study of *Blastocystis* Infection in an Urban Community in the Philippines

Maria Luz B. Belleza,¹ Jessa Louise C. Cadacio,² Maridel P. Borja,³ Juan Antonio A. Solon,¹ Mildred A. Padilla,⁴ Pilarita N. Tongol-Rivera,¹ and Windell L. Rivera^{5,6}

¹Department of Parasitology, College of Public Health, University of the Philippines, Ermita, 1000 Manila, Philippines

²College of Medicine, University of the Philippines, Ermita, 1000 Manila, Philippines

³Department of Epidemiology and Biostatistics, College of Public Health, University of the Philippines, 1000 Manila, Philippines

⁴Department of Veterinary Paraclinical Sciences, College of Veterinary Medicine, University of the Philippines, Los Baños, 4031 Laguna, Philippines

⁵Institute of Biology, College of Science, University of the Philippines, Diliman, 1101 Quezon City, Philippines

⁶Molecular Protozoology Laboratory, Natural Sciences Research Institute, University of the Philippines, Diliman, 1101 Quezon City, Philippines

Correspondence should be addressed to Windell L. Rivera; wrivera@science.upd.edu.ph

Received 1 March 2015; Accepted 29 April 2015

Academic Editor: Pam R. Factor-Litvak

Copyright © 2015 Maria Luz B. Belleza et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Blastocystis has been considered as the most common intestinal parasite in humans and has an augmented impact on public health. However, the prevalence of this parasite in the Philippines has not been determined. To contribute to a better understanding of the epidemiology of this infection, a cross-sectional study aimed at providing the first documented data on the prevalence and correlates, sociodemographic factors, hygiene practices, source of water supply, and dog ownership, associated with *Blastocystis* infection was carried out in randomly selected communities at Pateros, Metro Manila. Fecal samples from respondents were collected and cultured in diphasic agar medium for 3–7 days and examined using light microscopy. Of the 1,271 respondents, 12.98% (95% CI: 11.13–14.83) were detected positive for *Blastocystis*. Among the correlates of *Blastocystis* infection, dog ownership was found significantly associated as confirmed by multivariate analysis. Therefore, this factor should be considered in information to create awareness about *Blastocystis* and to prevent and control *Blastocystis* infection in particular and diarrheal diseases in general. Further studies using molecular approaches to distinguish subtype and to determine genetic characteristics of isolates from humans and dogs are recommended to analyze their relationship and provide more conclusive evidence of cross-transmission.

1. Introduction

Blastocystis sp. is currently the most common intestinal protist found in human feces and is considered an emerging parasite with a worldwide distribution [1, 2]. The accepted mode of transmission of the parasite is through the fecal-oral route [3]. Its pathogenic role in humans remains uncertain since *Blastocystis* infections are both symptomatic and asymptomatic [1]. Nevertheless, it is being associated with various nonspecific gastrointestinal symptoms including diarrhea, abdominal pain, flatulence, anorexia, nausea, and

vomiting [4]. The parasite may also be linked to irritable bowel syndrome and inflammatory bowel disease [5–8].

The prevalence of *Blastocystis* infection is generally higher in developing than in industrialized countries partly because of poor sanitary conditions, consumption of contaminated food or water, and close animal contact [9–12]. *Blastocystis* infection is being linked with demographic factors such as age, gender, and level of education as well as exposure factors such as hygiene, source of water supply, and exposure to animals [13–16]. Moreover, higher risks of infection and high prevalence have been identified in food and animal handlers,

providing conclusive evidence on its zoonotic potential [17–21]. In the Philippines, isolates of *Blastocystis* subtypes from humans were classified in the same corresponding subtypes as isolates from chickens, monkeys, and pigs in close contact with humans [22].

Diarrhea is still one of the leading causes of morbidity and is included in the top ten notifiable diseases in the Philippines [23–25]. Recent developments associating *Blastocystis* with diarrhea and showing the protist's zoonotic potential have become the bases of this community-based study. This study aimed to determine the prevalence of *Blastocystis* infection in humans using culture and light microscopy techniques and to identify which factors are associated with *Blastocystis* infection using univariate and multivariate analyses. The independent variables studied were age, gender, level of education, hygiene practices, water supply, and dog ownership.

2. Methods

2.1. Study Area and Population. A cross-sectional study was conducted in Pateros, Metro Manila, Philippines, from April 2011 to February 2012 (Figure 1). This urban community, which is mainly residential, consists of ten villages (locally called *barangay*) with a total population of 60,688 and is located southeast of Metropolitan Manila [25]. Pateros has a median age of 24 years and a sex ratio of 97.5 males for every 100 females [25]. It has population proportion of young dependents (0 to 14 years) of 32.5% and has 64.1% of economically active population (15 to 64 years). One of the great concerns of local health officials is environmental sanitation, particularly potability of water sources, toilet and excreta disposal facilities, and environmental pollution. Diarrhea was reported among the top ten notifiable diseases and was cause of consultations in the health centers in Pateros [26]. The study employed a three-stage random sampling among permanent residents of households in five different villages. The study population consisted of individuals with age > 1 year and those who had not taken antiprotozoal or antidiarrheal medications two weeks prior to sample collection.

2.2. Data Collection. The study utilized pretested interview schedule and direct observation to collect information on the correlates of *Blastocystis* infection such as sociodemographic factors (age, gender, and level of education) and exposure factors (hand washing, excreta disposal, source of water supply, and dog ownership). Applicability of the questionnaire was determined using a pretest group with similar characteristics as the target population. For children who have reduced ability to judge, their parents or guardians responded on their behalf.

2.3. Sample Collection and Processing of Fecal Specimens. Containers prelabeled with individual's name and identification number and sticks were distributed to each participant. A total of 1,271 stool specimens from humans were collected using three-stage random sampling design (municipality, *barangay* or village, and then household) and transported

immediately to the Molecular Protozoology Laboratory, Natural Sciences Research Institute, University of the Philippines, Diliman, Quezon City, Philippines, for processing and examination. The presence of parasite and stool consistency (formed or diarrheic, mucoid or watery) was determined using gross examination.

2.4. Isolation and In Vitro Cultivation of *Blastocystis*. Approximately 50 mg stool samples were aseptically inoculated in diphasic medium (1.5% nonnutrient agar overlaid with buffer solution containing 137 mM NaCl, 19.6 mM Na₂HPO₄, 1.98 mM KH₂PO₄, and 3.78 mM L-asparagine) supplemented with 10% heat-inactivated horse serum (Gibco, Life Technologies, Carlsbad, CA, USA) and penicillin-streptomycin antibiotics [18]. Samples were incubated at 37°C for 3–7 days and examined for *Blastocystis* using light microscopy. Samples having the characteristic morphology of *Blastocystis* under unstained microscopic examination were considered positive for culture. Cultures were reported as negative when there was no observed parasite growth until the last day of incubation.

2.5. Data Management and Statistical Analysis. Data collected from individuals who were positive for culture and completed the questionnaire were computed, coded, and analyzed using STATA Standard Edition version 11.0 for Windows (StataCorp, College Station, TX, USA). Prevalence of *Blastocystis* infection for humans was estimated at 95% confidence interval (95% CI). Factors having p value ≤ 0.25 in univariate analysis were considered as potential confounders. A percent change in estimate of odds ratio (OR) ≥ 10 was used as basis of significant confounding effect. Identification of association between the studied factors and *Blastocystis* infection was conducted using multivariate analysis.

2.6. Ethical Issues. Prior to data collection, the study protocol was reviewed and approved by the Ethics Committee of the College of Public Health, University of the Philippines, Ermita, Manila, Philippines, and permission for field work was secured from the City Mayor through the Municipality Health Officer.

3. Results

3.1. Demographic and Socioeconomic Profiles. A total of 1,271 individuals aged 1 to 70 and above participated in this study with a median age of 24.7 ± 19.9 years. The majority of participants were females (60.82%) and most of them belong to age brackets 15–29 (26.12%) and 30–44 (19.35%). Overall, the studied population came from an average socioeconomic background with more than half having more than 6 years of formal education. Among the exposure factors studied (Table 1), hygiene practices showed that most of the respondents wash their hands with soap and water immediately after using toilet (79.94%). Regarding excreta disposal, seven out of 10 (78.99%) had family owned toilets. Lastly, only a few respondents owned dogs (12.27%).




FIGURE 1: Map showing the location of the villages in Pateros, Metro Manila, Philippines, involved in the study.

3.2. Prevalence and Distribution of *Blastocystis* Infection. The overall prevalence of *Blastocystis* infection in humans was 12.98% (95% CI: 11.13%–14.83%). Both males and females had almost equal prevalence (12.65%, 95% CI: 9.75%–15.38% and 13.20%, 95% CI: 10.81%–15.59%). Figure 2 shows *Blastocystis* prevalence (%) according to age of the subjects in years. High school educational level on the other hand had the highest prevalence when grouped according to education (15.78%) (Table 1). Lastly, one for every four dog owners (25%, 95% CI: 18.18%–32.18%) was found positive with *Blastocystis*.

3.3. Statistical Analysis. Potential confounders ($p \leq 0.25$) such as sociodemographic factors (age and level of education), hygiene practices (hand washing and excreta disposal), water supply, and dog ownership, identified using univariate analysis, were included in multivariate analysis. Among the tested potential confounders, there was no significant confounding effect to association between *Blastocystis* infection and the tested predictors (% change in estimate of OR ≥ 10). Results of multivariate analysis showed that only dog ownership was significantly associated with *Blastocystis* infection (Table 2). The odds of having *Blastocystis* infection were 2.6 times higher among dog owners than nondog owners (OR = 2.6, 90% CI: 1.9–3.7, $p = 0.000$). Associations of other factors such as sociodemographic factors (OR = 1.0, 90% CI: 0.7–1.3, OR = 3.0, 90% CI: 1.3–5.1, and OR = 1.8, 90% CI: 1.2–2.6), hygiene practices (OR = 1.6, 90% CI: 1.0–2.5 and OR = 1.7, 90% CI: 1.0–2.5), and water supply (OR = 1.4, 90% CI: 1.0–2.0) to *Blastocystis* infection were weak and not statistically significant.




FIGURE 2: Graph showing *Blastocystis* prevalence (%) versus age (years).

4. Discussion

Infection with *Blastocystis* is a common health problem in many tropical and subtropical areas of the world, especially in developing countries. In the Philippines, studies on the prevalence and correlates of *Blastocystis* sp. are not well documented. This study was the first investigation on the

TABLE 1: Prevalence of *Blastocystis* infection among permanent residents of Pateros, Philippines, according to sociodemographic and exposure factors.

	<i>n</i> (%)	% infected	95% CI
Overall	1271	12.98	11.13–14.83
<i>Sociodemographic factors</i>			
Gender			
Male	498 (39.18)	12.65	9.75–15.38
Female	773 (60.18)	13.20	10.81–15.59
Age (in years)			
1–4	165 (12.98)	7.88	3.75–12.01
5–14	239 (18.80)	15.06	10.51–19.61
15–29	332 (26.12)	13.25	9.60–16.91
30–44	246 (19.35)	14.23	9.85–18.61
45–59	195 (15.34)	12.8	8.11–17.53
60–69	60 (4.72)	8.33	1.27–15.39
70 and above	34 (2.68)	20.59	6.78–34.40
Level of education			
College	331 (26.04)	10.27	6.99–13.55
High school	431 (33.91)	15.78	12.33–19.23
Elementary	306 (24.08)	12.09	8.43–15.75
No education	203 (15.97)	12.81	8.19–17.41
<i>Exposure factors</i>			
Hygiene practices			
Hand washing			
Wash hands with soap and water immediately after using toilet	1016 (79.94)	14.17	12.03–16.32
Wash hands with water only after using toilet	162 (12.75)	8.64	4.30–12.99
Wash hands with soap and water but delays washing for more than 5 minutes after using toilet	93 (7.32)	7.52	2.13–12.92
Excreta disposal			
Family owned toilet	1003 (78.99)	14.04	11.89–16.20
Communal toilet	267 (21.01)	8.99	5.54–12.43
Water supply			
Public water system	1085 (85.37)	12.53	10.56–14.51
Communal faucet	186 (14.63)	15.59	10.35–20.82
Dog ownership			
Nondog owner	1115 (87.73)	11.30	9.43–13.16
Dog owner	156 (12.27)	25.00	18.18–32.18

n: number of examined.

prevalence and epidemiology of *Blastocystis* in an urban community.

This study determined the prevalence of *Blastocystis* using culture and light microscopy techniques and its association with various factors using univariate and multivariate analyses. Culture method was preferred because of its higher sensitivity and specificity compared to direct fecal smear microscopy [27] and stool polymerase chain reaction (PCR) [28, 29]. Santos and Rivera [28] considered *in vitro* culture as the gold standard in detecting *Blastocystis* cells and reported sensitivity of the following methods: 19.4% for direct fecal smear method, 19.4% for PCR from stool, and 66.7% for PCR from *Blastocystis* culture. However, *in vitro* culture is a selective technique; it is affected by the composition of medium used and the protocol applied in cultivation [28]. Thus, some isolates of *Blastocystis* are refractory to *in vitro* culture. Stensvold et al. [30] reported 100% sensitivity and specificity for culture when compared with formol-ethyl

acetate concentration (FECT), trichrome staining, and xenic *in vitro* culture using PCR. Roberts et al. [31] observed 82.6% sensitivity and 100% specificity for culture. In the same year, conventional polymerase chain reaction (PCR) was found to be the most effective [31]. However, factors like requirement for special equipment (PCR machine), high cost, and need for intensive labor limited its use in this study. Compared with PCR, culture method is a cost-effective method for *Blastocystis* detection in stool, and it can also yield valid prevalence estimates. Lastly, culture method has high detection rate, since *Blastocystis* are allowed to grow and propagate, even starting with low infection.

Results show prevalence of 12.98% (~13%) in the study population. High prevalence rates were found among individuals aged 5–59 (79.61%) and those who owned dogs (25%). With regard to age, a possible reason may be an increased exposure of individuals to the parasite. Studies in endemic areas of Nicaragua [32], Bangladesh [33], and Brazil [34]

TABLE 2: Final models for various factors and association with *Blastocystis* infection.

	Adjusted OR (90% CI)	<i>p</i> value
Final Model 1: association between sociodemographic factors and <i>Blastocystis</i>		
<i>Sociodemographic factors</i>		
Gender		
Male*	1.0 (—)	—
Female	1.0 (0.7–1.3)	0.919
Age (in years)		
1–4*	1.0 (—)	—
5–59	3.0 (1.3–5.1)	0.020
60 and above	2.0 (0.8–4.5)	0.186
Level of education		
College*	1.0 (—)	—
High school	1.8 (1.2–2.6)	0.179
Elementary	1.4 (0.9–2.1)	0.040
No education	3.0 (1.7–5.4)	0.002
Final Model 2: association between hygiene practices and <i>Blastocystis</i>		
<i>Hygiene practices</i>		
Hand washing		
With soap and water immediately*	1.0 (—)	—
With water only or delayed washing	1.6 (1.0–2.5)	0.091
Excreta disposal		
Family owned toilet*	1.0 (—)	—
Communal faucet	1.7 (1.1–2.5)	0.036
Final Model 3: association between source of water supply and <i>Blastocystis</i> infection		
<i>Water supply</i>		
Public water system*	1.0 (—)	—
Communal toilet	1.4 (1.0–2.0)	0.135
Final Model 4: association between dog ownership and <i>Blastocystis</i> infection		
<i>Dog ownership</i>		
Nondog owner*	1.0 (—)	—
Dog owner	2.6 (1.9–3.7)	0.000

*Reference.

showed peak prevalence of *Blastocystis* in age group 5–14. The determined prevalence rate in this study was close to the 12% reported prevalence of the Department of Parasitology in University of the Philippines Manila among prescreened clients referred from hospitals and travel agencies (unpublished data). This study shows that stool culture method is more sensitive because it allows even few *Blastocystis* to grow and multiply.

Several factors that may be associated with risk of *Blastocystis* infection, namely, sociodemographic factors (age, gender, and level of education), hygiene practices (hand washing and excreta disposal), source of water supply, and dog ownership, were analyzed in this study. Multivariate analysis identified significant association of dog ownership with *Blastocystis* infection. Dog ownership as a potential risk factor may be attributed to the zoonotic potential of *Blastocystis*. Doyle et al. [35] observed that individuals who had close contact with animals, mainly pets, could be found positive for *Blastocystis* infection. In another study, Salim et al. [20] observed animal handlers in Malaysia and reported that exposure of animal handlers to their animals was associated with *Blastocystis* infection ($p = 0.0000313$).

5. Conclusion

The prevalence of *Blastocystis* was 12.98% in Pateros, Metro Manila. Such data is indicative of the probability of acquiring this parasite in this community. Among the correlates studied, dog ownership was significantly associated with *Blastocystis* infection. The borderline confidence interval of this factor showed positive direction to association with *Blastocystis* infection. This factor could be considered to have important role in the transmission of *Blastocystis* infection, and understanding it provides better interventions in its prevention and control.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors are grateful to the following colleagues for providing comments for unpublished results and technical

assistance: Kim L. Cochon and Maylin C. Palatino (University of the Philippines, Manila, Philippines) and Davin Edric V. Adao (University of the Philippines, Diliman, Quezon City, Philippines). This work was supported financially by the Philippine Department of Science and Technology-Science Education Institute and the Natural Sciences Research Institute, University of the Philippines, Diliman, Quezon City, Philippines.

References

- [1] K. S. W. Tan, "New insights on classification, identification, and clinical relevance of *Blastocystis* spp.," *Clinical Microbiology Reviews*, vol. 21, no. 4, pp. 639–665, 2008.
- [2] M. A. Alfellani, C. R. Stensvold, A. Vidal-Lapiedra, E. S. U. Onuoha, A. F. Fagbenro-Beyioku, and C. G. Clark, "Variable geographic distribution of *Blastocystis* subtypes and its potential implications," *Acta Tropica*, vol. 126, no. 1, pp. 11–18, 2013.
- [3] H. Yoshikawa, K. Yoshida, A. Nakajima, K. Yamanari, S. Iwatani, and I. Kimata, "Fecal-oral transmission of the cyst form of *Blastocystis hominis* in rats," *Parasitology Research*, vol. 94, no. 6, pp. 391–396, 2004.
- [4] K. S. W. Tan, H. Mirza, J. D. W. Teo, B. Wu, and P. A. MacAry, "Current views on the clinical relevance of *Blastocystis* spp.," *Current Infectious Disease Reports*, vol. 12, no. 1, pp. 28–35, 2010.
- [5] N. Abe, "Molecular and phylogenetic analysis of *Blastocystis* isolates from various hosts," *Veterinary Parasitology*, vol. 120, no. 3, pp. 235–242, 2004.
- [6] D. J. Sheehan, B. G. Raucher, and J. C. McKittrick, "Association of *Blastocystis hominis* with signs and symptoms of human disease," *Journal of Clinical Microbiology*, vol. 24, no. 4, pp. 548–550, 1986.
- [7] P. Poirier, I. Wawrzyniak, C. P. Vivarès, F. Delbac, and H. El Alaoui, "New insights into *Blastocystis* spp.: a potential link with irritable bowel syndrome," *PLoS Pathogens*, vol. 8, no. 3, Article ID e1002545, 2012.
- [8] K. F. Boorom, H. Smith, L. Nimri et al., "Oh my aching gut: irritable bowel syndrome, *Blastocystis*, and asymptomatic infection," *Parasites & Vectors*, vol. 1, no. 1, article 40, 2008.
- [9] L. I. Lee, T. T. Chye, B. M. Karmacharya, and S. K. Govind, "*Blastocystis* sp.: waterborne zoonotic organism, a possibility?" *Parasites and Vectors*, vol. 5, no. 1, article 130, 2012.
- [10] L.-H. Li, X.-N. Zhou, Z.-W. Du et al., "Molecular epidemiology of human *Blastocystis* in a village in Yunnan province, China," *Parasitology International*, vol. 56, no. 4, pp. 281–286, 2007.
- [11] F. Eroglu, A. Genc, G. Elgun, and I. S. Koltas, "Identification of *Blastocystis hominis* isolates from asymptomatic and symptomatic patients by PCR," *Parasitology Research*, vol. 105, no. 6, pp. 1589–1592, 2009.
- [12] T. K. Graczyk, C. K. Shiff, L. Tamang, F. Munsaka, A. M. Beitin, and W. J. Moss, "The association of *Blastocystis hominis* and *Endolimax nana* with diarrheal stools in Zambian school-age children," *Parasitology Research*, vol. 98, no. 1, pp. 38–43, 2005.
- [13] A. Duda, D. J. Stenzel, and P. F. L. Boreham, "Detection of *Blastocystis* sp. in domestic dogs and cats," *Veterinary Parasitology*, vol. 76, no. 1–2, pp. 9–17, 1998.
- [14] L. F. Nimri, "Evidence of an epidemic of *Blastocystis hominis* infections in preschool children in Northern Jordan," *Journal of Clinical Microbiology*, vol. 31, no. 10, pp. 2706–2708, 1993.
- [15] P. Taamasri, M. Mungthin, R. Rangsin, B. Tongupprakarn, W. Areekul, and S. Leelayoova, "Transmission of intestinal blastocystosis related to the quality of drinking water," *Southeast Asian Journal of Tropical Medicine and Public Health*, vol. 31, no. 1, pp. 112–117, 2000.
- [16] K. Suresh, H. V. Smith, and T. C. Tan, "Viable *Blastocystis* cysts in Scottish and Malaysian sewage samples," *Applied and Environmental Microbiology*, vol. 71, no. 9, pp. 5619–5620, 2005.
- [17] H. Yoshikawa, Z. Wu, K. Pandey et al., "Molecular characterization of *Blastocystis* isolates from children and rhesus monkeys in Kathmandu, Nepal," *Veterinary Parasitology*, vol. 160, no. 3–4, pp. 295–300, 2009.
- [18] H. Yoshikawa, Z. Wu, I. Kimata et al., "Polymerase chain reaction-based genotype classification among human *Blastocystis hominis* populations isolated from different countries," *Parasitology Research*, vol. 92, no. 1, pp. 22–29, 2004.
- [19] H. Yoshikawa, Z. Wu, I. Nagano, and Y. Takahashi, "Molecular comparative studies among *Blastocystis* isolates obtained from humans and animals," *Journal of Parasitology*, vol. 89, no. 3, pp. 585–594, 2003.
- [20] H. R. Salim, G. S. Kumar, S. Vellayan et al., "*Blastocystis* in animal handlers," *Parasitology Research*, vol. 85, no. 12, pp. 1032–1033, 1999.
- [21] U. Parkar, R. J. Traub, S. Vitali et al., "Molecular characterization of *Blastocystis* isolates from zoo animals and their animal-keepers," *Veterinary Parasitology*, vol. 169, no. 1–2, pp. 8–17, 2010.
- [22] W. L. Rivera, "Phylogenetic analysis of *Blastocystis* isolates from animal and human hosts in the Philippines," *Veterinary Parasitology*, vol. 156, no. 3–4, pp. 178–182, 2008.
- [23] Health and Welfare, *Philippine Yearbook*, 2011, http://www.census.gov.ph/sites/default/files/2011PY_Health%20and%20Welfare.pdf.
- [24] National Statistics Office, *The Philippines in Figures, Manila, Philippines*, National Statistics Office, 2006.
- [25] Department of Health. National Epidemiology Center, *The 2009 Philippine Health Statistics*, 2009, <http://www.doh.gov.ph/sites/default/files/PHILIPPINE%20HEALTH%20STATISTICS%202009.0.pdf>.
- [26] Pateros Health Department Annual Report, 2010.
- [27] S. Leelayoova, P. Taamasri, R. Rangsin, T. Naaglor, U. Thathaisong, and M. Mungthin, "In-vitro cultivation: a sensitive method for detecting *Blastocystis hominis*," *Annals of Tropical Medicine and Parasitology*, vol. 96, no. 8, pp. 803–807, 2002.
- [28] H. J. Santos and W. L. Rivera, "Comparison of direct fecal smear microscopy, culture, and polymerase chain reaction for the detection of *Blastocystis* sp. in human stool samples," *Asian Pacific Journal of Tropical Medicine*, vol. 6, no. 10, pp. 780–784, 2013.
- [29] S. Termmathurapoj, S. Leelayoova, P. Aimpun et al., "The usefulness of short-term *in vitro* cultivation for the detection and molecular study of *Blastocystis hominis* in stool specimens," *Parasitology Research*, vol. 93, no. 6, pp. 445–447, 2004.
- [30] C. R. Stensvold, H. V. Nielsen, K. Mølbak, and H. V. Smith, "Pursuing the clinical significance of *Blastocystis*—diagnostic limitations," *Trends in Parasitology*, vol. 25, no. 1, pp. 23–29, 2009.
- [31] T. Roberts, J. Barratt, J. Harkness, J. Ellis, and D. Stark, "Comparison of microscopy, culture, and conventional polymerase chain reaction for detection of *Blastocystis* sp. in clinical stool samples," *American Journal of Tropical Medicine and Hygiene*, vol. 84, no. 2, pp. 308–312, 2011.

- [32] A. Tellez Sierra, L. Cortez Ruiz, A. Aust Kettis, G. Huldts, J. Jonsson, and H. Schroder, "Amebiasis in Nicaragua: class specific serum antibody responses," *Archives of Medical Research*, vol. 23, no. 2, pp. 261–264, 1992.
- [33] M. M. Hossain, I. Ljungstrom, R. I. Glass, L. Lundin, B. J. Stoll, and G. Huldts, "Amoebiasis and giardiasis in Bangladesh: parasitological and serological studies," *Transactions of the Royal Society of Tropical Medicine and Hygiene*, vol. 77, no. 4, pp. 552–554, 1983.
- [34] L. L. Braga, A. A. M. Lima, C. L. Sears et al., "Seroepidemiology of *Entamoeba histolytica* in a slum in northeastern Brazil," *The American Journal of Tropical Medicine and Hygiene*, vol. 55, no. 6, pp. 693–697, 1996.
- [35] P. W. Doyle, M. M. Helgason, R. G. Mathias, and E. M. Proctor, "Epidemiology and pathogenicity of *Blastocystis hominis*," *Journal of Clinical Microbiology*, vol. 28, no. 1, pp. 116–121, 1990.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

