

Research Article

Noncoercive Perturbed Densely Defined Operators and Application to Parabolic Problems

Teffera M. Asfaw

Department of Mathematics, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

Correspondence should be addressed to Teffera M. Asfaw; tefferam@yahoo.com

Received 30 June 2015; Accepted 9 August 2015

Academic Editor: Naseer Shahzad

Copyright © 2015 Teffera M. Asfaw. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let X be a real locally uniformly convex reflexive separable Banach space with locally uniformly convex dual space X^* . Let $T : X \supseteq D(T) \rightarrow 2^{X^*}$ be maximal monotone and $S : X \supseteq D(S) \rightarrow X^*$ quasibounded generalized pseudomonotone such that there exists a real reflexive separable Banach space $W \subset D(S)$, dense and continuously embedded in X . Assume, further, that there exists $d \geq 0$ such that $\langle v^* + Sx, x \rangle \geq -d\|x\|^2$ for all $x \in D(T) \cap D(S)$ and $v^* \in Tx$. New surjectivity results are given for noncoercive, not everywhere defined, and possibly unbounded operators of the type $T + S$. A partial positive answer for Nirenberg's problem on surjectivity of expansive mapping is provided. Leray-Schauder degree is applied employing the method of elliptic superregularization. A new characterization of linear maximal monotone operator $L : X \supseteq D(L) \rightarrow X^*$ is given as a result of surjectivity of $L + S$, where S is of type (M) with respect to L . These results improve the corresponding theory for noncoercive and not everywhere defined operators of pseudomonotone type. In the last section, an example is provided addressing existence of weak solution in $X = L^p(0, T; W_0^{1,p}(\Omega))$ of a nonlinear parabolic problem of the type $u_t - \sum_{i=1}^n (\partial/\partial x_i) a_i(x, t, u, \nabla u) = f(x, t)$, $(x, t) \in Q$; $u(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$; $u(x, 0) = 0$, $x \in \Omega$, where $p > 1$, Ω is a nonempty, bounded, and open subset of \mathbb{R}^N , $a_i : \Omega \times (0, T) \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ ($i = 1, 2, \dots, n$) satisfies certain growth conditions, and $f \in L^{p'}(Q)$, $Q = \Omega \times (0, T)$, and p' is the conjugate exponent of p .

1. Introduction—Preliminaries

In what follows, X is a real reflexive separable locally uniformly convex Banach space with locally uniformly convex dual space X^* . The norm of the space X , and any other normed spaces herein, will be denoted by $\|\cdot\|$. For $x \in X$ and $x^* \in X^*$, the pairing $\langle x^*, x \rangle$ denotes the value $x^*(x)$. Let X and Y be real Banach spaces. For a multivalued mapping $T : X \rightarrow 2^Y$, we define the domain $D(T)$ of T by $D(T) = \{x \in X : Tx \neq \emptyset\}$ and the range $R(T)$ of T by $R(T) = \bigcup_{x \in D(T)} Tx$. We also denote the graph of T by $G(T) = \{(x, Tx) : x \in D(T)\}$. A mapping $T : X \supseteq D(T) \rightarrow Y$ is “demicontinuous” if it is continuous from the strong topology of $D(T)$ to the weak topology of Y . A multivalued mapping $T : X \supseteq D(T) \rightarrow 2^Y$ is “bounded” if it maps bounded subsets of $D(T)$ to bounded subsets of Y . It is “compact” if it is strongly continuous and maps bounded subsets of $D(T)$ to relatively compact subset of Y . It is “finitely continuous” if it is upper semicontinuous

from each finite dimensional subspace F of X to the weak topology of Y . It is “quasibounded” if for every $M > 0$ there exists $K(M) > 0$ such that $[x, w^*] \in G(T)$ with $\|x\| \leq M$ and $\langle w^*, x \rangle \leq M\|x\|$ imply $\|w^*\| \leq K(M)$. It is “strongly quasibounded” if for every $M > 0$ there exists $K(M) > 0$ such that $[x, w^*] \in G(T)$ with $\|x\| \leq M$ and $\langle w^*, x \rangle \leq M$ imply $\|w^*\| \leq K(M)$. In what follows, a mapping will be called “continuous” if it is strongly continuous.

Let $\psi : [0, \infty) \rightarrow [0, \infty)$ be continuous strictly increasing function such that $\psi(0) = 0$ and $\psi(t) \rightarrow \infty$ as $t \rightarrow \infty$. The duality mapping corresponding to ψ denoted by $J_\psi : X \rightarrow 2^{X^*}$ is defined by

$$J_\psi(x) = \{x^* \in X^* : \langle x^*, x \rangle = \|x^*\| \|x\|, \|x^*\| = \psi(\|x\|)\}. \quad (1)$$

It is well-known that, for each $x \in X$, the Hahn-Banach Theorem implies $J_\psi(x) \neq \emptyset$. Since X and X^* are locally

uniformly convex, J_ψ is single-valued, bounded monotone of type (S_+) and bicontinuous. If $\psi(t) = t$ for $t \geq 0$, then J_ψ is denoted by J and is called the normalized duality mapping.

An operator $T : X \supset D(T) \rightarrow 2^{X^*}$ is said to be “monotone” if, for every $x \in D(T)$, $y \in D(T)$, and every $u^* \in Tx$, $v^* \in Ty$, we have $\langle u^* - v^*, x - y \rangle \geq 0$. A monotone mapping $T : X \supset D(T) \rightarrow 2^{X^*}$ is “maximal monotone” if $R(T + \lambda J) = X^*$ for every $\lambda > 0$; that is, T is maximal monotone if and only if T is monotone and $\langle u^* - u_0^*, x - x_0 \rangle \geq 0$ for every $(x, u^*) \in G(T)$ implies $x_0 \in D(T)$ and $u_0^* \in Tx_0$. If T is maximal monotone, the operator $T_t : X \rightarrow X^*$, $t \in (0, \infty)$, defined by $T_t x = (T^{-1} + tJ^{-1})^{-1}x$, is bounded, continuous, maximal monotone and such that $T_t x \rightarrow T^{(0)}x$ as $t \rightarrow 0^+$, for every $x \in D(T)$, where $\|T^{(0)}x\| = \inf\{\|y^*\| : y^* \in Tx\}$. The “resolvent” $J_t : X \rightarrow D(T)$, defined by $J_t x = x - tJ^{-1}(T_t x)$, is continuous and $T_t x \in T(J_t x)$ for every $x \in X$. Moreover, $\lim_{t \rightarrow 0} J_t x = x$ for all $x \in \text{co}D(T)$, where $\text{co}D(T)$ is the convex hull of the set $D(T)$. An operator $A : X \supseteq D(A) \rightarrow 2^{X^*}$ is called “coercive” if either $D(A)$ is bounded or there exists a function $\psi : [0, \infty) \rightarrow (-\infty, \infty)$ such that $\psi(t) \rightarrow \infty$ as $t \rightarrow \infty$ and $\langle y^*, x \rangle \geq \psi(\|x\|)\|x\|$ for all $x \in D(A)$ and $y^* \in Ax$. For an operator $A : X \supseteq D(A) \rightarrow 2^{X^*}$ and $x \in D(A)$, we denote $|Ax| = \inf\{\|v^*\| : v^* \in Ax\}$. It is called weakly coercive if either $D(A)$ is bounded or $|Ax| \rightarrow \infty$ as $\|x\| \rightarrow \infty$.

The following definitions are used throughout the paper. In arbitrary Banach space X , Browder and Hess [1] introduced the definitions of pseudomonotone and generalized pseudomonotone operators. The original definition for single-valued pseudomonotone, generalized pseudomonotone, and operators of type (M) with domain all of X , is due to Brézis [2].

Definition 1. An operator $S : X \supseteq D(S) \rightarrow X^*$ is called

- (i) “generalized pseudomonotone” if, for each sequence $\{x_n\}$ in $D(S)$ with $x_n \rightarrow x_0$ and $Sx_n \rightarrow v_0^*$ as $n \rightarrow \infty$ such that $\limsup_{n \rightarrow \infty} \langle Sx_n, x_n - x_0 \rangle \leq 0$, then $x_0 \in D(S)$, $Sx_0 = v_0^*$, and $\langle Sx_n, x_n \rangle \rightarrow \langle Sx_0, x_0 \rangle$ as $n \rightarrow \infty$.
- (ii) “type (M) ” if, for each sequence $\{x_n\}$ in $D(S)$ with $x_n \rightarrow x_0$ in X and $Sx_n \rightarrow v_0^*$ as $n \rightarrow \infty$ such that $\limsup_{n \rightarrow \infty} \langle Sx_n, x_n - x_0 \rangle \leq 0$, then $x_0 \in D(S)$ and $Sx_0 = v_0^*$.
- (iii) “ α -expansive” if there exists $\alpha > 0$ such that $\|v^* - u^*\| \geq \alpha\|x - y\|$ for all $x \in D(S)$, $y \in D(S)$, $v^* \in Sx$, and $u^* \in Sy$. It is called expansive if $\alpha = 1$.

We notice here that the definition of single-valued expansive mapping is due to Nirenberg [3]. In order to enlarge the class of single-valued operators, the multivalued version is introduced in (iii) of Definition 1. It is not hard to notice that every uniformly monotone operator is expansive. Furthermore, in a Hilbert space $X = H$, if $T : H \supseteq D(T) \rightarrow 2^H$ is monotone, we see that, for each $\lambda > 0$, $T + \lambda I$ is multivalued expansive with domain $D(T)$.

The following definition gives a larger class of operators of monotone type, which can be found in Kartsatos and Skrypnik [4].

Definition 2. Let $T : X \supseteq D(T) \rightarrow 2^{X^*}$ be maximal monotone and $A : X \supseteq D(A) \rightarrow X^*$. Let $L \subseteq D(M) \cap D(A)$ be a linear subspace of X . Then A is said to be

- (i) “quasibounded with respect to T ” if, for each $M > 0$, there exists $K(M) > 0$ such that

$$\begin{aligned} \langle Au + u^*, u \rangle &\leq M, \\ \|u\| &\leq M, \end{aligned} \quad (2)$$

where $u \in L$ and $u^* \in Tu$, then $\|Au\| \leq K(M)$,

- (ii) “generalized (S_+) with respect to T ” if, for each $\{u_n\}$ in L with $u_n^* \in Tu_n$, $u_n \rightarrow u_0$ in X and $Au_n \rightarrow h_0^*$ in X^* as $n \rightarrow \infty$ such that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle Au_n, u_n - u_0 \rangle &\leq 0, \\ \langle u_n^* + Au_n, u_n \rangle &\leq 0 \end{aligned} \quad (3)$$

for all n , then $u_n \rightarrow u_0 \in D(A)$ and $Au_0 = h_0^*$,

- (iii) “generalized pseudomonotone with respect to T ” if, for each $\{u_n\}$ in L with $u_n^* \in Tu_n$, $u_n \rightarrow u_0$ in X and $Au_n \rightarrow h_0^*$ in X^* as $n \rightarrow \infty$ such that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle Au_n, u_n - u_0 \rangle &\leq 0, \\ \langle u_n^* + Au_n, u_n \rangle &\leq 0 \end{aligned} \quad (4)$$

for all n , then $u_0 \in D(A)$, $Au_0 = h_0^*$, and $\langle Su_n, u_n \rangle \rightarrow \langle Su_0, u_0 \rangle$ as $n \rightarrow \infty$,

- (iv) “of type (M) with respect to T ” if, for each $\{u_n\}$ in L with $u_n^* \in Tu_n$, $u_n \rightarrow u_0$ in X and $Au_n \rightarrow h_0^*$ in X^* as $n \rightarrow \infty$ such that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle Au_n, u_n - u_0 \rangle &\leq 0, \\ \langle u_n^* + Au_n, u_n \rangle &\leq 0 \end{aligned} \quad (5)$$

for all n , then $u_0 \in D(A)$ and $Au_0 = h_0^*$.

By Definition 2, it is not difficult to see that $0 \in D(T)$ and A is quasibounded implying that A is quasibounded with respect to T . Furthermore, it follows that the class of generalized (S_+) operators with respect to T includes the class of operators of type (S_+) .

For basic definitions and further properties of mappings of monotone type, the reader is referred to Barbu [5], Brézis et al. [6], Brézis [2], Browder and Hess [1], Pascali and Sburlan [7], Browder [8], and Zeidler [9]. For results concerning perturbations of maximal monotone operators by bounded and everywhere defined pseudomonotone type operators, the reader is referred to Browder and Hess [1], Brézis [2], Browder [10], Brézis and Nirenberg [11], Kenmochi [12–14], Guan et al. [15], Le [16], Guan and Kartsatos [15, 17],

and Kartsatos and Skrypnik [4] and the references therein. For recent degree theory and applications for solvability of operator inclusions involving bounded pseudomonotone perturbations of maximal monotone operators under general coercivity and Leray-Schauder type boundary conditions, we cite the paper due to Asfaw and Kartsatos [18]. Existence results concerning noncoercive operators of the type $T + S$, where $T : X \supseteq D(T) \rightarrow 2^{X^*}$ is maximal monotone and $S : X \rightarrow 2^{X^*}$ is bounded pseudomonotone, can be found in the paper due to Asfaw [19]. For applications of the theory of perturbed monotone type operators to variational and hemivariational inequality problems, the reader is referred to the papers due to Carl and Le [20], Carl et al. [21], Carl [22], and Carl and Motreanu [23] and the references therein. For a separable reflexive Banach space X and a nonempty, closed, and convex subset K of X , Asfaw and Kartsatos [24] gave existence results for locally defined operators of the type $T + S$, where $T : X \supseteq D(T) \rightarrow 2^{X^*}$ is maximal monotone and $S : K \rightarrow X^*$ is demicontinuous and generalized pseudomonotone under coercivity condition on S .

The main contribution of the paper is to obtain surjectivity results for *noncoercive* and *not everywhere defined* operators of the type

- (i) $T + S$, where $S : X \supseteq D(S) \rightarrow X^*$ is quasibounded demicontinuous generalized pseudomonotone such that
 - (a) there exists a real reflexive separable Banach space $W \subseteq D(S)$, dense and continuously embedded in X ;
 - (b) there exists $d \geq 0$ such that $\langle v^* + Sx, x \rangle \geq -d\|x\|^2$ for all $x \in D(T) \cap D(S)$ and $v^* \in Tx$;
 - (c) there exist $\alpha > d$ and $\mu \geq 0$ such that $\|v^* + Sx\| \geq \alpha\|x\| - \mu$ for all $x \in D(T) \cap D(S)$ and $v^* \in Tx$,
- (ii) $L + S$, where $S : X \supseteq D(S) \rightarrow X^*$ is quasibounded demicontinuous of type (M) with $D(L) \subseteq D(S)$ such that (b) and (c) of (i) are satisfied.

In Section 2, we proved surjectivity results for $T + S$ and $L + S$ satisfying conditions (i) and (ii), respectively. In Theorem 6, we provide a surjectivity result for operators of the type $T + S$, where T and S satisfy condition (i). Theorem 6 is new and improves the existing surjectivity results for an operator S , which is single-valued, everywhere defined, bounded, and coercive pseudomonotone. In particular, for a single-valued pseudomonotone operator S , Theorem 6 improves the surjectivity results due to Browder and Hess [1], Kenmochi [12–14], Le [16], Guan and Kartsatos [17], Asfaw and Kartsatos [18], and Asfaw [19, 25] because the results in these references require S to be everywhere defined, bounded, and coercive while Theorem 6 used S to be densely defined, quasibounded, and noncoercive. Moreover, Browder (cf. Zeidler [9, Theorem 32. A, pages 866–872]) gave the main theorem for perturbations of maximal monotone operator by a single-valued, bounded, demicontinuous, and coercive operator S with $D(S) = C$, a nonempty, closed, and convex subset of X . In view of this, Theorem 6 gives an analogous

result, where $D(S)$ is dense in X , possibly, neither closed nor convex, and S is weakly coercive. It is also known, due to Browder and Hess [1], that every pseudomonotone operator S from X into X^* with $D(S) = X$ is generalized pseudomonotone. It is also true that S is demicontinuous provided that it is bounded, single-valued, and everywhere defined. Consequently, the arguments used in the proof of Theorem 6 give analogous conclusion if $S : X = D(S) \rightarrow X^*$ is bounded pseudomonotone and T and S satisfy the given hypotheses. As a consequence of Corollary 7, a partial positive answer for Nirenberg’s problem on surjectivity of densely defined demicontinuous generalized pseudomonotone expansive mapping is provided. In addition, Theorem 8 provides surjectivity result for operators of the type $T + S$, where T and S satisfy condition (ii). As a result of Theorem 8, a new characterization of linear maximal monotone operator is proved when the space X is separable. It is well known due to Brézis (cf. Zeidler [9, Theorem 32. L, pages 897–899]) that a linear monotone operator L is maximal monotone if and only if L is closed and densely defined and the adjoint operator L^* is monotone. An interesting result in the present paper is that a linear monotone operator L is maximal monotone if and only if L is closed and densely defined, provided that X is separable. This result weakens the monotonicity condition on L^* used by Brézis (cf. Zeidler [9, Theorem 32. L, pages 897–899]). To the best of the author’s knowledge, Theorem 8 is a new result and Corollary 9 improves the well-known result of Brézis. In Section 3, we demonstrate the applicability of the results by proving existence of weak solution in $L^p(0, T; W_0^{1,p}(\Omega))$ of a nonlinear parabolic problem, where $p > 1$ and Ω is a nonempty, bounded, and open subset of \mathbb{R}^N .

The following important lemma is due to Brézis et al. [6].

Lemma 3. *Let B be a maximal monotone set in $X \times X^*$. If $(u_n, u_n^*) \in B$ such that $u_n \rightarrow u$, $u_n^* \rightarrow u^*$ as $n \rightarrow \infty$, and either*

$$\limsup_{n,m \rightarrow \infty} \langle u_n^* - u_m^*, u_n - u_m \rangle \leq 0 \tag{6}$$

or

$$\limsup_{n \rightarrow \infty} \langle u_n^* - u^*, u_n - u \rangle \leq 0, \tag{7}$$

then $(u, u^*) \in B$ and $\langle u_n^*, u_n \rangle \rightarrow \langle u^*, u \rangle$ as $n \rightarrow \infty$.

Browder and Ton [26] gave the following important embedding result.

Lemma 4. *Let X be a separable reflexive Banach space. Then there exists a real separable Hilbert space H and a compact injection $Q : H \rightarrow X$ such that $\overline{Q(H)} = X$.*

In this paper, we use the following fixed point result for compact operators, originally due to Leray and Schauder, which may be found in the book of Granas and Dugundji [27, Theorem 5.2, page 123].

Lemma 5. *Let C be a convex subset of a normed linear space X and let U be nonempty relatively open in C with $0 \in U$. Then each compact map $F : \overline{U} \rightarrow C$ satisfies that either*

(i) F has a fixed point in U

or

(ii) there exist $x \in \partial_C U$ and $\lambda \in (0, 1)$ such that $x = \lambda F(x)$, where $\partial_C U$ is the boundary of U with respect to the subspace topology on C .

2. Main Results

In this section, we prove the following new surjectivity result for maximal monotone perturbation of densely defined non-coercive generalized pseudomonotone operator in separable reflexive Banach spaces.

Theorem 6. Let $T : X \supseteq D(T) \rightarrow 2^{X^*}$ be maximal monotone with $0 \in T(0)$ and $S : X \supseteq D(S) \rightarrow X^*$ quasibounded demicontinuous generalized pseudomonotone. Suppose $W \subseteq D(S)$ is a real reflexive separable Banach space dense and continuously embedded in X . Assume, further, that there exist $\mu \geq 0$, $d \geq 0$, and $\alpha > d$ satisfying

$$\langle Sx, x \rangle \geq -d \|x\|^2 \quad (8)$$

for all $x \in D(S)$ and either

(i)

$$\|v^* + Sx\| \geq \alpha \|x\| - \mu \quad \forall x \in D(T) \cap D(S), v^* \in Tx \quad (9)$$

or

(ii) there exists $\phi : [0, \infty) \rightarrow (-\infty, \infty)$ such that $\phi(t) \rightarrow \infty$ as $t \rightarrow \infty$ and

$$\|v^* + Sx\| \geq \phi(\|x\|) \|x\| \quad \forall x \in D(T) \cap D(S), v^* \in Tx. \quad (10)$$

Then $T + S$ is surjective.

Proof. Let $\lambda > 0$ be fixed temporarily and T_λ the Yosida approximant of T . For each $\varepsilon > 0$, by using the inner product condition on S and monotonicity of T_λ ($T_\lambda(0) = 0$ for all $\lambda > 0$), we see that

$$\begin{aligned} & \langle T_\lambda x + Sx + \delta \|x\| Jx - f^*, x \rangle \\ & \geq \delta \|x\|^3 - d \|x\|^2 - \|f^*\| \|x\| \\ & = \|x\|^3 \left[\delta - \frac{d}{\|x\|} - \frac{\|f^*\|}{\|x\|^3} \right] > 0 \end{aligned} \quad (11)$$

for all $x \in D(S) \cap \partial B_{R_\delta}(0)$ for some $R_\delta > 0$. Let $G_\delta = B_{R_\delta}(0)$. Let H be a real separable Hilbert space and $Q : H \rightarrow W$ a compact injection such that $Q(H)$ is dense in W guaranteed by Lemma 4. Let $j : W \rightarrow X$ be the natural injection and let $Q^* : W^* \rightarrow H^*$ and $j^* : X^* \rightarrow W^*$ be adjoint of Q and j , respectively. It follows that $\psi = jQ : H \rightarrow X$ is a compact operator. Let $U = Q^{-1}(G_\delta \cap W)$. First we show that $G_\delta \cap W$ is open in W ; that is, $W \setminus (G_\delta \cap W) = W \cap (X \setminus G_\delta)$ is closed in

W . To this end, let $\{x_n\}$ be a sequence in $W \cap (X \setminus G_\delta)$ such that $x_n \rightarrow x_0$ in W as $n \rightarrow \infty$. Since W is continuously embedded in X , we get $x_n \rightarrow x_0$ in X as $n \rightarrow \infty$. Since $X \setminus G_\delta$ is closed in X , it follows that $x_0 \in X \setminus G_\delta$; that is, $x_0 \in W \cap (X \setminus G_\delta)$. This shows that $W \cap (X \setminus G_\delta)$ is closed in W ; that is, $G_\delta \cap W$ is open in W . The continuity of Q implies that U is open in H . Since W is continuously embedded in X , it follows that

$$\overline{G_\delta \cap W}^W \subseteq \overline{G_\delta \cap W}^X \subseteq \overline{G_\delta}, \quad (12)$$

where the closures are taken with respect to the spaces W and X , respectively. Since $\overline{G_\delta \cap W}^W \subseteq W$, we obtain that

$$\begin{aligned} (G_\delta \cap W) \cup \partial_W(G_\delta \cap W) &= \overline{G_\delta \cap W}^W \subseteq \overline{G_\delta} \cap W \\ &= (G_\delta \cap W) \cup (\partial G_\delta \cap W). \end{aligned} \quad (13)$$

Since the sets $G_\delta \cap W$ and $\partial_W(G_\delta \cap W)$ are disjoint, we conclude that

$$\partial_W(G_\delta \cap W) \subseteq \partial G_\delta \cap W. \quad (14)$$

For each $\lambda > 0$, let T_λ be the Yosida approximant of T . Let $J_1 x = \|x\| Jx$, $x \in X$. It is known that, for each $\delta > 0$, $T_\lambda + \delta J_1$ is bounded, continuous, monotone, and of type (S_+) . Let $\psi = jQ$ and $C_\varepsilon^{\lambda, \delta} : \bar{U} \rightarrow H$ be given by

$$C_\varepsilon^{\lambda, \delta}(v) = -\varepsilon^{-1} (\psi^* ((T_\lambda + S + \delta J_1) \psi(v) - f^*)), \quad v \in \bar{U}. \quad (15)$$

Since ψ is continuous, it follows that

$$\bar{U} = \overline{\psi^{-1}(G_\delta \cap W)} \subseteq \psi^{-1}(\overline{G_\delta \cap W}^W) \subseteq H \quad (16)$$

is closed subset of H . We show that $C_\varepsilon^{\lambda, \delta}$ is a compact operator. To this end, let $x_n \in \bar{U}$ such that $x_n \rightarrow x_0$ as $n \rightarrow \infty$. Since Q is continuous from H into X , we have $Qx_n \rightarrow Qx_0$ as $n \rightarrow \infty$. Since $x_n \in \bar{U}$, the sequence $\{Qx_n\}$ lies in W . Since $x_n \in H$ for all n and $x_0 \in \bar{U}$, it follows that $Qx_n \in W$ and $Qx_0 \in W$ for all n . Since S and T_λ are demicontinuous, it follows that $(T_\lambda + S)Qx_n \rightarrow (T_\lambda + S)Qx_0$ as $n \rightarrow \infty$. By the density of W in X , it is known that j^* is defined from W^* into H . As a result, for each $w \in W$, we see that

$$\begin{aligned} & \langle j^* (T_\lambda + S + \delta J_1) Qx_n - j^* (T_\lambda + S + \delta J_1) Qx_0, w \rangle \\ & = \langle (T_\lambda + S + \delta J_1) Qx_n, w \rangle \\ & \quad - \langle (T_\lambda + S + \delta J_1) Qx_0, w \rangle \end{aligned} \quad (17)$$

for all n . However, the right side expression goes to 0 as $n \rightarrow \infty$; that is, for each $w \in W$, it follows that

$$\begin{aligned} & \langle j^* (T_\lambda + S + \delta J_1) Qx_n - j^* (T_\lambda + S + \delta J_1) Qx_0, w \rangle \\ & \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (18)$$

On the other hand, by the density of W in X , for each $x \in X$, we get

$$\begin{aligned} & \langle j^*(T_\lambda + S + \delta J_1)Qx_n - j^*(T_\lambda + S + \delta J_1)Qx_0, x \rangle \\ & \rightarrow 0 \quad \text{as } n \rightarrow \infty; \end{aligned} \quad (19)$$

that is, $j^*(T_\lambda + S + \delta J_1)Qx_n \rightarrow j^*(T_\lambda + S + \delta J_1)Qx_0$ as $n \rightarrow \infty$. Since Q^* is compact linear, which is completely continuous and $(jQ)^* = Q^*j^*$, we arrive at $\psi^*(T_\lambda + S + \delta J_1)Qx_n \rightarrow \psi^*(T_\lambda + S + \delta J_1)Qx_0$ as $n \rightarrow \infty$. This shows that the mapping $C_\varepsilon^{\delta, \lambda}$ is continuous. Following similar argument as above, it is not difficult to show that $C_\varepsilon^{\delta, \lambda}$ maps any bounded subset of \bar{U} into relatively compact subset of H . As a result, we conclude that $C_\varepsilon^{\delta, \lambda}$ is a compact operator. Fix $\varepsilon > 0$. In order to use Lemma 5, it is enough to show that (i) of Lemma 5 does not hold; that is, for all $\mu \in (0, 1)$ and $x \in \partial_H U$, we have $x \neq \mu C_\varepsilon^{\delta, \lambda}(x)$. Suppose this is false; that is, there exist $x_0 \in \partial_H U$ and $\mu_0 \in (0, 1)$ such that $x_0 = \mu_0 C_\varepsilon^{\delta, \lambda}(x_0)$. This yields

$$\varepsilon x_0 + \mu_0 \psi^*((T_\lambda + S + \delta J_1)\psi x_0 - f^*) = 0. \quad (20)$$

We notice here that the continuity of Q , property of Q^{-1} , and definition of boundary of an open set imply that

$$\begin{aligned} \partial_H U &= \partial_H Q^{-1}(G_\delta \cap W) \subseteq Q^{-1}(\partial_W(G_\delta \cap W)) \\ &\subseteq Q^{-1}(\partial G_\delta \cap W) \end{aligned} \quad (21)$$

holds. Since $x_0 \in \partial_H U$, it follows that $Qx_0 \in \partial G \cap W$. By (11) and (20), we get

$$\begin{aligned} \frac{\varepsilon}{\mu_0} \|x_0\|^2 &= -\langle \psi^*((T_\lambda + S + \delta J_1)Qx_0 - f^*), x_0 \rangle \\ &= -\langle (T_\lambda + S + \delta J_1)Qx_0 - f^*, Qx_0 \rangle \leq 0, \end{aligned} \quad (22)$$

which implies $x_0 = 0$. But this is impossible because $0 \in G_\delta \cap W$. Therefore, by applying Lemma 5, for each $\varepsilon > 0$, $\lambda > 0$, and $\delta > 0$, we conclude that the compact operator $C_\varepsilon^{\delta, \lambda}$ has a fixed point $x_\varepsilon \in U$; that is,

$$\varepsilon x_\varepsilon + \psi^*((T_\lambda + S + \delta J_1)\psi x_\varepsilon - f^*) = 0. \quad (23)$$

Therefore, for each $\varepsilon_n \downarrow 0^+$, there exists $x_n \in \bar{U}$ such that

$$\varepsilon_n x_n + \psi^*((T_\lambda + S + \delta J_1)\psi x_n - f^*) = 0 \quad (24)$$

for all n . Since G_δ is bounded, the sequence $\{\psi x_n\}$ is bounded. Since T_λ and J_1 are bounded, it follows that the sequence $\{(T_\lambda + \delta J_1)\psi x_n\}$ is bounded. Since $W = \overline{Q(H)}^W$ and W is continuously embedded, we see that $W = \overline{Q(H)}^W \subseteq \overline{Q(H)}^X$, where the closures are with respect to the norms in W and X , respectively. As a result, the density of W in X implies that

$X = \overline{Q(H)}^X$. By using (11), the monotonicity of T_λ and J_1 , and property of ψ^* , we obtain that

$$\begin{aligned} & \langle S\psi x_n, \psi x_n \rangle \\ &= -\varepsilon_n \|x_n\|^2 \\ &\quad - \langle T_\lambda \psi x_n + \delta J_1 \psi x_n - (T_\lambda(0) + \delta J_1(0)), \psi x_n \rangle \\ &\quad + \langle T_\lambda(0) + \delta J_1(0) + f^*, \psi x_n \rangle \\ &\leq \|T_\lambda(0) + \delta J_1(0) + f^*\| \|\psi x_n\| \\ &\leq (|T(0)| + \|f^*\|) \|\psi x_n\| \end{aligned} \quad (25)$$

for all n . Since $Qx_n \in W$, it follows that $\psi x_n = jQx_n = Qx_n$ for all n . Consequently, we obtain that

$$\langle SQx_n, Qx_n \rangle \leq (|T(0)| + \|f^*\|) \|Qx_n\| \quad (26)$$

for all n . Since $\{Qx_n\}$ is bounded and S is quasibounded, we conclude that $\{SQx_n\}$ is bounded. Consequently, by using (24), it is not difficult to see that $\{\varepsilon_n \|x_n\|^2\}$ is bounded. If the sequence $\{x_n\}$ is bounded, then $\varepsilon_n x_n \rightarrow 0$ as $n \rightarrow \infty$. Otherwise, by using the boundedness of $\{\varepsilon_n \|x_n\|^2\}$, we assume without loss of generality that $\varepsilon_n x_n \rightarrow 0$ as $n \rightarrow \infty$, $Qx_n \rightarrow x_0$, and $SQx_n \rightarrow v_0^*$ as $n \rightarrow \infty$. Since $\overline{Q(H)}^X = X$, by choosing a sequence $\{z_m = Qy_m\}$ such that $z_m \rightarrow x_0$ as $m \rightarrow \infty$ and using (24) together with the monotonicity of $T_\lambda + \delta J_1$, we get

$$\begin{aligned} \langle SQx_n, Qx_n \rangle &= \langle SQx_n, Qy_m - Qy_m + Qx_n \rangle \\ &= \langle SQx_n, Qy_m \rangle + \langle SQx_n, Qx_n - Qy_m \rangle \\ &= \langle SQx_n, Qy_m \rangle + \langle f^*, Qx_n - Qy_m \rangle \\ &\quad - \langle (T_\lambda + \delta J_1)Qx_n - (T_\lambda + \delta J_1)Qy_m, Qx_n - Qy_m \rangle \\ &\quad - \langle (T_\lambda + \delta J_1)Qy_m, Qx_n - Qy_m \rangle \\ &\quad - \varepsilon_n \langle x_n, x_n - y_m \rangle \leq \langle SQx_n, Qy_m \rangle \\ &\quad - \langle (T_\lambda + \delta J_1)Qy_m, Qx_n - Qy_m \rangle + \varepsilon_n \langle x_n, y_m \rangle \\ &\quad + \langle f^*, Qx_n - Qy_m \rangle \end{aligned} \quad (27)$$

for all n and m . Fixing m and letting $n \rightarrow \infty$ in (27), we obtain that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \langle SQx_n, Qx_n \rangle \\ &\leq \langle v_0^*, Qy_m \rangle \\ &\quad - \langle (T_\lambda + \delta J_1)Qy_m - f^*, x_0 - Qy_m \rangle. \end{aligned} \quad (28)$$

Since $T_\lambda + \delta J_1$ is demicontinuous, letting $m \rightarrow \infty$, we arrive at

$$\limsup_{n \rightarrow \infty} \langle SQx_n, Qx_n \rangle \leq \langle v_0^*, x_0 \rangle; \quad (29)$$

that is,

$$\limsup_{n \rightarrow \infty} \langle SQx_n, Qx_n - x_0 \rangle \leq 0. \quad (30)$$

Since S is generalized pseudomonotone, we conclude that $x_0 \in D(S)$, $Sx_0 = v_0^*$, and $\langle SQx_n, Qx_n \rangle \rightarrow \langle v_0^*, x_0 \rangle$ as $n \rightarrow \infty$. For any $y \in Q(H)$, applying the monotonicity of $T_\lambda + \delta J_1$, we arrive at

$$\begin{aligned} \langle T_\lambda y + \delta J_1 y - f^*, y - x_0 \rangle &= \liminf_{n \rightarrow \infty} \langle T_\lambda y + \delta J_1 y \\ &\quad - f^*, y - Qx_n \rangle \geq \liminf_{n \rightarrow \infty} [\langle T_\lambda y + \delta J_1 y \\ &\quad - (T_\lambda Qx_n + \delta J_1 Qx_n), y - Qx_n \rangle] \\ &\quad + \liminf_{n \rightarrow \infty} [\langle T_\lambda Qx_n + \delta J_1 Qx_n - f^*, y - Qx_n \rangle] \\ &\geq \liminf_{n \rightarrow \infty} \langle T_\lambda Qx_n + \delta J_1 Qx_n - f^*, y - Qx_n \rangle. \end{aligned} \quad (31)$$

Moreover, from (24), we obtain that

$$\begin{aligned} \langle (T_\lambda + \delta J_1) Qx_n - f^*, y - Qx_n \rangle &= -\varepsilon_n \langle x_n, Q^{-1}y - x_n \rangle - \langle SQx_n, y - Qx_n \rangle \\ &= \varepsilon_n \|x_n\|^2 - \varepsilon_n \langle x_n, Q^{-1}y \rangle - \langle SQx_n, y - Qx_n \rangle \\ &\geq -\varepsilon_n \langle x_n, Q^{-1}y \rangle - \langle SQx_n, y - Qx_n \rangle \end{aligned} \quad (32)$$

for all n . As a result, we arrive at

$$\begin{aligned} \liminf_{n \rightarrow \infty} \langle (T_\lambda + \delta J_1) Qx_n - f^*, y - Qx_n \rangle &\geq -\langle Sx_0, y - x_0 \rangle. \end{aligned} \quad (33)$$

From (31) and (33), we obtain

$$\langle (T_\lambda + \delta J_1) y + Sx_0 - f^*, y - x_0 \rangle \geq 0 \quad (34)$$

for all $y \in Q(H)$. By the density of $Q(H)$ in X and the continuity of $T_\lambda + \delta J_1$, we conclude that

$$\langle T_\lambda y + \delta J_1 y + Sx_0 - f^*, y - x_0 \rangle \geq 0 \quad (35)$$

for all $y \in X$. Since, for any $y \in X$, $x_t = tx_0 + (1-t)y \in X$ for all $t \in [0, 1]$, using x_t in place of y , we obtain that

$$\langle (T_\lambda + \delta J_1) x_t + Sx_0 - f^*, (1-t)(y - x_0) \rangle \geq 0 \quad (36)$$

for all $t \in [0, 1]$; that is,

$$\langle (T_\lambda + \delta J_1) x_t + Sx_0 - f^*, y - x_0 \rangle \geq 0 \quad (37)$$

for all $t \in [0, 1]$. Since $T_\lambda + \delta J_1$ is continuous and $x_t \rightarrow x_0$ as $t \rightarrow 1^-$, we have $T_\lambda x_t + \delta J_1 x_t \rightarrow T_\lambda x_0 + \delta J_1 x_0$ as $t \rightarrow 1^-$. Letting $t \rightarrow 1^-$, we arrive at

$$\langle (T_\lambda + \delta J_1) x_0 + Sx_0 - f^*, y - x_0 \rangle \geq 0 \quad (38)$$

for all $y \in X$. Since $y \in X$ is arbitrary, setting $y + x_0$ in place of y yields

$$\langle (T_\lambda + \delta J_1) x_0 + Sx_0 - f^*, y \rangle \geq 0 \quad (39)$$

for all $y \in X$. Therefore, for each $\lambda > 0$ (by fixing $\delta > 0$ temporarily), we see that there exists $x_\lambda \in D(S) \cap \overline{G_\delta}$ such

that $T_\lambda x_\lambda + \delta J_1 x_\lambda + Sx_\lambda = f^*$. Thus, for each $\lambda_n \downarrow 0^+$, there exists $y_n \in D(S) \cap \overline{G_\delta}$ such that

$$T_{\lambda_n} y_n + \delta J_1 y_n + Sy_n - f^* = 0 \quad (40)$$

for all n . Since $\overline{G_\delta}$ and J_1 are bounded, it follows that $\{y_n\}$ and $\{J_1 y_n\}$ are bounded. Since S is quasibounded, it is not hard to see that $\{Sy_n\}$ is bounded, which implies the boundedness of $\{T_{\lambda_n} y_n\}$. Assume without loss of generality that $y_n \rightarrow y_0$, $Sy_n \rightarrow v_0^*$ and $T_{\lambda_n} y_n \rightarrow u_0^*$ as $n \rightarrow \infty$. Since $S + \delta J_1$ is generalized pseudomonotone with domain $D(S)$, it follows that

$$\liminf_{n \rightarrow \infty} \langle Sy_n + \delta J_1 y_n, y_n - y_0 \rangle \geq 0. \quad (41)$$

Consequently, from (40), we arrive at

$$\limsup_{n \rightarrow \infty} \langle T_{\lambda_n} y_n, y_n - y_0 \rangle \leq 0. \quad (42)$$

Let J_{λ_n} be the Yosida resolvent of T . It is well known that $J_{\lambda_n} y_n \in D(T)$, $J_{\lambda_n} y_n = x_n - \lambda_n J^{-1}(T_{\lambda_n} y_n)$, and $T_{\lambda_n} y_n \in T(J_{\lambda_n} y_n)$ for all n . Since $y_n \rightarrow y_0$ and $\{T_{\lambda_n} y_n\}$ is bounded, it follows that $J_{\lambda_n} y_n \rightarrow y_0$ as $n \rightarrow \infty$. Thus, we have

$$\begin{aligned} \langle T_{\lambda_n} y_n, J_{\lambda_n} y_n - y_0 \rangle &= \langle T_{\lambda_n} y_n, J_{\lambda_n} y_n - y_n \rangle \\ &\quad + \langle T_{\lambda_n} y_n, y_n - y_0 \rangle \\ &= -\langle T_{\lambda_n} y_n, \lambda_n J^{-1}(T_{\lambda_n} y_n) \rangle \\ &\quad + \langle T_{\lambda_n} y_n, y_n - y_0 \rangle \\ &= -t_n \|T_{\lambda_n} y_n\|^2 \\ &\quad + \langle T_{\lambda_n} y_n, y_n - y_0 \rangle \\ &\leq \langle T_{\lambda_n} y_n, y_n - y_0 \rangle \end{aligned} \quad (43)$$

for all n . Consequently, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \langle T_{\lambda_n} y_n, J_{\lambda_n} y_n - y_0 \rangle &\leq \lim_{n \rightarrow \infty} \langle T_{\lambda_n} y_n, y_n - y_0 \rangle \\ &\leq 0. \end{aligned} \quad (44)$$

By the maximality of T , applying Lemma 3, we obtain $x_0 \in D(T)$, $v_0^* \in Tx_0$, and $\langle T_{\lambda_n} y_n, J_{\lambda_n} y_n \rangle \rightarrow \langle v_0^*, x_0 \rangle$ as $n \rightarrow \infty$, which implies

$$\limsup_{n \rightarrow \infty} \langle Sy_n, y_n - y_0 \rangle \leq 0. \quad (45)$$

The generalized pseudomonotonicity of S implies $y_0 \in D(S)$ and $Sy_0 = h_0^*$. As a result, letting $n \rightarrow \infty$ in (40), we conclude that $v_0^* + Sy_0 + \delta J_1 y_0 = f^*$. This implies that, for each $\delta_n \downarrow 0^+$, there exist $z_n \in D(T) \cap D(S)$ and $v_n^* \in Tz_n$ such that

$$v_n^* + Sz_n + \delta_n J_1 z_n = f^* \quad (46)$$

for all n . Next we will show that $\{z_n\}$ is bounded. Assume without loss of generality that $\|z_n\| \rightarrow \infty$ as $n \rightarrow \infty$. By

the inner product condition on S and monotonicity of T with $0 \in T(0)$, we get

$$\delta_n \|z_n\|^3 \leq d \|z_n\|^2 + \|f^*\| \|z_n\| \tag{47}$$

for all n ; that is, dividing this inequality by $\|z_n\|$ for all large n , we get $\delta_n \|z_n\|^2 \leq d \|z_n\| + \|f^*\|$ for all large n . By using condition (i) and (46), we get that

$$\begin{aligned} -\mu + \alpha \|z_n\| &\leq \|v_n^* + Sz_n\| \leq \delta_n \|z_n\|^2 + \|f^*\| \\ &\leq d \|z_n\| + 2 \|f^*\| \end{aligned} \tag{48}$$

for all n . This gives $(\alpha - d)\|z_n\| \leq 2\|f^*\| + \mu$ for all n . Consequently, the boundedness of $\{z_n\}$ follows. Since S is quasibounded and $0 \in D(T)$, it is not hard to see that $\{Sz_n\}$ is bounded. Consequently, the boundedness of $\{v_n^*\}$ follows. Assuming that $z_n^* \rightarrow z_0^*$, $v_n^* \rightarrow v_0^*$, and $Sz_n \rightarrow h_0^*$ as $n \rightarrow \infty$ and using the arguments used in the first half of the proof of Theorem 6, we conclude that $z_0 \in D(T) \cap D(S)$, $v_0^* \in Tz_0$, $Sz_0 = h_0^*$, and $v_0^* + Sz_0 = f^*$; that is, for each $f^* \in X^*$, the inclusion problem $Tu + Su \ni f^*$ is solvable in $D(T) \cap D(S)$. Since $f^* \in X^*$ is arbitrary, we obtain the surjectivity of $T + S$. The proof using condition (ii) can be completed following similar arguments. The details are omitted here. This completes the proof. \square

It is worth mentioning that Theorem 6 is a new result because the perturbed operator $T + S$ is noncoercive and S is densely defined such that $D(S)$ contains a dense real separable reflexive Banach space. Under the conditions on $T + S$, the result was unknown earlier even for coercive operator $T + S$. The analog of Theorem 6 for single multivalued, finitely continuous, coercive, and quasibounded generalized pseudomonotone operator S such that $D(S)$ contains a dense linear subspace is due to Browder and Hess [1]. If S is quasimonotone with weakly closed graph or graph of T is weakly closed and S is monotone of type (M) , the arguments used in the proof of Theorem 6 can be easily carried out to conclude the surjectivity of $T + S$. The reader is referred to Gupta [28] for a result for $T + S$, where graph of T is assumed to be weakly closed and $S : X \supseteq D(S) \rightarrow 2^{X^*}$ is quasibounded, finitely continuous coercive operator of type (M) such that $D(S)$ contains a dense linear subspace. Theorem 6 improves and gives unifications of the existing surjectivity results due to Le [16], Asfaw and Kartsatos [18, 24], Asfaw [19], and Kenmochi [12–14] for maximal monotone perturbations of coercive bounded pseudomonotone operators with domain, all of X . In addition, it can be easily seen that the proof of Theorem 6 can go through if the quasiboundedness of S is omitted and T is assumed to be strongly quasibounded with $0 \in T(0)$. Another observation is that the condition $\langle v^* + Sx, x \rangle \geq -d\|x\|^2$ for all $x \in D(T) \cap D(S)$ and $v^* \in Tx$ can be replaced by a stronger condition $\langle v^* + Sx, x \rangle \geq -d\|x\|$ for all $x \in D(T) \cap D(S)$ and $v^* \in Tx$, and the weak coercivity condition (i) can be relaxed to satisfy $|Tx + Sx| \rightarrow \infty$ as $\|x\| \rightarrow \infty$. On the other hand, one can easily see that weak coercivity condition on $T + S$ is automatically satisfied if $T + S$ is α -expansive. Consequently, the following corollary is immediate.

Corollary 7. *Let $T : X \supseteq D(T) \rightarrow 2^{X^*}$ be strongly quasibounded maximal monotone with $0 \in T(0)$ and let $S : X \supseteq D(S) \rightarrow X^*$ be demicontinuous generalized pseudomonotone. Suppose $W \subseteq D(S)$ is a real reflexive separable Banach space dense and continuously embedded in X . Assume, further, that $T + S$ is α -expansive mapping and there exists $d \geq 0$ such that $\alpha > d$ and $\langle Sx, x \rangle \geq -d\|x\|^2$ for all $x \in D(S)$. Then $T + S$ is surjective.*

Proof. Since $0 \in T(0)$ and T is monotone, by the condition on S , it follows that $\langle v^* + Sx, x \rangle \geq -d\|x\|^2$ for all $x \in D(T) \cap D(S)$. Furthermore, by the expansiveness of $T + S$, for some $u_0 \in D(T) \cap D(S)$ and $v_0^* \in Tu_0$, we arrive at

$$\begin{aligned} \|v^* + Sx\| &\geq \|v^* + Sx - (v_0^* + Su_0)\| - \|v_0^* + Su_0\| \\ &\geq \alpha \|x - u_0\| - \|v_0^* + Su_0\| \\ &\geq \alpha \|x\| - \alpha \|u_0\| - \|v_0^* + Su_0\| \\ &= \alpha \|x\| - A_0, \end{aligned} \tag{49}$$

where $A_0 = \alpha\|u_0\| + \|v_0^* + Su_0\|$, for all $x \in D(T) \cap D(S)$ and $v^* \in Tx$. This shows that $T + S$ satisfies conditions of Theorem 6. By applying similar arguments as in the last part of the proof of Theorem 6 and using the strong quasiboundedness of T instead of quasiboundedness of S , we conclude that $T + S$ is surjective. The details are omitted here. \square

It is worth noticing here that Corollary 7 gives a partial positive answer for Nirenberg’s problem on the surjectivity of expansive mapping in a real separable reflexive Banach space. More precisely, Corollary 7 gives surjectivity of densely defined demicontinuous generalized pseudomonotone expansive mapping. To the best of the authors knowledge, this result was unknown. For related surjectivity results for continuous expansive mappings in a real Hilbert space, we cite the papers by Kartsatos [29] and Xiang [30]. For range result for single continuous quasimonotone expansive mapping defined from arbitrary reflexive Banach space into its dual space X^* , the reader is referred to the paper due to Asfaw [25].

The content of the following theorem addresses the solvability of operator equations involving operators of the type $L + S$, where $L : X \supseteq D(L) \rightarrow X^*$ is linear, densely defined, monotone, and closed, and $S : X \supseteq D(S) \rightarrow X^*$ is quasibounded demicontinuous of type (M) such that $D(L)$ lies in $D(S)$.

Theorem 8. *Let $L : X \supseteq D(L) \rightarrow X^*$ be closed, densely defined, and linear monotone, and let $S : X \supseteq D(S) \rightarrow X^*$ be quasibounded demicontinuous of type (M) with respect to L such that $D(L)$ lies in $D(S)$. Assume, further, that there exist $\mu \geq 0$ and $\alpha > d \geq 0$ such that*

$$\langle Lx + Sx, x \rangle \geq -d\|x\|^2 \tag{50}$$

for all $x \in D(L)$ and either

(i)

$$\|Lx + Sx\| \geq \alpha \|x\| - \mu \quad \forall x \in D(L) \tag{51}$$

or

(ii) *there exists $\phi : [0, \infty) \rightarrow (-\infty, \infty)$ such that $\phi(t) \rightarrow \infty$ as $t \rightarrow \infty$ and*

$$\|Lx + Sx\| \geq \phi(\|x\|) \|x\| \quad \forall x \in D(L). \quad (52)$$

Then $L + S$ is surjective.

Proof. Fix $f^* \in X^*$. Let $Y = D(L)$ and let $\|\cdot\|_Y$ be the graph norm on Y given by

$$\|x\|_Y = \|x\|_X + \|Lx\|_{X^*}, \quad x \in Y. \quad (53)$$

It is well-known that Y equipped with the graph norm becomes a real reflexive separable Banach space. By Lemma 4, let H be a Hilbert space and let $Q : H \rightarrow Y$ be a compact injection such that $Q(H)$ is dense in Y . Let $j : Y \rightarrow X$ be the natural embedding of Y into X and let $j^* : X^* \rightarrow Y^*$ be its adjoint. It follows that $\psi = jQ$ is a compact injection from H into X . By using the graph norm on Y , it follows that Y is dense and continuously embedded in X . Moreover, by the inner product condition on $L + S$, for each $\delta > 0$, there exists $R_\delta > 0$ such that

$$\begin{aligned} \langle Lx + Sx + \delta J_1 x, x \rangle &\geq \delta \|x\|^3 - d \|x\|^2 \\ &= \|x\|^3 \left[\delta - \frac{d}{\|x\|} \right] > 0 \end{aligned} \quad (54)$$

for all $x \in D(L) \cap D(S) \cap \partial B_{R_\delta}(0)$. Let $G_\delta = B_{R_\delta}(0)$. By using the arguments used in the first half of the proof of Theorem 6, we see that $G_\delta \cap Y$ is open in Y and $\partial_Y(G_\delta \cap Y) \subseteq \partial G_\delta \cap Y$. Let $U = \psi^{-1}(G_\delta \cap Y)$. Since $j : Y \rightarrow X$ and $Q : H \rightarrow Y$ are continuous, it follows that U is open in H . Since the operator $j^*Lj : Y \rightarrow Y^*$ is linear and monotone, it is continuous. By the arguments used in the proof of Theorem 6, using Y in place of X and the closed convex subset \bar{U} of H , it follows that the mapping $C_\varepsilon : \bar{U} \rightarrow H$ defined by

$$C_\varepsilon(v) = -\varepsilon^{-1} (\psi^*(L + S + \delta J_1) Jv - f^*), \quad v \in \bar{U} \quad (55)$$

is compact. In addition, we see that

$$\begin{aligned} \partial_H U &= \partial_H \psi^{-1}(G_\delta \cap Y) \subseteq \psi^{-1}(\partial_Y(G_\delta \cap Y)) \\ &\subseteq \psi^{-1}(\partial G_\delta \cap Y). \end{aligned} \quad (56)$$

Following the argument as in the proof of Theorem 6, it is not difficult to see that $x \neq \lambda C_\varepsilon(x)$ for all $x \in \partial_H U$ and all $\lambda \in (0, 1)$. Consequently, by Lemma 5, we obtain that, for each $\varepsilon > 0$, C_ε has a fixed point in U . Therefore, for each $\varepsilon_n \downarrow 0^+$, there exists $x_n \in U$ such that

$$\varepsilon_n x_n + \psi^*(L + S + \delta J_1) \psi x_n = \psi^* f^* \quad (57)$$

for all n ; that is,

$$\begin{aligned} \langle \varepsilon_n x_n, x \rangle + \langle Q^*(j^*(L + S + \delta J_1) jQx_n), x \rangle \\ = \langle Q^* j^*(f^*), x \rangle \quad \forall x \in H. \end{aligned} \quad (58)$$

Since $j : Y \rightarrow X$ and $Q : H \rightarrow Y$, by the definition of Q^* and j^* , we see that

$$\langle \varepsilon_n x_n, x \rangle + \langle (L + S + \delta J_1) Qx_n, Qx \rangle = \langle f^*, Qx \rangle \quad \forall x \in H. \quad (59)$$

Since $\psi x_n \in G_\delta \cap Y$ and G_δ is bounded in X , it follows that the sequence $\{jx_n\} = \{Qx_n\}$ is bounded in X . From (57), using the monotonicity $L(L(0) = 0)$, boundedness of $\{Qx_n\}$, and quasiboundedness of S , we get the boundedness of the sequence $\{SQx_n\}$. This gives

$$\begin{aligned} \varepsilon_n \|x_n\|^2 &= -\langle Q^*(L + S + \delta J_1) Qx_n - Q^* f^*, x_n \rangle \\ &\leq -\langle SQx_n + \delta J_1 Qx_n - f^*, Qx_n \rangle \\ &\leq \|SQx_n + \delta J_1 Qx_n - f^*\| \|Qx_n\| \quad \forall n. \end{aligned} \quad (60)$$

As a result, we get the boundedness of $\{\varepsilon_n \|x_n\|^2\}$. If $\{x_n\}$ is bounded, then $\varepsilon_n x_n \rightarrow 0$ as $n \rightarrow \infty$. If $\{x_n\}$ is unbounded, by passing into a subsequence, we see that

$$\varepsilon_n \|x_n\| = \frac{\varepsilon_n \|x_n\|^2}{\|x_n\|} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (61)$$

In all cases, we assume without loss of generality that $\varepsilon_n x_n \rightarrow 0$ as $n \rightarrow \infty$. As a result, we get

$$\begin{aligned} \langle LQx_n, Qx \rangle &= -\langle \varepsilon_n x_n, x \rangle - \langle SQx_n + \delta J_1 Qx_n, Qx \rangle \\ &\quad + \langle f^*, Qx \rangle \\ &\leq \|\varepsilon_n x_n + SQx_n + \delta J_1 Qx_n + f^*\| \|Qx\| \\ &\leq \mu \|Qx\|, \end{aligned} \quad (62)$$

where μ is an upper bound for the sequence $\{\|\varepsilon_n x_n + SQx_n + \delta J_1 Qx_n + f^*\|\}$. Since $Q(H)$ is dense in Y , for each $y \in Y$, there exists a sequence $\{y_m\}$ in H such that $Qy_m \rightarrow y$ as $n \rightarrow \infty$. This gives

$$\begin{aligned} |\langle LQx_n, y \rangle| &= \lim_{m \rightarrow \infty} |\langle LQx_n, Qy_m \rangle| \leq \lim_{m \rightarrow \infty} \mu \|Qy_m\| \\ &= \mu \|y\|. \end{aligned} \quad (63)$$

By similar argument, the density of Y in X implies that

$$|\langle LQx_n, x \rangle| \leq \mu \|x\| \quad \forall x \in X. \quad (64)$$

By using the uniform boundedness principle, we conclude that $\{LQx_n\}$ is bounded. Assume without loss of generality that $Qx_n \rightarrow x_0$ in X , $SQx_n \rightarrow v_0^*$, and $LQx_n \rightarrow h^*$ in X^* as $n \rightarrow \infty$. Since L is closed linear, it follows that $x_0 \in Y$ and $h^* = Lx_0$. By following the arguments used in the first half of the proof of Theorem 6 along with (57), we get

$$\limsup_{n \rightarrow \infty} \langle SQx_n - f^*, Qx_n - x_0 \rangle \leq 0. \quad (65)$$

On the other hand, from (58), by using $x_n \in H$ in place of x , we see that

$$\langle LQx_n + SQx_n - f^*, Qx_n \rangle \leq 0 \quad \forall n. \quad (66)$$

Since S is of type (M) with respect to L , it follows that $S - f^*$ is also of type (M) with respect to L , which yields $x_0 \in D(S)$ and $v_0^* = Sx_0$. Finally, letting $n \rightarrow \infty$ in (57), we get $\psi^*(L + S + \delta J_1)x_0 = \psi^* f^*$; that is, $Q^* j^*(L + S + \delta J_1)x_0 = Q^* j^* f^*$. Since $Q(H)$ and Y are dense in Y and X , respectively, it follows that j^* and Q^* are one to one. Therefore, we arrive at $Lx_0 + Sx_0 + \delta J_1 x_0 = f^*$. Consequently, for each $\delta_n \downarrow 0^+$, there exists $y_n \in D(L)$ such that

$$Ly_n + Sy_n + \delta_n J_1 y_n = f^* \quad \forall n. \tag{67}$$

Since L is closed and S is of type (M) , by weak coercivity condition on $L + S$, the same arguments used in the second half of the proof of Theorem 6 can be carried over to conclude the solvability of the equation $Lx + Sx \ni f^*$ in $D(L)$. Since $f^* \in X^*$ is arbitrary, we conclude that $L + S$ is surjective. The details are omitted here. \square

The following corollary gives a characterization of linear maximal monotone operator in separable reflexive Banach space.

Corollary 9. *Let X be a real separable reflexive Banach space and let $L : X \supseteq D(L) \rightarrow X^*$ be linear operator. Then the following two statements are equivalent:*

- (i) L is maximal monotone,
- (ii) L is monotone, densely defined, and closed.

Proof. The proof of (i) implies (ii) follows by the well-known result due to Brézis (cf. Zeidler [9, Theorem 32. L, page 897]). Next we prove (ii) implies (i). Let $\lambda > 0$. It is sufficient to show that $R(L + \lambda J) = X^*$. To this end, we will use Theorem 8. Since L is linear and monotone, that is, $\langle Lx, x \rangle \geq 0$ for all $x \in D(L)$, and J is monotone, it follows that $\langle Lx + \lambda Jx, x \rangle \geq \lambda \|x\|^2$ for all $x \in D(L)$. Therefore, for each $\lambda > 0$, it follows that

$$\|Lx + \lambda Jx\| \geq \lambda \|x\| \quad \forall x \in D(L). \tag{68}$$

By using J in place of S in Theorem 8, we conclude that $R(L + \lambda J) = X^*$ for any $\lambda > 0$. Thus, L is maximal monotone. \square

It is worth noticing that Brézis proved (i) in arbitrary reflexive Banach space provided that L^* is monotone and (ii) holds. As a result, Corollary 9 is an improvement of the result of Brézis when X is separable. It is important to mention that Gupta [28] gave surjectivity result for graph weakly closed maximal monotone perturbations of quasibounded, finitely continuous multivalued coercive operator S of type (M) such that $D(S)$ contains a dense linear subspace of X . However, the result in Theorem 8 is for noncoercive operator S along with weak coercivity of $L + S$. It is also important to mention here that the results in Theorems 6 and 8 are new even in the case where the operator S is coercive but not everywhere defined. In conclusion, Theorems 6 and 8 gave improvements over the existing theory for maximal monotone perturbations of coercive and everywhere defined operators of pseudomonotone type.

3. Example and Discussion

In this section, we demonstrate the existence of weak solution in $X = L^p(0, T; W_0^{1,p}(\Omega))$ for the parabolic problem of the type

$$\begin{aligned} \frac{\partial u}{\partial t} - \sum_{i=1}^N \frac{\partial}{\partial x_i} a_i(x, t, u, \nabla u) &= f(x, t) \quad (x, t) \in Q \\ u(x, t) &= 0 \quad (x, t) \in \partial\Omega \times (0, T) \\ u(x, 0) &= 0 \quad x \in \Omega, \end{aligned} \tag{69}$$

where $p > 1$, p' is conjugate exponent of p , $Q = \Omega \times (0, T)$, Ω is a nonempty, bounded, and open subset of \mathbb{R}^N , and $f \in L^{p'}(Q)$ such that the following conditions are satisfied:

- (A₁) $a_i(x, t, s, \xi)$ ($i = 1, 2, \dots, N$) satisfies the Carathéodory conditions; that is, for each $(s, \xi) \in \mathbb{R} \times \mathbb{R}^N$, the function $(x, t) \mapsto a_i(x, t, s, \xi)$ is measurable and, for almost all $(x, t) \in \Omega \times (0, T)$, the function $(s, \xi) \mapsto a_i(x, t, s, \xi)$ is continuous.
- (A₂) there exists positive constants μ_1 and μ_2 such that

$$\begin{aligned} \sum_{i=1}^N (a_i(x, t, s, \xi) - a_i(x, t, s, \eta)) (\xi_i - \eta_i) \\ \geq \mu_1 |\xi - \eta|^p, \end{aligned} \tag{70}$$

$$|a_i(x, t, s, \xi)| \leq \mu_2 [|s|^{p_0} + |\xi|^{p-1}] + g(x, t)$$

for all $(x, t, s, \xi) \in \Omega \times (0, T) \times \mathbb{R} \times \mathbb{R}^N$, where $\xi = (\xi_i) \in \mathbb{R}^N$ and $\eta = (\eta_i) \in \mathbb{R}^N$, $g \in L^{p'}(Q)$, $p_0 < p - 1 + 2(p - 1)/N$, and $1 < p < N$.

Let $X = L^p(0, T; V)$, $V = W_0^{1,p}(\Omega)$, and $L : X \supseteq D(L) \rightarrow X^*$ be defined by $Lu = u'$, where u' is understood in the sense of distributions; that is,

$$\begin{aligned} \int_0^T u'(t) \psi(t) dt &= - \int_0^T u(t) \psi'(t) dt, \\ \psi &\in C_0^\infty(0, T), \end{aligned} \tag{71}$$

where $D(L) = \{u \in X : u' \in X^*, u(0) = 0\}$. We notice that

$$\begin{aligned} \langle Lu, \phi \rangle &= \int_0^T \langle u'(t), \phi(t) \rangle_V dt, \\ u &\in D(L), \phi \in X. \end{aligned} \tag{72}$$

Let $A : X \supseteq D(A) \rightarrow X^*$ be defined by

$$\begin{aligned} \langle Au, \phi \rangle &= \sum_{i=1}^n \int_Q a_i(x, t, u, \nabla u) \frac{\partial \phi(x, t)}{\partial x_i} dx dt, \\ \phi &\in X, u \in D(A), \end{aligned} \tag{73}$$

where

$$D(A) = \{u \in X : u \in L^{\bar{p}}(Q)\}, \quad \bar{p} = \frac{p_0 p}{p - 1}. \tag{74}$$

It is known from Kartsatos and Skrypnik [4] that A is quasibounded demicontinuous generalized (S_+) with respect to L such that $D(A)$ contains $D(L)$. Moreover, it is well-known that L is linear, closed, and densely defined maximal monotone. The operator A is densely defined, that is, not everywhere defined, and coercive. Since $p_0 < p - 1$ in (A_2) , operator A may be unbounded. Therefore, by Theorem 8 using A in place of S , for each $f \in L^{p'}(Q)$, we conclude that the equation $Au + Lu = f^*$ is solvable in $D(L)$, where $f^* : X \rightarrow \mathbb{R}$ is given by $\langle f^*, \phi \rangle = \int_Q f(x, t)\phi(x, t)dxdt$. Therefore, the parabolic problem (69) admits at least one weak solution in $D(L)$.

Since Ω is bounded and $g \in L^{p'}(Q)$, it is well known that A is bounded, continuous, everywhere defined, and coercive provided that $p_0 = p - 1$ in (A_2) . More precisely, these conditions on A are satisfied if condition (A_2) is replaced by

(A_3) : there exists positive constants μ_1 and μ_2 such that

$$\sum_{i=1}^N (a_i(x, t, s, \xi) - a_i(x, t, s, \eta)) (\xi_i - \eta_i) \geq \mu_1 |\xi - \eta|^p, \quad (75)$$

$$|a_i(x, t, s, \xi)| \leq \mu_2 [|s|^{p-1} + |\xi|^{p-1}] + g(x, t)$$

for all $(x, t, s, \xi) \in \Omega \times (0, T) \times \mathbb{R} \times \mathbb{R}^N$, where $\xi = (\xi_i) \in \mathbb{R}^N$ and $\eta = (\eta_i) \in \mathbb{R}^N$, $g \in L^{p'}(Q)$, and $1 < p < N$.

Abstract existence results concerning nonlinear parabolic problems of the type in (69) under conditions (A_1) and (A_3) have been intensively studied by many researchers. For some of the basic and relevant references, the reader is referred to the papers by Browder and Hess [1], Brézis [2], Le [16], Kenmochi [12–14], Guan and Kartsatos [17], Asfaw and Kartsatos [18], Asfaw [19, 25], and the references therein. For further examples and applications of perturbed everywhere defined pseudomonotone type operators to inclusion, variational inequality, and evolution problems, the reader is referred to the papers of Landes and Mustonen [31], Kobayashi and Ôtani [32], and Mustonen [33] and books of Kinderlehrer and Stampacchia [34], Browder [8], and Naniecicz and Panagiotopoulos [35] and the references therein. The method of sub-supersolution is employed in the papers by Carl and Le [20], Carl et al. [21], Carl [22], Carl and Motreanu [23], and Le [36, 37] to study existence and properties of solution(s) for evolution inclusion problems of the type

$$u \in X : u' + A(u) \ni f^* \quad \text{in } X^*, \quad u(0) = u_0, \quad (76)$$

where $X = L^p(0, T; W_0^{1,p}(\Omega))$, $p > 1$, Ω is a nonempty, bounded, and open subset of \mathbb{R}^N , and A is noncoercive but still everywhere defined operator of pseudomonotone type. For further relevant information about sub-supersolution arguments concerning evolution type problems, the reader is referred to the recent book on nonsmooth analysis due to Carl et al. [38] and the references therein. Finally, it is important to indicate the readers that the results in this paper can be conveniently applied to address nonlinear parabolic problems of type (76) as well as elliptic problems of the type

$u \in Y : -\Delta_p u + B(u) \ni g^*, g^* \in Y^*$, where $Y = W_0^{1,p}(\Omega)$ and A and B are possibly noncoercive and densely defined and satisfy conditions of either Theorem 6 or Theorem 8.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The author would like thank the editor and anonymous referee for forwarding important suggestions. Furthermore, the author would like to thank the Virginia Tech's Open Access Sub-vention fund program for covering the full article processing fee.

References

- [1] F. E. Browder and P. Hess, "Nonlinear mappings of monotone type in Banach spaces," *Journal of Functional Analysis*, vol. 11, no. 3, pp. 251–294, 1972.
- [2] H. Brézis, "Équations et inéquations non linéaires dans les espaces vectoriels en dualité," *Annales de l'Institut Fourier*, vol. 18, no. 1, pp. 115–175, 1968.
- [3] L. Nirenberg, *Topics in Nonlinear Functional Analysis*, Lecture Notes, Courant Institute of Mathematical Sciences, New York University, New York, NY, USA, 1974.
- [4] A. G. Kartsatos and I. V. Skrypnik, "Topological degree theories for densely defined mappings involving operators of type (S_+) ," *Advances in Differential Equations*, vol. 4, no. 3, pp. 413–456, 1999.
- [5] V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff International Publishing, Leyden, The Netherlands, 1975.
- [6] H. Brézis, M. G. Crandall, and A. Pazy, "Perturbations of nonlinear maximal monotone sets in banach space," *Communications on Pure and Applied Mathematics*, vol. 23, no. 1, pp. 123–144, 1970.
- [7] D. Pascali and S. Sburlan, *Nonlinear Mappings of Monotone Type*, Sijthoff and Noordhoff, Bucharest, Romania, 1978.
- [8] F. E. Browder, "Nonlinear operators and nonlinear equations of evolution in Banach spaces," *Proceedings of Symposia in Pure Mathematics*, vol. 1, pp. 1–308, 1976.
- [9] E. Zeidler, *Nonlinear Functional Analysis and Its Applications*, Springer, New York, NY, USA, 1990.
- [10] F. E. Browder, "Pseudo-monotone operators and nonlinear elliptic boundary value problems on unbounded domains," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 74, no. 7, pp. 2659–2661, 1977.
- [11] H. Brézis and L. Nirenberg, "Characterizations of the ranges of some nonlinear operators and applications to boundary value problems," *Annali della Scuola Normale Superiore di Pisa*, vol. 5, pp. 225–326, 1978.
- [12] N. Kenmochi, "Nonlinear operators of monotone type in reflexive Banach spaces, and nonlinear perturbations," *Hiroshima Mathematical Journal*, vol. 4, no. 1, pp. 229–263, 1974.
- [13] N. Kenmochi, "Pseudomonotone operators and nonlinear elliptic boundary value problems," *Journal of the Mathematical Society of Japan*, vol. 27, pp. 121–149, 1975.

- [14] N. Kenmochi, "Monotonicity and compactness methods for nonlinear variational inequalities," in *Handbook of Differential Equations, IV*, pp. 203–298, Elsevier/North-Holland, Amsterdam, The Netherlands, 2007.
- [15] Z. Guan, A. G. Kartsatos, and I. V. Skrypnik, "Ranges of densely defined generalized pseudomonotone perturbations of maximal monotone operators," *Journal of Differential Equations*, vol. 188, no. 1, pp. 332–351, 2003.
- [16] V. K. Le, "A range and existence theorem for pseudomonotone perturbations of maximal monotone operators," *Proceedings of the American Mathematical Society*, vol. 139, no. 5, pp. 1645–1658, 2011.
- [17] Z. Guan and A. G. Kartsatos, "Ranges of generalized pseudomonotone perturbations of maximal monotone operators in reflexive Banach spaces," *Contemporary Mathematics*, vol. 204, pp. 107–123, 1997.
- [18] T. M. Asfaw and A. G. Kartsatos, "A Browder topological degree theory for multivalued pseudomonotone perturbations of maximal monotone operators in reflexive Banach spaces," *Advances in Mathematical Sciences and Applications*, vol. 22, no. 1, pp. 91–148, 2012.
- [19] T. M. Asfaw, "New surjectivity results for perturbed weakly coercive operators of monotone type in reflexive Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 113, pp. 209–229, 2015.
- [20] S. Carl and V. K. Le, "Quasilinear parabolic variational inequalities with multi-valued lower order terms," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 65, no. 5, pp. 845–864, 2014.
- [21] S. Carl, V. K. Le, and D. Motreanu, "Existence, comparison, and compactness results for quasilinear variational-hemivariational inequalities," *International Journal of Mathematics and Mathematical Sciences*, vol. 2005, no. 3, pp. 401–417, 2005.
- [22] S. Carl, "Existence and comparison results for noncoercive and nonmonotone multivalued elliptic problems," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 65, no. 8, pp. 1532–1546, 2006.
- [23] S. Carl and D. Motreanu, "General comparison principle for quasilinear elliptic inclusions," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 70, no. 2, pp. 1105–1112, 2009.
- [24] T. M. Asfaw and A. G. Kartsatos, "New results for perturbations of locally defined generalized pseudomonotone operators in separable reflexive Banach spaces," *Advances in Mathematical Sciences and Applications*, vol. 24, pp. 1–10, 2014.
- [25] T. M. Asfaw, "New variational inequality and surjectivity theories for perturbed noncoercive operators and application to nonlinear problems," *Advances in Mathematical Sciences and Applications*, vol. 24, pp. 611–668, 2014.
- [26] F. E. Browder and B. A. Ton, "Nonlinear functional equations in Banach spaces and elliptic super-regularization," *Mathematische Zeitschrift*, vol. 105, no. 3, pp. 177–195, 1968.
- [27] A. Granas and J. Dugundji, *Fixed Point Theory*, Springer Monographs in Mathematics, Springer, New York, NY, USA, 2003.
- [28] C. P. Gupta, "On an operator equation involving mappings of monotone type," *Proceedings of the American Mathematical Society*, vol. 53, no. 1, pp. 143–148, 1975.
- [29] A. G. Kartsatos, "On the connection between the existence of zeros and the asymptotic behavior of resolvents of maximal monotone operators in reflexive Banach spaces," *Transactions of the American Mathematical Society*, vol. 350, no. 10, pp. 3967–3987, 1998.
- [30] T. Xiang, "Notes on expansive mappings and a partial answer to Nirenberg's problem," *Electronic Journal of Differential Equations*, vol. 2013, no. 2, pp. 1–16, 2013.
- [31] R. Landes and V. Mustonen, "On pseudo-monotone operators and nonlinear noncoercive variational problems on unbounded domains," *Mathematische Annalen*, vol. 248, no. 3, pp. 241–246, 1980.
- [32] J. Kobayashi and M. Ôtani, "Topological degree for $(S)_+$ -mappings with maximal monotone perturbations and its applications to variational inequalities," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 59, no. 1-2, pp. 147–172, 2004.
- [33] V. Mustonen, "On elliptic operators in divergence form; old and new with applications," in *Proceedings of the International Conference on Function Spaces and Differential Operators and Nonlinear Analysis (FSDONA '04)*, pp. 188–200, 2004.
- [34] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, vol. 88 of *Classics in Applied Mathematics*, Academic Press, New York, NY, USA, 1980.
- [35] Z. Naniewicz and P. Panagiotopoulos, *Mathematical Theory of Hemivariational Inequalities and Applications*, vol. 188 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 1995.
- [36] V. K. Le, "On variational inequalities with maximal monotone operators and multivalued perturbing terms in Sobolev spaces with variable exponents," *Journal of Mathematical Analysis and Applications*, vol. 388, no. 2, pp. 695–715, 2012.
- [37] V. K. Le, "On variational and quasi-variational inequalities with multivalued lower order terms and convex functionals," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 94, pp. 12–31, 2014.
- [38] S. Carl, V. K. Le, and D. Motreanu, *Nonsmooth Variational Problems and their Inequalities: Comparison Principles and Applications*, Springer Monographs in Mathematics, Springer, New York, NY, USA, 2007.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

