



## POSTER PRESENTATION

## Open Access

# KIR-HLA footprints and NK cell-mediated recognition of HIV-1

CF Thobakgale<sup>1\*</sup>, J Carlson<sup>2</sup>, J Wright<sup>1</sup>, N van Teijlingen<sup>3</sup>, M Jaggernath<sup>1</sup>, P Goulder<sup>4</sup>, BD Walker<sup>3</sup>, M Carrington<sup>5</sup>, D Heckerman<sup>2</sup>, M Altfeld<sup>3</sup>, T Ndung'u<sup>1</sup>

From AIDS Vaccine 2012

Boston, MA, USA. 9-12 September 2012

## Background

Increasing data suggest an important role of KIR+ NK cells in the control of HIV-1, however the precise mechanisms on how NK cells recognize HIV-1-infected cells remain poorly understood. KIRs can bind to HLA class I molecules, but the binding affinity of these interactions is dependent on the HLA class I presented epitope. Recent reports have suggested that sequence variations within HIV-1 epitopes presented by HLA class I can affect the binding of inhibitory KIRs expressed on NK cells, potentially modulating NK cell responses to infected cells. Here we investigated whether HIV-1 might adapt to the combined KIR/HLA genotypes on a population level to identify areas within HIV-1 that might be targeted by KIR+ NK cells.

## Methods

HIV-1 Gag was sequenced in 390 untreated chronically clade C infected individuals from KwaZulu-Natal, South Africa. All study subjects were HLA class I and KIR typed. Phylogenetically-corrected logistic regression analysis of KIR/HLA associated Gag sequence polymorphisms was performed and q-values were used for multiple test correction.

## Results

A total of 93 sequence polymorphisms significantly associated with the combined HLA/KIR genotypes were identified ( $p < 0.05$ ), 6 of them with a false-positive rate of less than 20% ( $q < 0.2$ ). These significant associations were independent of previously identified KIR or HLA-linked polymorphisms.

## Conclusion

This study identified several sequence polymorphisms within HIV-1 Gag that were significantly associated with the expression of combined KIR/HLA genotypes at the population level, indicating adaptation of HIV-1 to NK cell mediated immune pressure. KIR/HLA class I binding studies in the context of the sequence polymorphisms and studies for NK cell function are ongoing to determine the consequence of these sequence changes for NK cell-mediated recognition of HIV-1-infected cells.

## Author details

<sup>1</sup>University of KwaZulu-Natal, Durban, South Africa. <sup>2</sup>Microsoft Research, Redmond, WA, USA. <sup>3</sup>Ragon Institute of MIT, MGH and Harvard, Boston, MA, USA. <sup>4</sup>University of Oxford, Oxford, UK. <sup>5</sup>Cancer and Inflammation Program, Laboratory of Experimental Immunology, Frederick, MD, USA.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P170

Cite this article as: Thobakgale et al.: KIR-HLA footprints and NK cell-mediated recognition of HIV-1. *Retrovirology* 2012 **9**(Suppl 2):P170.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>University of KwaZulu-Natal, Durban, South Africa  
Full list of author information is available at the end of the article