

Metagenomic analysis of orange colored protrusions from the muscle of Queen Conch *Lobatus gigas* (Linnaeus, 1758)

Jaison H. Cuartas¹, Juan F. Alzate², Claudia X. Moreno-Herrera¹ and Edna J. Marquez¹

¹Facultad de Ciencias, Laboratorio de Biología Molecular y Celular, Universidad Nacional de Colombia, Medellín, Antioquia, Colombia

²Facultad de Medicina, Centro Nacional de Secuenciación Genómica, Universidad de Antioquia, Medellín, Antioquia, Colombia

ABSTRACT

The endangered marine gastropod, *Lobatus gigas*, is an important fishery resource in the Caribbean region. Microbiological and parasitological research of this species have been poorly addressed despite its role in ecological fitness, conservation status and prevention of potential pathogenic infections. This study identified taxonomic groups associated with orange colored protrusions in the muscle of queen conchs using histological analysis, 454 pyrosequencing, and a combination of PCR amplification and automated Sanger sequencing. The molecular approaches indicate that the etiological agent of the muscle protrusions is a parasite belonging to the subclass Digenea. Additionally, the scope of the molecular technique allowed the detection of bacterial and fungi clades in the assignment analysis. This is the first evidence of a digenean infection in the muscle of this valuable Caribbean resource.

Subjects Biodiversity, Genomics, Marine Biology, Parasitology

Keywords Microbiology, Parasitic infection, 454 pyrosequencing, Bioinformatics, Trematode

Submitted 20 January 2017

Accepted 10 January 2018

Published 15 February 2018

Corresponding author

Edna J. Marquez,
ejmarque@unal.edu.co

Academic editor

Mauricio Rodríguez-Lanetty

Additional Information and
Declarations can be found on
page 12

DOI 10.7717/peerj.4307

© Copyright
2018 Cuartas et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

INTRODUCTION

The queen conch, *Lobatus gigas*, is an endangered marine gastropod of great socioeconomic, cultural and ecological importance in the Caribbean region. This species was included in Appendix II of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) in 1992 and the Red List of the International Union for Conservation of Nature (IUCN) in 1994. Despite these regulations, natural stocks of this species continue to decline (*Theile, 2001; Aldana, 2003*), likely by the loss of breeding habitats and detrimental human activities such as overfishing (*Glazer & Quintero, 1998; Aldana, 2003*).

Compared with studies in basic biology (*Randall, 1964*), fisheries (*Brownell & Stevely, 1981; Theile, 2001; Prada et al., 2009*), and genetics (*Mitton, Berg Jr & Orr, 1989; Tello-Cetina, Rodríguez-Gil & Rodríguez-Romero, 2005; Zamora-Bustillos et al., 2011; Márquez et al., 2013*), parasitological and microbial studies of *L. gigas* are less explored (*Acosta et al., 2009; Aldana et al., 2011; Rodríguez, Hariharan & Nimrod, 2011; Pérez et al., 2014*). So far, only one parasitic infection, with *Apicomplexa* coccidian protozoon, has been reported in *L. gigas* (*Baqueiro et al., 2007; Aldana et al., 2009; Aldana et al., 2011; Gros, Frenkiel &*

Aldana, 2009; Volland et al., 2010). Similarly, only three published studies report the association between *L. gigas* and bacteria of the family Vibrionaceae (*Acosta et al., 2009*), the phyla Firmicutes, Proteobacteria, Actinobacteria (*Pérez et al., 2014*) as well as potential bacterial pathogens (*Rodríguez, Hariharan & Nimrod, 2011*). Two recent investigations have also studied the symbiotic association of *L. gigas* with dinoflagellates of the genus *Symbiodinium* (*Banaszak, Ramos & Goulet, 2013; García Ramos & Banaszak, 2014*).

Moreover, an unknown etiological agent sporadically produces orange colored protrusions in the muscle of *L. gigas* in the Colombian San Andres archipelago. However, it remains to be elucidated whether such lesions are caused by different agents and posteriorly colonized by pigment-producing microorganisms or digenean infections as found in other marine gastropods. Specifically, the infections of *Cercaria parvicaudata* and *Renicola roscovita* have been reported to produce orange/lemon colored sporocysts in different tissues of *Littorina* snails (*Stunkard, 1950; Galaktionov & Skirnisson, 2000*), whereas *Renicola thaidus* has been found infecting *Nucella lapillus* (*Galaktionov & Skirnisson, 2000*). These trematodes, *C. parvicaudata*, *R. roscovita* and *R. thaidus* are considered synonymous based on morphological similarities and cercariae size parameters (*Werding, 1969*). Similarly, lemon-cream to orange colored sporocysts are produced by the congeners *Renicola* sp. “polychaetophila” and *Renicola* sp. “martini” in infections of the gonad and digestive glands in *Cerithidea californica* (*Hechinger & Miura, 2014*).

This work studied the presence of parasites, bacteria and fungi in orange colored protrusions in the muscle of Colombian Caribbean queen conchs. This was achieved by using histological analysis and molecular approaches based on 454 FLX and capillary automated sequencing using an ABI PRISM 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA). This 454 FLX next-generation platform (Roche, Basel, Switzerland) permits high-throughput identification of hundreds of samples at reasonable cost and time consumption (*Mardis, 2008*). This approach allows functional analysis of sequencing data sets for comparative analysis of microbiome diversity of orange colored protrusions found in the muscle of *L. gigas* by using metagenomic taxonomical classifiers (*Huson et al., 2007; Huson et al., 2011*). This information is required for queen conch conservation and management strategies of potential pathogenic infections for human beings.

MATERIALS AND METHODS

Orange colored protrusions were taken from three pieces of frozen muscle from one specimen of *L. gigas* processed for food trading in the Colombian Caribbean, San Andres archipelago (between 12°–16°N and 78°–82°W). These samples were provided by the Gobernación del Archipiélago de San Andrés, Providencia y Santa Catalina, through the scientific cooperation agreement #083/2012.

Since the etiological agent of these orange colored protrusions was unknown, we used three approaches to elucidate the origin of these lesions: (1) histological analysis, (2) 454 pyrosequencing of one whole genome shotgun library and (3) automated capillary sequencing (Sanger) of PCR amplified products to confirm the results provided by the metagenomic analysis. For histological analysis, samples from orange colored muscle

were fixed in 10% neutral phosphate-buffered formalin. The samples were prepared for histological examination by paraffin wax techniques and stained with hematoxylin and eosin following standard protocols ([García del Moral, 1993](#); [Prophet et al., 1995](#)).

Due to scarcity of samples, the orange protrusions were pooled and ground with liquid nitrogen to extract the genomic DNA using the commercial DNAeasy Blood & Tissue Kit (Qiagen, Hilden, Germany), according to manufacturer recommendations. Sample pooling was performed to obtain high-quantity and high-quality DNA required for the generation of the genomic library. Purified DNA from the pooled sample was sequenced using the 454 Whole Genome Shotgun strategy according to standard protocols recommended by 454 GS FLX platform (Roche, Basel, Switzerland) at the Centro Nacional de Secuenciación Genómica, Universidad de Antioquia ([Margulies et al., 2005](#)). The obtained raw reads were end polished of low-quality regions with the toolkit PRINSEQ lite ([Schmieder & Edwards, 2011](#)) and assembled using MIRA3 v3.4 software ([Chevreux, Wetter & Suhai, 1999](#)).

Classification of assembled contigs was carried out using the BLAST algorithm against nucleotide and protein non-redundant databases of the NCBI with further computation of the taxonomic position of the assembled dataset with MEGAN software v5.5.3 ([Huson et al., 2011](#)). This metagenomic software uses a Lowest Common Ancestor-based algorithm that assigns each contig to taxa such that the taxonomical level of the assigned taxon reflects the level of conservation of the sequence ([Huson et al., 2007](#)). Then, species-specific and widely conserved sequences were assigned to particular taxa as described by [Huson et al. \(2007\)](#). The contigs were classified using a bit-score threshold of 50, retaining only those hits that were within 10% of the best hit for a contig. Additionally, the *E*-value confidence criterion was set at $1E-15$, even though a threshold value of $1E-04$ is considered a good match ([De Wit et al., 2012](#)). Only contig alignment lengths above 100 nucleotides for BLASTN comparison or 100 amino acids for BLASTX comparisons were included in the assignment analysis. These analyses, comparing DNA or protein sequences, were carried out independently.

Furthermore, protein analysis assignments were classified to the proper taxonomic level according to [Monzoorul Haque et al. \(2009\)](#), who empirically proposes identity thresholds for restricting the assignments. Alignments having identities in ranges of 61–65%, 56–60%, 51–55% and 41–50% were conservatively restricted in the level of family, order, class and phylum, respectively. The identity threshold of 66–100% was used for restricting the assignment of contigs to either species or genus or family levels. Additionally, the taxonomic level within this identity range was distinguished by the difference between the two alignment parameters, the percentage of identities and positives.

Moreover, a 1,000 bp fragment of the mitochondrial *cytochrome c oxidase I* gene was amplified by PCR following conditions reported by [Leung et al. \(2009\)](#) and primers described by [Bowles et al. \(1993\)](#) (JB3: 5'-TTTTTTGGGCATCCTGAGGTTTAT-3') and [Králová-Hromadová \(2008\)](#) (trem.cox1.rnrl: 5' AATCATGATGCAAAAGGTA-3'). This sequence was used instead of ribosomal genes since the bioinformatic analysis indicated a high enrichment of molluscan and some fungi ribosomal sequences, which limited the amplification of the helminth sequences of 18S and 28S ribosomal genes (data not shown). The *cytochrome c oxidase I* amplicon was sequenced by automated Sanger method using

an ABI PRISM 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA) and compared by BLASTN against the NCBI nucleotide database to look for sequence matches of reported organisms.

Finally, a Bayesian tree was constructed using the sequence obtained from orange colored protrusions and published sequences of Platyhelminthes. Bayesian tree construction was performed using MrBayes (MB) V3.2 (Ronquist *et al.*, 2012) setting the GTR+I+G4 substitution model estimated by the software IQ-TREE, with 1,000,000 generations sampled every 1,000 generations and the other analysis parameters as default values. The convergence of the Markov Chain Monte Carlo iterations was assessed with the Potential Scale Reduction Factor (PSRF = 1; Gelman & Rubin, 1992) and the standard deviation of split frequencies (0.001).

RESULTS

Assembly and metagenomic approach

The massive shotgun sequencing generated 515,368 reads with an average length of 279 bp that were cleaned and then assembled using MIRA software into 5,180 contigs. Taxonomic classification of the contigs was carried out using the software MEGAN. For this analysis, all the contigs were compared with the NCBI's non-redundant protein database using the software BLASTX. With this strategy, 1,588 (30.7%) contigs were assigned to taxa (Bacteria: 412; Eukaryota: 1,157; other: 19), 866 (16.7%) were unassigned and 2,726 (52.6%) presented no hits. As expected, the Eukaryota group was dominant due to the origin of the sample. Furthermore, the group Gastropoda was frequently found in this analysis (186 hits), although many sequences remained unclassified due to the poor representation of these organisms in the public databases. Many bacterial sequences were also identified; 19 were assigned to the fungi group and 22 sequences were assigned to the Trematoda category. No viral or protozoa sequences were detected.

Following the MEGAN pipeline, with nucleotide comparisons using BLASTN and the nt/nr database, results were poorly classified. One contig was assigned to the root, 462 (8.9%) to particular taxa (Bacteria: 267; Eukaryota: 191; other: 4), 32 (0.6%) were unassigned and 4,685 (90.5%) had no hits. At the nucleotide level, most of the sequences were left unclassified. This reflects the lack of sequences in the databases of closely related organisms to the ones reported in this research.

Bacteria and fungi associated with orange colored protrusions

The ranges for the confidence criterion represented by the *E*-value, similarity and identity for protein comparisons are shown in Table 1. Bacteria assignments included the class *Gammaproteobacteria* and the phylum *Firmicutes*, which includes the orders *Bacillales* and *Lactobacillales* (Fig. 1). Specifically, the class *Gammaproteobacteria* showed 322 assignments for *Psychrobacter*, exhibiting hits to several types of proteins with strains of *Psychrobacter* sp. (Fig. 1, Table 1). The identity criterion for *Psychrobacter* sp. ranged from 67% to 100% (Table 1). Similarly, nucleotide sequences showed hits for different genomic regions of *Psychrobacter* sp. strains and congeners, displaying identities ranging from 72% to 99% and alignment lengths from 104 bp to 1,061 bp (Table 2).

Table 1 Diversity content in bacteria and fungi clades found in a pooled sample of orange colored protrusions from *L. gigas* muscle using translated contig sequences and the taxonomic classifier MEGAN.

Contig	Organism	Gene	Ranges				Assignment ^a
			E-value	Positives (%)	Identities (%)	Length (aa)	
8	<i>Psychrobacter</i> sp.	binding protein, kinase, transporter, adaptor, hypothetical proteins, membrane protein, glycosylase	0.000E+00; 1.325E-30	96; 100	96; 100	113; 259	Species
11	<i>Psychrobacter</i> sp.	dehydrogenase, hypothetical proteins, catalase, cytoplasmic protein, propionase, transferase, chaperone, deaminase, membrane protein	0.000E+00; 4.451E-73	94; 100	90; 99	105; 306	Genus
2	<i>Psychrobacter</i> sp.	channel protein, hypothetical protein	1.189E-107; 4.390E-77	77; 83	67; 73	183; 242	Family
2	<i>Carnobacterium jeotgali</i> ¹	replication initiator, phosphorylase	5.120E-93; 4.347E-87	99; 100	99; 100	135; 138	Species
2	<i>Carnobacterium</i> sp. ¹	hypothetical protein; integrase	2.691E-127; 4.938E-58	100; 100	100; 100	109; 183	Species
1	<i>Carnobacterium</i> sp. ¹	integrase	1.940E-61	86	80	129	Genus
1	<i>Carnobacterium</i> sp. ¹	hypothetical protein	2.200E-69	94	86	158	Family
2	<i>Lactobacillus jensenii</i> ¹	hypothetical protein	2.022E-78; 2.550E-60	73; 85	68; 81	166; 169	Genus
1	<i>Enterococcus faecalis</i> ¹	hypothetical protein	3.700E-45	68	44	172	Phylum
7	<i>Brochothrix thermosphacta</i> ^b	kinase, transcriptional regulator, transporter, ribosomal protein, reductase, hypothetical proteins	3.514E-162; 9.715E-69	100; 100	100; 100	180; 248	Species
6	<i>Brochothrix thermosphacta</i> ^b	dehydrogenase, hypothetical proteins, transposase, transferase	1.918E-148; 1.431E-53	76; 100	71; 99	120; 243	Genus
1	<i>Planococcus antarcticus</i> ^b	hypothetical protein	2.650E-55	90	80	106	Family
1	<i>Bacillus cytotoxicus</i> ^b	synthetase	7.050E-164	83	71	223	Family
1	<i>Lactococcus lactis</i> subsp. <i>Lactis</i> ¹	replication protein	6.020E-91	81	62	164	Family
1	<i>Staphylococcus aureus</i> ^b	hypothetical protein	2.050E-32	71	64	104	Family
1	<i>Fusarium oxysporum</i>	glutamine-rich protein	1.064E-16	56	41	243	Phylum
1	<i>Fusarium oxysporum</i>	glutamine-rich protein	2.900E-16	85	82	243	Genus
1	<i>Neurospora tetrasperma</i>	hypothetical protein	1.426E-58	92	90	107	Genus

Notes.

^aThe assignments were classified to the taxonomic level according to [Monzoorul Haque et al. \(2009\)](#), ^bBacillales, ¹Lactobacillales.




Figure 1 Phylogenetic diversity of translated contigs from orange colored protrusions of *Lobatus gigas* computed by MEGAN. The nodes of the cladogram represent the assigned taxa and the numbers indicate the relative abundance of assigned contigs.

Full-size [DOI: 10.7717/peerj.4307/fig-1](https://doi.org/10.7717/peerj.4307/fig-1)

On the other hand, 18 assignments for the order *Lactobacillales* (Fig. 1) showed hits for diverse proteins exhibiting similarities and identities up to 100% for *Carnobacterium jeotgali* and *Carnobacterium* sp. (Table 1). We also found hits for proteins of *Lactobacillus jensenii* and *Enterococcus faecalis* displaying identities above 68% and 44%, respectively (Table 1). Similarly, nucleotide analysis showed matches for genome regions and plasmids of *Carnobacterium* sp., displaying identities above 82% (Table 2). Additionally, the single hits for a plasmid and a genome fragment of *Enterococcus casseliflavus* and *Enterococcus faecalis* exhibited identities of 81% and 73%, respectively (Table 2).

A total of 28 contigs were assigned to different *Bacillales* bacteria within the phylum *Firmicutes* (Fig. 1); specifically, *Brochothrix thermosphacta* showed hits for several proteins exhibiting identities up to 100% (Table 1). *Planococcus antarcticus*, *Bacillus cytotoxicus*, *Lactococcus lactis* subsp. *Lactis* and *Staphylococcus aureus* showed identity ranges from 62% to 80% (Table 1). Furthermore, the nucleotide analysis showed hits for diverse bacteria belonging to genus *Listeria*, *Bacillus* and *Paenibacillus* (Table 2).

Only three out of 19 assignments to fungi clades satisfied the selection parameters; two hits supported the taxonomical levels of phylum and genus for *Fusarium oxysporum* and one hit classified to the genus taxonomical level for *Neurospora tetrasperma* (Table 1). In addition, nucleotide analysis (Table 2) showed three assignments for *Mrakia frigida* (rRNA genes, two hits) and *Togninia minima* (putative polyubiquitin protein mRNA, one hit).

A parasite associated with orange colored protrusions

The histological approach showed a tissue lesion characterized by the aggregation of hemocytes (cells endowed with phagocytic and immune-related functions) inside isolated foci surrounded by smooth muscle fibers and a basophilic tissue contiguous to a lamellated

Table 2 Diversity content in bacteria and fungi clades found in a pooled sample of orange colored protrusions from *L. gigas* muscle using nucleotide contig sequences and the taxonomic classifier MEGAN.

Contig	Organism	Gene	Ranges		
			E-value	Identities (%)	Length (bp)
1	<i>Psychrobacter</i> sp.	pRWF101	0.000E+00	98	993
1	<i>Psychrobacter</i> sp.	gf	0.000E+00	99	815
6	<i>Psychrobacter</i> sp.	p, gf	0.000E+00; 7.247E-59	90; 96	176; 679
1	<i>Psychrobacter</i> sp.	gf	0.000E+00	88	1,162
7	<i>Psychrobacter</i> sp.	p, gf	0.000E+00; 3.877E-131	80; 85	520; 914
3	<i>Psychrobacter</i> sp.	gf	4.648E-143; 2.430E-35	76; 79	288; 748
1	<i>Psychrobacter cryohalolentis</i>	gf	0.000E+00	93	1,047
3	<i>Psychrobacter cryohalolentis</i>	p, gf	5.595E-138; 6.961E-40	92; 95	112; 352
4	<i>Psychrobacter cryohalolentis</i>	gf	3.316E-153; 1.041E-103	83; 84	419; 537
3	<i>Psychrobacter cryohalolentis</i>	gf	1.626E-92; 3.574E-74	75; 78	524; 700
1	<i>Psychrobacter arcticus</i>	gf	0.000E+00	91	646
1	<i>Psychrobacter arcticus</i>	gf	5.090E-120	94	307
1	<i>Psychrobacter arcticus</i>	gf	8.830E-47	91	148
11	<i>Psychrobacter arcticus</i>	gf	0.000E+00; 3.900E-148	81; 88	611; 1,061
11	<i>Psychrobacter arcticus</i>	gf	0.000E+00; 6.195E-25	80; 89	104; 603
7	<i>Psychrobacter arcticus</i>	gf	1.962E-172; 1.181E-24	72; 79	224; 978
1	<i>Carnobacterium</i> sp. ¹	gf	0.000E+00	94	2,422
1	<i>Carnobacterium</i> sp. ¹	pWNCR9	0.000E+00	92	1,466
1	<i>Carnobacterium</i> sp. ¹	gf	0.000E+00	96	1,244
1	<i>Carnobacterium</i> sp. ¹	gf	0.000E+00	98	1,029
3	<i>Carnobacterium</i> sp. ¹	p, gf	0.000E+00; 1.117E-155	95; 98	321; 624
6	<i>Carnobacterium</i> sp. ¹	p, gf	0.000E+00; 8.843E-57	82; 88	264; 897
1	<i>Enterococcus casseliflavus</i> ¹	pTnpA	4.680E-85	81	464
1	<i>Enterococcus faecalis</i> ¹	gf	3.310E-37	73	537
1	<i>Listeria grayi</i> ^b	23S rRNA	0.00E+00	90	1,263
1	<i>Listeria welshimeri</i> ^b	23S rRNA	1.150E-176	88	539
1	<i>Listeria monocytogenes</i> ^b	gf	3.850E-103	74	902
1	<i>Listeria innocua</i> ^b	gf	7.870E-109	76	754
1	<i>Bacillus megaterium</i> ^b	gf	5.410E-170	78	908
1	<i>Bacillus toyonensis</i> ^b	gf	1.430E-66	74	622
1	<i>Bacillus cereus</i> ^b	gf	2.570E-17	78	174
1	<i>Paenibacillus larvae</i> ^b	pPL374	1.110E-170	100	335
1	Uncultured compost bacterium ^b	16S rRNA	0.000E+00	99	436
1	<i>Mrakia frigida</i>	25S rRNA	0.000E+00	100	1,429
1	<i>Mrakia frigida</i>	18S rRNA	0.000E+00	99	1,793
1	<i>Togninia minima</i>	protein mRNA	3.407E-28	90	3,261

Notes.^bBacillales, ¹Lactobacillales.

gf, genome fragment; p, plasmid; rRNA, ribosomal fragment.




Figure 2 Histological sections of orange-colored protrusions in the muscle of *Lobatus gigas*. The lesions showed hemocytes stained purplish-blue and smooth muscle fibers pink-red in color. (A) Presence of lamellated membrane (arrowhead) (40 \times). (B) Granulation process (arrowhead) (100 \times).

Full-size  DOI: [10.7717/peerj.4307/fig-2](https://doi.org/10.7717/peerj.4307/fig-2)

membrane (Fig. 2A). Additionally, some lesions exhibit interstitial immunocyte inclusions with morphology similar to a granulation process (Fig. 2B). Although the histological approach did not allow for detection of key features for identification, the microscopic images showed structures around 0.55 mm in diameter, which are compatible with immature developmental stages of a trematode.

Furthermore, the metagenomic analysis assigned 22 contigs to the trematode parasites clade. Specifically, these contigs had hits to an endonuclease-reverse transcriptase of *Schistosoma mansoni* (17) and *Schistosoma japonicum* (4), showing identities above 46% and 42%, respectively. Similarly, in the nucleotide analysis, seven contigs showed identities above 71% for different regions of two chromosomes of *S. mansoni* (Table 3).

We successfully amplified and sequenced a 740 bp region that confirmed the presence of trematode DNA in the *L. gigas* tissue (GenBank accession number KR092371). Moreover, this sequence clustered in a basal position to the suborder Xiphidiata (Trematoda: Digenea), which encompasses *Renicola* and *Helicometrina* genera (posterior probability: 0.98; Fig. 3). Additionally, the BLASTN results showed hits for some members of Xiphidiata, such as *Helicometrina labrisomid* (query coverage: 89%; identity: 77%), *Renicola cerithidicola* (query coverage: 70%; identity: 78%), *Synthesium pontoporiae* (query coverage: 42%; identity: 83%) and *Haematoloechus* sp. (query coverage: 39%; identity: 77%).

DISCUSSION

In this study, three approaches, including histological analysis, 454 pyrosequencing and automated Sanger amplification of the *cytochrome c oxidase I* gene, were used to explore the potential causal agent of orange colored protrusions in the muscle of *L. gigas*. Identification by histology was limited since no characteristic structures were detected in the sample. Also, several contigs had no hits for proteins (~52%) or nucleotide sequences (~90%), indicating a lack of information on such sequences in reference databases. This explanation is plausible since the current protein sequence reference databases cover only a small

Table 3 Diversity content in the trematoda clade found in a pooled sample of orange colored protrusions from *L. gigas* muscle using contig sequences and the taxonomic classifier MEGAN.

Contig	Organism	Gene	Ranges				Assignment ^a
			E-value	Positives (%)	Identities (%)	Length	
Translated contig sequences							
2	<i>Schistosoma japonicum</i>	endonuclease-reverse transcriptase	3.637E-61; 6.062E-42	75; 76	64; 64	141; 165	Family
2	<i>Schistosoma japonicum</i>	endonuclease-reverse transcriptase	2.102E-122; 1.995E-56	61; 63	42; 43	262; 489	Phylum
5	<i>Schistosoma mansoni</i>	endonuclease-reverse transcriptase	3.147E-152; 4.075E-45	77; 81	61; 67	155; 345	Family
5	<i>Schistosoma mansoni</i>	endonuclease-reverse transcriptase	2.497E-172; 2.424E-61	71; 74	56; 59	204; 346	Order
4	<i>Schistosoma mansoni</i>	endonuclease-reverse transcriptase	0.000E+00; 9.203E-89	69; 70	52; 52	275; 695	Class
3	<i>Schistosoma mansoni</i>	endonuclease-reverse transcriptase	1.343E-66; 1.719E-51	65; 68	46; 50	193; 262	Phylum
Nucleotide contig sequences							
1	<i>Schistosoma mansoni</i>	chromosome fragment W	1.320E-20	80	80	161	–
5	<i>Schistosoma mansoni</i>	chromosome fragments	1.640E-55; 7.006E-27	71; 73	71; 73	649; 763	–
1	<i>Schistosoma mansoni</i>	chromosome fragment 4	1.990E-19	77	77	199	–

Notes.

^aThe assignments were classified to the taxonomic level according to [Monzoorul Haque et al. \(2009\)](#).

fraction of the biodiversity believed to be present in the environment ([Wu et al., 2009](#)). Despite these limitations, the alignment lengths of the contigs (≥ 100 nucleotides or amino acids) and the bit-scores (50) used in this research ensure a reasonable level of confidence in the taxonomic assignments ([Huson et al., 2007](#)).

Bacteria and fungi associated with orange colored protrusions

The scope of the massive sequencing approach allowed the detection of some bacteria previously reported as microbiota associated with *L. gigas* ([Acosta et al., 2009](#); [Pérez et al., 2014](#)), as well as new reports. For instance, *Psychrobacter* sp. was found in the *L. gigas* muscle in both nucleotide and protein analyses ([Tables 1](#) and [2](#)). This outcome corroborates previous studies that found *Psychrobacter* sp. in environmental ([Acosta et al., 2009](#); [Pérez et al., 2014](#)) and tissue ([Pérez et al., 2014](#)) samples from *L. gigas*.

However, this study also found bacteria and fungi that have not been reported so far in *L. gigas*. Specifically, homologous protein and nucleotide sequences of species (e.g., *Carnobacterium jeotgali*), family and genus of *Carnobacterium* sp. were detected in the *L. gigas* muscle ([Tables 1](#) and [2](#)). *Carnobacterium* strains have been reported to inhabit live fish and a variety of seafood, dairy and meat ([Leisner et al., 2007](#)).

In addition, this research found homologous protein and nucleotide sequences of genus *Bacillus* and *Enterococcus* in the affected tissue of *L. gigas*. *Bacillus* species have ubiquitous distribution, inhabiting different environments such as soils, rocks, vegetation, foods and




Figure 3 Bayesian tree obtained from *cytochrome c oxidase I* gene sequences of orange-colored protrusions from the muscle of *L. gigas* (OPML) and GenBank Platyhelminthes sequences. *P. macrorchis* (JN592039.1), *M. sebastis* (NC_009055.1), *E. multilocularis* (AB018440.2), *E. granulosus* (AF297617.1), *T. saginata* (AY195858.1), *T. solium* (AY211880.1), *T. pisiformis* (GU569096.1), *S. mansoni* (AF216698.1), *S. japonicum* (AF215860.1), *S. chasanensis* (KU757308.1), *P. saginatum* (KX097855.1), *F. hepatica* (AF216697.1), *F. gigantica* (KF543342.1), *H. labrisomi* (KJ996009.1), *R. pinguis* (KU563724.1), *R. sterna* (KU563723.1), *R. lari* (KU563727.1), *R. cerithidicola* (KF512573.1), *R. buchanani* (KF512572.1), *Renicola* sp. 'polychaetophila' (KF512551.1) and *Renicola* sp. 'martini' (KF512560.1).

Full-size DOI: 10.7717/peerj.4307/fig-3

waters (Nicholson, 2002). Similarly, the ubiquitous nature of enterococci determines their frequently being found in foods as contaminants, although their predominant habitat is human and animal gastrointestinal tracts (Giraffa, 2002). However, they also occur in soil, surface waters, vegetables and fermented foods such as sausages, meat and cheese (Giraffa, 2002; Foulquié et al., 2006).

Furthermore, another bacteria present in the sample was *Brochothrix thermosphacta*, as it was assigned to bacterial species or genus taxonomical levels according to Monzoorul Haque et al. (2009). This bacterium, closely related to *Listeria*, is a non-proteolytic food spoilage organism in prepacked meats and fish products (Gardner, 1981; Lannelongue et al., 1982; Pin, García de Fernando & Ordóñez, 2002). In addition, some *Listeria* hits were detected in the nucleotide analysis, although the identity values did not allow species identification. This result is concordant with studies that have isolated *Listeria* members from freshwater and marine environments (Colburn et al., 1990; El Marrakchi, Boum'handi & Hamama, 2005).

Metagenomic analysis also showed some fungi assignments related to *Fusarium*, *Neurospora*, *Togninia* and *Mrakia*. Both *Fusarium* and *Neurospora* exhibit wide distribution,

including humid tropical and subtropical marine environments (Steele, 1967; Turner, Perkins & Fairfield, 2001; Babu et al., 2010; Summerell et al., 2010; Jebaraj et al., 2012; Saravanan & Sivakumar, 2013; Kumar, Gousia & Latha, 2015). Specifically, some *Fusarium* species are associated with infections in crustaceans and cultivated fishes (Hatai, 2012), whereas other species are endosymbionts of some seaweeds (Suryanarayanan, 2012), corals (Raghukumar & Ravindran, 2012) and some sea sponges (Höller et al., 2000; Wang, Li & Zhu, 2008; Liu et al., 2010; Paz et al., 2010).

In contrast, *Togninia* and *Mrakia* show more restricted distributions. For instance, *Togninia* comprises pathogenic fungi responsible for the development of wood diseases, and some strains have been isolated from submerged wood from streams, lakes, ponds, reservoirs and ditches (Hu, Cai & Hyde, 2012). Likewise, several *Mrakia* species have been isolated from icy environments, including meltwaters from glaciers and permafrost in Antarctica (Hua et al., 2010; Pathan et al., 2010; Carrasco et al., 2012; Zhang et al., 2012; Tsuji et al., 2013a; Tsuji et al., 2013b), Argentina (Brizzio et al., 2007; De Garcia, Brizzio & Broock, 2012), the Qinghai-Tibet Plateau (Su et al., 2016), Italy (Turchetti et al., 2008; Branda et al., 2010; Thomas-Hall et al., 2010) and the Arctic (Pathan et al., 2010).

Considering that several of the new bacteria reports are related to food microorganisms, we hypothesized that they might grow under environmental or freezing conditions instead of being native microbiota. Fungi findings suggest an environmental source; however, since some species of *Fusarium* and *Neurospora* produce orange spores (Davis & Perkins, 2002; Hatai, 2012), the colored protrusions found in *L. gigas* may be due to an opportunistic or primary fungal infection. Thus, the role of bacteria and fungi in the muscle of *L. gigas* and their relationship with the lesion, native microbiota or the environment remains to be explored.

A parasite associated with orange colored protrusions

Histology showed evidence of a membrane, that is consistent with a syncytium, enclosing a multicellular parasite, a mollusk inflammatory response elicited by hemocytes (De Vico & Carella, 2012). Moreover, such membranes are also compatible with the wall layers of the life cycle stage of Platyhelminthes, suggesting a possible infection by trematodes that infect other mollusks (Cake, 1976; Sorensen & Minchella, 2001). This finding was supported by the metagenomics analysis that showed sequences homologous to an endonuclease-reverse transcriptase of some species of trematodes like *Schistosoma* (Table 3). This result is expected since highly repetitive sequences, such non-LTR retrotransposons with an estimated copy number going up to 24,000, are more likely to be detected in whole genome shotgun amplification (DeMarco et al., 2005). Since databases of protein and nucleotide sequences are currently enriched with *Schistosoma*, but exhibit a poor representation of most members of Trematoda, these assignments require cautioned interpretation. Moreover, the lack of information in reference databases for most of the sequences from the studied sample (~90% of nucleotide sequences and ~52% of proteins) explains the relatively low number of hits for the parasite compared with Trematoda, bacteria and fungi taxa. Although these assignments are biased by the nucleotide and protein sequences available in the NCBI

databases, it supports the histological finding that the protrusions may be caused by a trematode.

The Bayesian tree supported the last result due to clustering of the sample in a basal position to the suborder Xiphidiata (Trematoda: Digenea), which includes *Renicola* species that produce colored pigments (Stunkard, 1950; Galaktionov & Skirnisson, 2000). The phylogenetic relationships with Xiphidiata were consistent with the BLASTN analysis that revealed genetic similitudes between the sequence found in this study and *Renicola*, *Helicometrina*, *Synthesium* and *Haematoloechus*, although its genetic distance with other members of these genera remains to be determined due to the lack of information for *cytochrome c oxidase I* and endonuclease-reverse transcriptase sequences of these taxa.

The molecular findings let us hypothesize about the structures approximately 0.55 mm in diameter found in the snail muscle tissue, although histology did not allow detection of key features for its identification. According to the life cycle described for Xiphidiata, the microscopic life cycle stage found in the muscle of *L. gigas* could be sporocysts, which are described to preferentially infect gonads and digestive glands, but can also disperse to other tissues in the form of more sporocysts or rediae (Cribb et al., 2003). Based on the structure's size, other stages such as metacercaria seems unlikely at least in *Renicola*, as they exhibit 0.12 to 0.16 mm in diameter (Stunkard, 1964). However, the evidence presented here is not enough to conclude which parasitic stage was observed within the colored lesions.

In conclusion, this study found evidence of a trematode infection, as well the presence of fungi and bacteria in the protruded muscle of *L. gigas*, which provides novel information for the parasitology and microbiology of this species. This first insight of a trematode infection in *L. gigas* is a baseline to expand the toolset to identify these organisms, the trematode life cycle, environmental conditions that trigger its appearance and epidemiological aspects regarding the host and possible effects on human health.

ACKNOWLEDGEMENTS

The authors thank the Laboratorio de Patología Animal and Centro de Secuenciación Genómica, Universidad de Antioquia, for assistance and services provided. The authors also thank to the anonymous reviewers for their comments, which substantially improved the final version of this article.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was funded by the Universidad Nacional de Colombia, Sede Medellín-the Gobernación del Archipiélago de San Andrés, Providencia y Santa Catalina, Grant 201010011420 (Scientific Cooperation Agreement # 083/2012), Convocatoria Nacional para el Programa Jóvenes Investigadores Colciencias No.525-2012-Identificación de parásitos asociados al caracol pala *Strombus gigas* en la Reserva de Biosfera Seaflower (Jaison H. Cuartas) and Programa de sostenibilidad 2016-2017, Grupo de Parasitología, Vicerrectoría de investigación, Universidad de Antioquia. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Universidad Nacional de Colombia.

Sede Medellín-the Gobernación del Archipiélago de San Andrés.

Providencia y Santa Catalina: 201010011420.

Scientific Cooperation Agreement: #083/2012.

Convocatoria Nacional para el Programa Jóvenes Investigadores Colciencias No.525-2012-Identificación de parásitos asociados al caracol pala *Strombus gigas* en la Reserva de Biosfera Seaflower (Jaison H. Cuartas).

Programa de sostenibilidad, Vicerrectoría de investigación, Universidad de Antioquia.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Jaison H. Cuartas, Juan F. Alzate, Claudia X. Moreno-Herrera and Edna J. Marquez conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.

Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The muscle samples of *Lobatus gigas* were provided by the Gobernación del Archipiélago de San Andrés, Providencia y Santa Catalina, through the scientific cooperation agreement # 083/2012.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

GenBank accession [KR092371](#).

Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as a [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4307#supplemental-information>.

REFERENCES

- Acosta EA, Gómez E, Romero M, Cadavid GE, Moreno CX. 2009. Molecular identification of bacterial populations associated to Queen Conch (*Strombus gigas*) from Colombian Caribe. *Acta Biologica Colombiana* 14:83–96.

- Aldana D. 2003.** *El caracol Strombus gigas: conocimiento integral para su manejo sustentable en el Caribe.* In: *Programa iberoamericano de ciencia y tecnología para el desarrollo.* Yucatán: CYTED.
- Aldana D, Frenkiel L, Brulé T, Montero J, Baqueiro E. 2011.** Occurrence of Apicomplexa-like structures in the digestive gland of *Strombus gigas* throughout the Caribbean. *Journal of Invertebrate Pathology* **106**:174–178 DOI [10.1016/j.jip.2010.09.004](https://doi.org/10.1016/j.jip.2010.09.004).
- Aldana D, Glazer R, Delgado G, Baqueiro E, Montero J. 2009.** Occurrence of Apicomplexa infecting Queen Conch, *Strombus gigas*, from offshore and nearshore in Florida. *Proceedings of the Gulf and Caribbean Fisheries Institute* **61**:491–497.
- Babu R, Varadharajan D, Soundarapandian P, Balasubramanian R. 2010.** Fungi diversity in different coastal marine ecosystem along South East Coast of India. *International Journal for Microbiological Research* **1**:175–178.
- Banaszak AT, García Ramos M, Goulet TL. 2013.** The symbiosis between the gastropod *Strombus gigas* and the dinoflagellate *Symbiodinium*: an ontogenic journey from mutualism to parasitism. *Journal of Experimental Marine Biology and Ecology* **449**:358–365 DOI [10.1016/j.jembe.2013.10.027](https://doi.org/10.1016/j.jembe.2013.10.027).
- Baqueiro E, Frenkiel L, Zetina A, Aldana D. 2007.** Coccidian (Apicomplexa) parasite infecting *Strombus gigas* Linné, 1758 digestive gland. *Journal of Shellfish Research* **26**:319–321 DOI [10.2983/0730-8000\(2007\)26\[319:CAPISG\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2007)26[319:CAPISG]2.0.CO;2).
- Bowles J, Hope M, Tiu WU, Liu X, McManus DP. 1993.** Nuclear and mitochondrial genetic markers highly conserved between Chinese and Philippine *Schistosoma japonicum*. *Acta Tropica* **55**:217–229 DOI [10.1016/0001-706X\(93\)90079-Q](https://doi.org/10.1016/0001-706X(93)90079-Q).
- Branda E, Turchetti B, Diolaiuti G, Pecci M, Smiraglia C, Buzzini P. 2010.** Yeast and yeast-like diversity in the southernmost glacier of Europe (Calderone Glacier, Apennines, Italy). *FEMS Microbiology Ecology* **72**:354–369 DOI [10.1111/j.1574-6941.2010.00864.x](https://doi.org/10.1111/j.1574-6941.2010.00864.x).
- Brizzio S, Turchetti B, De García V, Libkind D, Buzzini P, Van Broock M. 2007.** Extracellular enzymatic activities of basidiomycetous yeasts isolated from glacial and subglacial waters of northwest Patagonia (Argentina). *Canadian Journal of Microbiology* **53**:519–525 DOI [10.1139/W07-010](https://doi.org/10.1139/W07-010).
- Brownell WN, Stevely JM. 1981.** The biology, fisheries, and management of the Queen Conch, *Strombus gigas*. *Marine Fisheries Review* **43**(7):1–12.
- Cake EW. 1976.** A key to larval cestodes of shallow-water, benthic mollusks of the northern Gulf of Mexico. In: *Proceedings of the helminthological society of Washington, vol. 43.* 160–171.
- Carrasco M, Rozas JM, Barahona S, Alcaíno J, Cifuentes V, Baeza M. 2012.** Diversity and extracellular enzymatic activities of yeasts isolated from King George Island, the sub-Antarctic region. *BMC Microbiology* **12**:251–260 DOI [10.1186/1471-2180-12-251](https://doi.org/10.1186/1471-2180-12-251).
- Chevreur B, Wetter T, Suhai S. 1999.** Genome sequence assembly using trace signals and additional sequence information. In: *Computer science and biology: proceedings of the German conference on bioinformatics (GCB).* 45–56.

- Colburn KG, Kaysner CA, Abeyta Jr C, Wekell MM. 1990. *Listeria* species in a California coast estuarine environment. *Applied and Environmental Microbiology* 56:2007–2011.
- Cribb TH, Bray RA, Olson PD, Littlewood DTJ. 2003. Life cycle evolution in the Digenea: a new perspective from phylogeny. *Advances in Parasitology* 54:197–254 DOI 10.1016/S0065-308X(03)54004-0.
- Davis RH, Perkins DD. 2002. *Neurospora*: a model of model microbes. *Nature Reviews. Genetics* 3:397–403 DOI 10.1038/nrg797.
- De Garcia V, Brizzio S, Van Broock MR. 2012. Yeasts from glacial ice of Patagonian Andes, Argentina. *FEMS Microbiology Ecology* 82:540–550 DOI 10.1111/j.1574-6941.2012.01470.x.
- DeMarco R, Machado AA, Bisson-Filho AW, Verjovski-Almeida S. 2005. Identification of 18 new transcribed retrotransposons in *Schistosoma mansoni*. *Biochemical and Biophysical Research Communications* 333:230–240 DOI 10.1016/j.bbrc.2005.05.080.
- De Vico G, Carella F. 2012. Morphological features of the inflammatory response in molluscs. *Research in Veterinary Science* 93:1109–1115 DOI 10.1016/j.rvsc.2012.03.014.
- De Wit P, Pespeni MH, Ladner JT, Barshis DJ, Seneca F, Jaris H, Therkildsen NO, Morikawa M, Palumbi SR. 2012. The simple fool's guide to population genomics via RNA-Seq: an introduction to high-throughput sequencing data analysis. *Molecular Ecology Resources* 12:1058–1067 DOI 10.1111/1755-0998.12003.
- El Marrakchi A, Boum'handi N, Hamama A. 2005. Performance of a new chromogenic plating medium for the isolation of *Listeria monocytogenes* from marine environments. *Letters in Applied Microbiology* 40:87–91 DOI 10.1111/j.1472-765X.2004.01651.x.
- Foulquié MR, Sarantinopoulos P, Tsakalidou E, De Vuyst L. 2006. The role and application of enterococci in food and health. *International Journal of Food Microbiology* 106:1–24 DOI 10.1016/j.ijfoodmicro.2005.06.026.
- Galaktionov KV, Skirnisson K. 2000. Digeneans from intertidal molluscs of SW Iceland. *Systematic Parasitology* 47:87–101 DOI 10.1023/A:1006426117264.
- García del Moral R. 1993. *Laboratorio de Anatomía Patológica*. Madrid: McGraw-Hill, Interamericana de España.
- García Ramos M, Banaszak AT. 2014. Symbiotic association between symbiodinium and the gastropod *Strombus gigas*: larval acquisition of symbionts. *Marine Biotechnology* 16:193–201 DOI 10.1007/s10126-013-9536-x.
- Gardner GA. 1981. *Brochothrix thermosphacta* (*Microbacterium thermosphactum*) in the spoilage of meats: a review. In: Roberts TA, Hobbs GA, Christian JHB, Skovgaard N, eds. *Psychrotrophic microorganisms in spoilage and pathogenicity*. London: Academic Press, 139–173.
- Gelman A, Rubin DB. 1992. Inference from iterative simulation using multiple sequences. *Statistical Science* 7:457–472 DOI 10.1214/ss/1177011136.
- Giraffa G. 2002. Enterococci from foods. *FEMS Microbiology Reviews* 26:163–171 DOI 10.1016/S0168-6445(02)00094-3.

- Glazer R, Quintero I. 1998.** Observations on the sensitivity of Queen Conch to water quality: implications for coastal development. In: *Proceedings of the Gulf and Caribbean Fisheries Institute*, 78–93.
- Gros O, Frenkiel L, Aldana D. 2009.** Structural analysis of the digestive gland of the Queen Conch *Strombus gigas* Linnaeus, 1758 and its intracellular parasites. *Journal of Molluscan Studies* 75:59–68 DOI 10.1093/mollus/eyn041.
- Hatai K. 2012.** Diseases of fish and shellfish caused by marine fungi. In: Raghukumar C, ed. *Biology of marine fungi*. Berlin: Springer-Verlag Berlin Heidelberg, 15–52.
- Hechinger RF, Miura O. 2014.** Two “new” renicolid trematodes (Trematoda: Digenea: Renicolidae) from the California horn snail, *Cerithidea californica* (Haldeman, 1840) (Gastropoda: Potamididae). *Zootaxa* 3784:559–574 DOI 10.11646/zootaxa.3784.5.5.
- Höller U, Wright AD, Matthée GF, König GM, Draeger S, Aust H-J, Schulz B. 2000.** Fungi from marine sponges: diversity, biological activity and secondary metabolites. *Mycological Research* 104:1354–1365 DOI 10.1017/S0953756200003117.
- Hu D-M, Cai L, Hyde KD. 2012.** Three new ascomycetes from freshwater in China. *Mycologia* 104:1478–1489 DOI 10.3852/11-430.
- Hua MX, Chi Z, Liu GL, Buzdar MA, Chi ZM. 2010.** Production of a novel and cold-active killer toxin by *Mrakia frigida* 2E00797 isolated from sea sediment in Antarctica. *Extremophiles* 14:515–521 DOI 10.1007/s00792-010-0331-6.
- Huson DH, Auch AF, Qi J, Schuster SC. 2007.** MEGAN analysis of metagenomic data. *Genome Research* 17:377–386 DOI 10.1101/gr.5969107.
- Huson DH, Mitra S, Ruscheweyh HJ, Weber N, Schuster SC. 2011.** Integrative analysis of environmental sequences using MEGAN4. *Genome Research* 21:1552–1560 DOI 10.1101/gr.120618.111.
- Jebaraj CS, Forster D, Kauff F, Stoeck T. 2012.** Molecular diversity of fungi from marine oxygen-deficient environments (ODEs). In: Raghukumar C, ed. *Biology of marine fungi*. Berlin, Heidelberg: Springer-Verlag, 1899–1208.
- Králová-Hromadová I, Spakulová M, Horácková E, Turceková L, Novobilský A, Beck R, Koudela B, Marinculić A, Rajský D, Pybus M. 2008.** Sequence analysis of ribosomal and mitochondrial genes of the giant liver fluke *Fascioloides magna* (Trematoda: Fasciolidae): intraspecific variation and differentiation from *Fasciola hepatica*. *Journal of Parasitology* 94:58–67 DOI 10.1645/GE-1324.1.
- Kumar KA, Gousia SK, Latha JNL. 2015.** Evaluation of biological activity of secondary metabolites of *Neurospora crassa* from Machilipatnam Sea Water. *Research Journal of Microbiology* 10:377–384 DOI 10.3923/jm.2015.377.384.
- Lannelongue M, Hanna MO, Finne G, Nickelson IIR, Vanderzant C. 1982.** Storage characteristics of finfish fillets (*Archosargus probatocephalus*) packaged in modified gas atmospheres containing carbon dioxide. *Journal of Food Protection* 45:440–444 DOI 10.4315/0362-028X-45.5.440.
- Leisner JJ, Laursen BG, Prévost H, Drider D, Dalgaard P. 2007.** *Carnobacterium*: positive and negative effects in the environment and in foods. *FEMS Microbiology Reviews* 31:592–613 DOI 10.1111/j.1574-6976.2007.00080.x.

- Leung TLF, Donald KM, Keeney DB, Koehler AV, Peoples RC, Poulin R. 2009.** Trematode parasites of Otago Harbour (New Zealand) soft-sediment intertidal ecosystems: life cycles, ecological roles and DNA barcodes. *New Zealand Journal of Marine and Freshwater Research* **43**:857–865 DOI [10.1080/00288330909510044](https://doi.org/10.1080/00288330909510044).
- Liu WC, Li CQ, Zhu P, Yang JL, Cheng KD. 2010.** Phylogenetic diversity of culturable fungi associated with two marine sponges: *Haliclona simulans* and *Gelliodes carnosa*, collected from the Hainan Island coastal waters of the South China Sea. *Fungal Diversity* **42**:1–15 DOI [10.1007/s13225-010-0022-8](https://doi.org/10.1007/s13225-010-0022-8).
- Mardis ER. 2008.** Next-generation DNA sequencing methods. *Annual Review of Genomics and Human Genetics* **9**:387–402 DOI [10.1146/annurev.genom.9.081307.164359](https://doi.org/10.1146/annurev.genom.9.081307.164359).
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bembien LA, Berka J, Braverman MS, Chen Y-J, Chen Z, Dewell SB, Du L, Fierro JM, Gomes XV, Godwin BC, He W, Helgesen S, Ho CH, Irzyk GP, Jando SC, Alenquer MLI, Jarvie TP, Jirage KB, Kim J-B, Knight JR, Lanza JR, Leamon JH, Lefkowitz SM, Lei M, Li J, Lohman KL, Lu H, Makhijani VB, McDade KE, McKenna MP, Myers EW, Nickerson E, Nobile JR, Plant R, Puc BP, Ronan MT, Roth GT, Sarkis GJ, Simons JF, Simpson JW, Srinivasan M, Tartaro KR, Tomasz A, Vogt KA, Volkmer GA, Wang SH, Wang Y, Weiner MP, Yu P, Begley RF, Rothberg JM. 2005.** Genome sequencing in microfabricated high-density picolitre reactors. *Nature* **437**:376–380 DOI [10.1038/nature04726](https://doi.org/10.1038/nature04726).
- Márquez E, Landínez-García RM, Ospina-Guerrero SP, Segura JA, Prada M, Castro E, Correa JL, Borda C. 2013.** In: *Genetic analysis of Queen Conch Strombus gigas from the Southwest Caribbean*. Gulf and Caribbean Fisheries Institute, 410–416.
- Mitton JB, Berg Jr CJ, Orr KS. 1989.** Population structure, larval dispersal, and gene flow in the Queen Conch, *Strombus gigas*, of the Caribbean. *Biological Bulletin* **177**:356–362 DOI [10.2307/1541595](https://doi.org/10.2307/1541595).
- Monzoorul Haque M, Ghosh TS, Komanduri D, Mande SS. 2009.** SOrt-ITEMS: sequence orthology based approach for improved taxonomic estimation of metagenomic sequences. *Bioinformatics* **25**:1722–1730 DOI [10.1093/bioinformatics/btp317](https://doi.org/10.1093/bioinformatics/btp317).
- Nicholson WL. 2002.** Roles of *Bacillus* endospores in the environment. *Cellular and Molecular Life Sciences* **59**:410–416 DOI [10.1007/s00018-002-8433-7](https://doi.org/10.1007/s00018-002-8433-7).
- Pathan AAK, Bhadra B, Begum Z, Shivaji S. 2010.** Diversity of yeasts from puddles in the vicinity of Midre Lovénbreen glacier, arctic and bioprospecting for enzymes and fatty acids. *Current Microbiology* **60**:307–314 DOI [10.1007/s00284-009-9543-3](https://doi.org/10.1007/s00284-009-9543-3).
- Paz Z, Komon-Zelazowska M, Druzhinina IS, Aveskamp MM, Shnaiderman A, Aluma Y, Carmeli S, Ilan M, Yarden O. 2010.** Diversity and potential antifungal properties of fungi associated with a Mediterranean sponge. *Fungal Diversity* **42**:17–26 DOI [10.1007/s13225-010-0020-x](https://doi.org/10.1007/s13225-010-0020-x).
- Pérez OM, Posada M, Cadavid GE, Moreno CX. 2014.** Assessment of the bacterial community diversity associated with the queen conch *Strombus gigas* (Linnaeus, 1758) from the Caribbean coast of Colombia using denaturing gradient gel electrophoresis and culturing. *Aquaculture Research* **45**:773–786 DOI [10.1111/are.12016](https://doi.org/10.1111/are.12016).

- Pin C, García de Fernando GD, Ordóñez JA. 2002.** Effect of modified atmosphere composition on the metabolism of glucose by *Brochothrix thermosphacta*. *Applied and Environmental Microbiology* **68**:4441–4447 DOI [10.1128/AEM.68.9.4441](https://doi.org/10.1128/AEM.68.9.4441).
- Prada M, Castro E, Taylor E, Puentes V, Appeldoorn R, Daves N. 2009.** Non-detrimental findings for the Queen Conch (*Strombus gigas*) in Colombia. In: *NDF workshop case studies*. 34.
- Prophet E, Mills B, Arrington J, Sobón L. 1995.** *Métodos histotecnológicos*. Washington, D.C.: Instituto de Patología de las Fuerzas Armadas de los Estados Unidos de América (AFIP), Registro de Patología de los Estados Unidos de América (ARP), 280.
- Raghukumar C, Ravindran J. 2012.** Fungi and their role in corals and coral reef ecosystems. In: Raghukumar C, ed. *Biology of marine fungi*. Berlin, Heidelberg: Springer-Verlag, 89–113.
- Randall JE. 1964.** Contributions to the biology of the Queen Conch, *Strombus gigas*. *Bulletin of Marine Science of the Gulf and Caribbean* **14**:246–295.
- Rodriguez AI, Hariharan H, Nimrod S. 2011.** Occurrence and antimicrobial drug resistance of potential bacterial pathogens from shellfish, including Queen Conchs (*Strombus gigas*) and Whelks (*Cittarium pica*) in Grenada. *Webmed Central Microbiology* **2**:1–11.
- Ronquist F, Teslenko M, Vander Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012.** MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**:539–542 DOI [10.1093/sysbio/sys029](https://doi.org/10.1093/sysbio/sys029).
- Saravanan R, Sivakumar T. 2013.** Biodiversity and biodegradation potentials of fungi isolated from marine systems of East Coast of Tamil Nadu, India. *International Journal of Current Microbiology and Applied Sciences* **2**:192–201.
- Schmieder R, Edwards R. 2011.** Quality control and preprocessing of metagenomic datasets. *Bioinformatics* **27**:863–864 DOI [10.1093/bioinformatics/btr026](https://doi.org/10.1093/bioinformatics/btr026).
- Sorensen RE, Minchella DJ. 2001.** Snail-trematode life history interactions: past trends and future directions. *Parasitology* **123**(Suppl):S3–S18 DOI [10.1017/S0031182001007843](https://doi.org/10.1017/S0031182001007843).
- Steele CW. 1967.** Fungus populations in marine waters and coastal sands of the Hawaiian, Line, and Phoenix Islands. *Pacific Science* **21**:317–331.
- Stunkard HW. 1950.** Further observations on *Cercaria parvicaudata* Stunkard and Shaw, 1931. *Biological Bulletin* **99**:136–142 DOI [10.2307/1538757](https://doi.org/10.2307/1538757).
- Stunkard HW. 1964.** Studies on the trematode genus *Renicola*: Observations on the life-history, specificity, and systematic position. *Biological Bulletin* **126**:467–489 DOI [10.2307/1539314](https://doi.org/10.2307/1539314).
- Su Y, Jiang X, Wu W, Wang M, Hamid MI, Xiang M, Liu X. 2016.** Genomic, transcriptomic and proteomic analysis provide insights into the cold adaptation mechanism of the obligate psychrophilic fungus *Mrakia psychrophila*. *G3: Genes|Genomes|Genetics* **6**:3603–3613 DOI [10.1534/g3.116.033308](https://doi.org/10.1534/g3.116.033308).

- Summerell BA, Laurence MH, Liew ECY, Leslie JF. 2010.** Biogeography and phylogeography of *Fusarium*: a review. *Fungal Diversity* **44**:3–13 DOI [10.1007/s13225-010-0060-2](https://doi.org/10.1007/s13225-010-0060-2).
- Suryanarayanan TS. 2012.** Fungal endosymbionts of seaweeds. In: Raghukumar C, ed. *Biology of marine fungi*. Berlin: Springer-Verlag Berlin Heidelberg, 53–70.
- Tello-Cetina JA, Rodríguez-Gil LA, Rodríguez-Romero F. 2005.** Genética poblacional del caracol rosado *Strombus gigas* en la Península de Yucatán: implicaciones para su manejo y pesquería. *Ciencias Marinas* **31**:379–386 DOI [10.7773/cm.v31i2.56](https://doi.org/10.7773/cm.v31i2.56).
- Theile S. 2001.** Queen conch fisheries and their management in the Caribbean. TRAFFIC Europe report for the CITES Secretariat Report. Brussels, Belgium. Available at http://www.trafficj.org/publication/02_Queen_Conch.pdf.
- Thomas-Hall SR, Turchetti B, Buzzini P, Branda E, Boekhout T, Theelen B, Watson K. 2010.** Cold-adapted yeasts from Antarctica and the Italian Alps—description of three novel species: *Mrakia robertii* sp. nov. *Mrakia blollopis* sp. nov. and *Mrakiella niccombsii* sp. nov. *Extremophiles* **14**:47–59 DOI [10.1007/s00792-009-0286-7](https://doi.org/10.1007/s00792-009-0286-7).
- Tsuji M, Fujiu S, Xiao N, Hanada Y, Kudoh S, Kondo H, Tsuda S, Hoshino T. 2013a.** Cold adaptation of fungi obtained from soil and lake sediment in the Skarvsnes ice-free area, Antarctica. *FEMS Microbiology Letters* **346**:121–130 DOI [10.1111/1574-6968.12217](https://doi.org/10.1111/1574-6968.12217).
- Tsuji M, Singh SM, Yokota Y, Kudoh S, Hoshino T. 2013b.** Influence of initial pH on ethanol production by the antarctic basidiomycetous yeast *Mrakia blollopis*. *Bioscience, Biotechnology and Biochemistry* **77**:2483–2485 DOI [10.1271/bbb.130497](https://doi.org/10.1271/bbb.130497).
- Turchetti B, Buzzini P, Goretti M, Branda E, Diolaiuti G, D’Agata C, Smiraglia C, Vaughan-Martini A. 2008.** Psychrophilic yeasts in glacial environments of Alpine glaciers. *FEMS Microbiology Ecology* **63**:73–83 DOI [10.1111/j.1574-6941.2007.00409.x](https://doi.org/10.1111/j.1574-6941.2007.00409.x).
- Turner BC, Perkins DD, Fairfield A. 2001.** *Neurospora* from natural populations: a global study. *Fungal Genetics and Biology* **32**:67–92 DOI [10.1006/fgbi.2001.1247](https://doi.org/10.1006/fgbi.2001.1247).
- Volland JM, Gros O, Frenkiel L, Aldana D. 2010.** Apicomplexan parasite in the digestive gland of various species of the family Strombidae: *Strombus costatus* S. *gigas*, and *S. pugilis*. Gulf and Caribbean Fisheries Institute, 430–432.
- Wang G, Li Q, Zhu P. 2008.** Phylogenetic diversity of culturable fungi associated with the Hawaiian Sponges *Suberites zeteki* and *Gelliodes fibrosa*. *Antonie Van Leeuwenhoek* **93**:163–174 DOI [10.1007/s10482-007-9190-2](https://doi.org/10.1007/s10482-007-9190-2).
- Werding B. 1969.** Morphologie, Entwicklung und Ökologie digener Trematoden-Larven der Strandschnecke *Littorina littorea*. *Marine Biology* **3**:306–333 DOI [10.1007/BF00698861](https://doi.org/10.1007/BF00698861).
- Wu D, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova NN, Kunin V, Goodwin L, Wu M, Tindall BJ, Hooper SD, Pati A, Lykidis A, Spring S, Anderson IJ, D’haeseleer P, Zemla A, Singer M, Lapidus A, Nolan M, Copeland A, Han C, Chen F, Cheng J-F, Lucas S, Kerfeld C, Lang E, Gronow S, Chain P, Bruce D, Rubin EM, Kyrpides NC, Klenk H-P, Eisen JA. 2009.** A phylogeny-driven genomic encyclopaedia of Bacteria and Archaea. *Nature* **462**:1056–1060 DOI [10.1038/nature08656](https://doi.org/10.1038/nature08656).

- Zamora-Bustillos R, Rodríguez-Canul R, García de León FJ, Tello-Cetina J. 2011.** Diversidad genética de dos poblaciones del caracol *Strombus gigas* (Gastropoda: Strombidae) en Yucatán, México, con microsatélite. *Revista de Biología Tropical* **59**:1127–1134.
- Zhang X, Hua M, Song C, Chi Z. 2012.** Occurrence and diversity of marine yeasts in Antarctica environments. *Journal of Ocean University of China* **11**:70–74
[DOI 10.1007/s11802-012-1820-2](https://doi.org/10.1007/s11802-012-1820-2).