

Influence of whole-wheat consumption on fecal microbial community structure of obese diabetic mice

Jose F. Garcia-Mazcorro^{1,2}, Ivan Ivanov³, David A. Mills⁴ and Giuliana Noratto^{5,#}

¹ Faculty of Veterinary Medicine, Universidad Autónoma de Nuevo León, General Escobedo, Nuevo Leon, Mexico

² Research Group Medical Eco-Biology, Universidad Autónoma de Nuevo León, General Escobedo, Nuevo Leon, Mexico

³ Veterinary Physiology and Pharmacology, Texas A&M University, College Station, Texas, United States

⁴ Department of Food Science and Technology, University of California, Davis, Davis, California, United States

⁵ School of Food Science, Washington State University, Pullman, Washington, United States

Current Address: Nutrition and Food Science, Texas A&M University, College Station, Texas, United States

ABSTRACT

The digestive tract of mammals and other animals is colonized by trillions of metabolically-active microorganisms. Changes in the gut microbiota have been associated with obesity in both humans and laboratory animals. Dietary modifications can often modulate the obese gut microbial ecosystem towards a more healthy state. This phenomenon should preferably be studied using dietary ingredients that are relevant to human nutrition. This study was designed to evaluate the influence of whole-wheat, a food ingredient with several beneficial properties, on gut microorganisms of obese diabetic mice. Diabetic (db/db) mice were fed standard (obese-control) or whole-wheat isocaloric diets (WW group) for eight weeks; non-obese mice were used as control (lean-control). High-throughput sequencing using the MiSeq platform coupled with freely-available computational tools and quantitative real-time PCR were used to analyze fecal bacterial 16S rRNA gene sequences. Short-chain fatty acids were measured in caecal contents using quantitative high-performance liquid chromatography photo-diode array analysis. Results showed no statistical difference in final body weights between the obese-control and the WW group. The bacterial richness (number of Operational Taxonomic Units) did not differ among the treatment groups. The abundance of Ruminococcaceae, a family containing several butyrate-producing bacteria, was found to be higher in obese (median: 6.9%) and WW-supplemented mice (5.6%) compared to lean (2.7%, $p = 0.02$, Kruskal-Wallis test). Caecal concentrations of butyrate were higher in obese (average: 2.91 mmol/mg of feces) but especially in WW-supplemented mice (4.27 mmol/mg) compared to lean controls (0.97 mmol/mg), while caecal succinic acid was lower in the WW group compared to obese but especially to the lean group. WW consumption was associated with ~3 times higher abundances of *Lactobacillus* spp. compared to both obese and lean control mice. Analysis of weighted UniFrac distances revealed a distinctive clustering of lean microbial communities separately from both obese and

Submitted 28 October 2015
Accepted 27 January 2016
Published 15 February 2016

Corresponding authors
Jose F. Garcia-Mazcorro,
josegarcia_mex@hotmail.com
Giuliana Noratto,
gnoratto@tamu.edu

Academic editor
Adina Howe

Additional Information and
Declarations can be found on
page 16

DOI 10.7717/peerj.1702

© Copyright
2016 Garcia-Mazcorro et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

WW-supplemented mice ($p = 0.001$, ANOSIM test). Predictive metagenome analysis revealed significant differences in several metabolic features of the microbiota among the treatment groups, including carbohydrate, amino acids and vitamin metabolism ($p < 0.01$, Kruskal-Wallis test). However, obese and WW groups tended to share more similar abundances of gene families compared to lean mice. Using an in vivo model of obesity and diabetes, this study suggests that daily WW supplementation for eight weeks may not be enough to influence body weight or to output a lean-like microbiome, both taxonomically and metabolically. However, WW-supplementation was associated with several statistically significant differences in the gut microbiome compared to obese controls that deserve further investigation.

Subjects Bioinformatics, Food science and technology, Microbiology, Nutrition, Metabolic sciences

Keywords Fecal microbiota, High-throughput sequencing, Metabolic pathways, Obesity, Whole-wheat

INTRODUCTION

Obesity is an epidemic with catastrophic consequences for the health of millions of people around the globe. Different strategies can help reduce body weight including changes in exercise and dietary habits, yet many patients genuinely struggle to successfully decrease their body weight due to multiple interrelated factors ([Gupta, 2014](#)).

The mammalian digestive tract is a complex organ that has been constantly co-evolving with trillions of microorganisms (the gut microbiota) to combat pathogens and maximize food digestion for at least 600 million years. Despite its general resilience, the gut microbiota is still susceptible to changes in dietary and other life habits, some of which can lead to imbalances and consequently to disease ([Lozupone et al., 2012](#)). For instance, substantial evidence has been published showing an association between obesity and changes in gut microbial populations and its metabolism of dietary and endogenous compounds ([Delzenne & Cani, 2011](#)). Interestingly, the changes in gut microbial communities between lean and obese individuals are not irreversible ([Turnbaugh et al., 2008](#)) with diet being the most practical alternative to reestablish microbial equilibrium within the gut. Understanding changes in gut microorganisms in response to dietary modifications is essential to develop effective dietary strategies to help obese patients.

Growing evidence shows that the consumption of specific dietary ingredients or supplements such as probiotics, prebiotics, polyphenols, as well as whole-grains has the potential of modifying gut health parameters in obese individuals, both in humans and animal models ([Katcher et al., 2008](#); [Noratto et al., 2014](#); [Petschow et al., 2013](#); [Vitaglione et al., 2015](#)). Whole-Wheat (WW) is often recommended by medical nutritionists as part of a healthy diet for both overweighted and lean individuals. While several investigations have previously addressed the nutritional benefits of consuming WW ([Stevenson et al., 2012](#)), very few studies have researched the potential of either WW or its

individual nutrients to alter the gut microbiota of lean or obese individuals ([Neyrinck et al., 2011](#)) or as part of dietary management to treat obesity. One study investigated the effect of replacing refined wheat with whole-grain wheat for 12 weeks on body weight and fat mass in overweighted women ([Kristensen et al., 2011](#)). This short-period of 12 weeks was enough to significantly reduce percentage fat mass but no body weights. Here we show that an 8-week consumption period of an isocaloric WW diet did not significantly change body weights in obese-diabetic mice. Overall, obese mice under WW-supplemented diet showed similarities to obese controls with regards to gut microbial composition and predicted metabolic profile. The effect of WW was mostly observed on caecal concentrations of butyrate and succinate and a few bacterial groups such as *Lactobacillus*. The results may have implications in clinical dietary management of obesity using WW.

METHODS

Study design

The Institutional Animal Care Use Committee from Washington State University approved all experimental procedures (animal protocol approval number: 04436-001). Two strains of male mice were used in this study, BKS.Cg- + *Lepr^{db}/+Lepr^{db}/OlaHsd* obese diabetic (db/db), and lean BKS.Cg-*Dock7m +/+ Lepr^{db}/OlaHsd* (Harlan Laboratories, Kent, WA). Animals were purchased at 5–6 weeks of age and maintained in ventilated rack system with food and water provided *ad libitum* throughout the study. We received 11 mice for the lean group and 10 mice from all other groups. After 7 days of acclimatization, obese mice were randomly divided into two groups (n = 10 each) namely obese (AIN-93 G Purified Rodent Diet) and WW (whole-wheat supplemented diet). The wild type mice group (n = 11) was named lean (AIN-93 Diet). Diets were made by Dyets Inc. (Bethlehem, PA) ([Table 1](#)). Four or five mice per cage were housed in an environment-controlled room (23 °C, 12 hours dark-light cycle). All mice were visually inspected every day and body weight was recorded from all animals once a week.

Fecal collection and DNA extraction

Fresh distal colon contents (see qPCR analysis below) and fecal samples were obtained from all mice at the end of the study (8 weeks) and stored at –80 °C prior to DNA and 16S rRNA gene profiling analysis. Total DNA was extracted from at least two different fecal pellets weighting approximately 200 mg. Following bead-beating, the QIAamp DNA Stool Mini Kit (Qiagen Inc., Valencia, CA, USA) was used for DNA extraction following the manufacturer's instructions. DNA concentration and purity was determined using a NanoDrop Spectrophotometer (Thermo Scientific, Wilmington, DE, USA) and diluted to a working concentration of 5 ng/μL.

High-throughput sequencing of 16S rRNA genes

Amplification and sequencing were performed as described elsewhere ([Bokulich et al., 2014](#)). Briefly, the V4 semi-conserved region of bacterial 16S rRNA genes was amplified using primers F515 (5'-GTGCCAGCMGCCGCGGTAA-3') and R806

Table 1 Formulation of experimental diets (g/100 g).

Ingredients	Lean and obese diet	Wheat diet
Casein, high nitrogen	20	0.0
L-Cysteine	0.3	0.3
Whole-wheat meal	0.0	87.94
Soybean oil	7.0	7.0
Sucrose	10	0.0
Cornstarch	39.74	0.0
Dyetrose	13.2	0.0
t-Butylhydroquinone	0.0014	0.0014
Cellulose	5	0.0
Mineral mix #210025	3.5	3.5
Vitamin mix #310025	1.0	1.0
Choline bitartrate	0.25	0.25
Kcal/100 g	376.00	387.76

(5'-GGACTACHVGGGTWTCTAAT-3'), with the forward primer modified to contain a unique 8-nt barcode and a 2-nt linker sequence at the 5' terminus. Amplicons were combined into one pooled sample and submitted to the University of California Davis Genome Center DNA Technologies Core for Illumina paired-end library preparation, cluster generation, and 250-bp paired-end sequencing on an Illumina MiSeq instrument in one runs. For data analysis, raw Illumina fastq files were demultiplexed, quality filtered, and analyzed using the freely available Quantitative Insights into Microbial Ecology (QIIME) Virtual Box v.1.8.0 ([Caporaso et al., 2010](#)). Operational Taxonomic Units (OTUs) were assigned using two different approaches: first, using UCLUST v.1.2.22 ([Edgar, 2010](#)) as implemented in QIIME using the open-reference clustering algorithm described in ([Rideout et al., 2014](#)) for alpha and beta diversity analyses; and second, using the `pick_closed_reference_otus.py` QIIME script for further analysis using PICRUSt (see Predicted metabolic profiles below). The Greengenes 13_5 97% OTU representative 16S rRNA gene sequences was used as the reference sequence collection ([DeSantis et al., 2006](#)). Alfa and beta diversity analyses were performed using 3000 random sequences per sample (lowest number of sequences in a sample after demultiplexing, filtering and OTU picking). Raw sequences were uploaded into the Sequence Read Archive at NCBI (accession number: [PRJNA281761](#)). The `trim.seqs` command in MOTHUR ([Schloss et al., 2009](#)) was used for splitting original fastq files per sample for uploading to SRA.

Predicted metabolic profiles

OTUs from the `closed_reference` script were normalized and used to predict metagenome functional content using the online Galaxy version of Phylogenetic Investigation of Communities by Reconstruction of Unobserved States (PICRUSt) ([Langille et al., 2013](#)). PICRUSt uses existing annotations of gene content as well as 16S copy numbers from reference microbial genomes in the IMG database ([Markowitz et al., 2012](#)) and a

functional classification scheme to catalogue the predicted metagenome content. The current galaxy version supports three types of functional predictions; this current study used the popular KEGG Orthologs ([Kanehisa et al., 2012](#)).

Quantitative real-time PCR (qPCR) analysis

DNA was extracted from distal colon content using the ZR Fecal DNA MiniPrep™ kit following the manufacturer's protocol (Zymo Research, Irvine, CA, USA). qPCR was used to detect specific bacterial groups as described elsewhere ([Noratto et al., 2014](#)). [Table 2](#) shows the primers sequences used for all qPCR analyses.

Measurement of Short Chain Fatty Acids (SCFAs) in caecal contents

SCFAs were quantified as reported elsewhere ([Campos et al., 2012](#)). Briefly, samples were analyzed by an HPLC-PDA system using an Aminex HPX-87H strong cation-exchange resin column (300 × 7.8 mm) and fitted with an ion exchange microguard refill cartridge (Bio-Rad, Hercules, CA, USA). The HPLC-PDA system consisted of a Water 2695 Separation Module (Waters, Milford, MA), which was equipped with a Water 2996 Photodiode Array detector (PDA). Samples (20 μL) were eluted isocratically with 5 mM sulfuric acid at 0.6 mL/min, and the column temperature was held at 50 °C. Sodium butyrate, acetic acid, oxalic acid, and succinic acid were identified and quantified by comparing retention time and UV-Visible spectral data to standards.

Statistical analysis

ANOVA and the non-parametric alternative Kruskal-Wallis test were used to analyze final body weights and SCFAs concentrations, respectively. Multiple-comparisons were performed using Tukey and Mann-Whitney tests. The Bonferroni (for Mann-Whitney tests) and False Discovery Rate (for Tukey's tests) corrections were used to adjust for multiple comparisons. Analysis of Similarities (ANOSIM) was used to test for clustering of microbial communities using weighted and unweighted UniFrac distance matrices. QIIME v.1.8.0, R v.3.0.3 ([R Core Team, 2014](#)), PAST ([Hammer, Harper & Ryan, 2001](#)) and Excel were used for statistics and graphics. The Linear Discriminant Analysis (LDA) Effect Size (LEfSe) method was used to assess differences in microbial communities using a LDA score threshold of 3 ([Segata et al., 2011](#)). STAMP ([Parks & Beiko, 2010](#)) was used to visualize and analyze the PICRUSt data with ANOVA and False Discovery Rate. Unless otherwise noted, an alpha of 0.05 was considered to reject null hypotheses.

RESULTS

One mouse in the obese group died for reasons unrelated to this study. At the end of the study, there was a significant ($p < 0.01$, ANOVA) difference in body weight between the lean (average: 30.6 ± 2.2 g) and both the obese (46.1 ± 2.8 g) and WW groups (45.3 ± 5.8 g). WW consumption was not associated with a lower body weight compared to obese control group ($p = 0.96$, Tukey's test).

Table 2 Oligonucleotides used in this study for qPCR analyses.

qPCR primers	Sequence (5'-3')	Target	Reference
HDA1	ACTCCTACGGGAGGCAGCAGT	All bacteria (V2–V3 regions, position 339–539 in the <i>E. coli</i> 16S gene)	<i>Walter et al. (2000)</i>
HDA2	GTATTACCGCGGCTGCTGGCAC		
Bact834F	GGARCATGTGGTTTAATTCGATGAT	Bacteroidetes (Phylum)	<i>Guo et al. (2008)</i>
Bact1060R	AGCTGACGACAACCATGCAG		
928F-Firm	TGAAACTYAAAGGAATTGACG	Firmicutes (Phylum)	<i>Bacchetti et al. (2011)</i>
1040firmR	ACCATGCACCACCTGTC		
BifF	GCGTGCTTAACACATGCAAGTC	<i>Bifidobacterium</i> (genus)	<i>Penders et al. (2005)</i>
BifR	CACCCGTTTCCAGGAGCTATT		
<i>E. coli</i> F	CATGCCGCGTGTATGAAGAA	<i>E. coli</i>	<i>Huijsdens et al. (2002)</i>
<i>E. coli</i> R	CGGGTAAACGTCAATGAGCAA		
TuriciF	CAGACGGGGACAACGATTGGA	<i>Turibacter</i> (genus)	<i>Suchodolski et al. (2012)</i>
TuriciR	TACGCATCGTCGCCTTGGA		
RumiF	ACTGAGAGGTTGAACGGCCA	Ruminococcaceae (family)	<i>Garcia-Mazcorro et al. (2012)</i>
RumiR	CCTTTACACCCAGTAAWTCCGGA		
FaecaliF	GAAGGCGGCCTACTGGGCAC	<i>Faecalibacterium</i> (genus)	<i>Garcia-Mazcorro et al. (2012)</i>
FaecaliR	GTGCAGGCGAGTTGCAGCCT		
Eco1457-F	CATTGACGTTACCCGCAGAAGAAGC	Enterobacteriaceae (family)	<i>Bartosch et al. (2004)</i>
Eco1652-R	CTTACGAGACTCAAGCTTGC		
V1F	CAGCACGTGAAGGTGGGGAC	<i>Akkermansia muciniphila</i>	<i>Collado et al. (2007)</i>
V1R	CCTTGCGGTTGGCTTCAGAT		
PrevF	CACCAAGGCGACGATCA	<i>Prevotella</i> (genus)	<i>Larsen et al. (2010)</i>
PrevR	GGATAACGCCYGGACCT		
Bfr-F	CTGAACCAGCCAAGTAGCG	<i>Bacteroides fragilis</i>	<i>Liu et al. (2003)</i>
Bfr-R	CCGCAAACCTTTCACAACCTGACTTA		

Fecal microbiota composition

A total of 8686 different OTUs were detected using the open reference algorithm described by [Rideout et al. \(2014\)](#). On the other hand, the closed_reference method used to generate data for PICRUSt (see PICRUSt below) only yielded 1302 OTUs. Fecal microbial composition of all mice was mostly comprised by Firmicutes (average: 58.7% across all samples) and Bacteroidetes (average: 32.8%) ([Fig. 1](#)). Other less abundant Phyla were Actinobacteria (~4%), Proteobacteria (~3%), Verrucomicrobia (~0.8%) and others ([Fig. 1](#)). There was no statistical difference in relative abundance of the two most abundant phyla (Firmicutes and Bacteroidetes), partly because of the high variability among individual mice. The ratio Bacteroidetes/Firmicutes was lower in the lean (median: 38.7%) compared to the obese group (median: 85.3%) and the WW group (median: 75.4%) but this difference did not reach significance ($p = 0.12$, Kruskal-Wallis). Several statistical differences were found in low abundant phyla. Actinobacteria and Verrucomicrobia were higher in lean, Cyanobacteria, TM7 and Tenericutes were higher in the WW group, and Deferribacteres was higher in obese-control ($p < 0.01$, Kruskal-Wallis, [Fig. 1](#)).




Figure 1 Box plots. Composition of fecal microbiota at the phylum level in the Lean (n = 11), obese control (Obese, n = 9) and whole-wheat (Wheat, n = 10) group. Boxes represent the 25–75 quartiles, the median is shown with a vertical line inside the box. Values outside 1.5 times the box height are shown as circles; values outside 3 times the box height are shown as stars. The differences in relative abundance of Firmicutes, Bacteroidetes and Proteobacteria did not reach statistical significance (see main text for details).

LEfSe showed statistical significant differences for several microbial groups at lower taxonomic levels (Fig. 2). Among the bacterial groups that showed differences indicating an effect of WW-supplementation include the genus *Lactobacillus*, the class Gammaproteobacteria and the controversial S24-7 family <http://groups.google.com/forum/#!topic/qiime-forum/Ds75aZoVrFY> (Fig. 2). Other differences in bacterial abundances suggested that WW-supplementation did not generate a lean-like microbiome. For example, the genera *Bifidobacterium*, *Allobaculum* and *Akkermansia* were higher in lean compared to both obese and WW group (Fig. 2). Also, the family Ruminococcaceae was more similar between obese (median: 6.9%) and WW-supplemented mice (5.6%) compared to lean (2.7%). Despite these differences, overall the fecal microbial composition of obese-control and WW mice was more similar to each other compared to lean although WW-supplementation yielded a unique pattern of bacterial abundances that did not necessarily cluster together with all obese samples (Fig. 3).

Alpha diversity

There was no significant difference in number of OTUs and Chao1 diversity index. Interestingly, samples from the obese group showed a more disperse distribution of




Figure 2 Bar charts of most significant results using the LDA Effect Size method (LefSe). LefSe identifies those bacterial groups that showed statistical significance effect size and associate them with the class (in this study treatment group) with the highest median. (A) lean; (B) obese-control; (C) WW-supplemented obese mice. Dotted lines represent medians; straight lines represent averages.

OTUs (Fig. 4). Rarefied plots of number of OTUs showed that more than the 3000 sequences per sample used in this study are needed to fully describe the fecal microbiota of all mice.

Beta-diversity

Principal Coordinate Analysis (PCoA) of weighted and unweighted UniFrac metrics showed different clustering of microbial communities. Weighted (which takes phylogenetic information as well as sequence abundance into account) metrics clearly showed a different microbial structure in lean individuals compared to obese and WW groups (ANOSIM, $p = 0.001$) (Fig. 5). This was expected based on the clustering of lean subjects using relative abundance of sequence reads (Fig. 3). On the other hand, the qualitative (does not take sequence abundance into account) unweighted UniFrac analysis shows that the microbiota of the WW group clustered separately from the lean and obese groups (ANOSIM, $p = 0.001$) (Fig. 5).




Figure 3 Heatmap. Heatmap of relative abundance of the most abundant bacterial taxa at the family level (x axis, ordered by abundance) in Lean (n = 11), Obese (n = 9) and Wheat (n = 10) group. This figure shows that lean subjects clustered separately from obese and WW subjects. Clustering was performed using Bray-Curtis distances in R v.3.2.2.

Predicted metabolic profile

The taxa predicted by 16S RNA marker gene sequencing was used to predict the functional profile of the fecal microbiome in all three experimental groups. Using a $p < 0.01$ for ANOVA tests in STAMP, several statistical differences were found (Table 3). Overall, obese and WW groups tended to share more similar abundances of gene families compared to lean mice, an observation that supports the differences in bacterial abundances.

qPCR assessment of microbiota in distal colon contents

We performed qPCR analysis for bacterial groups of interest to health in distal colon contents. Similarly to the sequencing results from fecal samples, qPCR results revealed several differences in relative abundance for different bacterial groups (Fig. 6).




Figure 4 Dispersion in number of OTUs detected. (A) shows the relationship between the numbers of OTUs and the proportion of samples containing those OTUs for each treatment group. These plots show that more obese control samples contained higher numbers of OTUs compared to whole-wheat. Lines were used to illustrate 50% of the samples (vertical line) and 1000 OTUs (horizontal line). (B) shows boxplots to illustrate the distributions of the number of OTUs for each treatment group (boxes represent the 25–75 quartiles, the median is shown with a horizontal line inside the box).

SCFAs caecal concentrations

There was a statistically significant difference among the treatment groups for several SCFAs in caecal contents (Table 4). Butyrate concentrations were higher in the WW group compared to both the lean and the obese group ($p < 0.001$, Kruskal-Wallis). Also, WW consumption was associated with lower succinic acid concentrations ($p = 0.009$, Kruskal-Wallis).

DISCUSSION

Obesity is a worldwide epidemic disease that has been associated with changes in the gut microbiome in many different studies. Consumption of whole grains is often recommended by medical nutritionists as part of a healthy diet. To our knowledge, this is




Figure 5 Principal Coordinates Analysis plots. Principal Coordinates Analysis (PCoA) plots of weighted (A) and unweighted (B) UniFrac distance metrics. Please note that each plot gives contrasting results with regards to the clustering of samples.

the first study evaluating the in vivo effect of WW consumption on fecal bacterial community structure of obese diabetic mice, adding valuable information to the literature with regard to the use and development of dietary strategies to help obese patients.

Ley et al. (2005) showed that lean mice have more Bacteroidetes and less Firmicutes compared to obese mice, a finding that has been reported by several other research groups. However, it is important to note that these observations were division-wide (in other words, there was no specific subgroup such as families or genera that were present high or low in abundance) and, more importantly, that other researchers have found either no difference in Firmicutes and Bacteroidetes between obese and lean (*Duncan et al., 2008*) or more Bacteroidetes in obese compared to normal-weight individuals (*Zhang et al., 2009*). Interestingly, in this study sequencing showed no statistical difference in the abundance of both phyla Firmicutes and Bacteroidetes between lean and obese control; nonetheless, two important aspects must be taken into account. First, obese and WW mice were consistently more like each other compared to lean mice with regard to the abundance of both phyla. Also, contrary to the observations by *Ley et al. (2005)*, lean mice had more Firmicutes and less Bacteroidetes compared to both obese and WW mice, a difference that did not reach statistical significance. qPCR confirmed the sequencing results about the

Table 3 Metabolic features in Lean (n = 11), Obese (n = 9) and whole-wheat supplemented (n = 10) mice.

KEGG gene categories			Treatment Groups			p value
Level_1	Level_2	Level_3	Lean Mean ± st. dev.	Obese Mean ± st. dev.	Whole-wheat Mean ± st. dev.	
Cellular processes	Transport and catabolism	Peroxisome	0.15 ± 0.03 ^a	0.18 ± 0.04 ^{a,b}	0.23 ± 0.03 ^c	0.001
Environmental information processing	Signal transduction	Phosphatidylinositol signaling system	0.11 ± 0.01 ^a	0.08 ± 0.01 ^b	0.10 ± 0.01 ^{a,c}	0.002
	Signaling molecules and interaction	Ion channels	0.05 ± 0.01 ^a	0.02 ± 0.01 ^b	0.02 ± 0.01 ^{b,c}	<0.001
Genetic information processing	Replication and repair	Base excision repair	0.50 ± 0.06 ^a	0.41 ± 0.03 ^b	0.41 ± 0.04 ^{b,c}	0.001
Human diseases	Infectious diseases	Tuberculosis	0.18 ± 0.02 ^a	0.13 ± 0.01 ^b	0.13 ± 0.02 ^{b,c}	0.001
	Cancers	Pathways in cancer	0.07 ± 0.01 ^a	0.04 ± 0.00 ^b	0.05 ± 0.00 ^{b,c}	<0.001
	Neurodegenerative diseases	Amyotrophic Lateral Sclerosis (ALS)	0.02 ± 0.01 ^a	0.03 ± 0.01 ^b	0.04 ± 0.01 ^{b,c}	<0.001
	Cancers	Renal cell carcinoma	0.03 ± 0.01 ^a	0.01 ± 0.00 ^b	0.01 ± 0.01 ^{b,c}	<0.001
Metabolism	Carbohydrate metabolism	Fructose and mannose metabolism	1.16 ± 0.19 ^a	0.94 ± 0.08 ^b	0.86 ± 0.12 ^{b,c}	0.005
		Pentose phosphate pathway	0.92 ± 0.05 ^a	0.86 ± 0.07 ^{a,b}	0.78 ± 0.06 ^c	0.002
	Metabolism of cofactors and vitamins	Porphyrin and chlorophyll metabolism	0.55 ± 0.15 ^a	0.85 ± 0.13 ^b	0.66 ± 0.14 ^{a,c}	0.009
		Vitamin B6 metabolism	0.17 ± 0.02 ^a	0.20 ± 0.02 ^b	0.22 ± 0.01 ^{b,c}	<0.001
	Metabolism of other amino acids	Selenocompound metabolism	0.40 ± 0.02 ^a	0.37 ± 0.01 ^b	0.35 ± 0.01 ^{b,c}	<0.001
		Cyanoamino acid metabolism	0.23 ± 0.04 ^a	0.33 ± 0.02 ^b	0.30 ± 0.06 ^{b,c}	0.002
		Beta-alanine metabolism	0.18 ± 0.04 ^a	0.22 ± 0.04	0.28 ± 0.06 ^b	0.005
	Carbohydrate metabolism	C5-branched dibasic acid metabolism	0.23 ± 0.06 ^a	0.32 ± 0.02 ^b	0.31 ± 0.02 ^{b,c}	0.003
	Biosynthesis of other secondary metabolites	Phenylpropanoid biosynthesis	0.12 ± 0.03 ^a	0.20 ± 0.02 ^b	0.17 ± 0.05	0.003
	Metabolism of terpenoids and polyketides	Biosynthesis of ansamycins	0.14 ± 0.04 ^a	0.11 ± 0.02	0.09 ± 0.02 ^b	0.006
		Flavonoid biosynthesis	0.02 ± 0.01 ^a	0.01 ± 0.00 ^b	0.00 ± 0.00 ^{b,c}	<0.001
	Biosynthesis of other secondary metabolites	Stilbenoid, diarylheptanoid and gingerol biosynthesis	0.01 ± 0.01 ^a	0.00 ± 0.00 ^b	0.00 ± 0.00 ^{b,c}	0.007

Note:

This table only shows features that reached statistical significant differences (ANOVA, $p < 0.1$ adjusted for False Discovery Rate in STAMP). Different letters^(a,b,c) indicate statistical significant difference (Tukey's post-hoc tests, $p < 0.05$).

abundance of Firmicutes but not Bacteroidetes, maybe due to the use of fecal (sequencing) or colon (qPCR) contents for bacterial analysis. Regardless, differences in taxa abundance at the phylum level have little relevance when considering all their individual groups within. For instance, many bacterial groups at lower taxonomic levels deserve attention, like the mucin-degrader *Akkermansia* which has been shown to be




Figure 6 Boxplots. Quantitative real-time PCR (qPCR) results for selected bacterial groups. Results are expressed as relative abundance of 16S rRNA amplified DNA (all results were normalized to qPCR data for total bacteria). * $p < 0.05$ against lean; # $p < 0.05$ against Whole-Wheat (WW).

Table 4 Median (minimum-maximum) for all Short-Chain Fatty Acids (SCFAs). Results are expressed in mmol/mg of caecal contents.

SCFA	Lean	Obese	Whole-wheat	<i>p</i> value
Sodium butyrate	0.97 (0.15–2.65) ^a	2.91 (1.47–4.35) ^b	4.27 (3.05–6.26) ^{b,c}	<0.001
Acetic acid	10.2 (7.7–26.3)	12.0 (8.3–18.7)	15.4 (10.1–31.9)	0.208
Oxalic acid	15.14(6.68–18.91) ^a	14.60 (8.78–28.01)	9.96 (6.76–12.15) ^b	0.033
Succinic acid	39.84 (15.29–97.63) ^a	22.97 (3.86–71.18) ^{a,b}	3.12 (0.91–63.36) ^c	0.009

Note:

Different letters^(a,b,c) denote statistical significance. *p* values come from the Kruskal-Wallis test and multiple comparisons were performed using the Mann-Whitney test and corrected with the Bonferroni method.

inversely correlated with body weight in rodents and humans (Everard *et al.*, 2013). Accordingly, both sequencing and qPCR in this current study showed that obese mice had fewer *Akkermansia* and WW consumption surprisingly helped to decrease its abundance even further. Here it is important to note that a higher abundance in feces does not necessarily imply a higher abundance in the mucus. WW consumption was also associated with much more *Lactobacillus* spp., a bacterial genus frequently used in probiotic formulations, and the genus *Allobaculum* was practically absent in both obese and WW groups while lean individuals were heavily colonized by this group. These changes in bacterial abundances deserve more investigation.

Beta diversity metrics are useful to study similarities of microbiomes, which in turn have critical consequences for understanding health and disease processes. Lozupone *et al.* (2007) explains that quantitative beta-diversity measures (weighted UniFrac distances) are better for revealing community differences that are due to changes in relative taxon (OTUs) abundance, while qualitative (unweighted) are most informative

when communities differ by what can live in them. Most studies report either weighted or unweighted but few report both. In this study, weighted analysis showed a clear separation of lean samples from all samples from the obese and WW groups, suggesting that the numbers of OTUs are an important determinant to separate lean microbiomes from obese individuals with and without WW. In this study, the results of weighted analysis also show that animal genetics was the predominant factor to separate microbial communities. On the other hand, unweighted analysis showed opposite results: lean and obese samples clustered separately from all samples of the WW group, suggesting that WW helped create an environment that favored a phylogenetically different ecosystem. Importantly, the variation explained by the axes is much lower when using unweighted UniFrac. At this point, both methods should be considered for explaining the changes in gut microbiomes in investigations like this study ([Lozupone et al., 2007](#)). The discrepancy between the results of weighted and unweighted results suggests that an 8-week period of WW consumption helped change the overall environment in the intestinal lumen, thus modulating what can live and proliferate in it (unweighted results). Thus, the different environment could promote changes in the abundance of specific taxa (weighted results), as shown in this study for several bacterial groups. Given that the assessment of microbial diversity is a major component in microbial ecological studies and closely relates to our understanding of health and health deviations, we expect others to start inspecting and reporting both weighted and unweighted UniFrac distance metrics. The use of both metrics has been shown to be useful in various investigations ([Campbell et al., 2015](#); [Igarashi et al., 2014](#); [Wu et al., 2010](#)).

Microbial butyrate is essential for colon health and lower concentrations of this fatty acid are usually considered non-optimal for gut health ([Donohoe et al., 2011](#)). Nonetheless, studies have shown that obese individuals actually have higher fecal butyrate and other SCFAs compared to lean individuals ([Fernandes et al., 2014](#)), an observation that suggests that both lower and higher butyrate concentrations than normal may be associated with and perhaps aggravate disease. Similarly, obese mice in this current study (with and without WW supplementation) had higher butyrate concentrations in caecal contents compared to lean mice. Butyrate-producers are abundant in the mammalian gut and mainly belong to the family Ruminococcaceae within the Firmicutes ([Louis & Flint, 2009](#)). In this study both sequencing and qPCR revealed higher fecal Ruminococcaceae in obese and WW groups compared to lean individuals, thus potentially explaining the higher caecal butyrate concentrations. Another SCFA that deserves attention is succinic acid, which has been shown to increase in rats fed a high-fat diet ([Jakobsdottir et al., 2013](#)). In this current study, obese mice had lower concentrations of succinic acid and WW-supplementation seemingly helped to drastically decrease it. Unfortunately, far more attention has been paid to butyrate compared to succinate, propionate and other SCFA ([Cheng et al., 2013](#); [Reichardt et al., 2014](#)).

The assessment of microbial metabolic activity in complex ecosystems is hampered in part by the huge number of microorganisms and the cost of sequencing either whole genomes or transcriptomes. PICRUSt allows a prediction of the metabolic profile using

taxa predicted by 16S rRNA gene sequencing. PICRUSt is, however, not exempt of pitfalls: it only uses information for well-defined 16S sequences and the presence of a given set of genes does not tell anything about their functional activity depending on the specific environmental conditions. Supported by the similarities in abundance of most bacterial groups between obese-control and WW groups, this study showed that 8-week WW consumption was not enough to make a significant difference in the abundance of bacterial gene families.

Caveats

This study was designed to obtain preliminary information about the influence of WW consumption on gut microbial ecology of obese diabetic mice; therefore, we did not aim to determine the exact compound(s) behind the observed effects. Wheat is a fiber-rich grain and consumption of fiber alone is associated with changes in the gut microbiome and the immune system of the host (*Bermudez-Brito et al., 2015*). Aside from fiber, WW also contains other bioactive compounds (e.g. polyphenols) that may be responsible for specific effects on host metabolism, physiology and immune system. For instance, it has been recently shown that wheat-derived alkylresorcinols were capable of showing beneficial effects on diet-induced obese mice (*Oishi et al., 2015*). Interestingly, our group showed that carbohydrate-free polyphenol-rich juice from plum is capable of impeding body weight gain in obese Zucker rats (*Noratto et al., 2014*), a finding that was not observed with WW consumption in mice in this current study. More research is necessary to investigate the separate effect of the different nutrients in WW.

SUMMARY

In summary, this study suggests that an 8-week consumption of whole-wheat may not be enough to exert an effect on body weight and to output a lean-like microbiome using an in vivo model of obesity and diabetes. However, WW-supplementation was associated with several statistically significant changes compared to obese controls that deserve further investigations. These results may or may not apply to obesity in human patients. Also, our experimental scheme was not designed to address the effect of WW supplementation on lean mice; whether the observed changes in the gut microbiome and metabolite concentrations are irrespective of mice phenotype may warrant further research. The clinical relevance of this present work remains to be determined.

Future directions

In humans, obesity is a multifactorial disease that can be partly controlled with dietary modifications. This paper adds valuable information to the current literature with regard to the potential influence of WW consumption on the gut microbiota of obese diabetic mice. However, research is needed to investigate the effect of WW on obese human individuals.

ACKNOWLEDGEMENTS

The authors would like to express our deepest gratitude to the QIIME, Mothur and PICRUSt Developers and Help Forums for the creation of such exceptional

computational tools and all the support provided. The authors would also like to thank Indira Mohanty and Alejandra Mencia for their technical assistance in the analysis of SCFAs and qPCR.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the Washington Gran Commission (Grant number: 3057–4668). JFGM received financial support from CONACYT (Mexico) through the National System of Researchers (SNI, for initials in Spanish) program and PRODEP (Mexico). DAM received support from the Peter J. Shields Endowed Chair. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Washington Gran Commission: 3057–4668.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Jose F. Garcia-Mazcorro analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Ivan Ivanov analyzed the data, wrote the paper, reviewed drafts of the paper.
- David A. Mills performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Giuliana Noratto conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The Institutional Animal Care Use Committee from Washington State University approved all experimental procedures (animal protocol approval number: 04436–001).

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:
NCBI SRA: [PRJNA281761](https://www.ncbi.nlm.nih.gov/sra/PRJNA281761).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1702#supplemental-information>.

REFERENCES

- Bacchetti DG, Aldred N, Clare AS, Burgess JG. 2011. Improvement of phylum- and class-specific primers for real-time PCR quantification of bacterial taxa. *Journal of Microbiological Methods* 86(3):351–356 DOI 10.1016/j.mimet.2011.06.010.
- Bartosch S, Fite A, Macfarlane GT, McMurdo ME. 2004. Characterization of bacterial communities in feces from healthy elderly volunteers and hospitalized elderly patients by using real-time PCR and effects of antibiotic treatment on the fecal microbiota. *Applied Environmental Microbiology* 70(6):3575–3581 DOI 10.1128/AEM.70.6.3575-3581.2004.
- Bermudez-Brito M, Sahasrabudhe NM, Rösch C, Schols HA, Faas MM, de Vos P. 2015. The impact of dietary fibers on dendritic cell responses in vitro is dependent on the differential effects of the fibers on intestinal epithelial cells. *Molecular Nutrition and Food Research* 30(4):698–710 DOI 10.1002/mnfr.201400811.
- Bokulich NA, Thorngate JH, Richardson PM, Mills DA. 2014. Microbial biogeography of wine grapes is conditioned by cultivar, vintage, and climate. *Proceedings of the National Academy of Sciences of the United States of America* 111(1):E139–E148 DOI 10.1073/pnas.1317377110.
- Campbell AM, Fleisher J, Sinigalliano C, White JR, Lopez JV. 2015. Dynamics of marine bacteria community diversity of the coastal waters of the reefs, inlets, and wastewater outfalls of southeast Florida. *MicrobiologyOpen* 4(3):390–408 DOI 10.1002/mbo3.245.
- Campos D, Betalleluz-Pallardel I, Chirinos R, Aguilar-Galvez A, Noratto G, Pedreschi R. 2012. Prebiotic effects of yacon (*Smallanthus sonchifolius* Poepp. & Endl), a source of fructooligosaccharides and phenolic compounds with antioxidant activity. *Food Chemistry* 135(3):1592–1599 DOI 10.1016/j.foodchem.2012.05.088.
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turbnbaugh PJ, Walters WA, Widmann J, Yatsunencko T, Zaneveld J, Knight R. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7:335–336 DOI 10.1038/nmeth.f.303.
- Cheng KK, Wang GY, Zeng J, Zhang JA. 2013. Improved succinate production by metabolic engineering. *BioMed Research International* 2013:538790 DOI 10.1155/2013/538790.
- Collado MC, Derrien M, Isolauri E, de Vos WM, Salminen S. 2007. Intestinal integrity and *Akkermansia muciniphila*, a mucin-degrading member of the intestinal microbiota present in infants, adults, and the elderly. *Applied Environmental Microbiology* 73(23):7767–7770 DOI 10.1128/AEM.01477-07.
- Delzenne NM, Cani PD. 2011. Interaction between obesity and the gut microbiota: relevance in nutrition. *Annual Review of Nutrition* 31:15–31 DOI 10.1146/annurev-nutr-072610-145146.
- DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen GL. 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied Environmental Microbiology* 72(7):5069–5072 DOI 10.1128/AEM.03006-05.
- Donohoe DR, Garge N, Zhang X, Sun W, O’Connell TM, Bunker MK, Bultman SJ. 2013. The microbiome and butyrate regulate energy metabolism and autophagy in the mammalian colon. *Cell Metabolism* 13(5):517–526 DOI 10.1016/j.cmet.2011.02.018.
- Duncan SH, Lopley GE, Holtrop G, Ince J, Johnstone AM, Louis P, Flint HJ. 2008. Human colonic microbiota associated with diet, obesity and weight loss. *International Journal of Obesity* 32:1720–1724 DOI 10.1038/ijo.2008.155.

- Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26(19):2460–2461 DOI 10.1093/bioinformatics/btq461.
- Everard A, Belzer C, Geurts L, Ouwerkerk JP, Druart C, Bindels LB, Guiot Y, Derrien M, Muccioli GG, Delzenne NM. 2013. Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity. *Proceedings of the National Academy of Sciences of the United States of America* 110(22):9066–9071 DOI 10.1073/pnas.1219451110.
- Fernandes J, Su W, Rahat-Rozenbloom S, Wolever TM, Comelli EM. 2014. Adiposity, gut microbiota and faecal short chain fatty acids are linked in adult humans. *Nutrition & Diabetes* 4:e121 DOI 10.1038/nutd.2014.23.
- Garcia-Mazcorro JF, Suchodolski JS, Jones KR, Clark-Price SC, Dowd SE, Minamoto Y, Markel M, Steiner JM, Dossin O. 2012. Effect of the proton pump inhibitor omeprazole on the gastrointestinal bacterial microbiota of healthy dogs. *FEMS Microbiology Ecology* 80(3):624–636 DOI 10.1111/j.1574-6941.2012.01331.x.
- Guo X, Xia X, Tang R, Zhou J, Zhao H, Wang K. 2008. Development of a real-time PCR method for *Firmicutes* and *Bacteroidetes* in faeces and its application to quantify intestinal population of obese and lean pigs. *Letters in Applied Microbiology* 47(5):367–373 DOI 10.1111/j.1472-765X.2008.02408.x.
- Gupta H. 2014. Barriers to and facilitators of long term weight loss maintenance in adult UK people: a thematic analysis. *International Journal of Preventive Medicine* 5(12):1512–1520.
- Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: paleontological statistics software package for education and data analysis. *Paleontologia Electronica* 4:1–9.
- Huijsdens XW, Linskens RK, Mak M, Meuwissen SGM, Vandenbroucke-Grauls CMJE, Savelkoul PHM. 2002. Quantification of bacteria adherent to gastrointestinal mucosa by real-time PCR. *Journal of Clinical Microbiology* 40(12):4423–4427 DOI 10.1128/JCM.40.12.4423-4427.2002.
- Igarashi H, Maeda S, Ohno K, Horigome A, Odamaki T, Tsujimoto H. 2014. Effect of oral administration of metronidazole or prednisolone on fecal microbiota in dogs. *PLoS ONE* 9(9):e107909 DOI 10.1371/journal.pone.0107909.
- Jakobsdottir G, Xu J, Molin G, Ahrné S, Nyman M. 2013. High-fat diet reduces the formation of butyrate, but increases succinate, inflammation, live fat and cholesterol in rats, while dietary fibre counteracts these effects. *PLoS ONE* 8(11):e80476 DOI 10.1371/journal.pone.0080476.
- Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. 2012. KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Research* 40(D1):D109–D114 DOI 10.1093/nar/gkr988.
- Katcher HI, Legro RS, Kunselman AR, Gillies PJ, Demers LM, Bagshaw DM, Kris-Etherton PM. 2008. The effects of a whole grain-enriched hypocaloric diet on cardiovascular disease risk factors in men and women with metabolic syndrome. *The American Journal of Clinical Nutrition* 87(1):79–90.
- Kristensen M, Toubro S, Jensen MG, Ross AB, Riboldi G, Petronio M, Bügel S, Tetens I, Astrup A. 2011. Whole grain compared with refined wheat decreases the percentage of body fat following a 12-week, energy-restricted dietary intervention in postmenopausal women. *Journal of Nutrition* 142(4):710–716 DOI 10.3945/jn.111.142315.
- Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC, Burkpile DE, Vega Thurber RL, Knight R, Beiko RG, Huttenhower C. 2013. Predictive

- functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nature Biotechnology* 31:814–823 DOI 10.1038/nbt.2676.
- Larsen N, Vogensen FK, van den Berg FW, Nielsen DS, Andreasen AS, Pedersen BK, Al-Soud WA, Sørensen SJ, Hansen LH, Jakobsen M. 2010. Gut microbiota in human adults with type 2 diabetes differs from non-diabetic adults. *PLoS ONE* 5(2):e9085 DOI 10.1371/journal.pone.0009085.
- Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI. 2005. Obesity alters gut microbial ecology. *Proceedings of the National Academy of Sciences of the United States of America* 102(31):11070–11075 DOI 10.1073/pnas.0504978102.
- Liu C, Song Y, McTeague M, Vu AW, Wexler H, Finegold SM. 2003. Rapid identification of the species of the *Bacteroides fragilis* group by multiplex PCR assays using group- and species-specific primers. *FEMS Microbiology Letters* 222(1):9–16 DOI 10.1016/S0378-1097(03)00296-9.
- Louis P, Flint HJ. 2009. Diversity, metabolism and microbial ecology of butyrate-producing bacteria from the human large intestine. *FEMS Microbiology Letters* 294(1):1–8 DOI 10.1111/j.1574-6968.2009.01514.x.
- Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. 2012. Diversity, stability and resilience of the human gut microbiota. *Nature* 489:220–230 DOI 10.1038/nature11550.
- Lozupone CA, Hamady M, Kelley ST, Knight R. 2007. Quantitative and qualitative beta diversity measures lead to different insights into factors that structure microbial communities. *Applied Environmental Microbiology* 73(5):1576–1585 DOI 10.1128/AEM.01996-06.
- Markowitz VM, Chen IM, Palaniappan K, Chu K, Szeto E, Grechkin Y, Ratner A, Jacob B, Huang J, Williams P, Huntemann M, Anderson I, Mavromatis K, Ivanova NN, Kyrpides NC. 2012. IMG: the Integrated Microbial Genomes database and comparative analysis system. *Nucleic Acids Research* 40(D1):D115–D122 DOI 10.1093/nar/gkr1044.
- Neyrinck AM, Possemiers S, Druart C, van de Wiele T, De Backer F, Cani PD, Larondelle Y, Delzenne NM. 2011. Prebiotic effects of wheat arabinoxylan related to the increase in bifidobacteria, roseburia and bacteroides/prevotella in diet-induced obese mice. *PLoS ONE* 6(6):e20944 DOI 10.1371/journal.pone.0020944.
- Noratto G, Garcia-Mazcorro JF, Markel M, Martino HS, Minamoto Y, Steiner JM, Byrne D, Suchodolski JS, Mertens-Talcott SU. 2014. Carbohydrate-free peach and plum juice affects fecal microbial ecology in an obese animal model. *PLoS ONE* 9(7):e101723 DOI 10.1371/journal.pone.0101723.
- Oishi K, Yamamoto S, Itoh N, Nakao R, Yasumoto Y, Tanaka K, Kikuchi Y, Fukudome S, Okita K, Takano-Ishikawa Y. 2015. Wheat alkylresorcinols suppress high-fat, high-sucrose diet-induced obesity and glucose intolerance by increasing insulin sensitivity and cholesterol excretion in male mice. *Journal of Nutrition* 145(2):199–206 DOI 10.3945/jn.114.202754.
- Parks DH, Beiko RG. 2010. Identifying biologically relevant differences between metagenomic communities. *Bioinformatics* 26(6):715–721 DOI 10.1093/bioinformatics/btq041.
- Penders J, Vink C, Driessen C, London N, Thijs C, Stobberingh EE. 2005. Quantification of *Bifidobacterium* spp., *Escherichia coli* and *Clostridium difficile* in faecal samples of breast-fed and formula-fed infants by real-time PCR. *FEMS Microbiology Letters* 243(1):141–147.
- Petschow B, Doré J, Hibberd P, Dinan T, Reid G, Blaser M, Cani PD, Degnan FH, Foster J, Gibson G, Hutton J, Klaenhammer TR, Ley R, Nieuwdorp M, Pot B, Relman D, Serazin A, Sanders ME. 2013. Probiotics, prebiotics, and the host microbiome: the science of translation. *Annals of the New York Academy of Sciences* 1306:1–17 DOI 10.1111/nyas.12303.

- R Core Team.** 2014. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <http://www.R-project.org/> .
- Reichardt N, Duncan SH, Young P, Belenguer A, Leitch CM, Scott KP, Flint HJ, Louis P.** 2014. Phylogenetic distribution of three pathways for propionate production within the human gut microbiota. *ISME Journal* 8:1323–1335 DOI [10.1038/ismej.2014.14](https://doi.org/10.1038/ismej.2014.14).
- Rideout JR, He Y, Navas-Molina JA, Walters WA, Ursell LK, Gibbons SM, Chase J, McDonald D, Gonzalez A, Robbins-Pianka A, Clemente JC, Gilbert JA, Huse SM, Zhou HW, Knight R, Caporaso JG.** 2014. Subsampled open-reference clustering creates consistent, comprehensive OTU definitions and scales to billions of sequences. *PeerJ* 2:e545 DOI [10.7717/peerj.545](https://doi.org/10.7717/peerj.545).
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF.** 2009. Introducing mothur: open source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied Environmental Microbiology* 75(23):7537–7541 DOI [10.1128/AEM.01541-09](https://doi.org/10.1128/AEM.01541-09).
- Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C.** 2011. Metagenomic biomarker discovery and explanation. *Genome Biology* 12:R60 DOI [10.1186/gb-2011-12-6-r60](https://doi.org/10.1186/gb-2011-12-6-r60).
- Stevenson L, Phillips F, O’Sullivan K, Walton J.** 2012. Wheat bran: its composition and benefits to health, a European perspective. *International Journal of Food Sciences and Nutrition* 63(8):1001–1013 DOI [10.3109/09637486.2012.687366](https://doi.org/10.3109/09637486.2012.687366).
- Suchodolski JS, Markel ME, Garcia-Mazcorro JF, Unterer S, Heilmann RM, Dowd SE, Kachroo P, Ivanov I, Minamoto Y, Dillman EM, Steiner JM, Cook AK, Toresson L.** 2012. The fecal microbiome in dogs with acute diarrhea and idiopathic inflammatory bowel disease. *PLoS ONE* 7(12):e51907 DOI [10.1371/journal.pone.0051907](https://doi.org/10.1371/journal.pone.0051907).
- Turnbaugh PJ, Bäckhed F, Fulton L, Gordon JI.** 2008. Diet-induced obesity is linked to marked but reversible alterations in the mouse distal gut microbiome. *Cell Host & Microbe* 17(4):213–223 DOI [10.1016/j.chom.2008.02.015](https://doi.org/10.1016/j.chom.2008.02.015).
- Vitaglione P, Mennella I, Ferracane R, Rivellese AA, Giacco R, Ercolini D, Gibbons SM, La Storia A, Gilbert JA, Jonnalagadda S, Thielecke F, Gallo MA, Scalfi L, Fogliano V.** 2015. Whole-grain wheat consumption reduces inflammation in a randomized controlled trial on overweight and obese subjects with unhealthy dietary and lifestyle behaviors: role of polyphenolics bounds to cereal dietary fiber. *American Journal of Clinical Nutrition* 101(2):251–261 DOI [10.3945/ajcn.114.088120](https://doi.org/10.3945/ajcn.114.088120).
- Walter J, Tannock GW, Tilsala-Timisjarvi A, Rodtong S, Loach DM, Munro K, Alatossava T.** 2000. Detection and identification of gastrointestinal *Lactobacillus* species by using denaturing gel electrophoresis and species-specific PCR primers. *Applied Environmental Microbiology* 66(1):297–303.
- Wu GD, Lewis JD, Hoffmann C, Chen YY, Knight R, Bittinger K, Hwang J, Chen J, Berkowsky R, Nessel L, Li H, Bushman FD.** 2010. Sampling and pyrosequencing methods for characterizing bacterial communities in the human gut using 16S sequence tags. *BMC Microbiology* 10:206 DOI [10.1186/1471-2180-10-206](https://doi.org/10.1186/1471-2180-10-206).
- Zhang H, DiBaise JK, Zuccolo A, Kudrna D, Braidotti M, Yu Y, Parameswaran P, Crowell MD, Wing R, Rittmann BE, Krajmalnik-Brown R.** 2009. Human gut microbiota in obesity and after gastric bypass. *Proceedings of the National Academy of Sciences of the United States of America* 106(7):2365–2370 DOI [10.1073/pnas.0812600106](https://doi.org/10.1073/pnas.0812600106).