

RESEARCH NOTES COMPLETE LIFT OF A STRUCTURE SATISFYING

$$F^K - (-)^{K+1} F = 0$$

LOVEJOY S. DAS

Department of Mathematics
Kent State University
Tuscarawas Campus
New Philadelphia, Ohio 44663

(Received August 30, 1991 and in revised form January 12, 1992)

ABSTRACT. The idea of f -structure manifold on a differentiable manifold was initiated and developed by Yano [1], Ishihara and Yano [2], Goldberg [3] and among others. The horizontal and complete lifts from a differentiable manifold M^n of class C^∞ to its cotangent bundles have been studied by Yano and Patterson [4,5]. Yano and Ishihara [6] have studied lifts of an f -structure in the tangent and cotangent bundles. The purpose of this paper is to obtain integrability conditions of a structure satisfying $F^K - (-)^{K+1}F = 0$ and $F^W - (-)^{W+1}F \neq 0$ for $1 < W < K$, in the tangent bundle.

KEY WORDS AND PHRASES. Tangent bundle, Complete lift, F -structure, Integrability, Distributions.

1991 AMS SUBJECT CLASSIFICATION CODE. 53C15.

1. INTRODUCTION.

Let F be a nonzero tensor field of the type $(1,1)$ and of class C^∞ on an n dimensional manifold M^n such that [7]

$$F^K - (-)^{K+1} F = 0 \quad \text{and} \quad F^W - (-)^{W+1} F \neq 0 \quad \text{for} \quad 1 < W < K, \quad (1.1)$$

where K is a fixed positive integer greater than 2. The degree of the manifold being $K(K \geq 3)$. Such a structure on M^n has been called $F(K, -(-)^{K+1})$ - structure of rank r , where the rank $(F) = r$ and is constant on M^n . The case when K is odd and $K(\geq 3)$ has been considered in this paper.

Let the operators on M^n be defined as follows [7]

$$l = (-)^{K+1} F^{K+1} \quad \text{and} \quad m = I - (-)^{K+1} F^{K+1}, \quad (1.2)$$

where I denotes the identity operator on M^n . We will state the following two theorems. [7]

THEOREM (1.1). Let M^n be an $F(K, -(-)^{K+1})$ manifold then,

$$l + m = I, \quad l^2 = l \quad \text{and} \quad m^2 = m \quad (1.3)$$

For F satisfying (1.1), there exist complementary distributions L and M , corresponding to the projection operators l and m respectively. If the rank of F is constant and is equal to $r = r(F)$ then $\dim L = r$ and $\dim M = (n-r)$.

THEOREM (1.2). We have

$$a) \quad F1 = 1F = F \quad \text{and} \quad Fm = mF = 0 \tag{1.4)a}$$

$$b) \quad F^{K-1} 1 = 1 \quad \text{and} \quad F^{K-1} m = 0 \tag{1.4)b}$$

Then $F^{\frac{K-1}{2}}$ acts on L as an almost product structure and on M as a null operator.

2. COMPLETE LIFT ON $F(K, (-)^{K+1})$ - STRUCTURE IN TANGENT BUNDLE.

Let M be an n-dimensional differentiable manifold of class C^∞ and $T_p(M^n)$ the tangent space at a point p of M_n and

$$T(M^n) = \bigcup_{p \in M^n} T_p(M^n) \text{ is the tangent bundle over the manifold } M^n.$$

Let us denote by $T_s^r(M^n)$, the set of all tensor fields of class C^∞ and of type (r,s) in M^n and $T(M^n)$ be the tangent bundle over M^n . The complete lift F^C of an element of $T_1^1(M^n)$ with local components F_i^h has components of the form [5]

$$F^C : \begin{pmatrix} F_i^h & 0 \\ \delta_i^h & F_i^h \end{pmatrix}$$

Now we obtain the following results on the complete lift of F satisfying (1.1).

THEOREM (2.1). For $F \in T_1^1(M^n)$, the complete lift F^C of F is an $F(K, (-)^{K+1})$ - structure iff it is for F also. Then F is of rank r iff F^C is of rank 2r.

PROOF. Let $F, G \in T_1^1(M^n)$. Then we have [5]

$$(FG)^C = F^C G^C \tag{2.2}$$

Replacing G by F in (2.2) we obtain

$$\begin{aligned} (FF)^C &= F^C F^C \\ \text{or, } (F^2)^C &= (F^C)^2 \end{aligned} \tag{2.3}$$

Now putting $G = F^{K-1}$ in (2.2) since G is (1,1) tensor field therefore F^{K-1} is also (1,1) so we obtain $(FF^{K-1})^C = F^C (F^{K-1})^C$ which in view of (2.3) becomes

$$(F^K)^C = (F^C)^K \tag{2.4}$$

Taking complete lift on both sides of equation (1.1) we get

$$(F^K)^C - ((-)^{K+1} F)^C = 0$$

which in consequence of equation (2.4) gives

$$(F^C)^K - (-)^{K+1} F^C = 0 \tag{2.5}$$

Thus equation (1.1) and (2.5) are equivalent. The second part of the theorem follows in view of equation (2.1).

Let F satisfying (1.1) be an F-structure of rank r in M^n . Then the complete lifts 1^C of 1 and m^C of m are complementary projection tensors in $T(M^n)$. Thus there exist in $T(M^n)$ two complementary distributions L^C and M^C determined by 1^C and m^C respectively.

3. INTEGRABILITY CONDITIONS OF $F(K, (-)^{K+1})$ STRUCTURE IN TANGENT BUNDLE.

Let $F \in T_1^{-1}(M^n)$, then the Nijenhuis tensor N_F of F satisfying (1.1) is a tensor field of the type (1,2) given by [6]

$$N_F(X, Y) = [FX, FY] - F[FX, Y] - F[X, FY] + F^2[X, Y]. \tag{3.1a}$$

Let N^C be the Nijenhuis tensor of F^C in $T(M^n)$ of F in M^n , then we have

$$N^C(X^C, Y^C) = [F^C X^C, F^C Y^C] - F^C[F^C X^C, Y^C] - F^C[X^C, F^C Y^C] + (F^2)^C[X^C, Y^C] \tag{3.1b}$$

For any $X, Y \in T_0^{-1}(M^n)$ and $F \in T_1^{-1}(M^n)$ we have [5]

$$[X^C, Y^C] = [X, Y]^C \text{ and } (X+Y)^C = X^C + Y^C \tag{3.2a}$$

$$F^C X^C = (FX)^C \tag{3.2b}$$

From (1.4)a and (3.2)b we have

$$F^C m^C = (Fm)^C = 0 \tag{3.3}$$

THEOREM (3.1). The following identities hold

$$N^C(m^C X^C, m^C Y^C) = (F^C)^2 [m^C X^C, m^C Y^C], \tag{3.4}$$

$$m^C N^C(X^C, Y^C) = m^C [F^C X^C, F^C Y^C] \tag{3.5}$$

$$m^C N^C(1^C X^C, 1^C Y^C) = m^C [F^C X^C, F^C Y^C] \tag{3.6}$$

$$m^C N^C((F^C)^{K-2} X^C, (F^C)^{K-2} Y^C) = m^C [1^C X^C, 1^C Y^C] \tag{3.7}$$

PROOF. The proofs of (3.4) to (3.7) follow in view of equations (1.4), (3.1)a, and (3.3)

THEOREM (3.2). For any $X, Y \in T_0^{-1}(M^n)$, the following conditions are equivalent.

$$m^C N^C(X^C, Y^C) = 0, \tag{i}$$

$$m^C N^C(1^C X^C, 1^C Y^C) = 0, \tag{ii}$$

$$m^C N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0 \tag{iii}$$

PROOF. In consequence of equations (3.1)b and (1.4)a,b it can be easily proved that

$$N^C(1^C X^C, 1^C Y^C) = 0 \text{ iff } N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0 \text{ for all } X \text{ and } Y \in T_0^{-1}(M^n)$$

Now due to the fact that equations (3.5) and (3.6) are equal, the conditions (i), (ii) and (iii) are equivalent to each other.

THEOREM (3.3). The complete lift of M^C of the distribution M in $T(M^n)$ is integrable iff M is integrable in M^n .

PROOF. It is known that the distribution M is integrable in M^n iff [2]

$$l[mX, mY] = 0 \tag{3.8}$$

for any $X, Y \in T_0^{-1}(M^n)$.

Taking complete lift on both sides of equations (3.8) we get

$$1^C [m^C X^C, m^C Y^C] = 0 \tag{3.9}$$

where $1^C = (I - m)^C = I - m^C$, is the projection tensor complementary to m^C . Thus the

conditions (3.8) and (3.9) are equivalent.

THEOREM (3.4). For any $X, Y \in T_0^1(M^n)$, let the distribution M be integrable in M^n iff $N(mX, mY) = 0$.

Then the distribution m^C is integrable in $T(M^n)$ iff $1^C N^C(m^C X^C, m^C Y^C) = 0$ or equivalently, $N^C(m^C X^C, m^C Y^C) = 0$.

PROOF. By virtue of condition (3.4) we have

$$N^C(m^C X^C, m^C Y^C) = (F^C)^2 [m^C X^C, m^C Y^C].$$

Multiplying throughout by 1^C we get

$$1^C N^C(m^C X^C, m^C Y^C) = (F^C)^2 1^C [m^C X^C, m^C Y^C].$$

which in view of equation (3.9) becomes

$$1^C N^C(m^C X^C, m^C Y^C) = 0 \tag{3.10}$$

Making use of equation (3.3), we get

$$m^C N^C(m^C X^C, m^C Y^C) = 0 \tag{3.11}$$

Adding (3.10) and (3.11) we obtain

$$(1^C + m^C) N^C(m^C X^C, m^C Y^C) = 0$$

Since $1^C + m^C = I^C = I$ we have $N^C(m^C X^C, m^C Y^C) = 0$.

THEOREM (3.5). For any $X, Y \in T_0^1(M^n)$ let the distribution L be integrable in M^n that is $mN(X, Y) = 0$ then the distribution L^C is integrable in $T(M^n)$ iff any one of the conditions of theorem (3.2) is satisfied.

PROOF. The distribution L is integrable in M^n iff [2] holds i.e. $m[1X, 1Y] = 0$. Thus the distribution L^C is integrable in $T(M^n)$ iff $m^C[1^C X^C, 1^C Y^C] = 0$.

On making use of equation (3.7), the theorem follows.

We now define the following:

- (i) distribution L is integrable,
- (ii) an arbitrary vector field Z tangent to an integral manifold of L ,
- (iii) the operator F , such that $FZ = FZ$.

Hence by virtue of theorem (1.2) the induced structure F^* is an almost product structure on each integral manifold of L and F^* makes tangent spaces invariant of every integral manifold of L . Let us denote the vector valued 2-form $N^*(Z, W)$, the Nijenhuis tensor corresponding to the Nijenhuis tensor of the almost product structure induced form $F(K, -(-)^{K+1})$ structure, on each integral manifold of L and for any two $Z, W \in T_0^1(M^n)$ tangent to an integral manifold of L , then we have

$$N^*(Z, W) = [F^*Z, W^*] - F^*[F^*Z, W^*] - F^*[Z, F^*W] + F^{*2} [Z, W], \tag{3.12}$$

which in view of (3.1)b and (3.12) yields

$$N^C[1^C X^C, 1^C Y^C] = N^*(1^C X^C, 1^C Y^C) \tag{3.13}$$

DEFINITION (3.1). We say that $F(K, -(-)^{K+1})$ - structure is partially integrable if the distribution L is integrable and the almost product structure F induced from F^* on each integral manifold of L is also integrable.

THEOREM (3.6). For any $X, Y \in T_0^{-1}(M^n)$ let the $F(K, -(-)^{K+1})$ - structure be partially integrable in M^n i.e. $N(1X, 1Y) = 0$. Then the necessary and sufficient condition for $F(K, -(-)^{K+1})$ - structure to be partially integrable in $T(M^n)$ is that $N^C(1^C X^C, 1^C Y^C) = 0$ or equivalently $N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0$.

PROOF. In view of equation (1.4) and equation (3.1)b we can prove easily that $N^C(1^C X^C, 1^C Y^C) = 0$ iff $N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0$, for any $X, Y \in T_0^{-1}(M^n)$.

Now in view of equation (3.13) and theorem (3.5), the result follows immediately.

When both distributions L and M are integrable we can choose a local coordinate system such that all L and M are respectively represented by putting $(n - r)$ local coordinates constant and r -coordinates constant. We call such a coordinate system an adapted coordinate system. It can be supposed that in an adapted coordinate system the projection operators l and m have the components of the form

$$l = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}, \quad m = \begin{pmatrix} 0 & 0 \\ 0 & I_{n-r} \end{pmatrix}$$

respectively where I_r denotes the unit matrix of order 'r' and I_{n-r} is of order $(n - r)$.

Since F satisfies equation (1.4)a, the tensor has components of the form

$$F = \begin{pmatrix} F_r & 0 \\ 0 & 0 \end{pmatrix}$$

in an adapted coordinate system where F_r denotes $r \times r$ square matrix.

DEFINITION (3.2). We say that on $F(K, -(-)^{K+1})$ structure is integrable if

- (i) The structure $F(K, -(-)^{K+1})$ is partially integrable.
- (ii) The distribution M is integrable i.e. $N(mX, mY) = 0$.
- (iii) The components of the $F(K, -(-)^{K+1})$ structure are independent of the coordinates which are constant along the integral manifold of L in an adapted coordinate system.

THEOREM (3.7). For any $X, Y \in T_0^{-1}(M^n)$ let $F(K, -(-)^{K+1})$ - structure be integrable in M^n iff $N(X, Y) = 0$. Then the $F(K, -(-)^{K+1})$ - structure is integrable in $T(M^n)$ iff $N^C(X^C, Y^C) = 0$.

PROOF. In view of equations (3.1)a and (3.1)b we get



$$N^C(X^C, Y^C) = (N(X, Y))^C.$$

Since $F(K, -(-)^{K+1})$ is integrable in M^n thus the theorem follows.

REFERENCES

1. YANO, K., On a structure defined by a tensor field f of type (1,1) satisfying $f^3 + f = 0$, Tensor Vol. 14 (1963), pp. 99-109.
2. ISHIHARA, S. and YANO, K., On integrability conditions of a structure f satisfying $f^3 + f = 0$, Quarterly J. Math. Vol. 15 (1964), pp. 217-222.
3. GOLDBERG, S.I. and YANO, K., Globally framed f -manifolds, Illinois J. Math. Vol 15 (1971), pp. 456-476.

4. YANO, K. and PATTERSON, E. M., Vertical and Complete lifts from a manifold to its cotangent bundles, Journal Math. Soc. Japan. Vol. 19 (1967), pp. 91-113.
5. YANO, K. and PATTERSON, E. M., Horizontal lifts from a manifold to its cotangent bundles, Journal Math. Soc. Japan. Vol. 19 (1967), pp. 185-198.
6. YANO, K. and ISHIHARA, S., Tangent and Cotangent Bundles, Marcel Dekker, Inc. New York, (1973).
7. DAS, LOVEJOY S., On integrability conditions of $F(K, -(-)^{K+1})$ - structure on a differentiable manifold, Revista Mathematica. Argentina. Vol. 26 (1977).
8. DAS, LOVEJOY S. and UPADHYAY, M. D., On F - structure manifold, Kyung Pook Mathematical Journal. Korea. Vol. 18 (1978), pp. 277-283.
9. DAS, LOVEJOY S., On differentiable manifold with $F(K, -(-)^{K+1})$ - structure of rank 'r', Revista Mathematica. Argentina. Vol. 27 (1978), pp. 277-283.
10. DAS, LOVEJOY S., Prolongations of F - structure to the tangent bundle of order 2. International J. Math & Math Sci., (to appear).
11. HELGASON, S., Differential Geometry. Lie Groups and Symmetric Spaces, Academic Press, New York, (1970).



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

