

Research Article

Investigating the Mechanisms of Action of Depside Salt from *Salvia miltiorrhiza* Using Bioinformatic Analysis

Hua Li^{1,2} and Hongying Liu^{3,4}

¹Shanghai Institute of Cardiovascular Diseases, Zhongshan Hospital, Fudan University, Shanghai 200032, China

²Institutes of Biomedical Sciences, Fudan University, Shanghai 200032, China

³Institutes of Panvascular Medicine, Fudan University, Shanghai 200032, China

⁴Shanghai Green Valley Pharmaceutical Company, Shanghai 201203, China

Correspondence should be addressed to Hua Li; lihua199988@hotmail.com and Hongying Liu; liuhongying@green-valley.com

Received 26 June 2017; Accepted 5 September 2017; Published 19 November 2017

Academic Editor: Darren R. Williams

Copyright © 2017 Hua Li and Hongying Liu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Salvia miltiorrhiza is a traditional Chinese medicinal herb used for treating cardiovascular diseases. Depside salt from *S. miltiorrhiza* (DSSM) contains the following active components: magnesium lithospermate B, lithospermic acid, and rosmarinic acid. This study aimed to reveal the mechanisms of action of DSSM. After searching for DSSM-associated genes in GeneCards, Search Tool for Interacting Chemicals, SuperTarget, PubChem, and Comparative Toxicogenomics Database, they were subjected to enrichment analysis using Multifaceted Analysis Tool for Human Transcriptome. A protein-protein interaction (PPI) network was visualised; module analysis was conducted using the Cytoscape software. Finally, a transcriptional regulatory network was constructed using the TRRUST database and Cytoscape. Seventy-three DSSM-associated genes were identified. JUN, TNF, NFKB1, and FOS were hub nodes in the PPI network. Modules 1 and 2 were identified from the PPI network, with pathway enrichment analysis, showing that the presence of *NFKB1* and *BCL2* in module 1 was indicative of a particular association with the NF- κ B signalling pathway. JUN, TNF, NFKB1, FOS, and *BCL2* exhibited notable interactions among themselves in the PPI network. Several regulatory relationships (such as *JUN* \rightarrow *TNF/FOS*, *FOS* \rightarrow *NFKB1* and *NFKB1* \rightarrow *BCL2/TNF*) were also found in the regulatory network. Thus, DSSM exerts effects against cardiovascular diseases by targeting *JUN*, *TNF*, *NFKB1*, *FOS*, and *BCL2*.

1. Introduction

Salvia miltiorrhiza (also named Danshen) is widely used as a traditional Chinese medicinal herb to prevent and treat vascular diseases [1, 2]. Depside salt from *S. miltiorrhiza* (DSSM) is a new medicine that contains the active components magnesium lithospermate B (MLB), lithospermic acid (LA), and rosmarinic acid (RA) [3]. MLB plays protective roles in relieving atherosclerosis and combating myocardial ischaemia-reperfusion injury [4]. RA and LA also have beneficial effects on cardiovascular diseases, such as atherosclerosis and neointimal hyperplasia [5, 6]. Thus, understanding the mechanisms of action of DSSM is important for its better utilisation in a clinical setting.

In patients with unstable angina, DSSM can suppress platelet activation and aggregation as well as matrix metalloproteinase 9 (*MMP-9*) expression and secretion [7]. MLB protects against diabetic atherosclerosis by inducing the nuclear factor erythroid 2-related factor-2-antioxidant responsive element-NAD(P)H: quinone oxidoreductase-1 pathway [8]. Du et al. reported that MLB can be used to treat ischaemic heart diseases as it specifically inhibits transforming growth factor β -activated protein kinase 1-binding protein 1-p38 apoptosis signalling [9]. Kim et al. also demonstrated that RA inhibits adriamycin-induced cardiotoxicity by suppressing reactive oxygen species generation as well as extracellular signal-regulated kinase and c-Jun N-terminal kinase (JNK) activation [10]. Moreover, LA inhibits

foetal bovine serum-induced vascular smooth muscle cell (VSMC) proliferation by arresting cell cycle progression and suppressing cyclin D1 expression and lipopolysaccharide-induced VSMC migration by downregulating *MMP-9* expression; thus, LA may be used for preventing neointimal hyperplasia, restenosis, and atherosclerosis [11]. However, no comprehensive survey of DSSM-associated genes has been reported.

In this study, we searched for DSSM-associated genes in several common databases. Subsequently, we applied multiple bioinformatic methods to identify further DSSM-associated key genes; these methods included enrichment analysis, protein-protein interaction (PPI) network and module analyses, and transcriptional regulatory network analysis. This study may provide a better understanding of the mechanisms of action of DSSM.

2. Methods

2.1. Search for DSSM-Associated Genes. Several databases were used in this study. The human genomic database GeneCards (<http://www.genecards.org/>) was used [12], which provides concise genome, transcriptome, proteome, and function data of all predicted and known human genes. Search Tool for Interacting Chemicals (STICH, version 5.0, <http://stich.embl.de/>) was also used [13], which is a database that integrates the interactions between chemicals and proteins. SuperTarget (<http://insilico.charite.de/supertarget/>) [14] was also used, which is a database that includes drug-associated information correlated with drug metabolism, adverse drug effects, medical indications, Gene Ontology (GO) terms, and pathways for target proteins. The PubChem (<https://www.ncbi.nlm.nih.gov/pccompound/>) [15] database was also used, which provides information on chemical substances and corresponding biological activities and is linked to the National Institutes of Health PubMed Entrez. Moreover, the study features the Comparative Toxicogenomics Database (CTD, <http://ctdbase.org/>) [16], which is a database associated with environmental chemical-gene product interactions as well as the transportation and accumulation of chemical substances in the human body. In GeneCards [12], STICH (parameters set as organism = human, score > 0.4) [13], SuperTarget [14], PubChem [15], and CTD [16] databases, “magnesium lithospermate B”, “lithospermic acid” and “rosmarinic acid” were used as keywords to search for genes associated with them (which were combined into a set of DSSM-associated genes).

2.2. Functional and Pathway Enrichment Analyses. The GO (<http://www.geneontology.org>) database describes the associations of gene products with the categories of biological process (BP), molecular function (MF), and cellular component (CC), as well as with more specific subcategories [17]. The Kyoto Encyclopedia of Genes and Genomes (KEGG, <http://www.genome.ad.jp/kegg>) is a database used for annotating the functions of genes or other molecules [18]. The “BioCloud” online platform was developed for managing problems encountered in the analysis of high-throughput data. By applying the Multifaceted Analysis Tool for Human

Transcriptome (MATHT, <http://www.biocloudservice.com>) in the “BioCloud” online platform, GO functional and KEGG pathway enrichment analyses were conducted for DSSM-associated genes, setting a threshold of <0.05 for the false discovery rate (FDR).

2.3. PPI Network and Module Analyses. On combining the Search Tool for the Retrieval of Interacting Genes (STRING, version 10.0, <http://www.string-db.org/>, combined score > 0.4) [19], Biological General Repository for Interaction Datasets (BioGRID, version 3.4, <https://wiki.thebiogrid.org/>) [20] and Human Protein Reference Database (release 9, <http://www.hprd.org/>) [21] interaction databases, PPI pairs among DSSM-associated genes were predicted. Subsequently, the PPI network was visualised for DSSM-associated genes using the Cytoscape software (<http://www.cytoscape.org>) [22]. Using the CytoNCA plug-in [23] (version 2.1.6, <http://apps.cytoscape.org/apps/cytonca>) in Cytoscape, degree centrality (DC), betweenness centrality (BC), and closeness centrality of the nodes were analysed to obtain the hub proteins in the PPI network [24]. The parameter was set as “without weight.”

Based on the MCODE plug-in [25] (version 1.4.2; <http://apps.cytoscape.org/apps/mcode>; parameters set as degree cut-off = 2, maximum depth = 100, node score cut-off = 0.2, and *K*-core = 2) in Cytoscape, module analysis was conducted for the PPI network. Subsequently, KEGG pathway enrichment analysis was performed for the nodes of significant modules, with FDR < 0.05 as the cut-off criterion.

2.4. Transcriptional Regulatory Network Construction. Using the transcriptional regulatory relationships unravelled by a sentence-based text-mining (TRRUST, <http://www.grnpedia.org/trrust/>) [26] database, transcription factors (TFs) among DSSM-associated genes were searched and then their targets were screened. Finally, a transcriptional regulatory network was constructed using Cytoscape [22].

3. Results

3.1. Search for DSSM-Associated Genes. The numbers of MLB-, RA-, and LA-associated genes identified from GeneCards, STICH, SuperTarget, PubChem, and CTD databases are listed in Table 1. The MLB-, RA-, and LA-associated genes were combined into a group of 73 DSSM-associated genes.

3.2. Functional and Pathway Enrichment Analyses. The main BP, CC, and MF terms, as well as KEGG pathways enriched for DSSM-associated genes, are shown in Figure 1. The significantly enriched terms included inflammatory response (BP), cytosol (CC), protein binding (MF), and tumour necrosis factor (TNF) signalling pathway (KEGG pathway).

3.3. PPI Network and Module Analyses. The PPI network for DSSM-associated genes comprised 65 nodes and 431 edges (Figure 2). The 65 nodes included 21 MLB-associated genes, 56 RA-associated genes, and eight LA-associated genes. Among them, two genes [nitric oxide synthase 2 (*NOS2*) and nitric oxide synthase 3 (*NOS3*)] were associated with all MLB,




FIGURE 1: The main BP, CC, and MF terms as well as KEGG pathways enriched for DSSM-associated genes. BP, biological process; CC, cellular component; MF, molecular function; KEGG, Kyoto Encyclopedia of Genes and Genomes; DSSM, deposite salt from *Salvia miltiorrhiza*.

RA, and LA; four genes (interleukin 2, interleukin 1 beta, caspase 3, and v-rel avian reticuloendotheliosis viral oncogene homolog A) were associated with both MLB and RA; and four genes (xanthine dehydrogenase, aldo-keto reductase family 1 member B1, protein tyrosine phosphatase no-receptor type 1 and procollagen-lysine, and 2-oxoglutarate 5-dioxygenase) were associated with both MLB and LA. Jun protooncogene (JUN), TNF, nuclear factor kappa B subunit 1 (NFKB1), and Fos protooncogene (FOS) were noted as particular hub nodes according to their BC, closeness centrality, and DC scores (Table 2). Moreover, two modules were also identified from the PPI network: module 1 (21 nodes, score = 13.2) and module 2 (12 nodes, score = 7.8) (Figure 3). In addition, pathway enrichment analysis demonstrated that the nuclear factor kappa B (NF- κ B) signalling pathway [Figure 4; $p = 3.14E^{-06}$; involving *NFKB1* and B-cell CLL/lymphoma 2 (*BCL2*)] and leishmaniasis ($p = 4.72E^{-07}$) were particularly associated with the nodes in modules 1 and 2, respectively (Table 3 and Figure 5). JUN, TNF, NFKB1, FOS, and BCL2 also exhibited notable interactions among themselves in the PPI network.

3.4. Transcriptional Regulatory Network Analysis. Based on the TRRUST database, TFs among DSSM-associated genes

were searched for and their targets were screened. The constructed transcriptional regulatory network had 37 nodes (including eight TFs and 29 target genes) and 108 relationship pairs (such as *JUN* \rightarrow *TNF/FOS*, *FOS* \rightarrow *NFKB1*, and *NFKB1* \rightarrow *BCL2/TNF*) (Figure 6).

4. Discussion

In this study, a total of 73 DSSM-associated genes were identified from various databases. In the PPI network constructed on the basis of these genes and their interactions, JUN, TNF, NFKB1, and FOS were established as hub nodes according to their BC, closeness centrality, and DC scores. JUN, TNF, NFKB1, FOS, and BCL2 were also shown to interact among themselves in the PPI network. Moreover, two distinct modules (modules 1 and 2) of the PPI network were identified. In addition, several regulatory relationships (such as *JUN* \rightarrow *TNF/FOS*, *FOS* \rightarrow *NFKB1*, and *NFKB1* \rightarrow *BCL2/TNF*) were found to be involved in the transcriptional regulatory network.

Prolonged anti-TNF- α therapy has beneficial effects on the signs of subclinical cardiovascular disease in patients with severe psoriasis [27]. Studies have shown that, compared with

TABLE 1: MLB-associated genes, RA-associated genes, and LA-associated genes searched from the GeneCards, STICH, SuperTarget, PubChem, and CTD databases.

Database	MLB		Count	RA		Count	LA	
	Count	Symbol		Symbol	Symbol			
GeneCards	6	NOS2, NOS3, AKR1B1, XDH, PLOD1, PTPN1	33	SLC16A1, PTGS2, TNF, ALOX5, IL1B, TAT, JUN, BCL2, CXCL8, CCL11, LCK, FOS, CCL2, CREB1, ERVK-6, PIK3CG, NFKB1, PLCG1, IKKBK, SHC1, MAP2K1, CCL3, RELA, GRB2, XIAP, EEF1A1, CCR3, CXCL2, BIRC2, ZAP70, BIRC3, ITK, MIR155	8	NOS2, NOS3, AKR1B1, XDH, PLOD1, PTPN1, SPTLC1, SPTLC2		
STICH	8	IL15, IL2, IL4, KCNMA1, KCNU1, MAPK8, RRAGA, RRAGB	7	CCR3, FOS, IKKBK, IL2, LCK, PARG, PROCR	0	/		
SuperTarget	0	/	5	ALDR, CIR, HYALI, LCK, TYRO PROCR, PARG, TNF, ADAM17, SERPINE1, MAOB, MAOA, IL1B, DNMT1, COL1A1	0	/		
PubChem	0	/	10	PROCR, ADAM17, IL1B, CXCL1, IL6, TNF, SERPINE1, AIFM1, CASP3, COL1A1, COMT, DNMT1, FN1, GPT, MAOA, MAOB, NOS2, NOS3, OGG1, PARG, RELA	0	/		
CTD	7	IFNG, IL1B, CASP3, NFE2L2, SIRT1, CAT, RELA	21		1	XDH		

MLB, magnesium lithospermate B; RA, rosmarinic acid; LA, lithospermic acid, STICH, Search Tool for Interacting Chemicals; CTD, Comparative Toxicogenomics Database.






FIGURE 3: Modules 1 and 2 identified from the protein-protein interaction network. Green squares and purple circles represent magnesium lithospermate B-associated genes and rosmarinic acid-associated genes, respectively. Red diamonds represent genes associated with more than one drug. The larger nodes indicate genes with higher degrees.

the reduction of plasma inflammatory markers [39]. By increasing *BCL2* expression, *miR-21* promotes heart failure progression with preserved left ventricular ejection fraction and subsequently inhibits cardiac fibrosis [40]. Moreover,

Tang et al. reported that *miR-1* targets *BCL2* functions in mediating cardiomyocyte apoptosis [41]. In this study, pathway enrichment analysis showing the presence of *NFKB1* and *BCL2* in module 1 indicated a particular association with



(a)



(b)

FIGURE 5: Pathways enriched for the nodes involved in modules 1 (a) and 2 (b).

TABLE 3: Pathways enriched for the genes separately involved in module 1 and module 2 (top 10 listed).

Module	Term	Count	Genes	FDR
	hsa04668:TNF signaling pathway	12	<i>PIK3CG, FOS, IL6, CCL2, MAP2K1, JUN, CREB1, NFKB1, MAPK8, BIRC3, IKKBK, BIRC2</i>	1.18E - 12
	hsa05161:Hepatitis B	11	<i>PIK3CG, FOS, IL6, MAP2K1, GRB2, JUN, BCL2, CREB1, NFKB1, MAPK8, IKKBK</i>	2.20E - 09
	hsa04660:T cell receptor signaling pathway	10	<i>PIK3CG, FOS, MAP2K1, GRB2, JUN, LCK, ZAP70, NFKB1, IKKBK, IL2</i>	4.40E - 09
	hsa04380:Osteoclast differentiation	10	<i>PIK3CG, FOS, MAP2K1, GRB2, JUN, CREB1, LCK, NFKB1, MAPK8, IKKBK</i>	3.98E - 08
Module 1	hsa05200:Pathways in cancer	13	<i>PIK3CG, FOS, IL6, XIAP, MAP2K1, GRB2, JUN, BCL2, NFKB1, MAPK8, BIRC3, IKKBK, BIRC2</i>	9.52E - 08
	hsa05142:Chagas disease (American trypanosomiasis)	9	<i>PIK3CG, FOS, IL6, CCL2, JUN, NFKB1, MAPK8, IKKBK, IL2</i>	2.56E - 07
	hsa04722:Neurotrophin signaling pathway	9	<i>PIK3CG, MAP2K1, GRB2, JUN, BCL2, NFKB1, MAPK8, SHC1, IKKBK</i>	8.15E - 07
	hsa04510:Focal adhesion	10	<i>PIK3CG, XIAP, MAP2K1, GRB2, JUN, BCL2, MAPK8, SHC1, BIRC3, BIRC2</i>	2.33E - 06
	hsa04064:NF-kappa B signaling pathway	8	<i>XIAP, BCL2, LCK, ZAP70, NFKB1, BIRC3, IKKBK, BIRC2</i>	3.14E - 06
	hsa04621:NOD-like receptor signaling pathway	7	<i>IL6, CCL2, NFKB1, MAPK8, BIRC3, IKKBK, BIRC2</i>	8.05E - 06
	hsa05140:Leishmaniasis	7	<i>IL4, TNF, PTGS2, RELA, IFNG, IL1B, NOS2</i>	4.72E - 07
	hsa05142:Chagas disease (American trypanosomiasis)	7	<i>CCL3, TNF, RELA, SERPINE1, IFNG, IL1B, NOS2</i>	4.90E - 06
	hsa05146:Amoebiasis	6	<i>CASP3, TNF, RELA, IFNG, IL1B, NOS2</i>	3.72E - 04
	hsa05321:Inflammatory bowel disease (IBD)	5	<i>IL4, TNF, RELA, IFNG, IL1B</i>	2.36E - 03
Module 2	hsa05133:Pertussis	5	<i>CASP3, TNF, RELA, IL1B, NOS2</i>	4.47E - 03
	hsa05152:Tuberculosis	6	<i>CASP3, TNF, RELA, IFNG, IL1B, NOS2</i>	4.76E - 03
	hsa05132:Salmonella infection	5	<i>CCL3, RELA, IFNG, IL1B, NOS2</i>	6.71E - 03
	hsa04066:HIF-1 signaling pathway	5	<i>RELA, SERPINE1, IFNG, NOS3, NOS2</i>	1.30E - 02
	hsa04668:TNF signaling pathway	5	<i>CASP3, TNF, PTGS2, RELA, IL1B</i>	1.78E - 02
	hsa05145:Toxoplasmosis	5	<i>CASP3, TNF, RELA, IFNG, NOS2</i>	2.72E - 02

the NF- κ B signalling pathway, indicating that *NFKB1* and *BCL2* may be correlated with the effects of DSSM against cardiovascular diseases through this pathway.

5. Conclusion

In conclusion, a total of 73 DSSM-associated genes were identified by conducting a search of public databases. *JUN*,

TNF, *NFKB1*, *FOS*, and *BCL2* were also revealed as potential targets of DSSM for treating cardiovascular diseases. However, further experimental research should be performed to confirm these findings obtained by bioinformatic analysis.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

- kinase and extracellular signal-regulated kinase," *Biochemical Pharmacology*, vol. 70, no. 7, pp. 1066–1078, 2005.
- [11] L. Chen, W.-Y. Wang, and Y.-P. Wang, "Inhibitory effects of lithospermic acid on proliferation and migration of rat vascular smooth muscle cells," *Acta Pharmacologica Sinica*, vol. 30, no. 9, pp. 1245–1252, 2009.
 - [12] M. Safran, I. Dalah, J. Alexander et al., "GeneCards Version 3: the human gene integrator," *Database*, vol. 2010, Article ID baq020, 2010.
 - [13] D. Szklarczyk, A. Santos, C. Von Mering, L. J. Jensen, P. Bork, and M. Kuhn, "STITCH 5: Augmenting protein-chemical interaction networks with tissue and affinity data," *Nucleic Acids Research*, vol. 44, no. 1, pp. D380–D384, 2016.
 - [14] N. Hecker, J. Ahmed, J. von Eichborn et al., "SuperTarget goes quantitative: update on drug-target interactions," *Nucleic Acids Research*, vol. 40, no. 1, Article ID gkr912, pp. D1113–D1117, 2012.
 - [15] S. Kim, P. A. Thiessen, E. E. Bolton et al., "PubChem substance and compound databases," *Nucleic Acids Research*, vol. 44, no. 1, pp. D1202–D1213, 2016.
 - [16] A. P. Davis, C. J. Grondin, R. J. Johnson et al., "The Comparative Toxicogenomics Database: Update 2017," *Nucleic Acids Research*, vol. 45, no. 1, pp. D972–D978, 2017.
 - [17] S. Tweedie, M. Ashburner, K. Falls et al., "FlyBase: enhancing Drosophila Gene Ontology annotations," *Nucleic Acids Research*, vol. 37, no. 1, pp. D555–D559, 2009.
 - [18] M. Kanehisa and S. Goto, "KEGG: kyoto encyclopedia of genes and genomes," *Nucleic Acids Research*, vol. 28, no. 1, pp. 27–30, 2000.
 - [19] D. Szklarczyk, A. Franceschini, S. Wyder et al., "STRING v10: protein-protein interaction networks, integrated over the tree of life," *Nucleic Acids Research*, vol. 43, pp. D447–D452, 2015.
 - [20] A. Chatr-Aryamontri, B.-J. Breitkreutz, R. Oughtred et al., "The BioGRID interaction database: 2015 update," *Nucleic Acids Research*, vol. 43, no. 1, pp. D470–D478, 2015.
 - [21] T. S. Keshava Prasad, R. Goel, K. Kandasamy et al., "Human protein reference database—2009 update," *Nucleic Acids Research*, vol. 37, no. 1, pp. D767–D772, 2009.
 - [22] R. Saito, M. E. Smoot, K. Ono et al., "A travel guide to Cytoscape plugins," *Nature Methods*, vol. 9, no. 11, pp. 1069–1076, 2012.
 - [23] Y. Tang, M. Li, J. Wang, Y. Pan, and F.-X. Wu, "CytoNCA: a cytoscape plugin for centrality analysis and evaluation of protein interaction networks," *BioSystems*, vol. 127, pp. 67–72, 2015.
 - [24] X. He and J. Zhang, "Why do hubs tend to be essential in protein networks?" *PLoS Genetics*, vol. 2, no. 6, p. e88, 2006.
 - [25] G. D. Bader and C. Hogue, "An automated method for finding molecular complexes in large protein interaction networks," *BMC Bioinformatics*, vol. 4, 2003.
 - [26] H. Han, H. Shim, D. Shin et al., "TRRUST: A reference database of human transcriptional regulatory interactions," *Scientific Reports*, vol. 5, Article ID 11432, 2015.
 - [27] E. Herédi, J. Végh, L. Pogácsás et al., "Subclinical cardiovascular disease and its improvement after long-term TNF- α inhibitor therapy in severe psoriatic patients," *Journal of the European Academy of Dermatology and Venereology*, vol. 30, no. 9, pp. 1531–1536, 2016.
 - [28] D. H. Solomon, J. R. Curtis, K. G. Saag et al., "Cardiovascular risk in rheumatoid arthritis: comparing tnf- α blockade with nonbiologic DMARDs," *American Journal of Medicine*, vol. 126, no. 8, pp. 730.e9–730.e17, 2013.
 - [29] S. L. Westlake, A. N. Colebatch, J. Baird et al., "Tumour necrosis factor antagonists and the risk of cardiovascular disease in patients with rheumatoid arthritis: a systematic literature review," *Rheumatology*, vol. 50, no. 3, pp. 518–531, 2011.
 - [30] H. Zhang, Y. Park, J. Wu et al., "Role of TNF- α in vascular dysfunction," *Clinical Science*, vol. 116, no. 3, pp. 219–230, 2009.
 - [31] Y. Zhang, X. Yang, F. Bian, and et al., "TNF- α promotes early atherosclerosis by increasing transcytosis of LDL across endothelial cells: Crosstalk between NF- κ B and PPAR- γ ," *Journal of Molecular and Cellular Cardiology*, vol. 72, pp. 85–94, 2014.
 - [32] C. Ferrandi, R. Ballerio, P. Gaillard et al., "Inhibition of c-Jun N-terminal kinase decreases cardiomyocyte apoptosis and infarct size after myocardial ischemia and reperfusion in anaesthetized rats," *British Journal of Pharmacology*, vol. 142, no. 6, pp. 953–960, 2004.
 - [33] G. Milano, S. Morel, C. Bonny, and et al., "A peptide inhibitor of c-Jun NH2-terminal kinase reduces myocardial ischemia-reperfusion injury and infarct size in vivo," *American Journal of Physiology, Heart and Circulatory Physiology*, vol. 292, no. 4, pp. H1828–H1835, 2007.
 - [34] H. Yasumoto, S. Kim, Y. Zhan et al., "Dominant negative c-Jun gene transfer inhibits vascular smooth muscle cell proliferation and neointimal hyperplasia in rats," *Gene Therapy*, vol. 8, no. 22, pp. 1682–1689, 2001.
 - [35] S. Saliques, J.-R. Teyssier, C. Vergely et al., "Smoking and FOS expression from blood leukocyte transcripts in patients with coronary artery disease," *Atherosclerosis*, vol. 219, no. 2, pp. 931–936, 2011.
 - [36] X. Palomer, E. Capdevila-Busquets, G. Botteri, and et al., "miR-146a targets Fos expression in human cardiac cells," *Disease Models and Mechanisms*, vol. 8, no. 9, pp. 1081–1091, 2015.
 - [37] Y. N. Yang, J. Y. Zhang, and Y. T. Ma, "GW25-e3074 94 ATTG Insertion/Deletion Polymorphism of the NFKB1 Gene Is Associated with Coronary Artery Disease in Han and Uygur Women in China," *Genetic Testing and Molecular Biomarkers*, vol. 18, no. 6, pp. 430–438, 2014.
 - [38] H. Lai, Q. Chen, X. Li et al., "Association between genetic polymorphism in NFKB1 and NFKBIA and coronary artery disease in a Chinese Han population," *International Journal of Clinical and Experimental Medicine*, vol. 8, no. 11, pp. 21487–21496, 2015.
 - [39] V. Boccardi, M. R. Rizzo, R. Marfella et al., "-94 ins/del ATTG NFKB1 gene variant is associated with lower susceptibility to myocardial infarction," *Nutrition, Metabolism and Cardiovascular Diseases*, vol. 21, no. 9, pp. 679–684, 2011.
 - [40] S. Dong, W. Ma, B. Hao, and et al., "microRNA-21 promotes cardiac fibrosis and development of heart failure with preserved left ventricular ejection fraction by up-regulating Bcl-2," *International Journal of Clinical and Experimental Pathology*, vol. 7, no. 2, pp. 565–574, 2014.
 - [41] Y. Tang, J. Zheng, Y. Sun, Z. Wu, Z. Liu, and G. Huang, "MicroRNA-1 regulates cardiomyocyte apoptosis by targeting Bcl-2," *International Heart Journal*, vol. 50, no. 3, pp. 377–387, 2009.



Hindawi
Submit your manuscripts at
<https://www.hindawi.com>

