

RESEARCH

Open Access

Weighted differentiation composition operators from weighted bergman space to n th weighted space on the unit disk

Liang Zhang and Hong-Gang Zeng *

* Correspondence: zhgng@tju.edu.cn
Department of Mathematics,
Tianjin University, Tianjin 300072,
People's Republic of China**Abstract**

This paper characterizes the boundedness and compactness of the weighted differentiation composition operator from weighted Bergman space to n th weighted space on the unit disk of \mathbb{D} .

2000 Mathematics Subject Classification: Primary: 47B38; Secondary: 32A37, 32H02, 47G10, 47B33.

Keywords: weighted differentiation composition operators, weighted Bergman space, n th weighted space, boundedness, compactness

1. Introduction

Let \mathbb{D} be the open unit disk in the complex space \mathbb{C} , dA the Lebesgue measure on \mathbb{D} normalized so that $A(\mathbb{D}) = 1$. Let $H(\mathbb{D})$ be the space of all analytic functions on \mathbb{D} .

Let $\alpha > -1$, $p > 0$. f is said to belong to the weighted Bergman space, denoted by $A_\alpha^p (= A_\alpha^p(\mathbb{D}))$, if $f \in H(\mathbb{D})$ and

$$\|f\|^p = (\alpha + 1) \int_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\alpha dA < \infty.$$

When $0 < p < 1$, it is complete metric space; when $p \geq 1$, it is a Banach space.

Let $\mu(z)$ (weight) be a positive continuous function on \mathbb{D} and $n \in \mathbb{N}_0$. The n th weighted space on the unit disk, denoted by $\omega_\mu^{(n)}(\mathbb{D})$, consists of all $f \in H(\mathbb{D})$ such that

$$b_{\omega_\mu^{(n)}(\mathbb{D})}(f) = \sup_{z \in \mathbb{D}} \mu(z) |f^{(n)}(z)| < \infty.$$

For $n = 0$, the space becomes the weighted-type space $H_\mu^\infty(\mathbb{D})$; for $n = 1$, the Bloch-type space $\mathcal{B}_\mu(\mathbb{D})$; and for $n = 2$, the Zygmund-type $Z_\mu(\mathbb{D})$. For more details about these spaces, we recommend the readers to ([1,2]).

The expression $b_{\omega_\mu^{(n)}(\mathbb{D})}(f)$ defines a semi-norm on the n th weighted space $\omega_\mu^{(n)}(\mathbb{D})$, while the natural norm is given by

$$\|f\|_{\omega_\mu^{(n)}(\mathbb{D})} = \sum_{j=0}^{n-1} |f^{(j)}(0)| + b_{\omega_\mu^{(n)}(\mathbb{D})}(f).$$

With this norm, $\omega_\mu^{(n)}(\mathbb{D})$ becomes a Banach space. The little n th weighted space, denoted by $\omega_{\mu,0}^{(n)}(\mathbb{D})$, is a closed subspace of $\omega_\mu^{(n)}(\mathbb{D})$, consisting of those f for which

$$\lim_{|z| \rightarrow 1} \mu(z) |f^{(n)}(z)| = 0.$$

Let ϕ be a non-constant analytic self-map of \mathbb{D} , $u(z) \in H(\mathbb{D})$, and $m \in \mathbb{N}$. The weighted differentiation composition operator $D_{\phi,u}^m$ is defined by

$$D_{\phi,u}^m f(z) = u(z) f^{(m)}(\phi(z)),$$

for $z \in \mathbb{D}$, $f \in H(\mathbb{D})$. If $m = 1$, $u(z) = \phi'(z)$, then $D_{\phi,u}^m = DC_\phi$; if let $m = 1$, $u(z) = 1$, then $D_{\phi,u}^m = C_\phi D$.

Recently, there have been some interests in studying some particular cases of operators, such as DC_ϕ , $C_\phi D$ and $D_{\phi,u}^m$ between different function spaces. From those studies, they gave some sufficient and necessary conditions for these operators to be bounded and compact. Concerning these results, we also recommend the interested readers to ([3-9]).

In this paper, we characterize the boundedness and compactness of the operator $D_{\phi,u}^m$ from A_α^p to n th weighted space. For the case of $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$, we have the following results:

Theorem 1. Assume that $p > 0$, $\alpha > -1$, $n, m \in \mathbb{N}$, μ is a weight on \mathbb{D} , ϕ is a non-constant analytic self-map of \mathbb{D} , and $u \in H(\mathbb{D})$. Then,

(1a) $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is bounded if and only if for each $k \in \{0, 1, \dots, n\}$

$$\sup_{z \in \mathbb{D}} \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right|}{(1 - |\phi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} < \infty. \tag{1}$$

(1b) $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is compact if and only if $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is bounded and for each $k \in \{0, 1, \dots, n\}$

$$\lim_{|\phi(z)| \rightarrow 1} \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right|}{(1 - |\phi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} = 0. \tag{2}$$

For the case of $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_{\mu,0}^{(n)}$, our main results are the following:

Theorem 2. Assume that $p > 0$, $\alpha > -1$, $n, m \in \mathbb{N}$, μ is a weight on \mathbb{D} , ϕ is a non-constant analytic self-map of \mathbb{D} , and $u \in H(\mathbb{D})$. Then,

(2a) $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_{\mu,0}^{(n)}$ is bounded if and only if $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is bounded and for each $k \in \{0, 1, \dots, n\}$,

$$\lim_{|z| \rightarrow 1} \mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right| = 0. \tag{3}$$

(2b) $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_{\mu,0}^{(n)}$ is compact if and only if $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_{\mu,0}^{(n)}$ is bounded and for each $k \in \{0, 1, \dots, n\}$,

$$\lim_{|z| \rightarrow 1} \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|}{(1 - |\varphi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} = 0. \tag{4}$$

The organization of this paper is as follows: we give some lemmas in Section 2, and then prove Theorem 1 in Section 3 and Theorem 2 in Section 4, respectively.

Throughout this paper, we will use the symbol C to denote a finite positive number, and it may differ from one occurrence to the other.

2. Some Lemmas

Lemma 1 is a direct consequence of the well-known estimate in ([10], Proposition 1.4.10). Hence, we omit its proof.

Lemma 1. *Assume that $p > 0$, $\alpha > -1$, $n \in \mathbb{N}$, $n > 0$, and $w \in \mathbb{D}$. Then the function*

$$g_{w,n}(z) = \frac{(1 - |w|^2)^n}{(1 - \bar{w}z)^{n+\frac{2+\alpha}{p}}}$$

belongs to A_α^p . Moreover, $\sup_{w \in \mathbb{D}} \|g_{w,n}\|_{A_\alpha^p} < \infty$.

The next lemma comes from ([11]).

Lemma 2. *Assume that $p > 0$, $\alpha > -1$, $n \in \mathbb{N}$, and $z \in \mathbb{D}$. Then, there is a positive constant C independent of f such that*

$$|f^{(n)}(z)| \leq C \frac{\|f\|_{A_\alpha^p}}{(1 - |z|^2)^{n+\frac{2+\alpha}{p}}}.$$

Lemma 3. *Let $p > 0$, $\alpha > -1$, $m \in \mathbb{N}$, $a = m + 1 + \frac{\alpha+2}{p}$ and*

$$D_{n+1} = \begin{vmatrix} 1 & 1 & \dots & 1 \\ a & a+1 & \dots & a+n \\ \prod_{j=0}^{n-1} (a+j) & \prod_{j=0}^{n-1} (a+j+1) & \dots & \prod_{j=0}^{n-1} (a+j+n) \end{vmatrix}.$$

Then, $D_{n+1} = \prod_{j=1}^n j!$.

Proof. With $a = m + 1 + \frac{\alpha+2}{p}$ and replacing n by $n + 1$ in ([12], Lemma 2.3), the lemma easily follows. \square

The next lemma can be found in ([7], Lemma 4).

Lemma 4. *Assume $n \in \mathbb{N}$, $g, u \in H(\mathbb{D})$ and ϕ is an analytic self-map of \mathbb{D} . Then,*

$$(u(z)g(\varphi(z)))^{(n)} = \sum_{k=0}^n g^{(k)}(\varphi(z)) \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)),$$

where

$$B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) = \sum_{k_1, \dots, k_l} \frac{l!}{k_1! \dots k_l!} \prod_{j=1}^l \left(\frac{\varphi^{(j)}(z)}{j!} \right)^{k_j}, \tag{5}$$

and the sum in (5) is over all non-negative integers k_1, \dots, k_l satisfying $k_1 + k_2 + \dots + k_l = k$ and $k_1 + 2k_2 + \dots + lk_l = l$.

By a proof in a standard way ([1], Proposition 3.11), we can get the next lemma.

Lemma 5. Suppose $u \in H(\mathbb{D})$, $p > 0$, $\alpha > -1$, $n, m \in \mathbb{N}$ and ϕ is a non-constant analytic self-map of \mathbb{D} . Then the operator $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is compact if and only if $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_{\mu,0}^{(n)}$ is bounded and for any bounded sequence $\{f_k\}_{k \in \mathbb{N}}$ in A_α^p which converges to zero uniformly on compact subsets of \mathbb{D} as $k \rightarrow \infty$, we have $D_{\phi,u}^m f_k \rightarrow 0$ in $\omega_\mu^{(n)}$ as $k \rightarrow \infty$.

Lemma 6. Suppose $n \in \mathbb{N}$ and μ is a radial weight such that $\lim_{|z| \rightarrow 1} \mu(z) = 0$. A closed set K in $\omega_{\mu,0}^{(n)}$ is compact if and only if it is bounded and satisfies

$$\lim_{|z| \rightarrow 1} \sup_{f \in K} \mu(z) |f^{(n)}(z)| = 0.$$

Proof. The proof of this Lemma is followed by standard arguments similar to those outlined in ([13]). We omit the details. \square

3. The Proof of Theorem 1

(1a) Boundedness of $D_{\phi,u}^m$.

We will prove the sufficiency first. Suppose that the conditions in (1) hold. Then, for any $f \in A_\alpha^p$, from Lemma 2 and Lemma 4, we obtain

$$\begin{aligned} & \mu(z) \left| (D_{\phi,u}^m f)^{(n)}(z) \right| \\ &= \mu(z) \left| \sum_{k=0}^n f^{(k+m)}(\phi(z)) \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right| \\ &\leq \mu(z) \sum_{k=0}^n \left| f^{(k+m)}(\phi(z)) \right| \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right| \quad (6) \\ &\leq C \|f\|_{A_\alpha^p} \sum_{k=0}^n \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right|}{(1 - |\phi(z)|^2)^{k+m+\frac{\alpha+2}{p}}}. \end{aligned}$$

For $z = 0$ and every $d \in \{0, 1, \dots, n - 1\}$,

$$\begin{aligned} & \left| (D_{\phi,u}^m f)^{(d)}(0) \right| \\ &= \left| \sum_{k=0}^d f^{(k+m)}(\phi(0)) \sum_{l=k}^d C_d^l u^{(d-l)}(0) B_{l,k}(\phi'(0), \phi''(0), \dots, \phi^{(l-k+1)}(0)) \right| \quad (7) \\ &\leq C \|f\|_{A_\alpha^p} \sum_{k=0}^d \frac{\left| \sum_{l=k}^d C_d^l u^{(d-l)}(0) B_{l,k}(\phi'(0), \phi''(0), \dots, \phi^{(l-k+1)}(0)) \right|}{(1 - |\phi(0)|^2)^{k+m+\frac{\alpha+2}{p}}}. \end{aligned}$$

From (1), (6), and (7), we know that $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is bounded.

Conversely, suppose that $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is bounded. Then, there exists a constant C such that $\|D_{\phi,u}^m f\|_{\omega_\mu^{(n)}} \leq C \|f\|_{A_\alpha^p}$ for all $f \in A_\alpha^p$.

For a fixed $w \in \mathbb{D}$, and constants c_1, c_2, \dots, c_{n+1} , set

$$g_w(z) = \sum_{j=1}^{n+1} \frac{c_j}{\left(j + \frac{2+\alpha}{p}\right) \left(j + \frac{2+\alpha}{p} + 1\right) \dots \left(j + \frac{2+\alpha}{p} + m - 1\right)} \frac{(1 - |w|^2)^j}{(1 - \bar{w}z)^{j + \frac{2+\alpha}{p}}}. \quad (8)$$

Applying Lemma 1 and triangle inequality, it is easy to get that $g_w \in A_\alpha^p$ for every $w \in \mathbb{D}$. Moreover, we have that

$$\sup_{w \in \mathbb{D}} \|g_w\|_{A_\alpha^p} < \infty. \tag{9}$$

Now we show that for each $s \in \{m, m + 1, \dots, m + n\}$, there are constants c_1, c_2, \dots, c_{n+1} , such that,

$$g_w^{(s)}(w) = \frac{\overline{w}^s}{(1 - |w|^2)^{s + \frac{\alpha+2}{p}}}, \quad g_w^{(t)}(w) = 0, \quad t \in \{m, \dots, m + n\} \setminus \{s\} \tag{10}$$

Indeed, by differentiating function g_w for each $s \in \{m, m + 1, \dots, m + n\}$, the system in (10) becomes

$$\begin{aligned} c_1 + c_2 + \dots + c_{n+1} &= 0 \\ (m + 1 + \frac{\alpha+2}{p})c_1 + (m + 2 + \frac{\alpha+2}{p})c_2 + \dots + (m + n + 1 + \frac{\alpha+2}{p})c_{n+1} &= 0 \\ \dots\dots\dots \\ (m + 1 + \frac{\alpha+2}{p}) \dots (s + \frac{\alpha+2}{p})c_1 + \dots + (m + n + 1 + \frac{\alpha+2}{p}) \dots (n + s + \frac{\alpha+2}{p})c_{n+1} &= 1 \\ \dots\dots\dots \\ (m + 1 + \frac{\alpha+2}{p}) \dots (m + n + \frac{\alpha+2}{p})c_1 + \dots + (m + n + 1 + \frac{\alpha+2}{p}) \dots (m + 2n + \frac{\alpha+2}{p})c_{n+1} &= 0 \end{aligned} \tag{11}$$

By Lemma 3, the determinant of system (11) is different from zero, which implies the statement. For each $k \in \{0, 1, 2, \dots, n\}$, we choose the corresponding family of functions that satisfy (10) with $s = m + k$ and denote it by $g_{w,k}$. For each fixed $k \in \{0, 1, \dots, n\}$, the boundedness of the operator $D_{\varphi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$, along with Lemma 4 and (8) implies that for each $\phi(w) \neq 0$,

$$\begin{aligned} &\frac{\mu(w) |\varphi(w)|^{m+k} \left| \sum_{l=k}^n C_n^l u^{(n-l)}(w) B_{l,k}(\varphi'(w), \varphi''(w), \dots, \varphi^{(l-k+1)}(w)) \right|}{(1 - |\varphi(w)|^2)^{k+m+\frac{\alpha+2}{p}}} \\ &\leq C \sup_{w \in \mathbb{D}} \|D_{\varphi,u}^m(g_{\varphi(w),k})\|_{\omega_\mu^{(n)}} \leq C \|D_{\varphi,u}^m\|_{A_\alpha^p \rightarrow \omega_\mu^{(n)}}. \end{aligned} \tag{12}$$

From (12), it follows that for each $k \in \{0, 1, \dots, n\}$,

$$\begin{aligned} &\sup_{|\varphi(z)| > \frac{1}{2}} \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|}{(1 - |\varphi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} \\ &\leq C \|D_{\varphi,u}^m\|_{A_\alpha^p \rightarrow \omega_\mu^{(n)}}. \end{aligned} \tag{13}$$

Now we use the test functions

$$h_k(z) = z^{k+m}, \quad k = 0, 1, \dots, n.$$

For each $k \in \mathbb{N}$, it is easy to get that

$$h_k \in A_\alpha^p, \quad \|h_k\|_{A_\alpha^p} \leq 1. \tag{14}$$

By applying Lemma 4 to the $h_0(z) = z^m$, we get

$$\begin{aligned} &(D_{\varphi,u}^m h_0)^{(n)}(z) \\ &= h_0^{(m)}(\varphi(z)) \sum_{l=0}^n C_n^l u^{(n-l)}(z) B_{l,0}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l+1)}(z)) \\ &= m! \sum_{l=0}^n C_n^l u^{(n-l)}(z) B_{l,0}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l+1)}(z)), \end{aligned}$$

which along with boundedness of the operator $D_{\varphi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ implies that

$$m! \sup_{z \in \mathbb{D}} \mu(z) \left| \sum_{l=0}^n C_n^l u^{(n-l)}(z) B_{l,0}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l+1)}(z)) \right| \leq C \|D_{\varphi,u}^m\|_{A_\alpha^p \rightarrow \omega_\mu^{(n)}}. \quad (15)$$

Assume now that we have proved the following inequalities

$$\sup_{z \in \mathbb{D}} \mu(z) \left| \sum_{l=j}^n C_n^l u^{(n-l)}(z) B_{l,j}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-j+1)}(z)) \right| \leq C \|D_{\varphi,u}^m\|_{A_\alpha^p \rightarrow \omega_\mu^{(n)}}. \quad (16)$$

for $j \in \{0, 1, \dots, k-1\}$, $k \leq n$.

Apply Lemma 4 to the $h_k(z) = z^{m+k}$, and knowing that $z^{(s)} \equiv 0$ for $s > m+k$ and the boundedness of the operator $D_{\varphi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$, we get

$$\begin{aligned} & \left| (D_{\varphi,u}^m h_k)^{(n)}(z) \right| \\ & \geq \left| \sum_{j=0}^{k-1} (m+k) \dots (k-j+1) (\varphi(z))^{(k-j)} \sum_{l=j}^n C_n^l u^{(n-l)}(z) B_{l,j}(\varphi'(z), \dots, \varphi^{(l-j+1)}(z)) \right| \\ & \quad - \left| (m+k)! \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \dots, \varphi^{(l-k+1)}(z)) \right|. \end{aligned} \quad (17)$$

Using hypothesis (16), we can know that

$$\sup_{z \in \mathbb{D}} \mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right| \leq C \|D_{\varphi,u}^m\|_{A_\alpha^p \rightarrow \omega_\mu^{(n)}}. \quad (18)$$

for each $k \in \{0, 1, \dots, n\}$. Then, for each $k \in \{0, 1, \dots, n\}$,

$$\begin{aligned} & \sup_{|\varphi(z)| \leq \frac{1}{2}} \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|}{(1 - |\varphi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} \\ & \leq C \sup_{z \in D} \mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|. \\ & \leq C \|D_{\varphi,u}^m\|_{A_\alpha^p \rightarrow \omega_\mu^{(n)}} \end{aligned} \quad (19)$$

From (13) and (19), we know that (1) holds.

(1b) Compactness of $D_{\varphi,u}^m$.

Suppose $D_{\varphi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is bounded and (2) holds. Then, by (1a), (1) holds. Let $(f_i)_{i \in \mathbb{N}}$ be a sequence in A_α^p such that, $\sup_{i \in \mathbb{N}} \|f_i\|_{A_\alpha^p} \leq M$ and f_i converges to 0 uniformly on compact subsets of \mathbb{D} as $i \rightarrow \infty$. By the assumption, for any $\varepsilon > 0$, there is a $\delta \in (0, 1)$, such that, for each $k \in \{0, 1, \dots, n\}$ and $\delta < |\phi(z)| < 1$,

$$\frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|}{(1 - |\varphi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} < \varepsilon. \quad (20)$$

From Lemma 2 and (20), we have

$$\begin{aligned} & \sup_{i \in \mathbb{N}} \mu(z) \left| \sum_{k=0}^n f_i^{(k+m)}(\varphi(z)) \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \dots, \varphi^{(l-k+1)}(z)) \right| \\ & \leq C \sup_{i \in \mathbb{N}} \|f_i\|_{A_\alpha^p} \sum_{k=0}^n \sup_{z \in \mathbb{D}} \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|}{(1 - |\varphi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} \quad (21) \\ & \leq CM(n+1)\varepsilon. \end{aligned}$$

If $|\phi(z)| \leq r$, then by Cauchy's estimate and (19), we have

$$\begin{aligned} & \sup \mu(z) \left| \sum_{k=0}^n f_i^{(k+m)}(\varphi(z)) \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \dots, \varphi^{(l-k+1)}(z)) \right| \quad (22) \\ & \leq C \sum_{k=0}^n \sup_{|\varphi(z)| \leq r} |f_i^{(m+k)}(\varphi(z))| \rightarrow 0, (i \rightarrow \infty) \end{aligned}$$

For $j = 0, 1, \dots, n - 1$, we have

$$\left| (D_{\varphi, u}^m f_i)^{(j)}(0) \right| \rightarrow 0, (i \rightarrow \infty) \quad (23)$$

Applying (21), (22), and (23), we know that $\|D_{\varphi, u}^m f_i\|_{\omega_\mu^{(n)}} \rightarrow 0, (i \rightarrow \infty)$. From Lemma 5, $D_{\varphi, u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is compact.

Conversely, suppose that $D_{\varphi, u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is compact, then $D_{\varphi, u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is bounded. Let $(z_i)_{i \in \mathbb{N}}$ be a sequence in \mathbb{D} such that $|\phi(z_i)| \rightarrow 1, i \rightarrow \infty$. If such a sequence does not exist, then the condition in (2) is easily satisfied. Now, assume that when $\|\phi\|_\infty = 1$ and (2) does not hold, then there is $k \in \{0, 1, \dots, n\}$ and $\delta > 0$ such that

$$\frac{\mu(z_i) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z_i) B_{l,k}(\varphi'(z_i), \varphi''(z_i), \dots, \varphi^{(l-k+1)}(z_i)) \right|}{(1 - |\varphi(z_i)|^2)^{k+m+\frac{\alpha+2}{p}}} \geq \delta.$$

Let $g_i(z) = g_{\varphi(z_i), k}(z), i \in \mathbb{N}, k \in \{0, 1, \dots, n\}$ be as in Theorem 1. Then, $\sup_{i \in \mathbb{N}} \|g_i\|_{A_\alpha^p} \leq M$ and $g_i \rightarrow 0$ uniformly on compact subsets of \mathbb{D} as $i \rightarrow \infty$. By the assumption and Lemma 5, we have that for $k \in \{0, 1, \dots, n\}$

$$\lim_{i \rightarrow \infty} \|D_{\varphi, u}^m g_{\varphi(z_i), k}\|_{\omega_\mu^{(n)}} = 0. \quad (24)$$

On the other hand, from (12), we obtain

$$\begin{aligned} & \|D_{\varphi, u}^m g_{\varphi(z_i), k}\|_{\omega_\mu^{(n)}} \\ & \geq \frac{\mu(z_i) |\varphi(z_i)|^{k+m} \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z_i) B_{l,k}(\varphi'(z_i), \varphi''(z_i), \dots, \varphi^{(l-k+1)}(z_i)) \right|}{(1 - |\varphi(z_i)|^2)^{k+m+\frac{\alpha+2}{p}}} \quad (25) \\ & > \frac{\delta}{3} \end{aligned}$$

for large enough i . From (24) and (25), this is a contradiction. So, (2) holds.

Now the proof of Theorem 1 is completed.

4. The Proof of Theorem 2

(2a) Boundedness of $D_{\varphi,u}^m$.

First, suppose that $D_{\varphi,u}^m : A_{\alpha}^p \rightarrow \omega_{\mu}^{(n)}$ is bounded and (3) holds. For each polynomial $p(z)$, we obtain $|p^{(m+k)}(z)| \leq C_p, z \in D, C_p$ is a constant depending on p .

And

$$\begin{aligned} & \mu(z) \left| (D_{\varphi,u}^m p)^{(n)}(z) \right| \\ &= \mu(z) \left| \sum_{k=0}^n p^{(k+m)}(\varphi(z)) \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \dots, \varphi^{(l-k+1)}(z)) \right| \\ &\leq C_p \sum_{k=0}^n |\mu(z)| \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \dots, \varphi^{(l-k+1)}(z)) \right| \\ &\rightarrow 0, |z| \rightarrow 1. \end{aligned} \tag{26}$$

From (26), we have that for each polynomial $p(z), D_{\varphi,u}^m p \in \omega_{\mu,0}^{(n)}$. Since the set of polynomials is dense in A_{α}^p , we have that for each $f \in A_{\alpha}^p$, there is a sequence of polynomials $(p_k)_{k \in \mathbb{N}}$ such that $\|f - p_k\|_{A_{\alpha}^p} \rightarrow 0$ as $k \rightarrow \infty$. From the boundedness of $D_{\varphi,u}^m : A_{\alpha}^p \rightarrow \omega_{\mu}^{(n)}$, we have that

$$\|D_{\varphi,u}^m f - D_{\varphi,u}^m p_k\|_{\omega_{\mu}^{(n)}} \leq \|D_{\varphi,u}^m\|_{A_{\alpha}^p \rightarrow \omega_{\mu}^{(n)}} \|f - p_k\|_{A_{\alpha}^p} \rightarrow 0, k \rightarrow \infty.$$

Then, $D_{\varphi,u}^m f \in \omega_{\mu,0}^{(n)}$ from which the boundedness of $D_{\varphi,u}^m : A_{\alpha}^p \rightarrow \omega_{\mu,0}^{(n)}$ follows.

Conversely, suppose that $D_{\varphi,u}^m : A_{\alpha}^p \rightarrow \omega_{\mu,0}^{(n)}$ is bounded. It is clear that $D_{\varphi,u}^m : A_{\alpha}^p \rightarrow \omega_{\mu}^{(n)}$ is bounded. Then, taking the test functions $h_k(z) = z^{m+k}$ for each $k \in \{0, 1, \dots, n\}$, we obtain $D_{\varphi,u}^m z^{m+k} \in \omega_{\mu,0}^{(n)}$. By the proof of Theorem 1, for each $k \in \{0, 1, \dots, n\}$,

$$\lim_{|z| \rightarrow 1} \mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right| = 0.$$

(2a) is completed.

(2b) Compactness of $D_{\varphi,u}^m$.

First, assume that $D_{\varphi,u}^m : A_{\alpha}^p \rightarrow \omega_{\mu,0}^{(n)}$ is compact, so it is bounded and (3) holds. Hence, if $\|\phi\|_{\infty} < 1$,

$$\begin{aligned} & \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|}{\left(1 - |\varphi(z)|^2\right)^{k+m+\frac{\alpha+2}{p}}} \\ &\leq \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\varphi'(z), \varphi''(z), \dots, \varphi^{(l-k+1)}(z)) \right|}{\left(1 - \|\varphi\|_{\infty}^2\right)^{k+m+\frac{\alpha+2}{p}}} \\ &\rightarrow 0, (|z| \rightarrow 1) \end{aligned} \tag{27}$$

If $\|\phi\|_\infty = 1$, since $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_\mu^{(n)}$ is compact too and (2) holds, then for all $\varepsilon > 0$, there is an $r \in (0, 1)$, such that when $r < |\phi(z)| < 1$, for $k \in \{0, 1, \dots, n\}$,

$$\frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right|}{(1 - |\phi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} < \varepsilon. \tag{28}$$

By (3), we know there is a $\delta \in (0, 1)$, such that $\delta < |z| < 1$, for $k \in \{0, 1, \dots, n\}$,

$$\begin{aligned} & \mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right| \\ & < \varepsilon (1 - r^2)^{k+m+\frac{\alpha+2}{p}}. \end{aligned} \tag{29}$$

Then, when $\delta < |z| < 1$ and $r < |\phi(z)| < 1$ for $k \in \{0, 1, \dots, n\}$, we get

$$\frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right|}{(1 - |\phi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} < \varepsilon. \tag{30}$$

In addition, when $|\phi(z)| \leq r$ and $\delta < |z| < 1$, we have

$$\begin{aligned} & \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right|}{(1 - |\phi(z)|^2)^{k+m+\frac{\alpha+2}{p}}} \\ & < \frac{\mu(z) \left| \sum_{l=k}^n C_n^l u^{(n-l)}(z) B_{l,k}(\phi'(z), \phi''(z), \dots, \phi^{(l-k+1)}(z)) \right|}{(1 - r^2)^{k+m+\frac{\alpha+2}{p}}} < \varepsilon. \end{aligned} \tag{31}$$

Combining (30) and (31), we know (4) holds.

Conversely, assume $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_{\mu,0}^{(n)}$ is bounded and (4) holds. Taking the supremum in (6) for all f in the unit ball of $A_{\phi'}^p$ and using the condition (4), we have $\lim_{|z| \rightarrow 1} \sup_{\|f\|_{A_\alpha^p} \leq 1} \mu(z) \left| (D_{\phi,u}^m f)^{(n)}(z) \right| = 0$, from which by Lemma 6, the compactness of $D_{\phi,u}^m : A_\alpha^p \rightarrow \omega_{\mu,0}^{(n)}$ follows.

Now the proof of Theorem 2 is finished.

Acknowledgements

We are grateful to the referee(s) for many helpful comments on the manuscript. Hong-Gang Zeng is supported in part by the National Natural Science Foundation of China (Grand Nos. 10971153, 10671141).

Authors' contributions

LZ found the question and drafted the manuscript. HGZ joined in the discussion about the question and revised the paper. All authors read and approved the final manuscript

Competing interests

The authors declare that they have no competing interests.

Received: 29 January 2011 Accepted: 21 September 2011 Published: 21 September 2011

References

1. Cowen, CC, Maccluer, BD: Composition Operators on Spaces of Analytic Functions. CRC Press, Boca Raton, FL (1995)
2. Zhu, K: Operator Theory in Function Spaces. Marcel Dekker, New York (1990)
3. Hibscheweiler, RA, Portnoy, N: Composition followed by differentiation between Bergman and Hardy spaces. The Rocky Mountain Journal of Mathematics. **35**(3), 843-855 (2005). doi:10.1216/rmj/1181069709
4. Li, S, Stević, S: Composition followed by differentiation between Bloch-type spaces. J Comput Anal Appl. **9**(2), 195-205 (2007)

5. Ohno, S: Products of composition and differentiation between Hardy spaces. *Bull Aust Math Soc.* **73**(2), 235–243 (2006). doi:10.1017/S0004972700038818
6. Stević, S: Products of composition and differentiation operators on the weighted Bergman space. *Bull Belg Math Soc Simon Stevin.* **16**(4), 623–635 (2009)
7. Stević, S: Weighted differentiation composition operators from H^∞ and Bloch spaces to n th weighted-type spaces on the unit disk. *Appl Math Comput.* **216**(12), 3634–3641 (2010). doi:10.1016/j.amc.2010.05.014
8. Yang, WF: Products of composition and differentiation operators from $Q_k(p, q)$ spaces to Bloch-Type Spaces. *Abst Appl Anal* **2009**, 14. Article ID 741920
9. Zhou, XL: Products of differentiation, composition and multiplication from Bergman type spaces to Bers type spaces. *Int Trans Special Funct.* **18**(3), 223–231 (2007). doi:10.1080/10652460701210250
10. Rudin, W: *Function Theory in the Unit ball of C^n* , vol. 241 of *Fundamental Principles of Mathematical Science*. Springer, New York, USA (1980)
11. Flett, TM: The dual of an inequality and Littlewood and some related inequalities. *J Math Anal Appl.* **38**, 746–765 (1972). doi:10.1016/0022-247X(72)90081-9
12. Stević, S: composition operators from the weighted Bergman space to the n th weighted-type space on the unit disk. *Hindawi Publ Corp Discret Dyn Nat Soc* **2009**, 11. Article ID 742019
13. Madigan, K, Matheson, A: Compact composition operators on the Bloch Space. *Trans Am Math Soc.* **347**(7), 2679–2687 (1995). doi:10.2307/2154848

doi:10.1186/1029-242X-2011-65

Cite this article as: Zhang and Zeng: Weighted differentiation composition operators from weighted bergman space to n th weighted space on the unit disk. *Journal of Inequalities and Applications* 2011 **2011**:65.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
