

RESEARCH

Open Access

Positive solutions of fractional differential equations at resonance on the half-line

Yi Chen and Xianhua Tang*

*Correspondence:
tangxh@mail.csu.edu.cn
School of Mathematical Sciences
and Computing Technology,
Central South University, Changsha,
Hunan 410083, P.R. China

Abstract

This article deals with the differential equations of fractional order on the half-line. By the recent Leggett-Williams norm-type theorem due to O'Regan and Zima, we present some new results on the existence of positive solutions for the fractional boundary value problems at resonance on unbounded domains.

MSC: 26A33; 34A08; 34A34

Keywords: fractional order; half-line; coincidence degree; at resonance

1 Introduction

In this article, we are concerned with the fractional differential equation

$$\begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)), & t \in [0, +\infty), \\ u(0) = u'(0) = u''(0) = 0, & D_{0+}^{\alpha-1} u(0) = \lim_{t \rightarrow +\infty} D_{0+}^{\alpha-1} u(t), \end{cases} \quad (1.1)$$

where D_{0+}^{α} is the Riemann-Liouville fractional derivative, $3 < \alpha < 4$, and $f : [0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies the following condition:

(H) $f : [0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and for each $l > 0$, there exists $\phi_l \in C[0, +\infty) \cap L^1[0, +\infty)$ satisfying $\sup_{t \geq 0} |\phi_l(t)| < +\infty$ and $\phi_l(t) > 0$, $t > 0$ such that

$$|u| < l \text{ implies } |f(t, (1 + t^{\alpha-1})u)| \leq \phi_l(t), \quad \text{a.e. } t \geq 0.$$

The problem (1.1) happens to be at resonance in the sense that the kernel of the linear operator D_{0+}^{α} is not less than one-dimensional under the boundary value conditions.

Fractional calculus is a generalization of the ordinary differentiation and integration. It has played a significant role in science, engineering, economy, and other fields. Some books on fractional calculus and fractional differential equations have appeared recently (see [1–3]); furthermore, today there is a large number of articles dealing with the fractional differential equations (see [4–15]) due to their various applications.

In [8], the researchers dealt with the existence of solutions for boundary value problems of fractional order of the form

$$\begin{aligned} {}^C D_{0+}^{\alpha} y(t) &= f(t, y(t)), & t \in [0, +\infty), \\ y(0) &= y_0, & y \text{ is bounded in } [0, +\infty), \end{aligned}$$

where $1 < \alpha \leq 2$ and $f : [0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous. The results are based on the fixed point theorem of Schauder combined with the diagonalization method.

In [9], Su and Zhang studied the following fractional differential equations on the half-line using Schauder's fixed point theorem

$$\begin{aligned} D_{0+}^{\alpha} u(t) &= f(t, u(t), D_{0+}^{\alpha-1} u(t)), \quad t \in (0, +\infty), 1 < \alpha \leq 2, \\ u(0) &= 0, \quad \lim_{t \rightarrow \infty} D_{0+}^{\alpha-1} u(t) = u_{\infty}. \end{aligned}$$

Employing the Leray-Schauder alternative theorem, in [12], Zhao and Ge considered the fractional boundary value problem

$$\begin{aligned} D_{0+}^{\alpha} u(t) + f(t, u(t)) &= 0, \quad t \in (0, +\infty), 1 < \alpha < 2, \\ u(0) &= 0, \quad \lim_{t \rightarrow \infty} D_{0+}^{\alpha-1} u(t) = \beta u(\xi). \end{aligned}$$

However, the articles on the existence of solutions of fractional differential equations on the half-line are still few, and most of them deal with the problems under nonresonance conditions. And as far as we know, recent articles, such as [4, 6, 7], investigating resonant problems are on the finite interval.

Motivated by the articles [16–20], in this article we study the differential equations (1.1) under resonance conditions on the unbounded domains. Moreover, we have successfully established the existence theorem by the recent Leggett-Williams norm-type theorem due to O'Regan and Zima. To our best knowledge, there is no article dealing with the resonant problems of fractional order on unbounded domains by the theorem.

The rest of the article is organized as follows. In Section 2, we give the definitions of the fractional integral and fractional derivative, some results about fractional differential equations, and the abstract existence theorem. In Section 3, we obtain the existence result of the solution for the problem (1.1) by the recent Leggett-Williams norm-type theorem. Then, an example is given in Section 4 to demonstrate the application of our result.

2 Preliminaries

First of all, we present some fundamental facts on the fractional calculus theory which we will use in the next section.

Definition 2.1 ([1–3]) The Riemann-Liouville fractional integral of order $\nu > 0$ of a function $h : (0, \infty) \rightarrow \mathbb{R}$ is given by

$$I_{0+}^{\nu} h(t) = D_{0+}^{-\nu} h(t) = \frac{1}{\Gamma(\nu)} \int_0^t (t-s)^{\nu-1} h(s) ds, \tag{2.1}$$

provided that the right-hand side is pointwise defined on $(0, \infty)$.

Definition 2.2 ([1–3]) The Riemann-Liouville fractional derivative of order $\nu > 0$ of a continuous function $h : (0, \infty) \rightarrow \mathbb{R}$ is given by

$$D_{0+}^{\nu} h(t) = \frac{1}{\Gamma(n-\nu)} \left(\frac{d}{dt} \right)^n \int_0^t (t-s)^{n-\nu-1} h(s) ds, \tag{2.2}$$

where $n = [\nu] + 1$, provided that the right-hand side is pointwise defined on $(0, \infty)$.

Lemma 2.1 ([1, 9]) *Assume that $h(t) \in L^1(0, +\infty)$. If $v_1, v_2, \nu > 0$, then*

$$I_{0+}^{\nu_1} I_{0+}^{\nu_2} h(t) = I_{0+}^{\nu_1 + \nu_2} h(t), \quad D_{0+}^{\nu} I_{0+}^{\nu} h(t) = h(t). \tag{2.3}$$

Lemma 2.2 ([9]) *Assume that $D_{0+}^{\nu} h(t) \in L^1(0, +\infty)$, $\nu > 0$. Then we have*

$$I_{0+}^{\nu} D_{0+}^{\nu} h(t) = h(t) + C_1 t^{\nu-1} + C_2 t^{\nu-2} + \dots + C_N t^{\nu-N}, \quad t > 0, \tag{2.4}$$

for some $C_i \in \mathbb{R}$, $i = 1, 2, \dots, N$, where N is the smallest integer greater than or equal to ν .

Now, let us recall some standard facts and the fixed point theorem due to O'Regan and Zima, and these can be found in [16, 17, 21–23].

Let X, Z be real Banach spaces. Consider an operation equation

$$Lu = Nu,$$

where $L : \text{dom } L \subset X \rightarrow Z$ is a linear operator, $N : X \rightarrow Z$ is a nonlinear operator. If $\dim \text{Ker } L = \text{codim } \text{Im } L < +\infty$ and $\text{Im } L$ is closed in Z , then L is called a Fredholm mapping of index zero. And if L is a Fredholm mapping of index zero, there exist linear continuous projectors $P : X \rightarrow X$ and $Q : Z \rightarrow Z$ such that $\text{Ker } L = \text{Im } P$, $\text{Im } L = \text{Ker } Q$ and $X = \text{Ker } L \oplus \text{Ker } P$, $Z = \text{Im } L \oplus \text{Im } Q$. Then it follows that $L_P = L|_{\text{dom } L \cap \text{Ker } P} : \text{dom } L \cap \text{Ker } P \rightarrow \text{Im } L$ is invertible. We denote the inverse of this map by K_P . For $\text{Im } Q$ is isomorphic to $\text{Ker } L$, there exists an isomorphism $J : \text{Im } Q \rightarrow \text{Ker } L$.

It is known that the coincidence equation $Lu = Nu$ is equivalent to

$$u = (P + JQN)u + K_P(I - Q)Nu.$$

A nonempty convex closed set $C \subset X$ is called a cone if

- (i) $\kappa x \in C$ for all $x \in C$ and $\kappa \geq 0$;
- (ii) $x, -x \in C$ implies $x = 0$.

Note that C induces a partial order \leq in X by

$$x \leq y \quad \text{if and only if} \quad y - x \in C.$$

The following lemma is valid for every cone in a Banach space.

Lemma 2.3 ([17, 23]) *Let C be a cone in the Banach space X . Then for every $u \in C \setminus \{0\}$, there exists a positive number $\sigma(u)$ such that*

$$\|x + u\| \geq \sigma(u)\|x\|,$$

for all $x \in C$.

Let $\gamma : X \rightarrow C$ be a retraction, i.e., a continuous mapping such that $\gamma(x) = x$ for all $x \in C$. Denote

$$\Psi := P + JQN + K_P(I - Q)N,$$

and

$$\Psi_\gamma := \Psi \circ \gamma.$$

Theorem 2.1 ([16, 17]) *Let C be a cone in X and let Ω_1, Ω_2 be open bounded subsets of X with $\overline{\Omega}_1 \subset \Omega_2$ and $C \cap (\overline{\Omega}_2 \setminus \Omega_1) \neq \emptyset$. Assume that:*

- 1° L is a Fredholm operator of index zero;
- 2° $QN : X \rightarrow Z$ is continuous and bounded and $K_P(I - Q)N : X \rightarrow X$ is compact on every bounded subset of X ;
- 3° $Lu \neq \lambda Nu$ for all $u \in C \cap \partial\Omega_2 \cap \text{dom } L$ and $\lambda \in (0, 1)$;
- 4° γ maps subsets of $\overline{\Omega}_2$ into bounded subsets of C ;
- 5° $d_B([I - (P + JQN)\gamma]|_{\text{Ker } L}, \text{Ker } L \cap \Omega_2, 0) \neq 0$, where d_B stands for the Brouwer degree;
- 6° there exists $u_0 \in C \setminus \{0\}$ such that $\|u\| \leq \sigma(u_0)\|\Psi u\|$ for $u \in C(u_0) \cap \partial\Omega_1$, where $C(u_0) = \{u \in C : \mu u_0 \leq u \text{ for some } \mu > 0\}$ and $\sigma(u_0)$ is such that $\|u + u_0\| \geq \sigma(u_0)\|u\|$ for every $u \in C$;
- 7° $(P + JQN)\gamma(\partial\Omega_2) \subset C$;
- 8° $\Psi_\gamma(\overline{\Omega}_2 \setminus \Omega_1) \subset C$.

Then the equation $Lx = Nx$ has a solution in the set $C \cap (\overline{\Omega}_2 \setminus \Omega_1)$.

Let

$$X = \left\{ x \mid x \in C[0, +\infty), \lim_{t \rightarrow +\infty} \frac{x(t)}{1 + t^{\alpha-1}} \text{ exists} \right\}$$

with the norm

$$\|x\|_X = \sup_{t \geq 0} \frac{|x(t)|}{1 + t^{\alpha-1}},$$

and

$$Z = \left\{ z \mid z \in C[0, +\infty) \cap L^1[0, +\infty), \sup_{t \geq 0} |z(t)| < +\infty \right\},$$

equipped with the norm

$$\|z\|_Z = \sup_{t \geq 0} |z(t)| + \int_0^{+\infty} |z(t)| dt.$$

Remark 2.1 It is easy for us to prove that $(X, \|\cdot\|_X)$ and $(Z, \|\cdot\|_Z)$ are Banach spaces.

Set

$$\text{dom } L = \left\{ u \in X \mid D_{0+}^\alpha u(t) \in C[0, +\infty) \cap L^1[0, +\infty), u(0) = u'(0) = u''(0) = 0, \right. \\ \left. D_{0+}^{\alpha-1} u(0) = \lim_{t \rightarrow +\infty} D_{0+}^{\alpha-1} u(t) \right\}.$$

Define

$$L : \text{dom } L \rightarrow Z, \quad u \rightarrow D_{0+}^\alpha u(t), \tag{2.5}$$

and

$$N : X \rightarrow Z, \quad u \rightarrow f(t, u(t)). \tag{2.6}$$

Then the multi-point boundary value problem (1.1) can be written by

$$Lu = Nu, \quad u \in \text{dom } L.$$

Definition 2.3 $u \in X$ is called a solution of the problem (1.1) if $u \in \text{dom } L$ and u satisfied Equation (1.1).

Next, similar to the compactness criterion in [12, 24], we establish the following criterion, and it can be proved in a similar way.

Lemma 2.4 \mathcal{U} is a relatively compact set in X if and only if the following conditions are satisfied:

- (a) \mathcal{U} is uniformly bounded, that is, there exists a constant $R > 0$ such that for each $u \in \mathcal{U}$, $\|u\|_X \leq R$.
- (b) The functions from \mathcal{U} are equicontinuous on any compact subinterval of $[0, +\infty)$, that is, let J be a compact subinterval of $[0, +\infty)$, then $\forall \varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that for $t_1, t_2 \in J$, $|t_1 - t_2| < \delta$,

$$\left| \frac{u(t_1)}{1 + t_1^{\alpha-1}} - \frac{u(t_2)}{1 + t_2^{\alpha-1}} \right| < \varepsilon, \quad \forall u \in \mathcal{U}.$$

- (c) The functions from \mathcal{U} are equiconvergent, that is, given $\varepsilon > 0$, there exists $T = T(\varepsilon) > 0$ such that

$$\left| \frac{u(s_1)}{1 + s_1^{\alpha-1}} - \frac{u(s_2)}{1 + s_2^{\alpha-1}} \right| < \varepsilon,$$

for $s_1, s_2 > T, \forall u \in \mathcal{U}$.

3 Main results

In this section, we will present the existence theorem for the fractional differential equation on the half-line. In order to prove our main result, we need the following lemmas.

Lemma 3.1 Let $g \in Z$. Then $u \in X$ is the solution of the following fractional differential equation:

$$\begin{cases} D_{0+}^\alpha u(t) = g(t), & t \in [0, +\infty), \\ u(0) = u'(0) = u''(0) = 0, & D_{0+}^{\alpha-1} u(0) = \lim_{t \rightarrow +\infty} D_{0+}^{\alpha-1} u(t), \end{cases}$$

if and only if

$$u(t) = ct^{\alpha-1} + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g(s) ds, \quad c \in \mathbb{R},$$

and

$$\int_0^{+\infty} g(t) dt = 0.$$

Proof In view of Lemmas 2.1 and 2.2, we can certify the conclusion easily, so we omit the details here. \square

Lemma 3.2 *The operator L is a Fredholm mapping of index zero. Moreover,*

$$\text{Ker } L = \{u | u = ct^{\alpha-1}, t \geq 0, c \in \mathbb{R}\} \subset X, \tag{3.1}$$

and

$$\text{Im } L = \left\{ g \in Z \mid \int_0^{+\infty} g(t) dt = 0 \right\} \subset Z. \tag{3.2}$$

Proof It is obvious that Lemma 3.1 implies (3.1) and (3.2). Now, let us focus our minds on proving that L is a Fredholm mapping of index zero.

Define $Q : Z \rightarrow Z$

$$(Qg)(t) = e^{-t} \int_0^{+\infty} g(s) ds, \quad t \geq 0, \tag{3.3}$$

where $g \in Z$. Evidently, $\text{Ker } Q = \text{Im } L$, $\text{Im } Q = \{g | g = ce^{-t}, t \geq 0, c \in \mathbb{R}\}$, and $Q : Z \rightarrow Z$ is a continuous linear projector. In fact, for an arbitrary $g \in Z$, we have

$$Q^2g = Q(Qg) = Q\left(e^{-t} \int_0^{+\infty} g(s) ds\right) = Q(e^{-t}) \int_0^{+\infty} g(s) ds = e^{-t} \int_0^{+\infty} g(s) ds = Qg,$$

that is to say, $Q : Z \rightarrow Z$ is idempotent.

Let $g = g - Qg + Qg = (I - Q)g + Qg$, where $g \in Z$ is an arbitrary element. Since $Qg \in \text{Im } Q$ and $(I - Q)g \in \text{Ker } Q$, we obtain that $Z = \text{Im } Q + \text{Ker } Q$. Take $z_0 \in \text{Im } Q \cap \text{Ker } Q$, then z_0 can be written as $z_0 = ce^{-t}$, $c \in \mathbb{R}$, for $z_0 \in \text{Im } Q$. Since $z_0 \in \text{Ker } Q = \text{Im } L$, by (3.2), we get that $Q(z_0) = Q(ce^{-t}) = cQ(e^{-t}) = ce^{-t} = 0$, which implies that $c = 0$, and then $z_0 = 0$. Therefore, $\text{Im } Q \cap \text{Ker } Q = \{0\}$, thus, $Z = \text{Im } Q \oplus \text{Ker } Q = \text{Im } Q \oplus \text{Im } L$.

Now, $\dim \text{Ker } L = 1 = \dim \text{Im } Q = \text{codim } \text{Ker } Q = \text{codim } \text{Im } L < +\infty$, and observing that $\text{Im } L$ is closed in Z , so L is a Fredholm mapping of index zero. \square

Let $P : X \rightarrow X$ be defined by

$$(Pu)(t) = \left(\frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} u(s) ds \right) t^{\alpha-1}, \quad t \geq 0, u \in X. \tag{3.4}$$

It is clear that $P : X \rightarrow X$ is a linear continuous projector and

$$\text{Im } P = \{u | u = ct^{\alpha-1}, t \geq 0, c \in \mathbb{R}\} = \text{Ker } L.$$

Also, proceeding with the proof of Lemma 3.2, we can show that $X = \text{Im } P \oplus \text{Ker } P = \text{Ker } L \oplus \text{Ker } P$.

Consider the mapping $K_P : \text{Im } L \rightarrow \text{dom } L \cap \text{Ker } P$

$$(K_P g)(t) = -\left(\frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} g(s) ds\right) t^{\alpha-1} + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g(s) ds, \quad g \in \text{Im } L.$$

Note that

$$(K_P L)u = K_P(Lu) = u, \quad \forall u \in \text{dom } L \cap \text{Ker } P,$$

and

$$(LK_P)g = L(K_P g) = g, \quad \forall g \in \text{Im } L.$$

Thus, $K_P = (L_P)^{-1}$, where $L_P = L|_{\text{dom } L \cap \text{Ker } P} : \text{dom } L \cap \text{Ker } P \rightarrow \text{Im } P$.

Define the linear isomorphism $J : \text{Im } Q \rightarrow \text{Ker } L$ as

$$J(ce^{-t}) = ct^{\alpha-1}, \quad t \geq 0, c \in \mathbb{R}.$$

Thus, $JQN + K_P(I - Q)N : X \rightarrow X$ is given by

$$[JQN + K_P(I - Q)N]u(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)} \int_0^{+\infty} G(t,s)f(s,u(s)) ds, \quad t \geq 0, \tag{3.5}$$

where

$$G(t,s) = \begin{cases} 0, & t = 0; \\ \Gamma(\alpha) + \frac{1}{2} - e^{-s} - \int_0^t \frac{(t-\tau)^{\alpha-1}}{t^{\alpha-1}} e^{-\tau} d\tau + \frac{(t-s)^{\alpha-1}}{t^{\alpha-1}}, & t \neq 0 \text{ and } 0 \leq s \leq t < +\infty; \\ \Gamma(\alpha) + \frac{1}{2} - e^{-s} - \int_0^t \frac{(t-\tau)^{\alpha-1}}{t^{\alpha-1}} e^{-\tau} d\tau, & 0 < t \leq s < +\infty. \end{cases}$$

Then, it is easy to verify that

$$0 < \Gamma(\alpha) - \frac{1}{2} \leq G(t,s) \leq \Gamma(\alpha) + \frac{3}{2}. \tag{3.6}$$

Now, we state the main result on the existence of the positive solutions to the problem (1.1) in the following.

Theorem 3.1 *Let $f : [0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$ satisfy the condition (H). Assume that there exist six nonnegative functions $\alpha_i(t)$ ($i = 1, 2, 3$), $\beta_j(t)$ ($j = 1, 2$) and $\mu(t)$ such that*

$$f(t,u) \leq -\alpha_1(t)|f(t,u)| + \alpha_2(t) \frac{u}{1+t^{\alpha-1}} + \alpha_3(t), \quad t \geq 0, \tag{3.7}$$

and

$$-\mu(t) \frac{u}{1+t^{\alpha-1}} \leq f(t,u) \leq -\beta_1(t) \frac{u}{1+t^{\alpha-1}} + \beta_2(t), \quad t \geq 0, \tag{3.8}$$

where $0 \leq \frac{u}{1+t^{\alpha-1}} \leq R$, $R > R_0$, and R_0 is defined by (3.12), $\alpha_1(t)$ is bounded on $[0, +\infty)$, $\beta_1(t) > 0$, $t \geq 0$, $\alpha_2(t), \alpha_3(t), \beta_1(t), \beta_2(t) \in L^1[0, +\infty)$,

$$\alpha_0 := \inf_{t \geq 0} \alpha_1(t) > 0, \quad \int_0^{+\infty} \alpha_3(t) dt > 0, \quad \int_0^{+\infty} \beta_2(t) dt > 0, \quad (3.9)$$

$$\Gamma_0 := \frac{2}{\alpha_0} \sup_{t \geq 0} \frac{\alpha_2(t) + \alpha_0 e^{-t}(1 + t^{\alpha-1})/2}{\beta_1(t)} < +\infty, \quad (3.10)$$

and

$$\int_0^{+\infty} \mu(t) dt < \frac{\Gamma(\alpha) + 1}{2(\Gamma(\alpha) + 1/2)(\Gamma(\alpha) + 3/2)}, \quad e^t \mu(t) < \frac{1 + t^{\alpha-1}}{\Gamma(\alpha) + 3/2}. \quad (3.11)$$

Then the problem (1.1) has at least one positive solution in $\text{dom } L$.

Proof For the simplicity of notation, we denote

$$\varepsilon_1 := \frac{\Gamma(\alpha)}{\Gamma(\alpha) + 1/2} + \frac{\Gamma(\alpha) + 3/2}{\Gamma(\alpha) + 1} \int_0^{+\infty} \mu(s) ds < 1, \quad \beta_0 := \int_0^{+\infty} \frac{s^{\alpha-1} \beta_1(s)}{1 + s^{\alpha-1}} ds,$$

and

$$R_0 := \max \left\{ \frac{\Gamma_0}{\Gamma(\alpha)} \int_0^{+\infty} \beta_2(s) ds + \frac{2}{\alpha_0 \Gamma(\alpha)} \int_0^{+\infty} \alpha_3(s) ds, \frac{1}{\beta_0} \int_0^{+\infty} \beta_2(s) ds \right\}. \quad (3.12)$$

Consider the cone

$$C = \{u | u \in X, u(t) \geq 0, t \geq 0\}.$$

Set

$$\Omega_1 = \left\{ u \in X \mid \varepsilon_0 \|u\|_X < \frac{u(t)}{1 + t^{\alpha-1}} < R_1, t \geq 0 \right\}, \quad \Omega_2 = \{u \in X \mid \|u\|_X < R_2, t \geq 0\},$$

where $R_2 \in (R_0, R)$, $R_1 \in (0, R_2)$, $\varepsilon_0 \in (\varepsilon_1, 1)$. Clearly, Ω_1 and Ω_2 are an open bounded set of X .

Step 1: In view of Lemma 3.2, the condition 1° of Theorem 2.1 is fulfilled.

Step 2: By virtue of Lemma 2.4, we can get that $QN : X \rightarrow Z$ is continuous and bounded and $K_P(I - Q)N : X \rightarrow X$ is compact on every bounded subset of X , which ensures that the assumption 2° of Theorem 2.1 holds.

Step 3: Suppose that there exist $u^* \in C \cap \partial\Omega_2 \cap \text{dom } L$ and $\lambda^* \in (0, 1)$ such that $Lu^* = \lambda^* Nu^*$.

Since

$$u^* = (I - P)u^* + Pu^* = K_P L(I - P)u^* + Pu^* = K_P Lu^* + Pu^*,$$

we have

$$\frac{u^*(t)}{1 + t^{\alpha-1}} = -\frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} D_{0+}^\alpha u^*(s) ds \cdot \frac{t^{\alpha-1}}{1 + t^{\alpha-1}} + \frac{1}{\Gamma(\alpha)} \int_0^t \frac{(t-s)^{\alpha-1}}{1 + t^{\alpha-1}} D_{0+}^\alpha u^*(s) ds$$

$$\begin{aligned}
 & + \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} u^*(s) ds \cdot \frac{t^{\alpha-1}}{1+t^{\alpha-1}} \\
 & < \frac{2}{\Gamma(\alpha)} \int_0^{+\infty} |D_{0+}^\alpha u^*(s)| ds + \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} u^*(s) ds.
 \end{aligned} \tag{3.13}$$

From (3.7) and (3.8), we get that

$$\begin{aligned}
 D_{0+}^\alpha u^*(t) = \lambda^* f(t, u^*(t)) & \leq -\lambda^* \alpha_1(t) |f(t, u^*(t))| + \lambda^* \alpha_2(t) \frac{u^*(t)}{1+t^{\alpha-1}} + \lambda^* \alpha_3(t) \\
 & \leq -\alpha_1(t) |D_{0+}^\alpha u^*(t)| + \alpha_2(t) \frac{u^*(t)}{1+t^{\alpha-1}} + \alpha_3(t),
 \end{aligned} \tag{3.14}$$

and

$$D_{0+}^\alpha u^*(t) = \lambda^* f(t, u^*(t)) \leq -\lambda^* \beta_1(t) \frac{u^*(t)}{1+t^{\alpha-1}} + \lambda^* \beta_2(t). \tag{3.15}$$

On account of the fact that

$$\int_0^{+\infty} D_{0+}^\alpha u^*(s) ds = \int_0^{+\infty} D(D_{0+}^{\alpha-1} u^*(s)) ds = \lim_{t \rightarrow +\infty} D_{0+}^{\alpha-1} u^*(t) - D_{0+}^{\alpha-1} u^*(0) = 0,$$

and considering (3.14) and (3.15), we have

$$\begin{aligned}
 0 & = \int_0^{+\infty} D_{0+}^\alpha u^*(s) ds \\
 & \leq - \int_0^{+\infty} \alpha_1(s) |D_{0+}^\alpha u^*(s)| ds + \int_0^{+\infty} \alpha_2(s) \frac{u^*(s)}{1+s^{\alpha-1}} ds + \int_0^{+\infty} \alpha_3(s) ds,
 \end{aligned}$$

and

$$0 = \int_0^{+\infty} D_{0+}^\alpha u^*(s) ds \leq - \int_0^{+\infty} \lambda^* \beta_1(s) \frac{u^*(s)}{1+s^{\alpha-1}} ds + \int_0^{+\infty} \lambda^* \beta_2(s) ds.$$

Thus,

$$\int_0^{+\infty} |D_{0+}^\alpha u^*(s)| ds \leq \frac{1}{\alpha_0} \int_0^{+\infty} \alpha_2(s) \frac{u^*(s)}{1+s^{\alpha-1}} ds + \frac{1}{\alpha_0} \int_0^{+\infty} \alpha_3(s) ds,$$

and

$$\int_0^{+\infty} \beta_1(s) \frac{u^*(s)}{1+s^{\alpha-1}} ds \leq \int_0^{+\infty} \beta_2(s) ds.$$

By (3.9), (3.10) and (3.13), we obtain that

$$\begin{aligned}
 & \frac{u^*(t)}{1+t^{\alpha-1}} \\
 & < \frac{2}{\Gamma(\alpha)} \int_0^{+\infty} |D_{0+}^\alpha u^*(s)| ds + \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} u^*(s) ds \\
 & \leq \frac{2}{\alpha_0 \Gamma(\alpha)} \int_0^{+\infty} \alpha_2(s) \frac{u^*(s)}{1+s^{\alpha-1}} ds + \frac{2}{\alpha_0 \Gamma(\alpha)} \int_0^{+\infty} \alpha_3(s) ds + \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} u^*(s) ds
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} \left(\frac{2}{\alpha_0} \frac{\alpha_2(s) + \alpha_0 e^{-s}(1 + s^{\alpha-1})/2}{\beta_1(s)} \right) \beta_1(s) \frac{u^*(s)}{1 + s^{\alpha-1}} ds + \frac{2}{\alpha_0 \Gamma(\alpha)} \int_0^{+\infty} \alpha_3(s) ds \\
 &\leq \frac{\Gamma_0}{\Gamma(\alpha)} \int_0^{+\infty} \beta_1(s) \frac{u^*(s)}{1 + s^{\alpha-1}} ds + \frac{2}{\alpha_0 \Gamma(\alpha)} \int_0^{+\infty} \alpha_3(s) ds \\
 &\leq \frac{\Gamma_0}{\Gamma(\alpha)} \int_0^{+\infty} \beta_2(s) ds + \frac{2}{\alpha_0 \Gamma(\alpha)} \int_0^{+\infty} \alpha_3(s) ds \leq R_0 < R_2,
 \end{aligned}$$

which is a contradiction to $u^* \in C \cap \partial\Omega_2 \cap \text{dom}L$. Therefore, 3° is satisfied.

Step 4: Let $(\gamma u)(t) = |u(t)|$, then we can verify that $\gamma : X \rightarrow C$ is a retraction and 4° holds.

Step 5: Let $u \in \text{Ker}L \cap \overline{\Omega}_2$, then $u(t) = ct^{\alpha-1}$, $t \geq 0$, $c \in \mathbb{R}$. Inspired by Aijun and Wang [5], we set

$$H(ct^{\alpha-1}, \rho) = [I - \rho(P + JQN)\gamma](ct^{\alpha-1}) = \left(c - \rho|c| - \rho \int_0^{+\infty} f(s, |c|s^{\alpha-1}) ds \right) t^{\alpha-1},$$

where $c \in [-R_2, R_2]$ and $\rho \in [0, 1]$.

Define homeomorphism $J_1 : \text{Ker}L \cap \overline{\Omega}_2 \rightarrow \mathbb{R}$ by $J_1(ct^{\alpha-1}) = c$, then

$$\begin{aligned}
 d_B(H(ct^{\alpha-1}, \rho), \text{Ker}L \cap \Omega_2, 0) &= d_B(J_1H(J_1^{-1}c, \rho), J_1(\text{Ker}L \cap \Omega_2), J_1(0)) \\
 &= d_B(J_1H(J_1^{-1}c, \rho), J_1(\text{Ker}L \cap \Omega_2), 0).
 \end{aligned}$$

It is obvious that $J_1H(J_1^{-1}c, \rho) = 0$ implies that $c \geq 0$ by (3.8) and (3.11).

Take $c_0 \in J_1(\text{Ker}L \cap \partial\Omega_2)$, then $|c_0| = R_2$. Suppose that $J_1H(J_1^{-1}c, \rho) = 0$, $\rho \in (0, 1]$, then we have that $c_0 = R_2$. Also, in view of (3.8),

$$\begin{aligned}
 R_2 &= \rho \left(R_2 - \int_0^{+\infty} f(s, R_2 s^{\alpha-1}) ds \right) \\
 &\leq \rho \left(R_2 + R_2 \int_0^{+\infty} \beta_1(s) \frac{s^{\alpha-1}}{1 + s^{\alpha-1}} ds + \int_0^{+\infty} \beta_2(s) ds \right) \\
 &< \rho R_2 \leq R_2.
 \end{aligned}$$

It is a contradiction. Besides, if $\rho = 0$, then $R_2 = 0$, which is impossible. Hence, for $c \in J_1(\text{Ker}L \cap \partial\Omega_2)$, $J_1H(J_1^{-1}c, \rho) \neq 0$, $\rho \in [0, 1]$.

Therefore,

$$\begin{aligned}
 d_B([I - (P + JQN)\gamma]|_{\text{Ker}L}, \text{Ker}L \cap \Omega_2, 0) &= d_B(H(\cdot, 1), \text{Ker}L \cap \Omega_2, 0) \\
 &= d_B(J_1H(J_1^{-1}c, 1), J_1(\text{Ker}L \cap \Omega_2), 0) \\
 &= d_B(J_1H(J_1^{-1}c, 0), J_1(\text{Ker}L \cap \Omega_2), 0) \\
 &= d_B(I, J_1(\text{Ker}L \cap \Omega_2), 0) = 1 \neq 0,
 \end{aligned}$$

which shows that 5° is true.

Step 6: Let $u_0 = 1 + t^{\alpha-1} \in C \setminus \{0\}$, then we have

$$C(u_0) = \left\{ u \in C \mid \inf_{t \geq 0} \frac{u(t)}{1 + t^{\alpha-1}} > 0 \right\}.$$

And we can take $\sigma(u_0) = 1$.

Let $t_0 > 0$ such that

$$\frac{t_0^{\alpha-1}}{1+t_0^{\alpha-1}} > \frac{\Gamma(\alpha)+1/2}{\Gamma(\alpha)+1}.$$

For $u \in C(u_0) \cap \partial\Omega_1$, we have that

$$\|u\|_X \leq R_1 < R_2, \quad \frac{u(t)}{1+t^{\alpha-1}} \geq \varepsilon_0 \|u\|_X.$$

Therefore, combining (3.6), (3.8) and (3.11), we get that

$$\begin{aligned} & \frac{(\Psi u)(t_0)}{1+t_0^{\alpha-1}} \\ &= \frac{1}{1+t_0^{\alpha-1}} (Pu(t_0) + [JQN + K_P(I - Q)N]u(t_0)) \\ &= \frac{t_0^{\alpha-1}}{1+t_0^{\alpha-1}} \left(\frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} u(s) ds + \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} G(t_0, s) f(s, u(s)) ds \right) \\ &> \frac{\Gamma(\alpha)+1/2}{\Gamma(\alpha)+1} \left(\varepsilon_0 \|u\|_X \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} (1+s^{\alpha-1}) ds - \frac{\|u\|_X}{\Gamma(\alpha)} \int_0^{+\infty} G(t_0, s) \mu(s) ds \right) \\ &\geq \|u\|_X \frac{\Gamma(\alpha)+1/2}{\Gamma(\alpha)+1} \left(\varepsilon_0 \frac{\Gamma(\alpha)+1}{\Gamma(\alpha)} - \frac{\Gamma(\alpha)+3/2}{\Gamma(\alpha)} \int_0^{+\infty} \mu(s) ds \right) \\ &= \|u\|_X. \end{aligned}$$

Thus, $\|u\|_X \leq \sigma(u_0) \|\Psi u\|_X$ for all $u \in C(u_0) \cap \partial\Omega_1$. So, 6° holds.

Step 7: For $u \in \partial\Omega_2$, from (3.8) and (3.11), we have

$$\begin{aligned} (P + JQN)(\gamma u)(t) &= \left(\frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} |u(s)| ds + \int_0^{+\infty} f(s, |u(s)|) ds \right) t^{\alpha-1} \\ &\geq \frac{t^{\alpha-1}}{\Gamma(\alpha)} \int_0^{+\infty} [e^{-s} (1+s^{\alpha-1}) - \mu(s)] \frac{|u(s)|}{1+s^{\alpha-1}} ds \geq 0, \end{aligned}$$

which implies that $(P + JQN)\gamma(\partial\Omega_2) \subset C$. Hence, 7° holds.

Step 8: For $u \in \overline{\Omega_2} \setminus \Omega_1$, by (3.6), (3.8) and (3.11), we obtain that

$$\begin{aligned} \Psi_\gamma u(t) &= [P + JQN + K_P(I - Q)N] |u(t)| \\ &= \left(\frac{1}{\Gamma(\alpha)} \int_0^{+\infty} e^{-s} |u(s)| ds + \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} G(t, s) f(s, |u(s)|) ds \right) t^{\alpha-1} \\ &\geq \frac{t^{\alpha-1}}{\Gamma(\alpha)} \left(\int_0^{+\infty} [e^{-s} (1+s^{\alpha-1}) - G(t, s) \mu(s)] \frac{|u(s)|}{1+s^{\alpha-1}} ds \right) \\ &\geq \frac{t^{\alpha-1}}{\Gamma(\alpha)} \left(\int_0^{+\infty} \left[e^{-s} (1+s^{\alpha-1}) - \left(\Gamma(\alpha) + \frac{3}{2} \right) \mu(s) \right] \frac{|u(s)|}{1+s^{\alpha-1}} ds \right) \\ &\geq 0. \end{aligned}$$

Thus, $\Psi_\gamma(\overline{\Omega_2} \setminus \Omega_1) \subset C$, that is, 8° is satisfied.

Hence, applying Theorem 2.1, the problem (1.1) has a positive solution in the set $C \cap (\overline{\Omega_2} \setminus \Omega_1)$. □

4 Examples

To illustrate our main result, we will present an example.

Example 4.1

$$\begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)), & t \in [0, +\infty), \\ u(0) = u'(0) = u''(0) = 0, & D_{0+}^{\alpha-1} u(0) = \lim_{t \rightarrow +\infty} D_{0+}^{\alpha-1} u(t), \end{cases} \quad (4.1)$$

where $\alpha = 3.5$, and for $(t, u) \in \mathbb{R}^2$,

$$f(t, u) = -\beta_1(t) \frac{u}{1 + t^{\alpha-1}} + \beta_2(t),$$

and

$$\beta_1(t) = \frac{1}{40} e^{-t} (1 + t^{\alpha-1}), \quad \beta_2(t) = \frac{1}{1 + t^2}.$$

It is easy for us to certify that f satisfies the condition (H).

Noting that

$$f(t, u) \leq -\alpha_1(t) |f(t, u)| + \alpha_2(t) \frac{u}{1 + t^{\alpha-1}} + \alpha_3(t), \quad t \geq 0,$$

and

$$-\mu(t) \frac{u}{1 + t^{\alpha-1}} \leq f(t, u) \leq -\beta_1(t) \frac{u}{1 + t^{\alpha-1}} + \beta_2(t), \quad t \geq 0,$$

for $u \geq 0$, where

$$\alpha_1(t) = 2, \quad \alpha_2(t) = \beta_1(t), \quad \alpha_3(t) = 3\beta_2(t), \quad \mu(t) = \beta_1(t).$$

Evidently, $\mu(t)$ satisfies (3.11).

Meanwhile, by simple computation we can get that

$$\alpha_0 = 2, \quad \int_0^{+\infty} \alpha_3(t) dt = \frac{3\pi}{2}, \quad \int_0^{+\infty} \beta_2(t) dt = \frac{\pi}{2}, \quad \Gamma_0 = 41.$$

Thus, to sum up the points which we have just indicated, by Theorem 3.1, we can conclude that the problem (4.1) has at least one positive solution.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All the authors typed, read, and approved the final manuscript.

Acknowledgement

This project is supported by the Hunan Provincial Innovation Foundation For Postgraduate (NO. CX2011B079) and the National Natural Science Foundation of China (NO. 11171351).

References

1. Kilbas, AA, Srivastava, HM, Trujillo, JJ: Theory and Applications of Fractional Differential Equations. Elsevier, Amsterdam (2006)
2. Miller, KS, Ross, B: An Introduction to the Fractional Calculus and Fractional Differential Equations. Wiley, New York (1993)
3. Lakshmikantham, V, Leela, S, Vasundhara Devi, J: Theory of Fractional Dynamic Systems. Cambridge Academic Publishers, Cambridge (2009)
4. Bai, Z, Zhang, Y: The existence of solutions for a fractional multi-point boundary value problem. *Comput. Math. Appl.* **60**, 2364-2372 (2010)
5. Yang, A, Wang, H: Positive solutions of two-point boundary value problems of nonlinear fractional differential equation at resonance. *Electron. J. Qual. Theory Differ. Equ.* **71**, 1-15 (2011)
6. Jiang, W: The existence of solutions to boundary value problems of fractional differential equations at resonance. *Nonlinear Anal.* **74**, 1987-1994 (2011)
7. Bai, Z: Solvability for a class of fractional-point boundary value problem at resonance. *Comput. Math. Appl.* **62**, 1292-1302 (2011)
8. Arara, A, Benchohra, M, Hamidi, N, Nieto, JJ: Fractional order differential equations on an unbounded domain. *Nonlinear Anal.* **72**, 580-586 (2010)
9. Su, X, Zhang, S: Unbounded solutions to a boundary value problem of fractional order on the half-line. *Comput. Math. Appl.* **61**, 1079-1087 (2011)
10. Kou, C, Zhou, H, Yan, Y: Existence of solutions of initial value problems for nonlinear fractional differential equations on the half-axis. *Nonlinear Anal.* **74**, 5975-5986 (2011)
11. Liang, S, Zhang, J: Existence of three positive solutions of m -point boundary value problems for some nonlinear fractional differential equations on an infinite interval. *Comput. Math. Appl.* **61**, 3343-3354 (2011)
12. Zhao, X, Ge, W: Unbounded solutions for a fractional boundary value problems on the infinite interval. *Acta Appl. Math.* **109**, 495-505 (2010)
13. Yang, L, Chen, H: Unique positive solution for boundary value problem of fractional differential equations. *Appl. Math. Lett.* **23**, 1095-1098 (2010)
14. Chai, G: Positive solutions for boundary value problem of fractional differential equation with p -Laplacian operator. *Bound. Value Probl.* **2012**, 18 (2012)
15. Ahmad, B, Nieto, JJ: Riemann-Liouville fractional integro-differential equations with fractional nonlocal integral boundary conditions. *Bound. Value Probl.* **2011**, 36 (2011)
16. O'Regan, D, Zima, M: Leggett-Williams norm-type theorems for coincidences. *Arch. Math.* **87**, 233-244 (2006)
17. Franco, D, Infante, G, Zima, M: Second order nonlocal boundary value problems at resonance. *Math. Nachr.* **284**, 875-884 (2011)
18. Infante, G, Zima, M: Positive solutions of multi-point boundary value problems at resonance. *Nonlinear Anal.* **69**, 2458-2465 (2008)
19. Yang, L, Shen, C: On the existence of positive solution for a kind of multi-point boundary value problem at resonance. *Nonlinear Anal.* **72**, 4211-4220 (2010)
20. Yang, L, Shen, C: Positive solutions for second order four-point boundary value problems at resonance. *Topol. Methods Nonlinear Anal.* **38**, 1-16 (2011)
21. Mawhin, J: Equivalence theorems for nonlinear operator equations and coincidence degree theory for some mappings in locally convex topological vector spaces. *J. Differ. Equ.* **12**, 610-636 (1972)
22. Santanilla, J: Some coincidence theorems in wedges, cones, and convex sets. *J. Math. Anal. Appl.* **105**, 357-371 (1985)
23. Petryshyn, WV: On the solvability of $x \in Tx + \lambda Fx$ in quasiconvex cones with T and F k -set contractive. *Nonlinear Anal.* **5**, 589-591 (1981)
24. Agarwal, RP, O'Regan, D: Infinite Interval Problems for Differential, Difference and Integral Equations. Kluwer Academic, Dordrecht (2001)

doi:10.1186/1687-2770-2012-64

Cite this article as: Chen and Tang: Positive solutions of fractional differential equations at resonance on the half-line. *Boundary Value Problems* 2012 **2012**:64.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com