

RESEARCH

Open Access

Growth properties of Green-Sch potentials at infinity

Tao Zhao¹ and Alexander Yamada Jr.^{2*}

*Correspondence:

yamadaayu71@yahoo.com

²Matematiska Institutionen,
Stockholms Universitet, Stockholm,
106 91, SwedenFull list of author information is
available at the end of the article**Abstract**

This paper gives growth properties of Green-Sch potentials at infinity in a cone, which generalizes results obtained by Qiao-Deng. The proof is based on the fact that the estimations of Green-Sch potentials with measures are connected with a kind of densities of the measures modified by the measures.

MSC: 35J10; 35J25**Keywords:** stationary Schrödinger operator; Green-Sch potential; growth property; cone

1 Introduction and main results

Let \mathbf{R} and \mathbf{R}_+ be the set of all real numbers and the set of all positive real numbers, respectively. We denote by \mathbf{R}^n ($n \geq 2$) the n -dimensional Euclidean space. A point in \mathbf{R}^n is denoted by $P = (X, x_n)$, $X = (x_1, x_2, \dots, x_{n-1})$. The Euclidean distance of two points P and Q in \mathbf{R}^n is denoted by $|P - Q|$. Also $|P - O|$ with the origin O of \mathbf{R}^n is simply denoted by $|P|$. The boundary, the closure and the complement of a set S in \mathbf{R}^n are denoted by ∂S , \bar{S} , and S^c , respectively. For $P \in \mathbf{R}^n$ and $r > 0$, let $B(P, r)$ denote the open ball with center at P and radius r in \mathbf{R}^n .

We introduce a system of spherical coordinates (r, Θ) , $\Theta = (\theta_1, \theta_2, \dots, \theta_{n-1})$, in \mathbf{R}^n which are related to cartesian coordinates $(x_1, x_2, \dots, x_{n-1}, x_n)$ by

$$x_1 = r \left(\prod_{j=1}^{n-1} \sin \theta_j \right) \quad (n \geq 2), \quad x_n = r \cos \theta_1,$$

and if $n \geq 3$, then

$$x_{n-m+1} = r \left(\prod_{j=1}^{m-1} \sin \theta_j \right) \cos \theta_m \quad (2 \leq m \leq n-1),$$

where $0 \leq r < +\infty$, $-\frac{1}{2}\pi \leq \theta_{n-1} < \frac{3}{2}\pi$, and if $n \geq 3$, then $0 \leq \theta_j \leq \pi$ ($1 \leq j \leq n-2$).

The unit sphere and the upper half unit sphere in \mathbf{R}^n are denoted by S^{n-1} and S_+^{n-1} , respectively. For simplicity, a point $(1, \Theta)$ on S^{n-1} and the set $\{\Theta; (1, \Theta) \in \Omega\}$ for a set Ω , $\Omega \subset S^{n-1}$, are often identified with Θ and Ω , respectively. For two sets $\Xi \subset \mathbf{R}_+$ and $\Omega \subset S^{n-1}$, the set $\{(r, \Theta) \in \mathbf{R}^n; r \in \Xi, (1, \Theta) \in \Omega\}$ in \mathbf{R}^n is simply denoted by $\Xi \times \Omega$. In particular, the half space $\mathbf{R}_+ \times S_+^{n-1} = \{(X, x_n) \in \mathbf{R}^n; x_n > 0\}$ will be denoted by T_n .

By $C_n(\Omega)$, we denote the set $\mathbf{R}_+ \times \Omega$ in \mathbf{R}^n with the domain Ω on \mathbf{S}^{n-1} ($n \geq 2$). We call it a cone. Then T_n is a special cone obtained by putting $\Omega = \mathbf{S}_+^{n-1}$. We denote the sets $I \times \Omega$ and $I \times \partial\Omega$ with an interval on \mathbf{R} by $C_n(\Omega; I)$ and $S_n(\Omega; I)$. By $S_n(\Omega; r)$ we denote $C_n(\Omega) \cap S_r$. By $S_n(\Omega)$ we denote $S_n(\Omega; (0, +\infty))$, which is $\partial C_n(\Omega) - \{O\}$.

Let $C_n(\Omega)$ be an arbitrary domain in \mathbf{R}^n and \mathcal{A}_a denote the class of nonnegative radial potentials $a(P)$, i.e. $0 \leq a(P) = a(r)$, $P = (r, \Theta) \in C_n(\Omega)$, such that $a \in L_{loc}^b(C_n(\Omega))$ with some $b > n/2$ if $n \geq 4$ and with $b = 2$ if $n = 2$ or $n = 3$.

If $a \in \mathcal{A}_a$, then the stationary Schrödinger operator

$$Sch_a = -\Delta + a(P)I = 0,$$

where Δ is the Laplace operator and I is the identical operator, can be extended in the usual way from the space $C_0^\infty(C_n(\Omega))$ to an essentially self-adjoint operator on $L^2(C_n(\Omega))$ (see [1, Ch. 13]). We will denote it Sch_a as well. This last one has a Green-Sch function $G_\Omega^a(P, Q)$. Here $G_\Omega^a(P, Q)$ is positive on $C_n(\Omega)$ and its inner normal derivative $\partial G_\Omega^a(P, Q)/\partial n_Q \geq 0$, where $\partial/\partial n_Q$ denotes the differentiation at Q along the inward normal into $C_n(\Omega)$. We denote this derivative by $PI_\Omega^a(P, Q)$, which is called the Poisson-Sch kernel with respect to $C_n(\Omega)$.

We shall say that a set $E \subset C_n(\Omega)$ has a covering $\{r_j, R_j\}$ if there exists a sequence of balls $\{B_j\}$ with centers in $C_n(\Omega)$ such that $E \subset \bigcup_{j=0}^\infty B_j$, where r_j is the radius of B_j and R_j is the distance from the origin to the center of B_j .

For positive functions h_1 and h_2 , we say that $h_1 \lesssim h_2$ if $h_1 \leq Mh_2$ for some constant $M > 0$. If $h_1 \lesssim h_2$ and $h_2 \lesssim h_1$, we say that $h_1 \approx h_2$.

Let Ω be a domain on \mathbf{S}^{n-1} with smooth boundary. Consider the Dirichlet problem

$$\begin{aligned} (\Lambda_n + \lambda)\varphi &= 0 \quad \text{on } \Omega, \\ \varphi &= 0 \quad \text{on } \partial\Omega, \end{aligned}$$

where Λ_n is the spherical part of the Laplace opera Δ_n

$$\Delta_n = \frac{n-1}{r} \frac{\partial}{\partial r} + \frac{\partial^2}{\partial r^2} + \frac{\Lambda_n}{r^2}.$$

We denote the least positive eigenvalue of this boundary value problem by λ and the normalized positive eigenfunction corresponding to λ by $\varphi(\Theta)$, $\int_\Omega \varphi^2(\Theta) dS_1 = 1$. In order to ensure the existence of λ and a smooth $\varphi(\Theta)$. We put a rather strong assumption on Ω : if $n \geq 3$, then Ω is a $C^{2,\alpha}$ -domain ($0 < \alpha < 1$) on \mathbf{S}^{n-1} surrounded by a finite number of mutually disjoint closed hypersurfaces (e.g. see [2, pp.88-89] for the definition of $C^{2,\alpha}$ -domain).

For any $(1, \Theta) \in \Omega$, we have (see [3, pp.7-8])

$$\varphi(\Theta) \approx \text{dist}((1, \Theta), \partial C_n(\Omega)),$$

which yields

$$\delta(P) \approx r\varphi(\Theta), \tag{1.1}$$

where $P = (r, \Theta) \in C_n(\Omega)$ and $\delta(P) = \text{dist}(P, \partial C_n(\Omega))$.

Solutions of an ordinary differential equation

$$-Q''(r) - \frac{n-1}{r}Q'(r) + \left(\frac{\lambda}{r^2} + a(r)\right)Q(r) = 0, \quad 0 < r < \infty. \tag{1.2}$$

It is well known (see, for example, [4]) that if the potential $a \in \mathcal{A}_a$, then (1.2) has a fundamental system of positive solutions $\{V, W\}$ such that V is nondecreasing with (see [5–8])

$$0 \leq V(0+) \leq V(r) \quad \text{as } r \rightarrow +\infty,$$

and W is monotonically decreasing with

$$+\infty = W(0+) > W(r) \searrow 0 \quad \text{as } r \rightarrow +\infty.$$

We will also consider the class \mathcal{B}_a , consisting of the potentials $a \in \mathcal{A}_a$ such that there exists the finite limit $\lim_{r \rightarrow \infty} r^2 a(r) = k \in [0, \infty)$, and moreover, $r^{-1}|r^2 a(r) - k| \in L(1, \infty)$. If $a \in \mathcal{B}_a$, then the (sub)superfunctions are continuous (see [9]).

In the rest of paper, we assume that $a \in \mathcal{B}_a$ and we shall suppress this assumption for simplicity.

Denote

$$i_k^\pm = \frac{2 - n \pm \sqrt{(n-2)^2 + 4(k + \lambda)}}{2},$$

then the solutions to (1.2) have the asymptotic (see [10])

$$V(r) \approx r^{i_k^+}, \quad W(r) \approx r^{i_k^-} \quad \text{as } r \rightarrow \infty. \tag{1.3}$$

We denote the Green-Sch potential with a positive measure ν on $C_n(\Omega)$ by

$$G_\Omega^a \nu(P) = \int_{C_n(\Omega)} G_\Omega^a(P, Q) d\nu(Q).$$

Let ν be any positive measure $C_n(\Omega)$ such that $G_\Omega^a \nu(P) \not\equiv +\infty$ (resp. $G_\Omega^0 \nu(P) \not\equiv +\infty$) for $P \in C_n(\Omega)$. The positive measure ν' (rep. ν'') on \mathbf{R}^n is defined by

$$d\nu'(Q) = \begin{cases} W(t)\varphi(\Phi) d\nu(Q), & Q = (t, \Phi) \in C_n(\Omega; (1, +\infty)), \\ 0, & Q \in \mathbf{R}^n - C_n(\Omega; (1, +\infty)). \end{cases}$$

$$\left(d\nu''(Q) = \begin{cases} t^{i_0^-} \varphi(\Phi) d\nu(Q), & Q = (t, \Phi) \in C_n(\Omega; (1, +\infty)), \\ 0, & Q \in \mathbf{R}^n - C_n(\Omega; (1, +\infty)). \end{cases} \right)$$

Let $\epsilon > 0$, $0 \leq \alpha < n$, and λ be any positive measure on \mathbf{R}^n having finite total mass. For each $P = (r, \Theta) \in \mathbf{R}^n - \{O\}$, the maximal function $M(P; \lambda, \alpha)$ is defined by (see [11])

$$M(P; \lambda, \alpha) = \sup_{0 < \rho < \frac{r}{2}} \lambda(B(P, \rho)) V(\rho) W(\rho) \rho^{\alpha-2}.$$

The set

$$\{P = (r, \Theta) \in \mathbf{R}^n - \{O\}; M(P; \lambda, \alpha) V^{-1}(r) W^{-1}(r) r^{2-\alpha} > \epsilon\}$$

is denoted by $E(\epsilon; \lambda, \alpha)$.

Remark 1 If $\lambda(\{P\}) > 0$ ($P \neq O$), then $M(P; \lambda, \alpha) = +\infty$ for any positive number β . So we can find $\{P \in \mathbf{R}^n - \{O\}; \lambda(\{P\}) > 0\} \subset E(\epsilon; \lambda, \alpha)$.

About the growth properties of Green potentials at infinity in a cone, Qiao-Deng (see [12, Theorem 1]) has proved the following result.

Theorem A Let ν be a positive measure on $C_n(\Omega)$ such that $G_\Omega^0 \nu(P) \neq +\infty$ for any $P = (r, \Theta) \in C_n(\Omega)$. Then there exists a covering $\{r_j, R_j\}$ of $F(\epsilon; \nu'', \alpha) (\subset C_n(\Omega))$ satisfying

$$\sum_{j=0}^{\infty} \left(\frac{r_j}{R_j}\right)^{n-\alpha} < \infty,$$

such that

$$\lim_{r \rightarrow \infty, P \in C_n(\Omega) - F(\epsilon; \nu'', \alpha)} r^{-\alpha} \varphi^{\alpha-1}(\Theta) G_\Omega^0 \nu(P) = 0,$$

where

$$H(P; \nu'', \alpha) = \sup_{0 < \rho < \frac{r}{2}} \frac{\nu''(B(P, \rho))}{\rho^{n-\alpha}}$$

and

$$F(\epsilon; \nu'', \alpha) = \{P = (r, \Theta) \in \mathbf{R}^n - \{O\}; H(P; \nu'', \alpha) r^{n-\alpha} > \epsilon\}.$$

Now we state our first result.

Theorem 1 Let ν be a positive measure on $C_n(\Omega)$ such that

$$G_\Omega^a \nu(P) \neq +\infty \quad (P = (r, \Theta) \in C_n(\Omega)). \tag{1.4}$$

Then there exists a covering $\{r_j, R_j\}$ of $E(\epsilon; \nu', \alpha) (\subset C_n(\Omega))$ satisfying

$$\sum_{j=0}^{\infty} \left(\frac{r_j}{R_j}\right)^{2-\alpha} \frac{V(R_j) W(R_j)}{V(r_j) W(r_j)} < \infty, \tag{1.5}$$

such that

$$\lim_{r \rightarrow \infty, P \in C_n(\Omega) - E(\epsilon; \nu', \alpha)} V^{-1}(r) \varphi^{\alpha-1}(\Theta) G_\Omega^a \nu(P) = 0. \tag{1.6}$$

Remark 2 By comparison the condition (1.4) is fairly briefer and easily applied. Moreover, $E(\epsilon; \nu', 1)$ is a set of 1-finite view in the sense of [13, 14] (see [13, Definition 2.1] for the definition of 1-finite view). In the case $a = 0$, Theorem 1 (1.6) is just the result of Theorem A.

Corollary 1 Let ν be a positive measure on $C_n(\Omega)$ such that (1.4) holds. Then for a sufficiently large L and a sufficiently small ϵ we have

$$\{P \in C_n(\Omega; (L, +\infty)); G_\Omega^a \nu(P) \geq V(r)\varphi^{1-\alpha}(\Theta)\} \subset E(\epsilon; \mu', \alpha).$$

2 Some lemmas

Lemma 1 (see [15, 16])

$$G_\Omega^a(P, Q) \approx V(t)W(r)\varphi(\Theta)\varphi(\Phi) \tag{2.1}$$

$$\text{(resp. } G_\Omega^a(P, Q) \approx V(r)W(t)\varphi(\Theta)\varphi(\Phi)\text{)}, \tag{2.2}$$

for any $P = (r, \Theta) \in C_n(\Omega)$ and any $Q = (t, \Phi) \in C_n(\Omega)$ satisfying $0 < \frac{t}{r} \leq \frac{4}{5}$ (resp. $0 < \frac{r}{t} \leq \frac{4}{5}$);
 Further, for any $P = (r, \Theta) \in C_n(\Omega)$ and any $Q = (t, \Phi) \in C_n(\Omega; (\frac{4}{5}r, \frac{5}{4}r))$, we have

$$G_\Omega^0(P, Q) \lesssim \frac{\varphi(\Theta)\varphi(\Phi)}{t^{n-2}} + \Pi_\Omega(P, Q), \tag{2.3}$$

where

$$\Pi_\Omega(P, Q) = \min \left\{ \frac{1}{|P-Q|^{n-2}}, \frac{rt\varphi(\Theta)\varphi(\Phi)}{|P-Q|^n} \right\}.$$

Lemma 2 Let ν be a positive measure on $C_n(\Omega)$ such that there is a sequence of points $P_i = (r_i, \Theta_i) \in C_n(\Omega)$, $r_i \rightarrow +\infty$ ($i \rightarrow +\infty$) satisfying $G_\Omega^a \nu(P_i) < +\infty$ ($i = 1, 2, \dots; Q \in C_n(\Omega)$). Then, for a positive number l ,

$$\int_{C_n(\Omega; (l, +\infty))} W(t)\varphi(\Phi) d\nu(Q) < +\infty \tag{2.4}$$

and

$$\lim_{R \rightarrow +\infty} \frac{W(R)}{V(R)} \int_{C_n(\Omega; (0, R))} V(t)\varphi(\Phi) d\nu(Q) = 0. \tag{2.5}$$

Proof Take a positive number l satisfying $P_1 = (r_1, \Theta_1) \in C_n(\Omega)$, $r_1 \leq \frac{4}{5}l$. Then from (2.2), we have

$$V(r_1)\varphi(\Theta_1) \int_{S_n(\Omega; (l, +\infty))} W(t)\varphi(\Phi) d\mu(Q) \lesssim \int_{S_n(\Omega)} G_\Omega^a(P, Q) d\mu(Q) < +\infty,$$

which gives (2.4). For any positive number ϵ , from (2.4), we can take a number R_ϵ such that

$$\int_{S_n(\Omega; (R_\epsilon, +\infty))} W(t)\varphi(\Phi) d\mu(Q) < \frac{\epsilon}{2}.$$

If we take a point $P_i = (r_i, \Theta_i) \in C_n(\Omega)$, $r_i \geq \frac{5}{4}R_\epsilon$, then we have from (2.1)

$$W(r_i)\varphi(\Theta_i) \int_{S_n(\Omega; (0, R_\epsilon])} V(t)\varphi(\Phi) d\mu(Q) \lesssim \int_{S_n(\Omega)} G_\Omega^\alpha(P, Q) d\mu(Q) < +\infty.$$

If R ($R > R_\epsilon$) is sufficiently large, then

$$\begin{aligned} & \frac{W(R)}{V(R)} \int_{S_n(\Omega; (0, R))} V(t)\varphi(\Phi) d\mu(Q) \\ & \lesssim \frac{W(R)}{V(R)} \int_{S_n(\Omega; (0, R_\epsilon])} V(t)\varphi(\Phi) d\mu(Q) + \int_{S_n(\Omega; (R_\epsilon, R))} W(t)\varphi(\Phi) d\mu(Q) \\ & \lesssim \frac{W(R)}{V(R)} \int_{S_n(\Omega; (0, R_\epsilon])} V(t)\varphi(\Phi) d\mu(Q) + \int_{S_n(\Omega; (R_\epsilon, +\infty))} W(t)\varphi(\Phi) d\mu(Q) \\ & \lesssim \epsilon, \end{aligned}$$

which gives (2.5). □

Lemma 3 *Let λ be any positive measure on \mathbf{R}^n having finite total mass. Then $E(\epsilon; \lambda, \alpha)$ has a covering $\{r_j, R_j\}$ ($j = 1, 2, \dots$) satisfying*

$$\sum_{j=1}^{\infty} \left(\frac{r_j}{R_j} \right)^{2-\alpha} \frac{V(R_j)W(R_j)}{V(r_j)W(r_j)} < \infty.$$

Proof Set

$$E_j(\epsilon; \lambda, \beta) = \{P = (r, \Theta) \in E(\epsilon; \lambda, \beta) : 2^j \leq r < 2^{j+1}\} \quad (j = 2, 3, 4, \dots).$$

If $P = (r, \Theta) \in E_j(\epsilon; \lambda, \beta)$, then there exists a positive number $\rho(P)$ such that

$$\left(\frac{\rho(P)}{r} \right)^{2-\alpha} \frac{V(r)W(r)}{V(\rho(P))W(\rho(P))} \approx \left(\frac{\rho(P)}{r} \right)^{n-\alpha} \leq \frac{\lambda(B(P, \rho(P)))}{\epsilon}.$$

Since $E_j(\epsilon; \lambda, \beta)$ can be covered by the union of a family of balls $\{B(P_{j,i}, \rho_{j,i}) : P_{j,i} \in E_j(\epsilon; \lambda, \beta)\}$ ($\rho_{j,i} = \rho(P_{j,i})$). By the Vitali lemma (see [17]), there exists $\Lambda_j \subset E_j(\epsilon; \lambda, \beta)$, which is at most countable, such that $\{B(P_{j,i}, \rho_{j,i}) : P_{j,i} \in \Lambda_j\}$ are disjoint and $E_j(\epsilon; \lambda, \beta) \subset \bigcup_{P_{j,i} \in \Lambda_j} B(P_{j,i}, 5\rho_{j,i})$.

So

$$\bigcup_{j=2}^{\infty} E_j(\epsilon; \lambda, \beta) \subset \bigcup_{j=2}^{\infty} \bigcup_{P_{j,i} \in \Lambda_j} B(P_{j,i}, 5\rho_{j,i}).$$

On the other hand, note that

$$\bigcup_{P_{j,i} \in \Lambda_j} B(P_{j,i}, \rho_{j,i}) \subset \{P = (r, \Theta) : 2^{j-1} \leq r < 2^{j+2}\},$$

so that

$$\begin{aligned} \sum_{P_{j,i} \in \Lambda_j} \left(\frac{5\rho_{j,i}}{|P_{j,i}|} \right)^{2-\alpha} \frac{V(|P_{j,i}|)W(|P_{j,i}|)}{V(\rho_{j,i})W(\rho_{j,i})} &\approx \sum_{P_{j,i} \in \Lambda_j} \left(\frac{5\rho_{j,i}}{|P_{j,i}|} \right)^{n-\alpha} \leq 5^{n-\alpha} \sum_{P_{j,i} \in \Lambda_j} \frac{\lambda(B(P_{j,i}, \rho_{j,i}))}{\epsilon} \\ &\leq \frac{5^{n-\alpha}}{\epsilon} \lambda(C_n(\Omega; [2^{j-1}, 2^{j+2}])). \end{aligned}$$

Hence we obtain

$$\begin{aligned} \sum_{j=1}^{\infty} \sum_{P_{j,i} \in \Lambda_j} \left(\frac{\rho_{j,i}}{|P_{j,i}|} \right)^{2-\alpha} \frac{V(|P_{j,i}|)W(|P_{j,i}|)}{V(\rho_{j,i})W(\rho_{j,i})} &\approx \sum_{j=1}^{\infty} \sum_{P_{j,i} \in \Lambda_j} \left(\frac{\rho_{j,i}}{|P_{j,i}|} \right)^{n-\alpha} \\ &\leq \sum_{j=1}^{\infty} \frac{\lambda(C_n(\Omega; [2^{j-1}, 2^{j+2}]))}{\epsilon} \\ &\leq \frac{3\lambda(\mathbf{R}^n)}{\epsilon}. \end{aligned}$$

Since $E(\epsilon; \lambda, \beta) \cap \{P = (r, \Theta) \in \mathbf{R}^n; r \geq 4\} = \bigcup_{j=2}^{\infty} E_j(\epsilon; \lambda, \beta)$. Then $E(\epsilon; \lambda, \beta)$ is finally covered by a sequence of balls $\{B(P_{j,i}, \rho_{j,i}), B(P_1, 6)\}$ ($j = 2, 3, \dots; i = 1, 2, \dots$) satisfying

$$\sum_{j,i} \left(\frac{\rho_{j,i}}{|P_{j,i}|} \right)^{2-\alpha} \frac{V(|P_{j,i}|)W(|P_{j,i}|)}{V(\rho_{j,i})W(\rho_{j,i})} \approx \sum_{j,i} \left(\frac{\rho_{j,i}}{|P_{j,i}|} \right)^{n-\alpha} \leq \frac{3\lambda(\mathbf{R}^n)}{\epsilon} + 6^{n-\alpha} < +\infty,$$

where $B(P_1, 6)$ ($P_1 = (1, 0, \dots, 0) \in \mathbf{R}^n$) is the ball which covers $\{P = (r, \Theta) \in \mathbf{R}^n; r < 4\}$. \square

3 Proof of Theorem 1

For any point $P = (r, \Theta) \in C_n(\Omega; (R, +\infty)) - E(\epsilon; \nu', \alpha)$, where $R (\leq \frac{4}{5}r)$ is a sufficiently large number and ϵ is a sufficiently small positive number.

Write

$$G_{\Omega}^a \nu(P) = G_{\Omega}^a \nu(1)(P) + G_{\Omega}^a \nu(2)(P) + G_{\Omega}^a \nu(3)(P),$$

where

$$G_{\Omega}^a \nu(1)(P) = \int_{C_n(\Omega; (0, \frac{4}{5}r])} G_{\Omega}^a(P, Q) d\nu(Q),$$

$$G_{\Omega}^a \nu(2)(P) = \int_{C_n(\Omega; (\frac{4}{5}r, \frac{5}{4}r))} G_{\Omega}^a(P, Q) d\nu(Q),$$

and

$$G_{\Omega}^a \nu(3)(P) = \int_{C_n(\Omega; [\frac{5}{4}r, \infty))} G_{\Omega}^a(P, Q) d\nu(Q).$$

From (2.1) and (2.2) we obtain the following growth estimates:

$$G_{\Omega}^a \nu(1)(P) \lesssim \epsilon V(r)\varphi(\Theta), \tag{3.1}$$

$$G_{\Omega}^a \nu(3)(P) \lesssim \epsilon V(r)\varphi(\Theta). \tag{3.2}$$

By (2.3) and (3.1), we have

$$G_{\Omega}^{\alpha}v(2)(P) \leq G_{\Omega}^{\alpha}v(21)(P) + G_{\Omega}^{\alpha}v(22)(P),$$

where

$$G_{\Omega}^{\alpha}v(21)(P) = \varphi(\Theta) \int_{C_n(\Omega; (\frac{4}{5}r, \frac{5}{4}r))} V(t) dv'(Q)$$

and

$$G_{\Omega}^{\alpha}v(22)(P) = \int_{C_n(\Omega; (\frac{4}{5}r, \frac{5}{4}r))} \Pi_{\Omega}(P, Q) dv(Q).$$

Then by Lemma 2, we immediately get

$$G_{\Omega}^{\alpha}v(21)(P) \lesssim \epsilon V(r)\varphi(\Theta). \tag{3.3}$$

To estimate $G_{\Omega}^{\alpha}v(22)(P)$, take a sufficiently small positive number c independent of P such that

$$\Lambda(P) = \left\{ (t, \Phi) \in C_n \left(\Omega; \left(\frac{4}{5}r, \frac{5}{4}r \right) \right); |(1, \Phi) - (1, \Theta)| < c \right\} \subset B \left(P, \frac{r}{2} \right) \tag{3.4}$$

and divide $C_n(\Omega; (\frac{4}{5}r, \frac{5}{4}r))$ into two sets $\Lambda(P)$ and $\Lambda(P)$, where

$$\Lambda(P) = C_n \left(\Omega; \left(\frac{4}{5}r, \frac{5}{4}r \right) \right) - \Lambda(P).$$

Write

$$G_{\Omega}^{\alpha}v(22)(P) = G_{\Omega}^{\alpha}v(221)(P) + G_{\Omega}^{\alpha}v(222)(P),$$

where

$$G_{\Omega}^{\alpha}v(221)(P) = \int_{\Lambda(P)} \Pi_{\Omega}(P, Q) dv(Q)$$

and

$$G_{\Omega}^{\alpha}v(222)(P) = \int_{\Lambda(P)} \Pi_{\Omega}(P, Q) dv(Q).$$

There exists a positive c' such that $|P - Q| \geq c'r$ for any $Q \in \Lambda(P)$, and hence

$$\begin{aligned} G_{\Omega}^{\alpha}v(222)(P) &\lesssim \int_{C_n(\Omega; (\frac{4}{5}r, \frac{5}{4}r))} \frac{rt\varphi(\Theta)\varphi(\Phi)}{|P - Q|^n} dv(Q) \\ &\lesssim V(r)\varphi(\Theta) \int_{C_n(\Omega; (\frac{4}{5}r, \infty))} dv'(Q) \\ &\lesssim \epsilon V(r)\varphi(\Theta) \end{aligned} \tag{3.5}$$

from Lemma 2.

Now we estimate $G_{\Omega}^{\alpha}v(221)(P)$. Set

$$I_i(P) = \{Q \in \Lambda(P); 2^{i-1}\delta(P) \leq |P - Q| < 2^i\delta(P)\},$$

where $i = 0, \pm 1, \pm 2, \dots$

Since $P = (r, \Theta) \notin E(\epsilon; v', \alpha)$ and hence $v'(\{P\}) = 0$ from Remark 1, we can divide $G_{\Omega}^{\alpha}v(221)(P)$ into

$$G_{\Omega}^{\alpha}v(221)(P) = G_{\Omega}^A v(2211)(P) + G_{\Omega}^{\alpha}v(2212)(P),$$

where

$$G_{\Omega}^A v(2211)(P) = \sum_{i=-\infty}^{-1} \int_{I_i(P)} \Pi_{\Omega}(P, Q) dv(Q)$$

and

$$G_{\Omega}^{\alpha}v(2212)(P) = \sum_{i=0}^{\infty} \int_{I_i(P)} \Pi_{\Omega}(P, Q) dv(Q).$$

Since $\delta(Q) + |P - Q| \geq \delta(P)$, we have

$$t\varphi_{\Omega}(\Phi) \gtrsim \delta(Q) \gtrsim 2^{-1}\delta(P)$$

for any $Q = (t, \Phi) \in I_i(P)$ ($i = -1, -2, \dots$). Then by (1.1)

$$\begin{aligned} \int_{I_i(P)} \Pi_{\Omega}(P, Q) dv(Q) &\lesssim \int_{I_i(P)} \frac{1}{|P - Q|^{n-2} W(t)\varphi(\Phi)} dv'(Q) \\ &\lesssim \frac{r^{2-\alpha}}{W(r)} \varphi^{1-\alpha}(\Theta) \frac{v'(B(P, 2^i\delta(P)))}{\{2^i\delta(P)\}^{n-\alpha}} \\ &\lesssim \frac{r^{2-\alpha}}{W(r)} \varphi^{1-\alpha}(\Theta) M(P; v', \alpha) \quad (i = -1, -2, \dots). \end{aligned}$$

Since $P = (r, \Theta) \notin E(\epsilon; v', \alpha)$, we obtain

$$G_{\Omega}^{\alpha}v(2211)(P) \lesssim \epsilon V(r) \varphi^{1-\alpha}(\Theta). \tag{3.6}$$

By (3.4), we can take a positive integer $i(P)$ satisfying

$$2^{i(P)-1}\delta(P) \leq \frac{r}{2} < 2^{i(P)}\delta(P)$$

and $I_i(P) = \emptyset$ ($i = i(P) + 1, i(P) + 2, \dots$).

Since $r\varphi_{\Omega}(\Theta) \lesssim \delta(P)$ ($P = (r, \Theta) \in C_n(\Omega)$), we have

$$\begin{aligned} \int_{I_i(P)} \Pi_{\Omega}(P, Q) dv'(Q) &\lesssim r\varphi(\Theta) \int_{I_i(P)} \frac{t}{|P - Q|^n W(t)} dv'(Q) \\ &\lesssim \frac{r^{2-\alpha}}{W(r)} \varphi^{1-\alpha}(\Theta) \frac{v'(I_i(P))}{\{2^i\delta(P)\}^{n-\alpha}} \quad (i = 0, 1, 2, \dots, i(P)). \end{aligned}$$

Since $P = (r, \Theta) \notin E(\epsilon; v', \alpha)$, we have

$$\begin{aligned} \frac{v'(I_i(P))}{\{2^i \delta(P)\}^{n-\alpha}} &\lesssim v'(B(P, 2^i \delta(P))) V(2^i \delta(P)) W(2^i \delta(P)) \{2^i \delta(P)\}^{\alpha-2} \\ &\lesssim M(P; v', \alpha) \\ &\leq \epsilon V(r) W(r) r^{\alpha-2} \quad (i = 0, 1, 2, \dots, i(P) - 1) \end{aligned}$$

and

$$\frac{v'(I_i(P))}{\{2^i \delta(P)\}^{n-\alpha}} \lesssim v'(\Lambda(P)) V\left(\frac{r}{2}\right) W\left(\frac{r}{2}\right) \left(\frac{r}{2}\right)^{\alpha-2} \leq \epsilon V(r) W(r) r^{\alpha-2}.$$

Hence we obtain

$$G_\Omega^\alpha v(2212)(P) \lesssim \epsilon V(r) \varphi^{1-\alpha}(\Theta). \quad (3.7)$$

Combining (3.1)-(3.3) and (3.5)-(3.7), we finally obtain the result that if R is sufficiently large and ϵ is a sufficiently small, then $G_\Omega^\alpha v(P) = o(V(r)\varphi^{1-\alpha}(\Theta))$ as $r \rightarrow \infty$, where $P = (r, \Theta) \in C_n(\Omega; (R, +\infty)) - E(\epsilon; v', \alpha)$. Finally, there exists an additional finite ball B_0 covering $C_n(\Omega; (0, R])$, which together with Lemma 3, gives the conclusion of Theorem 1.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the manuscript and read and approved the final manuscript.

Author details

¹School of Mathematics and Information Science, Henan University of Economics and Law, Zhengzhou, 450046, P.R. China. ²Matematiska Institutionen, Stockholms Universitet, Stockholm, 106 91, Sweden.

Acknowledgements

The authors are very thankful to the anonymous referees for their valuable comments and constructive suggestions, which helped to improve the quality of the paper. This work is supported by the Academy of Finland Grant No. 176512.

Received: 19 September 2014 Accepted: 12 November 2014 Published online: 27 November 2014

References

- Escassut, A, Tutschke, W, Yang, CC: Some Topics on Value Distribution and Differentiability in Complex and P -Adic Analysis. Science Press, Beijing (2008)
- Gilbarg, D, Trudinger, NS: Elliptic Partial Differential Equations of Second Order. Springer, Berlin (1977)
- Miranda, C: Partial Differential Equations of Elliptic Type. Springer, Berlin (1970)
- Verzhbinskii, GM, Maz'ya, VG: Asymptotic behavior of solutions of elliptic equations of the second order close to a boundary. *I. Sib. Mat. Zh.* **12**(2), 874-899 (1971)
- Qiao, L, Deng, GT: A lower bound of harmonic functions in a cone and its application. *Sci. Sin., Math.* **44**(6), 671-684 (2014) (in Chinese)
- Qiao, L, Deng, GT: Minimally thin sets at infinity with respect to the Schrödinger operator. *Sci. Sin., Math.* **44**(12), 1247-1256 (2014) (in Chinese)
- Xue, GX: A remark on the a -minimally thin sets associated with the Schrödinger operator. *Bound. Value Probl.* **2014**, 133 (2014)
- Zhao, T: Minimally thin sets associated with the stationary Schrödinger operator. *J. Inequal. Appl.* **2014**, 67 (2014)
- Simon, B: Schrödinger semigroups. *Bull. Am. Math. Soc.* **7**(2), 447-526 (1982)
- Hartman, P: Ordinary Differential Equations. Wiley, New York (1964)
- Qiao, L, Ren, YD: Integral representations for the solutions of infinite order of the stationary Schrödinger equation in a cone. *Monatshefte Math.* **173**(4), 593-603 (2014)
- Qiao, L, Deng, GT: The Riesz decomposition theorem for superharmonic functions in a cone and its application. *Sci. Sin., Math.* **42**(8), 763-774 (2012) (in Chinese)
- Qiao, L: Integral representations for harmonic functions of infinite order in a cone. *Results Math.* **61**(4), 63-74 (2012)
- Qiao, L, Deng, GT: Integral representations of harmonic functions in a cone. *Sci. Sin., Math.* **41**(6), 535-546 (2011) (in Chinese)

15. Miyamoto, I, Yoshida, H: Two criteria of Wiener type for minimally thin sets and rarefied sets in a cone. *J. Math. Soc. Jpn.* **54**, 487-512 (2002)
16. Yoshida, H: Harmonic majorant of a radial subharmonic function on a strip and their applications. *Int. J. Pure Appl. Math.* **30**(2), 259-286 (2006)
17. Stein, EM: *Singular Integrals and Differentiability Properties of Functions*. Princeton University Press, Princeton (1970)

doi:10.1186/s13661-014-0245-9

Cite this article as: Zhao and Yamada Jr.: Growth properties of Green-Sch potentials at infinity. *Boundary Value Problems* 2014 **2014**:245.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
