



Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Amino-4-(4-chlorophenyl)-4H-chromeno[8,7-b]pyridine-3-carbonitrile

Abd El-Galil E. Amr,^{a,b,†} Ahmed M. El-Agrody,^c Nermien M. Sabry,^c Seik Weng Ng^{d,e} and Edward R. T. Tiekink^{d,*}

^aDrug Exploration & Development Chair (DEDC), College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, ^bApplied Organic Chemistry Department, National Research Center, Dokki 12622, Cairo, Egypt, ^cChemistry Department, Faculty of Science, King Khalid University, Abha 61413, PO Box 9004, Saudi Arabia, ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^eChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: edward.tiekink@gmail.com


Received 19 February 2013; accepted 22 February 2013

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.061; wR factor = 0.164; data-to-parameter ratio = 16.3.

The asymmetric unit of the title compound, $\text{C}_{19}\text{H}_{12}\text{ClN}_3\text{O}$, contains two molecules with similar conformations. The 14 non-H atoms comprising the 4H-chromeno[8,7-b]pyridine residue are essentially coplanar (r.m.s. deviations = 0.037 and 0.042 Å for the two molecules) and the main difference between them is seen in the twist about the bond linking the main residue to the attached chlorobenzene rings [dihedral angles = 79.01 (12) and 76.22 (11)° for the two molecules]. Zigzag supramolecular chains along the a -axis direction mediated by amino-pyridine $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds feature in the crystal packing; these are connected into a three-dimensional architecture by $\text{C}-\text{H}\cdots\pi$ interactions and $\text{Cl}\cdots\text{Cl}$ contacts [$\text{Cl}\cdots\text{Cl} = 3.3896$ (14) Å].

Related literature

For background to the chemistry and biological activity of 4H-pyran derivatives, see: Al-Ghamdi *et al.* (2012); El-Agrody *et al.* (2012). For the structure of the 2-chloro analogue, see: Wang *et al.* (2003).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{12}\text{ClN}_3\text{O}$	$V = 3165.2$ (6) Å ³
$M_r = 333.77$	$Z = 8$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 6.5311$ (8) Å	$\mu = 0.25$ mm ⁻¹
$b = 35.129$ (3) Å	$T = 295$ K
$c = 14.0903$ (14) Å	$0.30 \times 0.20 \times 0.05$ mm
$\beta = 101.740$ (11)°	

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector	20646 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2011)	7326 independent reflections
$T_{\min} = 0.833$, $T_{\max} = 1.000$	3471 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.068$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.164$	$\Delta\rho_{\max} = 0.21$ e Å ⁻³
$S = 1.01$	$\Delta\rho_{\min} = -0.32$ e Å ⁻³
7326 reflections	
450 parameters	

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 and Cg2 are the centroids of the $\text{N4}, \text{C20}-\text{C23}, \text{C28}$ and $\text{C33}-\text{C38}$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H1}\cdots\text{N4}^i$	0.84 (3)	2.34 (4)	3.172 (4)	174 (3)
$\text{N2}-\text{H2}\cdots\text{N5}^i$	0.89 (3)	2.61 (3)	3.308 (5)	136 (2)
$\text{N5}-\text{H3}\cdots\text{N1}$	0.87 (3)	2.15 (3)	3.014 (3)	173 (3)
$\text{C18}-\text{H18}\cdots\text{Cg1}^{\text{ii}}$	0.93	2.80	3.650 (3)	152
$\text{C24}-\text{H24}\cdots\text{Cg2}^{\text{iii}}$	0.93	2.74	3.668 (3)	174

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x - \frac{1}{2}, y - \frac{1}{2}, -z - \frac{1}{2}$; (iii) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012), *QMol* (Gans & Shalloway, 2001) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

† Additional correspondence author, e-mail: aamr1963@yahoo.com.

We thank the Research Center of the College of Pharmacy and Deanship of Scientific Research of King Saud University for supporting this study. We also thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR-MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB7046).

References

- Agilent (2011). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Al-Ghamdi, A. M., Abd EL-Wahab, A. H. F., Mohamed, H. M. & El-Agrody, A. M. (2012). *Lett. Drug. Des. Discov.* **9**, 459–470.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- El-Agrody, A. M., Khattab, E. S. A. E. H., Fouda, A. M. & Al-Ghamdi, A. M. (2012). *Med. Chem. Res.* **22**, 1339–1355.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Gans, J. & Shalloway, D. (2001). *J. Mol. Graph. Model.*, **19**, 557–559.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, X.-S., Shi, D.-Q. & Tu, S.-J. (2003). *Chin. J. Chem.* **21**, 1114–1117.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2013). E69, o462–o463 [doi:10.1107/S1600536813005217]

2-Amino-4-(4-chlorophenyl)-4*H*-chromeno[8,7-*b*]pyridine-3-carbonitrile

Abd El-Galil E. Amr, Ahmed M. El-Agrody, Nermien M. Sabry, Seik Weng Ng and Edward R. T. Tiekink

S1. Comment

Motivated by their biological activities and in continuation of an on-going programme on the chemistry of 4*H*-pyran derivatives (Al-Ghamdi *et al.*, 2012; El-Agrody *et al.*, 2012), the synthesis and crystal structure determination of (I) is reported.

Two independent molecules comprise the asymmetric unit of (I), Fig. 1. As illustrated in Fig. 2, where the O2-containing molecule is super-imposed upon the inverse of the O1-containing molecule, the molecules are virtually super-imposable with differences apparent in the relationship between the 4*H*-chromeno[8,7-*b*]pyridine residue and the attached benzene ring. For the O1-containing molecule, the r.m.s. deviation of the 14 non-hydrogen atoms comprising the fused ring system is 0.037 Å, the dihedral angle between this and the benzene ring is 79.01 (12)° and the twist between these groups is manifested in the C7—C12—C14—C15 torsion angle of -132.9 (3)°. The comparable values for the second molecule are 0.042 Å, 76.22 (11)° and 149.0 (3)°, respectively. The observed conformation is in accord with that established previously for the 2-chloro analogue (Wang *et al.*, 2003).


The most prominent feature of the crystal packing is the formation of supramolecular zigzag chains along the *a* axis mediated by (amino)N—H...N(pyridyl) hydrogen bonding, Fig. 3 and Table 1. Whereas the second N1-bound H2 atom forms a weak interaction to the N5 atom, Table 1, reinforcing the chain, the second N2-bound H4 atom does not form a significant intermolecular interaction. The chains are connected into a three-dimensional architecture by C—H... π interactions along with Cl2...Cl2^{*i*} contacts [Cl2...Cl2^{*i*} = 3.3896 (14) Å for *i*: 2 - *x*, 1 - *y*, 2 - *z*], Fig. 4.

S2. Experimental


A solution of 8-hydroxyquinoline (0.01 mol) in EtOH (30 ml) was treated with α -cyano-*p*-chlorocinnamionitrile (0.01 mol) and piperidine (0.5 ml). The reaction mixture was heated for 60 minutes by which time complete precipitation occurred. The solid product was collected by filtration and recrystallized from ethanol to give yellow prisms of the title compound, (I); *M*.pt: 522–523 K.

S3. Refinement


The C-bound H atoms were geometrically placed (C—H = 0.93–0.98 Å) and refined as riding with $U_{iso}(\text{H}) = 1.2U_{eq}(\text{C})$. The N-bound-H atoms were refined freely.

**Figure 1**


The molecular structures of the two independent molecules comprising the asymmetric unit in (I) showing displacement ellipsoids at the 35% probability level.

**Figure 2**

Overlay diagram of the two independent molecules in (I) with the inverted N1-containing molecule illustrated in red. The molecules are overlaid so that the pyridyl rings are superimposed.

**Figure 3**

A view of the supramolecular chain along the a axis in (I) sustained by $\text{N—H}\cdots\text{N}$ hydrogen bonding, shown as blue dashed lines.

**Figure 4**

A view in projection down the a axis of the crystal packing in (I). The $\text{N—H}\cdots\text{N}$, $\text{C—H}\cdots\pi$ and $\text{Cl}\cdots\text{Cl}$ interactions are shown as blue, purple and orange dashed lines, respectively.

2-Amino-4-(4-chlorophenyl)-4H-chromeno[8,7-b]pyridine-3-carbonitrile*Crystal data*C₁₉H₁₂ClN₃O $M_r = 333.77$ Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

 $a = 6.5311$ (8) Å $b = 35.129$ (3) Å $c = 14.0903$ (14) Å $\beta = 101.740$ (11)° $V = 3165.2$ (6) Å³ $Z = 8$ $F(000) = 1376$ $D_x = 1.401$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2452 reflections

 $\theta = 2.9$ – 27.5 ° $\mu = 0.25$ mm⁻¹ $T = 295$ K

Prism, yellow

 $0.30 \times 0.20 \times 0.05$ mm*Data collection*Agilent SuperNova Dual
diffractometer with an Atlas detectorRadiation source: SuperNova (Mo) X-ray
Source

Mirror monochromator

Detector resolution: 10.4041 pixels mm⁻¹ ω scanAbsorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2011) $T_{\min} = 0.833$, $T_{\max} = 1.000$

20646 measured reflections

7326 independent reflections

3471 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.068$ $\theta_{\max} = 27.6$ °, $\theta_{\min} = 3.0$ ° $h = -8 \rightarrow 8$ $k = -44 \rightarrow 45$ $l = -18 \rightarrow 18$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.061$ $wR(F^2) = 0.164$ $S = 1.01$

7326 reflections

450 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0527P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.21$ e Å⁻³ $\Delta\rho_{\min} = -0.32$ e Å⁻³Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0023 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.14716 (19)	0.04691 (3)	0.61417 (7)	0.0887 (4)
Cl2	0.8629 (2)	0.47842 (3)	0.89931 (6)	0.1067 (5)

O1	0.1506 (3)	0.25977 (5)	0.34064 (13)	0.0470 (5)
O2	0.6996 (3)	0.35767 (5)	0.38366 (13)	0.0464 (5)
N1	0.4684 (4)	0.27492 (6)	0.25590 (15)	0.0455 (6)
N2	-0.0484 (5)	0.28690 (7)	0.43039 (19)	0.0494 (7)
N3	-0.3636 (5)	0.21391 (8)	0.5043 (2)	0.0800 (10)
N4	1.0071 (4)	0.35491 (6)	0.28969 (16)	0.0475 (6)
N5	0.4811 (5)	0.31956 (7)	0.4397 (2)	0.0500 (7)
N6	0.2277 (5)	0.37488 (8)	0.5911 (2)	0.0873 (11)
C1	0.6186 (5)	0.28248 (8)	0.2095 (2)	0.0512 (8)
H1A	0.6696	0.3073	0.2126	0.061*
C2	0.7079 (5)	0.25603 (9)	0.15559 (19)	0.0528 (8)
H2A	0.8134	0.2632	0.1237	0.063*
C3	0.6363 (5)	0.21963 (9)	0.15116 (19)	0.0511 (8)
H3A	0.6953	0.2014	0.1171	0.061*
C4	0.4729 (5)	0.20941 (8)	0.19792 (18)	0.0445 (7)
C5	0.3886 (5)	0.17228 (8)	0.1979 (2)	0.0549 (9)
H5	0.4396	0.1529	0.1641	0.066*
C6	0.2346 (5)	0.16457 (8)	0.2465 (2)	0.0542 (8)
H6	0.1832	0.1399	0.2458	0.065*
C7	0.1495 (5)	0.19300 (7)	0.29835 (18)	0.0421 (7)
C8	0.2278 (5)	0.22901 (7)	0.29687 (18)	0.0403 (7)
C9	0.3921 (5)	0.23850 (7)	0.24918 (17)	0.0407 (7)
C10	-0.0007 (5)	0.25317 (8)	0.39165 (18)	0.0409 (7)
C11	-0.0810 (5)	0.21862 (8)	0.40245 (19)	0.0437 (7)
C12	-0.0248 (5)	0.18351 (7)	0.35028 (19)	0.0458 (7)
H12	-0.1477	0.1772	0.3003	0.055*
C13	-0.2384 (6)	0.21591 (8)	0.4580 (2)	0.0531 (8)
C14	0.0212 (5)	0.14887 (8)	0.4167 (2)	0.0453 (7)
C15	-0.1253 (6)	0.12104 (9)	0.4144 (2)	0.0702 (10)
H15	-0.2530	0.1233	0.3712	0.084*
C16	-0.0887 (7)	0.08930 (9)	0.4751 (3)	0.0751 (11)
H16	-0.1905	0.0706	0.4728	0.090*
C17	0.0984 (7)	0.08624 (9)	0.5375 (2)	0.0579 (9)
C18	0.2466 (6)	0.11359 (10)	0.5426 (2)	0.0690 (10)
H18	0.3736	0.1113	0.5863	0.083*
C19	0.2072 (6)	0.14497 (9)	0.4821 (2)	0.0637 (9)
H19	0.3086	0.1638	0.4858	0.076*
C20	1.1555 (5)	0.35407 (9)	0.2385 (2)	0.0552 (9)
H20	1.1859	0.3308	0.2131	0.066*
C21	1.2701 (5)	0.38601 (10)	0.2201 (2)	0.0584 (9)
H21	1.3715	0.3838	0.1827	0.070*
C22	1.2309 (5)	0.42014 (9)	0.2577 (2)	0.0562 (9)
H22	1.3058	0.4416	0.2461	0.067*
C23	1.0766 (5)	0.42312 (8)	0.31426 (19)	0.0464 (8)
C24	1.0277 (5)	0.45694 (8)	0.3591 (2)	0.0540 (9)
H24	1.0983	0.4793	0.3510	0.065*
C25	0.8789 (5)	0.45719 (7)	0.4137 (2)	0.0482 (8)
H25	0.8528	0.4797	0.4439	0.058*

C26	0.7618 (5)	0.42418 (7)	0.42611 (17)	0.0394 (7)
C27	0.8066 (5)	0.39158 (7)	0.38100 (18)	0.0381 (7)
C28	0.9661 (5)	0.38958 (7)	0.32597 (17)	0.0400 (7)
C29	0.5538 (5)	0.35562 (7)	0.43977 (18)	0.0404 (7)
C30	0.5022 (5)	0.38585 (7)	0.48884 (18)	0.0401 (7)
C31	0.5956 (5)	0.42540 (7)	0.48574 (18)	0.0405 (7)
H31	0.4841	0.4422	0.4525	0.049*
C32	0.3499 (6)	0.37995 (8)	0.5446 (2)	0.0537 (8)
C33	0.6695 (5)	0.44071 (7)	0.5878 (2)	0.0434 (7)
C34	0.5327 (6)	0.46012 (8)	0.6330 (2)	0.0602 (9)
H34	0.3988	0.4657	0.5987	0.072*
C35	0.5929 (7)	0.47155 (9)	0.7299 (3)	0.0708 (11)
H35	0.4991	0.4845	0.7599	0.085*
C36	0.7888 (7)	0.46368 (9)	0.7799 (2)	0.0644 (10)
C37	0.9283 (6)	0.44463 (8)	0.7368 (2)	0.0613 (9)
H37	1.0626	0.4395	0.7713	0.074*
C38	0.8671 (5)	0.43299 (8)	0.6410 (2)	0.0521 (8)
H38	0.9613	0.4197	0.6120	0.063*
H1	-0.032 (6)	0.3060 (10)	0.397 (2)	0.081 (12)*
H2	-0.157 (5)	0.2859 (8)	0.460 (2)	0.064 (11)*
H3	0.469 (5)	0.3081 (9)	0.384 (2)	0.073 (11)*
H4	0.369 (5)	0.3152 (8)	0.467 (2)	0.061 (10)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.1187 (10)	0.0705 (6)	0.0826 (6)	0.0147 (6)	0.0336 (6)	0.0305 (5)
Cl2	0.1573 (13)	0.1087 (8)	0.0572 (5)	-0.0307 (8)	0.0293 (6)	-0.0242 (5)
O1	0.0519 (15)	0.0330 (10)	0.0612 (11)	-0.0010 (10)	0.0238 (11)	-0.0079 (8)
O2	0.0496 (14)	0.0337 (10)	0.0638 (12)	-0.0065 (10)	0.0302 (11)	-0.0081 (9)
N1	0.0483 (18)	0.0410 (13)	0.0499 (13)	-0.0018 (12)	0.0161 (13)	-0.0042 (11)
N2	0.059 (2)	0.0390 (14)	0.0557 (15)	0.0046 (14)	0.0237 (15)	-0.0015 (12)
N3	0.074 (2)	0.085 (2)	0.090 (2)	-0.0125 (19)	0.038 (2)	-0.0017 (17)
N4	0.0521 (18)	0.0416 (13)	0.0539 (14)	0.0040 (13)	0.0228 (13)	0.0030 (11)
N5	0.058 (2)	0.0373 (13)	0.0627 (16)	-0.0117 (13)	0.0308 (16)	-0.0083 (12)
N6	0.093 (3)	0.0567 (18)	0.135 (3)	0.0027 (18)	0.077 (2)	0.0025 (18)
C1	0.053 (2)	0.0495 (17)	0.0538 (17)	-0.0009 (16)	0.0169 (17)	0.0005 (14)
C2	0.052 (2)	0.064 (2)	0.0446 (15)	0.0023 (18)	0.0150 (15)	0.0032 (15)
C3	0.057 (2)	0.0555 (19)	0.0419 (15)	0.0101 (17)	0.0140 (16)	-0.0029 (14)
C4	0.050 (2)	0.0438 (16)	0.0388 (14)	0.0091 (15)	0.0070 (14)	-0.0051 (12)
C5	0.064 (3)	0.0407 (16)	0.0620 (18)	0.0061 (17)	0.0180 (18)	-0.0129 (14)
C6	0.062 (2)	0.0406 (16)	0.0614 (18)	-0.0049 (17)	0.0148 (18)	-0.0123 (14)
C7	0.044 (2)	0.0355 (14)	0.0461 (15)	0.0006 (14)	0.0074 (14)	-0.0048 (12)
C8	0.043 (2)	0.0335 (14)	0.0431 (14)	0.0026 (14)	0.0069 (14)	-0.0067 (12)
C9	0.041 (2)	0.0410 (15)	0.0395 (14)	0.0021 (14)	0.0073 (14)	-0.0007 (12)
C10	0.041 (2)	0.0410 (15)	0.0413 (14)	0.0035 (15)	0.0103 (14)	-0.0009 (12)
C11	0.038 (2)	0.0440 (16)	0.0491 (16)	0.0006 (15)	0.0102 (15)	0.0001 (13)
C12	0.042 (2)	0.0415 (15)	0.0502 (16)	-0.0049 (15)	0.0015 (15)	-0.0017 (13)

C13	0.052 (2)	0.0469 (18)	0.0615 (19)	-0.0057 (17)	0.0152 (18)	0.0030 (14)
C14	0.043 (2)	0.0384 (15)	0.0537 (17)	-0.0019 (15)	0.0079 (16)	-0.0024 (13)
C15	0.061 (3)	0.057 (2)	0.085 (2)	-0.016 (2)	-0.004 (2)	0.0148 (18)
C16	0.079 (3)	0.054 (2)	0.089 (2)	-0.025 (2)	0.009 (2)	0.0132 (19)
C17	0.074 (3)	0.0490 (18)	0.0531 (18)	0.004 (2)	0.0177 (19)	0.0036 (15)
C18	0.065 (3)	0.068 (2)	0.067 (2)	-0.003 (2)	-0.0054 (19)	0.0121 (18)
C19	0.055 (2)	0.0527 (19)	0.077 (2)	-0.0133 (18)	-0.0028 (19)	0.0052 (17)
C20	0.059 (2)	0.0555 (19)	0.0569 (17)	0.0111 (18)	0.0257 (18)	0.0023 (15)
C21	0.053 (2)	0.072 (2)	0.0566 (18)	0.0007 (19)	0.0259 (17)	0.0088 (17)
C22	0.052 (2)	0.060 (2)	0.0598 (18)	-0.0096 (18)	0.0187 (17)	0.0118 (16)
C23	0.049 (2)	0.0456 (16)	0.0460 (15)	-0.0031 (16)	0.0127 (15)	0.0086 (13)
C24	0.065 (3)	0.0423 (17)	0.0570 (17)	-0.0141 (17)	0.0172 (18)	0.0074 (14)
C25	0.056 (2)	0.0326 (15)	0.0572 (17)	-0.0037 (15)	0.0137 (17)	-0.0016 (13)
C26	0.0399 (19)	0.0338 (14)	0.0451 (14)	-0.0006 (14)	0.0100 (14)	0.0026 (12)
C27	0.0391 (19)	0.0304 (13)	0.0464 (15)	-0.0025 (13)	0.0122 (14)	0.0030 (11)
C28	0.044 (2)	0.0383 (14)	0.0390 (14)	0.0003 (14)	0.0103 (14)	0.0042 (12)
C29	0.041 (2)	0.0349 (14)	0.0480 (15)	-0.0028 (14)	0.0147 (14)	-0.0002 (12)
C30	0.0402 (19)	0.0363 (14)	0.0468 (15)	-0.0034 (14)	0.0158 (14)	-0.0042 (12)
C31	0.0400 (19)	0.0319 (13)	0.0505 (15)	0.0037 (14)	0.0118 (14)	-0.0024 (12)
C32	0.058 (2)	0.0343 (15)	0.076 (2)	-0.0001 (16)	0.0303 (19)	-0.0032 (14)
C33	0.049 (2)	0.0294 (13)	0.0554 (16)	0.0000 (15)	0.0186 (16)	-0.0050 (12)
C34	0.053 (2)	0.0528 (18)	0.079 (2)	-0.0001 (18)	0.0229 (19)	-0.0214 (16)
C35	0.080 (3)	0.061 (2)	0.083 (2)	-0.015 (2)	0.046 (2)	-0.0294 (19)
C36	0.084 (3)	0.058 (2)	0.0573 (19)	-0.017 (2)	0.028 (2)	-0.0092 (16)
C37	0.075 (3)	0.0530 (18)	0.0532 (18)	-0.0008 (19)	0.0072 (18)	0.0023 (15)
C38	0.059 (2)	0.0430 (16)	0.0554 (18)	0.0070 (17)	0.0150 (17)	-0.0033 (14)

Geometric parameters (Å, °)

C11—C17	1.743 (3)	C14—C19	1.374 (4)
C12—C36	1.732 (3)	C15—C16	1.396 (4)
O1—C10	1.355 (3)	C15—H15	0.9300
O1—C8	1.389 (3)	C16—C17	1.357 (5)
O2—C29	1.358 (3)	C16—H16	0.9300
O2—C27	1.385 (3)	C17—C18	1.355 (4)
N1—C1	1.312 (3)	C18—C19	1.385 (4)
N1—C9	1.369 (3)	C18—H18	0.9300
N2—C10	1.367 (3)	C19—H19	0.9300
N2—H1	0.84 (3)	C20—C21	1.402 (4)
N2—H2	0.89 (3)	C20—H20	0.9300
N3—C13	1.147 (4)	C21—C22	1.356 (4)
N4—C20	1.321 (3)	C21—H21	0.9300
N4—C28	1.368 (3)	C22—C23	1.411 (4)
N5—C29	1.353 (3)	C22—H22	0.9300
N5—H3	0.87 (3)	C23—C28	1.409 (4)
N5—H4	0.91 (3)	C23—C24	1.412 (4)
N6—C32	1.145 (4)	C24—C25	1.357 (4)
C1—C2	1.400 (4)	C24—H24	0.9300

C1—H1A	0.9300	C25—C26	1.419 (4)
C2—C3	1.359 (4)	C25—H25	0.9300
C2—H2A	0.9300	C26—C27	1.370 (3)
C3—C4	1.410 (4)	C26—C31	1.502 (4)
C3—H3A	0.9300	C27—C28	1.421 (4)
C4—C9	1.414 (3)	C29—C30	1.347 (3)
C4—C5	1.416 (4)	C30—C32	1.403 (4)
C5—C6	1.354 (4)	C30—C31	1.522 (3)
C5—H5	0.9300	C31—C33	1.520 (4)
C6—C7	1.416 (4)	C31—H31	0.9800
C6—H6	0.9300	C33—C34	1.378 (4)
C7—C8	1.366 (4)	C33—C38	1.381 (4)
C7—C12	1.510 (4)	C34—C35	1.401 (5)
C8—C9	1.417 (4)	C34—H34	0.9300
C10—C11	1.343 (4)	C35—C36	1.358 (5)
C11—C13	1.417 (4)	C35—H35	0.9300
C11—C12	1.519 (4)	C36—C37	1.368 (5)
C12—C14	1.527 (4)	C37—C38	1.390 (4)
C12—H12	0.9800	C37—H37	0.9300
C14—C15	1.364 (4)	C38—H38	0.9300
C10—O1—C8	118.2 (2)	C17—C18—H18	120.3
C29—O2—C27	118.59 (19)	C19—C18—H18	120.3
C1—N1—C9	116.9 (2)	C14—C19—C18	121.4 (3)
C10—N2—H1	114 (2)	C14—C19—H19	119.3
C10—N2—H2	115 (2)	C18—C19—H19	119.3
H1—N2—H2	120 (3)	N4—C20—C21	124.2 (3)
C20—N4—C28	116.4 (2)	N4—C20—H20	117.9
C29—N5—H3	114 (2)	C21—C20—H20	117.9
C29—N5—H4	118.3 (19)	C22—C21—C20	119.0 (3)
H3—N5—H4	111 (3)	C22—C21—H21	120.5
N1—C1—C2	124.9 (3)	C20—C21—H21	120.5
N1—C1—H1A	117.5	C21—C22—C23	120.0 (3)
C2—C1—H1A	117.5	C21—C22—H22	120.0
C3—C2—C1	118.2 (3)	C23—C22—H22	120.0
C3—C2—H2A	120.9	C28—C23—C22	116.5 (3)
C1—C2—H2A	120.9	C28—C23—C24	118.9 (3)
C2—C3—C4	120.3 (3)	C22—C23—C24	124.6 (3)
C2—C3—H3A	119.9	C25—C24—C23	120.8 (3)
C4—C3—H3A	119.9	C25—C24—H24	119.6
C3—C4—C9	116.8 (3)	C23—C24—H24	119.6
C3—C4—C5	124.5 (3)	C24—C25—C26	122.0 (3)
C9—C4—C5	118.7 (3)	C24—C25—H25	119.0
C6—C5—C4	120.9 (3)	C26—C25—H25	119.0
C6—C5—H5	119.6	C27—C26—C25	117.0 (2)
C4—C5—H5	119.6	C27—C26—C31	122.0 (2)
C5—C6—C7	122.0 (3)	C25—C26—C31	120.9 (2)
C5—C6—H6	119.0	C26—C27—O2	123.4 (2)

C7—C6—H6	119.0	C26—C27—C28	123.0 (2)
C8—C7—C6	117.2 (3)	O2—C27—C28	113.7 (2)
C8—C7—C12	122.2 (2)	N4—C28—C23	123.8 (2)
C6—C7—C12	120.5 (2)	N4—C28—C27	118.0 (2)
C7—C8—O1	122.9 (2)	C23—C28—C27	118.2 (2)
C7—C8—C9	123.2 (2)	C30—C29—N5	127.9 (3)
O1—C8—C9	114.0 (2)	C30—C29—O2	122.4 (2)
N1—C9—C4	122.8 (3)	N5—C29—O2	109.7 (2)
N1—C9—C8	119.1 (2)	C29—C30—C32	116.8 (2)
C4—C9—C8	118.0 (2)	C29—C30—C31	124.2 (2)
C11—C10—O1	123.9 (2)	C32—C30—C31	119.0 (2)
C11—C10—N2	127.7 (3)	C26—C31—C33	114.4 (2)
O1—C10—N2	108.3 (2)	C26—C31—C30	109.2 (2)
C10—C11—C13	117.9 (3)	C33—C31—C30	110.4 (2)
C10—C11—C12	122.8 (2)	C26—C31—H31	107.5
C13—C11—C12	119.1 (3)	C33—C31—H31	107.5
C7—C12—C11	109.6 (2)	C30—C31—H31	107.5
C7—C12—C14	113.5 (2)	N6—C32—C30	178.9 (4)
C11—C12—C14	112.8 (2)	C34—C33—C38	118.0 (3)
C7—C12—H12	106.8	C34—C33—C31	120.1 (3)
C11—C12—H12	106.8	C38—C33—C31	121.7 (2)
C14—C12—H12	106.8	C33—C34—C35	120.8 (3)
N3—C13—C11	178.9 (4)	C33—C34—H34	119.6
C15—C14—C19	117.7 (3)	C35—C34—H34	119.6
C15—C14—C12	120.4 (3)	C36—C35—C34	119.7 (3)
C19—C14—C12	121.8 (3)	C36—C35—H35	120.2
C14—C15—C16	121.7 (4)	C34—C35—H35	120.2
C14—C15—H15	119.1	C35—C36—C37	120.8 (3)
C16—C15—H15	119.1	C35—C36—C12	119.4 (3)
C17—C16—C15	118.7 (3)	C37—C36—C12	119.8 (3)
C17—C16—H16	120.7	C36—C37—C38	119.3 (3)
C15—C16—H16	120.7	C36—C37—H37	120.3
C16—C17—C18	121.2 (3)	C38—C37—H37	120.3
C16—C17—C11	119.4 (3)	C33—C38—C37	121.4 (3)
C18—C17—C11	119.4 (3)	C33—C38—H38	119.3
C17—C18—C19	119.3 (3)	C37—C38—H38	119.3
C9—N1—C1—C2	-1.2 (4)	C28—N4—C20—C21	0.1 (5)
N1—C1—C2—C3	-0.6 (5)	N4—C20—C21—C22	1.0 (5)
C1—C2—C3—C4	1.5 (4)	C20—C21—C22—C23	-0.1 (5)
C2—C3—C4—C9	-0.7 (4)	C21—C22—C23—C28	-1.8 (4)
C2—C3—C4—C5	-179.6 (3)	C21—C22—C23—C24	178.1 (3)
C3—C4—C5—C6	178.5 (3)	C28—C23—C24—C25	0.9 (5)
C9—C4—C5—C6	-0.5 (5)	C22—C23—C24—C25	-178.9 (3)
C4—C5—C6—C7	0.7 (5)	C23—C24—C25—C26	-1.9 (5)
C5—C6—C7—C8	0.6 (5)	C24—C25—C26—C27	0.6 (4)
C5—C6—C7—C12	178.9 (3)	C24—C25—C26—C31	-179.0 (3)
C6—C7—C8—O1	176.7 (3)	C25—C26—C27—O2	-177.5 (2)

C12—C7—C8—O1	-1.6 (4)	C31—C26—C27—O2	2.2 (4)
C6—C7—C8—C9	-2.1 (4)	C25—C26—C27—C28	1.8 (4)
C12—C7—C8—C9	179.6 (3)	C31—C26—C27—C28	-178.6 (3)
C10—O1—C8—C7	3.9 (4)	C29—O2—C27—C26	-5.2 (4)
C10—O1—C8—C9	-177.2 (2)	C29—O2—C27—C28	175.5 (2)
C1—N1—C9—C4	2.1 (4)	C20—N4—C28—C23	-2.2 (4)
C1—N1—C9—C8	-179.1 (3)	C20—N4—C28—C27	179.7 (3)
C3—C4—C9—N1	-1.2 (4)	C22—C23—C28—N4	3.0 (4)
C5—C4—C9—N1	177.8 (3)	C24—C23—C28—N4	-176.8 (3)
C3—C4—C9—C8	180.0 (3)	C22—C23—C28—C27	-178.8 (3)
C5—C4—C9—C8	-1.0 (4)	C24—C23—C28—C27	1.3 (4)
C7—C8—C9—N1	-176.5 (3)	C26—C27—C28—N4	175.5 (3)
O1—C8—C9—N1	4.6 (4)	O2—C27—C28—N4	-5.2 (4)
C7—C8—C9—C4	2.3 (4)	C26—C27—C28—C23	-2.7 (4)
O1—C8—C9—C4	-176.6 (2)	O2—C27—C28—C23	176.6 (2)
C8—O1—C10—C11	-0.1 (4)	C27—O2—C29—C30	3.4 (4)
C8—O1—C10—N2	178.2 (2)	C27—O2—C29—N5	-174.5 (2)
O1—C10—C11—C13	179.8 (3)	N5—C29—C30—C32	-2.3 (5)
N2—C10—C11—C13	1.9 (5)	O2—C29—C30—C32	-179.7 (3)
O1—C10—C11—C12	-6.0 (5)	N5—C29—C30—C31	178.9 (3)
N2—C10—C11—C12	176.0 (3)	O2—C29—C30—C31	1.4 (5)
C8—C7—C12—C11	-3.7 (4)	C27—C26—C31—C33	126.5 (3)
C6—C7—C12—C11	178.1 (3)	C25—C26—C31—C33	-53.9 (3)
C8—C7—C12—C14	-130.8 (3)	C27—C26—C31—C30	2.3 (4)
C6—C7—C12—C14	51.0 (4)	C25—C26—C31—C30	-178.1 (2)
C10—C11—C12—C7	7.4 (4)	C29—C30—C31—C26	-4.0 (4)
C13—C11—C12—C7	-178.5 (3)	C32—C30—C31—C26	177.1 (3)
C10—C11—C12—C14	134.9 (3)	C29—C30—C31—C33	-130.6 (3)
C13—C11—C12—C14	-51.0 (4)	C32—C30—C31—C33	50.5 (4)
C7—C12—C14—C15	-132.9 (3)	C26—C31—C33—C34	149.0 (3)
C11—C12—C14—C15	101.7 (3)	C30—C31—C33—C34	-87.4 (3)
C7—C12—C14—C19	48.7 (4)	C26—C31—C33—C38	-36.7 (4)
C11—C12—C14—C19	-76.7 (4)	C30—C31—C33—C38	87.0 (3)
C19—C14—C15—C16	-0.7 (5)	C38—C33—C34—C35	-0.1 (4)
C12—C14—C15—C16	-179.2 (3)	C31—C33—C34—C35	174.5 (3)
C14—C15—C16—C17	-0.3 (5)	C33—C34—C35—C36	0.4 (5)
C15—C16—C17—C18	1.1 (5)	C34—C35—C36—C37	-0.1 (5)
C15—C16—C17—C11	179.8 (3)	C34—C35—C36—C12	179.2 (2)
C16—C17—C18—C19	-0.9 (5)	C35—C36—C37—C38	-0.6 (5)
C11—C17—C18—C19	-179.5 (2)	C12—C36—C37—C38	-179.9 (2)
C15—C14—C19—C18	1.0 (5)	C34—C33—C38—C37	-0.6 (4)
C12—C14—C19—C18	179.5 (3)	C31—C33—C38—C37	-175.1 (2)
C17—C18—C19—C14	-0.3 (5)	C36—C37—C38—C33	0.9 (5)

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the N4,C20–C23,C28 and C33–C38 rings, respectively.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H1···N4 ⁱ	0.84 (3)	2.34 (4)	3.172 (4)	174 (3)
N2—H2···N5 ⁱ	0.89 (3)	2.61 (3)	3.308 (5)	136 (2)
N5—H3···N1	0.87 (3)	2.15 (3)	3.014 (3)	173 (3)
C18—H18···Cg1 ⁱⁱ	0.93	2.80	3.650 (3)	152
C24—H24···Cg2 ⁱⁱⁱ	0.93	2.74	3.668 (3)	174

Symmetry codes: (i) $x-1, y, z$; (ii) $-x-3/2, y-1/2, -z-1/2$; (iii) $-x+2, -y+1, -z+1$.