

Research Article

Planck-Scale Dual-Curvature Lensing and Spacetime Noncommutativity

Giovanni Amelino-Camelia,¹ Leonardo Barcaroli,¹ Stefano Bianco,² and Laura Pensato¹

¹Dipartimento di Fisica, Università “La Sapienza” and Sez. Roma1 INFN, P.le A. Moro 2, 00185 Roma, Italy

²Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Am Mühlenberg 1, Golm, 14476 Potsdam, Germany

Correspondence should be addressed to Giovanni Amelino-Camelia; amelino@roma1.infn.it

Received 10 May 2017; Accepted 22 June 2017; Published 10 August 2017

Academic Editor: Sally Seidel

Copyright © 2017 Giovanni Amelino-Camelia et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

It was recently realized that Planck-scale momentum-space curvature, which is expected in some approaches to the quantum-gravity problem, can produce dual-curvature lensing, a feature which mainly affects the direction of observation of particles emitted by very distant sources. Several gray areas remain in our understanding of dual-curvature lensing, including the possibility that it might be just a coordinate artifact and the possibility that it might be in some sense a by-product of the better studied dual-curvature redshift. We stress that data reported by the IceCube neutrino telescope should motivate a more vigorous effort of investigation of dual-curvature lensing, and we observe that studies of the recently proposed “ ρ -Minkowski noncommutative spacetime” could be valuable from this perspective. Through a dedicated ρ -Minkowski analysis, we show that dual-curvature lensing is not merely a coordinate artifact and that it can be present even in theories without dual-curvature redshift.

1. Introduction

There has been considerable interest (see, e.g., [1–7]) over the last decade for the possibility that Planck-scale structures might have observably large implications for particle propagation over cosmological distances. We shall here focus on one of the scenarios for the quantum-gravity realm which can motivate such studies, the scenario such that the characteristic scale of quantum gravity (usually assumed to be of the order of the Planck scale) plays the role of scale of curvature of momentum space [8–11]. The most studied effect of this sort is “dual-curvature redshift” [12], an effect such that two ultrarelativistic (massless or effectively massless) particles of different energies emitted simultaneously at a distant source reach a detector at different times (earlier discussions of this effect, before the curved-momentum-space perspective emerged, can be found, e.g., in [1–5]). Recently it became clear that Planck-scale curvature of momentum space can also produce the effect of “dual-curvature lensing” [13–15] (or “dual-gravity lensing” (the effect is better known as dual-gravity lensing, but actually it is present also in absence of momentum-space geometrodynamics: momentum-space

curvature is sufficient for producing the effect. We also note that all models with Planck-scale-curved momentum space developed so far indeed do not have momentum-space geometrodynamics, but rather have a fixed background curved momentum-space geometry. We reserve the characterization of effects as “gravitational” only when geometrodynamics is present)), an effect such that two ultrarelativistic particles of different energies emitted by the same distant source reach a detector from different directions.

The understanding of dual-curvature lensing has not yet reached the level of our understanding of dual-curvature redshift. Most notably previous studies establish that dual-curvature redshift is a truly physical (observable) effect but leave the possibility that dual-curvature lensing might be just a coordinate artifact open. Moreover, these previous works do not clarify whether dual-curvature lensing is just some sort of by-product of dual-curvature redshift or instead one could have theories without dual-curvature redshift but with dual-curvature lensing.

This state of affairs became recently more disappointing since data reported by the IceCube telescope [16]

open a potentially powerful observational window on dual-curvature lensing, and actually a first exploratory analysis gave preliminarily encouraging results [17].

We here focus on the recently proposed “ ρ -Minkowski noncommutative spacetime” [14], using it as a conceptual laboratory for dual-curvature lensing. We show that ρ -Minkowski produces dual-curvature lensing, but dual-curvature redshift is absent. Most importantly we establish that in ρ -Minkowski dual-curvature lensing is a truly physical effect, not merely a coordinate artifact.

2. ρ -Minkowski and Its Relativistic Properties

We start by characterizing the relativistic properties of the ρ -Minkowski noncommutative spacetime. ρ -Minkowski is a 3D noncommutative spacetime, characterized by the following commutation relations among spacetime coordinates:

$$\begin{aligned} [x_i, x_0] &= \rho \epsilon_{ij} x_j, \\ [x_i, x_j] &= 0, \end{aligned} \quad (1)$$

where $i \in \{1, 2\}$.

As usual for this sort of noncommutative spacetime, in order to have an explicit description of the relativistic properties one must adopt an ordering convention for the basis of exponentials. We find that it is convenient to adopt time-to-the-right ordering, so that elements of the basis of exponentials are written as follows:

$$e^{ip_j x_j} e^{ip_0 x_0}. \quad (2)$$

Relying again on experience gained working with other similar noncommutative spacetimes, we infer the form of the law of composition of momenta by studying the properties of products of elements of the basis of exponentials. We have that

$$\begin{aligned} e^{ip_j x_j} e^{ip_0 x_0} e^{iq_l x_l} e^{iq_0 x_0} \\ \approx (1 + ip_j x_j) (1 + ip_0 x_0) (1 + iq_l x_l) (1 + iq_0 x_0) \\ \approx (1 + i(p_j + q_j - \rho p_0 \epsilon_{jl} q_l) x_j) (1 + i(p_0 + q_0) x_0), \end{aligned} \quad (3)$$

where we used (1) and we focused on the leading correction in ρ .

Result (3) suggests that, at leading order in ρ , the law of composition of momenta in ρ -Minkowski should take the form:

$$\begin{aligned} (p \oplus q)_0 &= p_0 + q_0, \\ (p \oplus q)_i &= p_i + q_i - \rho p_0 \epsilon_{ij} q_j. \end{aligned} \quad (4)$$

We shall here establish that ρ -Minkowski allows the formulation of a DSR-relativistic theory, in the sense first introduced in [18, 19] (also see [20, 21]), that is, a relativistic theory with two nontrivial relativistic invariant, also a momentum scale, in this case ρ^{-1} , in addition to the speed-of-light scale (which we set to unity by choice of dimensions but

can be easily brought back into the picture by dimensional analysis). The label “DSR” refers to the fact that this class of relativistic theories was at first called “doubly special reactivity.”

The DSR-compatibility of the setup requires that the action of boosts on momenta is introduced by requiring that the deformed momentum-conservation law is covariant. In order to ensure this it suffices to require that a vanishing composed momentum for observer Alice, $p \oplus q = 0$, also vanishes for observer Bob, boosted with respect to Alice:

$$\left\{ N_i^{[p \oplus q]}, p \oplus q \right\}_{|_{p \oplus q = 0}} = 0. \quad (5)$$

Taking the description of boosts in the rather similar κ -Minkowski noncommutative spacetime as guidance [22–24], in the case of ρ -Minkowski we are led to the following description of the action of boosts on composed momenta, when given in terms of the action of boosts on single-particle momenta:

$$N_i^{[p \oplus q]} = N_i^{[p]} + N_i^{[q]} - \rho p_0 \epsilon_{ij} N_j^{[q]}. \quad (6)$$

For the action of boosts on single-particle momenta we introduce a 6-parameter *ansatz*:

$$\begin{aligned} \{N_i, p_0\} &= p_i + \rho A p_0 p_i, \\ \{N_i, p_j\} &= \delta_{ij} p_0 + \rho \left(B p_0^2 + C |\vec{p}|^2 \right) \delta_{ij} \\ &\quad + \rho \left(D p_0^2 + E |\vec{p}|^2 \right) \epsilon_{ij} + \rho F \epsilon_{il} p_l p_j. \end{aligned} \quad (7)$$

The parameters appearing in (7) can be of course determined by making use of (6) and (7) in the compatibility condition (5), finding that the compatibility condition is satisfied by the following form of the action of boosts on single-particle momenta:

$$\begin{aligned} \{N_i, p_0\} &= p_i, \\ \{N_i, p_j\} &= \delta_{ij} p_0 + \frac{\rho}{2} \left(\epsilon_{ij} |\vec{p}|^2 - \epsilon_{ik} p_k p_j \right). \end{aligned} \quad (8)$$

Building on these starting points it is easy to complete the description of the relativistic symmetries of ρ -Minkowski by simply insisting that all relevant Jacobi identities are satisfied. This leads one to the conclusion that the only other needed deformation is in the Poisson bracket between two boosts,

$$\{N_i, N_j\} = -R \epsilon_{ij} + \frac{\rho}{2} \epsilon_{ij} N_i p_i, \quad (9)$$

while all other remaining Poisson brackets are undeformed.

We conclude that the full description of the relativistic symmetries of ρ -Minkowski is given in terms of the following Poisson brackets:

$$\begin{aligned}
\{p_0, p_i\} &= 0, \\
\{p_i, p_j\} &= 0, \\
\{R, p_0\} &= 0, \\
\{R, p_i\} &= \epsilon_{ij} p_j, \\
\{R, N_i\} &= \epsilon_{ij} N_j, \\
\{N_i, p_0\} &= p_i, \\
\{N_i, p_j\} &= \delta_{ij} p_0 + \frac{\rho}{2} (\epsilon_{ij} |\vec{p}|^2 - \epsilon_{ik} p_k p_j), \\
\{N_i, N_j\} &= -\epsilon_{ij} R + \frac{\rho}{2} \epsilon_{ij} N_a p_a.
\end{aligned} \tag{10}$$

Notably the ordinary special-relativistic mass Casimir,

$$C = p_0^2 - |\vec{p}|^2, \tag{11}$$

is still a Casimir also of the deformed symmetry algebra (10).

3. Dual-Curvature Lensing with Commuting Coordinates

As for most other studies of the implications of Planck-scale momentum-space curvature, we shall here focus on the implications of the associated DSR-deformed relativistic symmetries at the level of the trajectories of particles in the classical limit. Previous related studies [25] have shown that while the noncommutativity of coordinates plays evidently a crucial role in establishing the form of the DSR-relativistic symmetries, the truly physical content of the analysis of classical trajectories is the same whether one uses noncommuting coordinates or commuting ones. In these cases it turns out to be convenient to perform computations both using commuting and using noncommuting coordinates, since this facilitates establishing which properties are truly physical (properties which are coordinate artifacts will change from one formulation to the other). We therefore study dual-curvature lensing in this section using commuting coordinates, while in the next section we perform the same study using coordinates with ρ -Minkowski Poisson brackets.

The symplectic structure adopted for this section is

$$\begin{aligned}
\{\mathbf{x}_i, \mathbf{x}_0\} &= 0, \\
\{\mathbf{x}_i, \mathbf{x}_j\} &= 0, \\
\{p_0, \mathbf{x}_0\} &= 1, \\
\{p_i, \mathbf{x}_0\} &= 0, \\
\{p_0, \mathbf{x}_i\} &= 0, \\
\{p_i, \mathbf{x}_j\} &= -\delta_{ij}.
\end{aligned} \tag{12}$$

It is useful to start by noticing that in terms of this symplectic structure the relativistic-symmetry generators introduced in the previous section admit the following representation:

$$\begin{aligned}
P_\mu &= p_\mu, \\
\mathcal{R} &= \epsilon_{ij} \mathbf{x}_i p_j, \\
\mathcal{N}_i &= \mathbf{x}_i p_0 - \mathbf{x}_0 p_i + \frac{\rho}{2} \epsilon_{ia} (\mathbf{x}_a |\vec{p}|^2 - p_a \vec{\mathbf{x}} \cdot \vec{p}).
\end{aligned} \tag{13}$$

From this representation one finds that the Poisson brackets between rotations and the spacetime coordinates are the standard ones, while the Poisson brackets between boosts and spacetime coordinates are deformed:

$$\begin{aligned}
\{\mathcal{N}_i, \mathbf{x}_0\} &= \mathbf{x}_i \{p_0, \mathbf{x}_0\} = \mathbf{x}_i, \\
\{\mathcal{N}_i, \mathbf{x}_j\} &= -\mathbf{x}_0 \{p_i, \mathbf{x}_j\} \\
&\quad + \frac{\rho}{2} \epsilon_{ia} (\mathbf{x}_a \{|\vec{p}|^2, \mathbf{x}_j\} - p_a \vec{\mathbf{x}} \cdot \{\vec{p}, \mathbf{x}_j\}) \\
&= \mathbf{x}_0 \delta_{ij} + \frac{\rho}{2} \epsilon_{ia} (p_a \mathbf{x}_j - 2 \mathbf{x}_a p_j) + \frac{\rho}{2} \epsilon_{ij} \mathbf{x}_i p_l.
\end{aligned} \tag{14}$$

We can of course derive the equations of motion using the mass Casimir (11) as Hamiltonian of evolution in an affine parameter, finding

$$\begin{aligned}
\dot{\mathbf{x}}_0 &= \{\mathcal{C}, \mathbf{x}_0\} = \{p_0^2 - |\vec{p}|^2, \mathbf{x}_0\} = 2p_0, \\
\dot{\mathbf{x}}_i &= \{\mathcal{C}, \mathbf{x}_i\} = \{p_0^2 - |\vec{p}|^2, \mathbf{x}_i\} = 2p_i, \\
\dot{p}_\mu &= \{\mathcal{C}, p_\mu\} = \{p_0^2 - |\vec{p}|^2, p_\mu\} = 0,
\end{aligned} \tag{15}$$

where \dot{q} denotes the derivative of q with respect to the affine parameter. Momenta are constant of motion and a generic solution of the equations for the world line can be written as

$$\mathbf{x}_i - \bar{\mathbf{x}}_i = \frac{p_i}{p_0} (\mathbf{x}_0 - \bar{\mathbf{x}}_0), \tag{16}$$

where $\bar{\mathbf{x}}_\mu$ is a point of the world line.

Our analysis of dual-curvature lensing starts by considering an observer Alice that emits from its spacetime origin two massless particles of different energies. The relevant two massless particles have spatial momenta $\vec{p}_s = (p_{1s}, 0)$ and $\vec{p}_h = (p_{1h}, 0)$, with $p_{1s} > 0$, $p_{1h} > 0$, and the one with momentum \vec{p}_h is “hard” while the one with momentum \vec{p}_s is “soft,” meaning that $p_{1s} \ll p_{1h}$. From (16), we have that Alice describes the two particles with the same world line

$$\begin{aligned}
\mathbf{x}_{1,s}^A &= \mathbf{x}_{1,h}^A = \mathbf{x}_0^A, \\
\mathbf{x}_{2,s}^A &= \mathbf{x}_{2,h}^A = 0.
\end{aligned} \tag{17}$$

Alice is the observer at the source. The main feature of dual-curvature lensing will be manifest in the description given by a distant detector, with direction-discriminating

capabilities, of these two particles emitted at Alice. The next step of our analysis is to consider an observer Bob, which is related to Alice by a pure translation with parameters $b^\mu = (b, -b, 0)$. Since Bob and Alice are related by a pure translation with parameters $b^\mu = (b, -b, 0)$, one has of course the following relationship between Alice's coordinates and Bob's coordinates:

$$\begin{aligned} \mathbf{x}_0^B &= \mathbf{x}_0^A + b^\mu \{p_\mu, \mathbf{x}_0^A\} = \mathbf{x}_0^A + b, \\ \mathbf{x}_1^B &= \mathbf{x}_1^A + b^\mu \{p_\mu, \mathbf{x}_1^A\} = \mathbf{x}_1^A + b, \\ \mathbf{x}_2^B &= \mathbf{x}_2^A + b^\mu \{p_\mu, \mathbf{x}_2^A\} = \mathbf{x}_2^A. \end{aligned} \quad (18)$$

This leads us immediately to Bob's description of the world lines of the two particles emitted by Alice:

$$\begin{aligned} \mathbf{x}_{1,s}^B &= \mathbf{x}_{1,h}^B = \mathbf{x}_0^B, \\ \mathbf{x}_{2,s}^B &= \mathbf{x}_{2,h}^B = 0, \end{aligned} \quad (19)$$

for which we also used the fact that momenta are invariant under pure translations.

Evidently the map between Alice and Bob, purely translated with respect to Alice, remains trivial and unaffected by the curvature of momentum space. We shall not report explicitly the directional analysis for Bob, but the triviality of the map clearly implies that there is no dual-curvature lensing in the picture involving these two observers, Alice and Bob. We shall find that a truly physical feature (not a coordinate artifact) of dual-curvature lensing is found in ρ -Minkowski when the source and the detector are not just purely translated but rather are related by a composition of a translation and a boost. It is therefore appropriate to consider at this point a third observer Camilla, purely boosted with respect to Bob. Observer Camilla is therefore obtained from Alice by first acting with a translation with parameters $b^\mu = (b, -b, 0)$ and then with a boost with parameters $\xi^i = (\xi^1, 0)$

$$\begin{aligned} \mathbf{x}_\mu^C &= T_{\xi^i} (T_{b^\mu} (\mathbf{x}_\mu^A)) = T_{\xi^i} (\mathbf{x}_\mu^B), \\ p_\mu^C &= T_{\xi^i} (T_{b^\mu} (p_\mu^A)) = T_{\xi^i} (p_\mu^B). \end{aligned} \quad (20)$$

As a first step toward writing the world lines of the soft and the hard particle in Camilla's reference frame, we give the transformation laws for momentum between Camilla's frame and Alice's frame, specialized to the case of our interest (in which $p_\mu^A = (p_0^A, p_1^A, 0)$):

$$\begin{aligned} p_0^C &= p_0^A + \xi^1 \{ \mathcal{N}_1, p_0^A \} = p_0^A + \xi^1 p_1^A, \\ p_1^C &= p_1^A + \xi^1 \{ \mathcal{N}_1, p_1^A \} = p_1^A + \xi^1 p_0^A, \\ p_2^C &= p_2^A + \xi^1 \{ \mathcal{N}_1, p_2^A \} = \xi^1 \left(\frac{p}{2} (p_1^A)^2 \right). \end{aligned} \quad (21)$$

Here one sees that the ρ -Minkowski deformation of boost transformations is such that while for Alice the spatial parts of the momenta of the soft and hard particles are both directed along the \mathbf{x}^1 -axis, in Camilla's frame the momenta

have a component also along the \mathbf{x}^2 -axis, proportional to the energies of the particles in Alice's frame.

Our next step is to determine which coordinates Camilla assigns to the origin of Alice, so we consider $\bar{\mathbf{x}}_\mu^C = T_{\xi^i} (T_{b^\mu} (\bar{\mathbf{x}}_\mu^A = (0, 0, 0)))$, finding that

$$\begin{aligned} \mathbf{x}_0^C &= b + \xi^1 b, \\ \mathbf{x}_1^C &= b + \xi^1 b, \\ \mathbf{x}_2^C &= 0, \end{aligned} \quad (22)$$

which is valid both for the soft particle and for the hard particle. We note that after computing the Poisson brackets we used that $p_{2h}^A = 0$ and that $T_{b^{2h}} = 0$.

We are now ready to give the world lines of the particles according to Camilla. For the soft particle Camilla has

$$\begin{aligned} \mathbf{x}_{1,s}^C &= \mathbf{x}_0^C, \\ \mathbf{x}_{2,s}^C &= 0, \end{aligned} \quad (23)$$

where we neglected the deformation effects (the soft particle is a low-energy particle, and the deformation here of interest grows with energy), while Camilla's description of the world line of the hard particle is

$$\begin{aligned} \mathbf{x}_{1,h}^C &= \mathbf{x}_0^C, \\ \mathbf{x}_{2,h}^C &= \frac{\rho}{2} p_{0,h}^C \xi^1 \mathbf{x}_0^C. \end{aligned} \quad (24)$$

This establishes that both the soft particle and the hard particles go through the spacetime origin of Camilla's reference frame, but we are evidently most interested in what measurement results Camilla would get for the direction of the two particles. For this purpose it is important to notice that a good measurement procedure for directions involves the setup of a rigid nonpoint-like detector: we establish the direction of a particle by seeing that the particle went through two (or more) points of the rigid extended detector. Since we are thinking of possible applications of our results to the analysis of IceCube-telescope data, it is nice to notice that IceCube establishes directions exactly in this way. From a relativist perspective the notion of an extended detector is a bit cumbersome, since it is standard practice to abstract the notion of a point-like detector in association with the equally abstract notion of relativistic observer. We can however still rely on these standard abstractions by viewing a rigid extended detector as a network of point-like detectors in rigid motion. The results will be more transparent by introducing a different relativistic observer for each of the point-like detectors composing the extended detector, so that each of these many observers has a point-like detector in the spatial origin of its reference frame.

These conceptual considerations translate in the technical challenge for our computations of introducing two more observers which we will simply denote with D and D' . Both the soft particle and the hard particle go through the origin of Camilla's reference frame, so if we find that the soft (hard)

particle also goes through D (D') then we will have enough information for establishing the direction of the soft (hard) particle. It is appropriate to take D as an observer, at rest with respect to Camilla, whose origin of the reference frame coincides with the point which for Camilla has coordinates $\mathbf{x}_{\mu,D}^C = (\delta, \delta, 0)$. The observer D is related to Camilla by a translation with parameters $\delta^\mu = (\delta, -\delta, 0)$, so that

$$\begin{aligned}\mathbf{x}_0^D &= \mathbf{x}_{0,D}^C + \delta^\mu \{p_\mu, \mathbf{x}_{0,D}^C\} = \mathbf{x}_{0,D}^C + \delta, \\ \mathbf{x}_1^D &= \mathbf{x}_{1,D}^C + \delta^\mu \{p_\mu, \mathbf{x}_{1,D}^C\} = \mathbf{x}_{1,D}^C + \delta, \\ \mathbf{x}_2^D &= \mathbf{x}_{2,D}^C + \delta^\mu \{p_\mu, \mathbf{x}_{2,D}^C\} = \mathbf{x}_{2,D}^C.\end{aligned}\quad (25)$$

This straightforwardly leads us to the description given by observer D of the world lines of the soft particle and of the hard particle:

$$\begin{aligned}\mathbf{x}_{1,s}^D &= \mathbf{x}_0^D, \\ \mathbf{x}_{2,s}^D &= 0, \\ \mathbf{x}_{1,h}^D &= \mathbf{x}_0^D, \\ \mathbf{x}_{2,h}^D &= \frac{\rho}{2} p_{0,h}^D \xi^1 (\mathbf{x}_0^D - \delta),\end{aligned}\quad (26)$$

where $p_{0,h}^D = p_{0,h}^C$, since observer Camilla and observer D are connected by a pure translation. Notice that these equations for the world lines show in particular that the soft particle goes through the origin of observer D but the hard particle does not.

Next it is convenient to consider another observer D' , also at rest with respect to Camilla, whose origin of the reference frame coincides with the point which for Camilla has coordinates $\mathbf{x}_{\mu,D'}^C = (\delta, +\delta, +(\rho/2)p_{0,h}^C \xi^1 \delta)$. The observer D' is related to Camilla by a translation with parameters $\delta^\mu = (\delta, -\delta, -(\rho/2)p_{0,h}^C \xi^1 \delta)$, so that we have

$$\begin{aligned}\mathbf{x}_0^{D'} &= \mathbf{x}_{0,D'}^C + \delta^\mu \{p_\mu, \mathbf{x}_{0,D'}^C\} = \mathbf{x}_{0,D'}^C + \delta, \\ \mathbf{x}_1^{D'} &= \mathbf{x}_{1,D'}^C + \delta^\mu \{p_\mu, \mathbf{x}_{1,D'}^C\} = \mathbf{x}_{1,D'}^C + \delta, \\ \mathbf{x}_2^{D'} &= \mathbf{x}_{2,D'}^C + \delta^\mu \{p_\mu, \mathbf{x}_{2,D'}^C\} = \mathbf{x}_{2,D'}^C + \frac{\rho}{2} p_{0,h}^C \xi^1 \delta.\end{aligned}\quad (27)$$

Therefore according to observer D' the world lines of the soft particle and of the hard particle are described by

$$\begin{aligned}\mathbf{x}_{1,s}^{D'} &= \mathbf{x}_0^{D'}, \\ \mathbf{x}_{2,s}^{D'} &= \frac{\rho}{2} p_{0,h}^{D'} \xi^1 \delta, \\ \mathbf{x}_{1,h}^{D'} &= \mathbf{x}_0^{D'}, \\ \mathbf{x}_{2,h}^{D'} &= \frac{\rho}{2} p_{0,h}^{D'} \xi^1 \mathbf{x}_0^{D'},\end{aligned}\quad (28)$$

where $p_{0,h}^{D'} = p_{0,h}^D = p_{0,h}^C$, since the observers Camilla, D , and D' are all connected by a pure translation. Notice that these

equations for the world lines show in particular that the hard particle goes through the origin of observer D but the soft particle does not.

So we found that the soft particle goes through D and Camilla, while the hard particle goes through D' and Camilla. This is the dual-curvature lensing that we were looking for: the particles are emitted by the same source but their detection manifests a different direction of propagation. From the fact that the soft particle goes through D and Camilla and the hard particle goes through D' and Camilla one easily finds that the angle θ characterizing the difference in their observed directions of propagation is given by

$$\tan \theta \approx \frac{\rho}{2} p_{0,h}^C \xi^1. \quad (29)$$

4. Dual-Curvature Lensing with ρ -Minkowski Commuting Coordinates

As announced earlier, we shall redo, now assuming ρ -Minkowski Poisson brackets, the analysis done in the previous section with “commutative” Poisson brackets. We shall find the same physical result for dual-curvature lensing, but several details of the derivation will be different.

So we now adopt the following symplectic structure:

$$\begin{aligned}\{x_i, x_0\} &= \rho \epsilon_{ij} x_j, \\ \{x_i, x_j\} &= 0, \\ \{p_0, x_0\} &= 1, \\ \{p_0, x_i\} &= 0, \\ \{p_i, x_0\} &= 0, \\ \{p_i, x_j\} &= -\delta_{ij} + \rho \epsilon_{ij} p_0.\end{aligned}\quad (30)$$

The fact that the physical results cannot depend on the choice between this symplectic structure and the symplectic structure used in the previous section is clear upon noticing that these two symplectic structures are connected by a simple momentum-dependent redefinition of spacetime coordinates:

$$\begin{aligned}\mathbf{x}_0 &= x_0, \\ \mathbf{x}_i &= x_i - \rho \epsilon_{ij} x_j p_0,\end{aligned}\quad (31)$$

where \mathbf{x}_μ are the “commutative” coordinates used in the previous section, while x_μ are used in this section.

In terms of the x_μ coordinates the representation of the generators of the relativistic symmetries is

$$\begin{aligned}P_\mu &= p_\mu, \\ \mathcal{R} &= \epsilon_{ij} x_i p_j - \rho p_0 \vec{x} \cdot \vec{p}, \\ \mathcal{N}_i &= x_i p_0 - x_0 p_i \\ &\quad + \frac{\rho}{2} \epsilon_{ij} \left(x_j |\vec{p}|^2 - p_j \vec{x} \cdot \vec{p} - 2x_j p_0^2 \right).\end{aligned}\quad (32)$$

Accordingly we then have that

$$\begin{aligned}
\{\mathcal{R}, x_0\} &= \{\epsilon_{ij}x_i p_j - \rho p_0 x_i p_i, x_0\} = 0, \\
\{\mathcal{R}, x_i\} &= \{\epsilon_{ab}x_a p_b - \rho p_0 x_a p_a, x_i\} = \epsilon_{ab}x_a \{p_b, x_i\} \\
&\quad - \rho p_0 x_a \{p_a, x_i\} = \epsilon_{ab}x_a (-\delta_{bi} + \rho \epsilon_{bi} p_0) \\
&\quad + \rho p_0 x_a \delta_{ai} = \epsilon_{ia}x_a + \rho(-x_i p_0 + p_0 x_i) = \epsilon_{ia}x_a, \\
\{\mathcal{N}_i, x_0\} &= \left\{ x_i p_0 - x_0 p_i \right. \\
&\quad \left. + \frac{\rho}{2} \epsilon_{ia} \left(x_a |\vec{p}|^2 - p_a \vec{x} \cdot \vec{p} - 2x_a p_0^2 \right), x_0 \right\} \\
&= x_i \{p_0, x_0\} + p_0 \{x_i, x_0\} - \frac{\rho}{2} \epsilon_{ia} 2x_a \{p_0^2, x_0\} = x_i \\
&\quad - \rho \epsilon_{ij} x_j p_0, \\
\{\mathcal{N}_i, x_j\} &= \left\{ x_i p_0 - x_0 p_i \right. \\
&\quad \left. + \frac{\rho}{2} \epsilon_{ia} \left(x_a |\vec{p}|^2 - p_a \vec{x} \cdot \vec{p} - 2x_a p_0^2 \right), x_j \right\} \\
&= -x_0 \{p_i, x_j\} - p_i \{x_0, x_j\} + \frac{\rho}{2} \epsilon_{ia} \left(x_a \{|\vec{p}|^2, x_j\} \right. \\
&\quad \left. - x_b \{p_a p_b, x_j\} \right) = -x_0 (-\delta_{ij} + \rho \epsilon_{ij} p_0) + \rho \epsilon_{jk} x_k p_i \\
&\quad + \frac{\rho}{2} \epsilon_{ia} (-2x_a p_j + x_b (\delta_{bj} p_a + \delta_{aj} p_b)) = \delta_{ij} x_0 \\
&\quad + \rho \left(\frac{1}{2} (\epsilon_{ia} p_a x_j + \epsilon_{ij} x_a p_a) + \epsilon_{jk} x_k p_i - \epsilon_{ia} x_a p_j \right. \\
&\quad \left. - \epsilon_{ij} p_0 x_0 \right).
\end{aligned} \tag{33}$$

It is equally straightforward to see that the equations of motion take the form

$$\begin{aligned}
\dot{x}_0 &= \{\mathcal{C}, x_0\} = \{p_0^2 - |\vec{p}|^2, x_0\} = 2p_0, \\
\dot{x}_i &= \{\mathcal{C}, x_i\} = \{p_0^2 - |\vec{p}|^2, x_i\} = -2p_a \{p_a, x_i\} \\
&= 2p_i + 2\rho \epsilon_{ij} p_j p_0, \\
\dot{p}_\mu &= \{\mathcal{C}, x_0\} = \{p_0^2 - |\vec{p}|^2, x_0\} = 0.
\end{aligned} \tag{34}$$

This implies that momenta are constant of motion and that a generic world line can be written as

$$x_i - \bar{x}_i = \left(\frac{p_i}{p_0} + \rho \epsilon_{ij} p_j \right) (x_0 - \bar{x}_0), \tag{35}$$

where p_μ is the momentum of the particle and \bar{x}_μ is a point of the world line.

Of course we shall focus again on the same two massless particles emitted at Alice already considered in the previous section. With the choice of coordinates made in this section

one has that Alice's description of the relevant two world lines is

$$\begin{aligned}
x_{1,s}^A &= x_0^A, \\
x_{2,s}^A &= 0, \\
x_{1,h}^A &= x_0^A, \\
x_{2,h}^A &= -\rho p_0 x_0^A,
\end{aligned} \tag{36}$$

where p_0 denotes the energy of the hard particle ($p_0 \equiv p_{0,h}$) while in the equation for $x_{2,s}^A$ we neglected a term going with the energy of the soft particle (we are assuming again that for the soft particle the ρ -dependent effects will be negligible). Notice that even though the two particles have $p_{2,s} = p_{2,s} = 0$, in the coordinatization used in this section one gets that x_2 changes over time (visible in (36) for the hard particle, neglected for the soft particle). This is just a coordinate artifact due to the symplectic structure (30), and indeed even in this coordinatization the particles reach Bob (the observer already introduced in the previous section, which is purely translated with respect to Alice along the x_1 direction). In order to verify this important fact we start by characterizing the transformation between Alice's coordinates and Bob coordinates, using the coordinatization of this section, taking into account the fact that again the translation parameters connecting Alice and Bob are $b^\mu = (b, -b, 0)$:

$$\begin{aligned}
x_0^B &= x_0^A + b^\mu \{p_\mu, x_0^A\} = x_0^A + b, \\
x_1^B &= x_1^A + b^\mu \{p_\mu, x_1^A\} = x_1^A + b, \\
x_2^B &= x_2^A + b^\mu \{p_\mu, x_2^A\} = x_2^A - \rho b p_0^A.
\end{aligned} \tag{37}$$

Equipped with this one easily arrives at Bob's of the world lines of the soft particle and of the hard particle:

$$\begin{aligned}
x_{1,s}^B &= x_0^B, \\
x_{2,s}^B &= 0, \\
x_{1,h}^B &= x_0^B, \\
x_{2,h}^B &= -\rho p_{0,h}^B x_0^B.
\end{aligned} \tag{38}$$

This shows that indeed both the soft particle and the hard particle go through Bob's spacetime origin. With the “ ρ -Minkowski coordinates” we are using in this section there is a coordinate artifact apparently attributing to the particles a motion also along the x_2 direction, but physically our two particles just move along the 1 direction and indeed they are emitted at Alice and they are detected at Bob, with Bob purely translated with respect to Alice along the 1 direction.

The next step toward our goal of characterizing dual-curvature lensing with ρ -Minkowski coordinates is again the introduction of observer Camilla, the observer Camilla which is purely boosted with respect to Bob (and therefore

obtained from Alice by performing first a translation and then a boost):

$$\begin{aligned}x_{\mu}^C &= T_{\xi^i} (T_{b^{\mu}} (x_{\mu}^A)) = T_{\xi^i} (x_{\mu}^B), \\p_{\mu}^C &= T_{\xi^i} (T_{b^{\mu}} (p_{\mu}^A)) = T_{\xi^i} (p_{\mu}^B),\end{aligned}\quad (39)$$

where $b^{\mu} = (b, -b, 0)$ and $\xi^i = (\xi^1, 0)$.

The transformation laws for momenta between Alice's frame and Camilla's frame, specialized to the case of our interest, in which the spatial momentum of the particles does not have a component along the x_2 direction, take the following form:

$$\begin{aligned}p_0^C &= p_0^A + \xi^1 \{ \mathcal{N}_1, p_0^A \} = p_0^A + \xi^1 p_1^A, \\p_1^C &= p_1^A + \xi^1 \{ \mathcal{N}_1, p_1^A \} = p_1^A + \xi^1 p_0^A, \\p_2^C &= p_2^A + \xi^1 \{ \mathcal{N}_1, p_2^A \} = \xi^1 \left(\frac{\rho}{2} (p_1^A)^2 \right).\end{aligned}\quad (40)$$

Then proceeding just with the same steps of derivation discussed in the previous section one easily arrives at the description of the world lines according to Camilla:

$$\begin{aligned}x_{1,s}^C &= x_0^C, \\x_{2,s}^C &= 0, \\ \text{hard } x_{1,h}^C &= x_0^C, \\x_{2,h}^C &= -\rho p_{0,h}^C x_0^C + \frac{\xi^1}{2} \rho p_{0,h}^C x_0^C.\end{aligned}\quad (41)$$

Importantly both the soft particle and the hard particle go through the spacetime origin of Camilla's reference frame.

Toward our goal of describing dual-curvature lensing with ρ -Minkowski coordinates, we are left with only the final step, the one involving the observers D and D' . The steps of derivation are the same as in the previous section. Some formulas take a different form but the conclusions for the physical feature of dual-curvature lensing are unchanged. One easily finds that according to observer D the world lines take the form

$$\begin{aligned}x_{1,s}^D &= x_0^D, \\x_{2,s}^D &= 0, \\x_{1,h}^D &= x_0^D, \\x_{2,h}^D &= -\rho p_{0,h}^D x_0^D + \frac{\rho}{2} \xi^1 p_{0,h}^D x_0^D - \frac{\rho}{2} \xi^1 \delta p_{0,h}^D,\end{aligned}\quad (42)$$

which again show that the soft particle goes through the spacetime origin of observer D , while the hard particle does not.

Observer D' describes the same world lines as follows:

$$\begin{aligned}x_{1,s}^{D'} &= x_0^{D'}, \\x_{2,s}^{D'} &= \frac{\rho}{2} p_{0,h}^{D'} \xi^1 \delta, \\ \text{hard } x_{1,h}^{D'} &= x_0^{D'}, \\x_{2,h}^{D'} &= -\rho p_{0,h}^{D'} x_0^{D'} + \frac{\rho}{2} \xi^1 p_{0,h}^{D'} x_0^{D'},\end{aligned}\quad (43)$$

which again show that the hard particle goes through the spacetime origin of observer D , while the soft particle does not.

Finally reasoning just as done at the end of the previous section one finds from these results that the angle θ characterizing the difference in their observed directions of propagation is given by

$$\tan \theta \simeq \frac{\rho}{2} p_{0,h}^C \xi^1. \quad (44)$$

This is the same result found at the end of the previous section. While some intermediate steps of derivation do depend on the choice of coordinatization, the result for dual-curvature lensing is the same independently on whether one uses "commuting coordinates," as done in the previous section, or " ρ -Minkowski coordinates," as done in this section.

5. Closing Remarks

We here made some significant steps toward the understanding of dual-curvature lensing, most notably established as a truly physical effect. The noncommutative ρ -Minkowski turned out to be ideally suited for investigating dual-curvature lensing at the classical-mechanics level, and it should naturally provide a good option for exploring quantum manifestations of dual-curvature lensing, in the sense of studies of dual-curvature redshift such as the one reported in [26]. Of course, ultimately the most significant applications will concern phenomenology. In this respect it is noteworthy that for ρ -Minkowski we found dual-curvature lensing amounting to a mismatch of directions given by an angle θ of order $\rho p_0 \xi$, with ρ being the noncommutativity scale, p_0 being the energy difference between the two particles, and ξ being the boost between the (distant) observers at the source and at the detector. We expect that the quantification of dual-curvature lensing might be different in different spacetimes, so, in order to give guidance to the relevant phenomenology, it would be important to find other quantum spacetimes, in addition to ρ -Minkowski, to be used in the exploration of the realm of possibilities for dual-curvature lensing.





Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

- [1] G. Amelino-Camelia, J. Ellis, N. E. Mavromatos, D. V. Nanopoulos, and S. Sarkar, "Tests of quantum gravity from observations of big gamma-ray bursts," *Nature*, vol. 393, pp. 763–765, 1998.

- [2] R. Gambini and J. Pullin, “Nonstandard optics from quantum space-time,” *Physical Review D*, vol. 59, no. 12, Article ID 124021, 1999.
- [3] J. Alfaro, H. A. Morales-Técolt, and L. F. Urrutia, “Quantum Gravity Corrections to Neutrino Propagation,” *Physical Review Letters*, vol. 84, pp. 2318–2321, 2000.
- [4] U. Jacob and T. Piran, “Neutrinos from gamma-ray bursts as a tool to explore quantum-gravity-induced Lorentz violation,” *Nature Physics*, vol. 3, pp. 87–90, 2007.
- [5] G. Amelino-Camelia and L. Smolin, “Prospects for constraining quantum gravity dispersion with near term observations,” *Physical Review D*, vol. 80, Article ID 084017, 2009.
- [6] G. Amelino-Camelia, “Quantum-spacetime phenomenology,” *Living Reviews in Relativity*, vol. 16, no. 1, 5 pages, 2013.
- [7] D. Mattingly, “Modern tests of Lorentz invariance,” *Living Reviews in Relativity*, vol. 8, no. 1, 5 pages, 2005.
- [8] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman, and L. Smolin, “Principle of relative locality,” *Physical Review D*, vol. 84, no. 8, 2011.
- [9] J. M. Carmona, J. L. Cortés, D. Mazón, and F. Mercati, “Locality and the relativity principle beyond special relativity,” *Physical Review D*, vol. 84, Article ID 085010, 2011.
- [10] S. Mignemi and A. Samsarov, “Relative-locality effects in Snyder spacetime,” *Physics Letters A*, vol. 381, no. 19, pp. 1655–1658, 2017.
- [11] L. Freidel, R. G. Leigh, and D. Minic, “Modular spacetime,” *International Journal of Modern Physics D*, vol. 24, no. 12, Article ID 1544028, 2015.
- [12] G. Amelino-Camelia, L. Barcaroli, G. Gubitosi, and N. Loret, “Dual redshift on Planck-scale-curved momentum spaces,” *Classical and Quantum Gravity*, vol. 30, no. 23, 2013.
- [13] L. Freidel and L. Smolin, “Gamma ray burst delay times probe the geometry of momentum space,” 2011, <https://arxiv.org/abs/1103.5626>.
- [14] G. Amelino-Camelia, L. Barcaroli, and N. Loret, “Modeling transverse relative locality,” *International Journal of Theoretical Physics*, vol. 51, no. 11, pp. 3359–3375, 2012.
- [15] G. Amelino-Camelia, M. Arzano, S. Bianco, and R. J. Buoncore, “The DSR-deformed relativistic symmetries and the relative locality of 3D quantum gravity,” *Classical and Quantum Gravity*, vol. 30, no. 6, 12 pages, 2013.
- [16] IceCube Collaboration, “Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector,” *Science*, vol. 342, no. 6161, Article ID 1242856, 2013.
- [17] G. Amelino-Camelia, L. Barcaroli, DG. Amico, N. Loret, and G. Rosati, “IceCube and GRB neutrinos propagating in quantum spacetime,” *Physics Letters B*, vol. 761, pp. 318–325, 2016.
- [18] G. Amelino-Camelia, “Relativity in space-times with short-distance structure governed by an observer-independent (Planckian) length scale,” *International Journal of Modern Physics D*, vol. 11, 2002.
- [19] G. Amelino-Camelia, “Testable scenario for Relativity with minimum-length,” *Physics Letters B*, vol. 510, no. 1–4, Article ID 0012238, pp. 255–263, 2001.
- [20] J. Kowalski-Glikman and S. Nowak, “Non-commutative space-time of doubly special relativity theories,” *International Journal of Modern Physics D*, vol. 12, no. 2, pp. 299–315, 2003.
- [21] J. Magueijo and L. Smolin, “Generalized Lorentz invariance with an invariant energy scale,” *Physical Review D*, vol. 67, no. 4, Article ID 044017, 2003.
- [22] S. Majid and H. Ruegg, “Bicrossproduct structure of κ -Poincare group and non-commutative geometry,” *Physics Letters B*, vol. 334, no. 3–4, pp. 348–354, 1994.
- [23] J. Lukierski, H. Ruegg, and W. J. Zakrzewski, “Classical and Quantum Mechanics of Free κ -Relativistic Systems,” *Annals of Physics*, vol. 243, no. 1, pp. 90–116, 1995.
- [24] J. Kowalski-Glikman, “De Sitter space as an arena for doubly special relativity,” *Physics Letters B*, vol. 547, no. 3–4, Article ID 0207279, p. 291, 2002.
- [25] G. Amelino-Camelia, A. Marciano, M. Matassa, and G. Rosati, “Deformed Lorentz symmetry and relative locality in a curved/expanding spacetime,” *Physical Review D*, vol. 86, Article ID 124035, 2012.
- [26] G. Amelino-Camelia, V. Astuti, and G. Rosati, “Relative locality in a quantum spacetime and the pregeometry of κ -Minkowski,” *The European Physical Journal C*, vol. 73, no. 8, p. 2521, 2013.



Hindawi

Submit your manuscripts at
<https://www.hindawi.com>

