

A NEW CLASS OF COMPOSITION OPERATORS

S. PATTANAYAK, C.K. MOHAPATRA and A.K. MISHRA

School of Mathematical Sciences
Sambalpur University
Jyoti Vihar 768019, INDIA

(Received May 30, 1984 and in revised form June 14, 1985)

ABSTRACT. A new class of composition operators $P_\phi: H^2(T) \rightarrow H^2(T)$, with $\phi: T \rightarrow \bar{D}$ is introduced. Sufficient conditions on ϕ for P_ϕ to be bounded and Hilbert-Schmidt are obtained. Properties of P_ϕ with $\phi(e^{it}) = ae^{it} + be^{-it}$ for different values of the parameters a and b have been investigated. This paper concludes with a discussion on the compactness of P_ϕ .

KEY WORDS AND PHRASES. H^p Space, Composition operator, Hilbert Schmidt operator, Compact operator.

1980 AMS SUBJECT CLASSIFICATION CODES. 47B37, 47B99, 46J15.

1. PRELIMINARIES.

For a complex valued function f analytic in $D = \{z: |z| < 1\}$ and for $1 \leq p < \infty$ set

$$M_p(r, f) = \left(\int_0^{2\pi} |f(re^{i\theta})|^p \frac{d\theta}{2\pi} \right)^{\frac{1}{p}}, \quad 1 \leq p < \infty$$

and

$$M(r, f) = \sup_{0 < \theta < 2\pi} |f(re^{i\theta})|.$$

The function f is said to be in $H^p(D)$, if $\lim_{r \rightarrow 1^-} M_p(r, f) < \infty$. Similarly, let

$H^p(T)$, $T = \{z: |z| = 1\}$, be the class of functions in $L^p(T)$ such that

$$\int_0^{2\pi} f(e^{i\theta}) e^{in\theta} d\theta = 0, \quad n = 1, 2, 3, \dots$$

It is known [1,2] that for f in $H^p(D)$ $\lim_{r \rightarrow 1^-} f(re^{i\theta}) = f_*(e^{i\theta})$ exists for almost all

θ and f_* belongs to $H^p(T)$. Conversely, the Poisson integral of a function in $H^p(T)$ is in $H^p(D)$. Also, if f in $H^p(D)$ has the sequence $\{a_n\}$ as its Taylor coefficients then, f_* has the same sequence as its Fourier coefficients and vice versa. This correspondence establishes an isometrical isomorphism between $H^p(D)$ and $H^p(T)$. Thus, these

two spaces are interchangeably used and are usually referred to as the Hardy Space $H^p[1,2]$.

In the sequel we came across another space familiarly known as the weighted Hardy space [3]. Let $\rho(n)$ be a sequence of positive numbers. An analytic function $f: D \rightarrow \mathbb{C}$, given by $f(z) = \sum a_n z^n$, is said to be in the class $H^2(\rho)$, if $\|f\|_\rho^2 = \sum |a_n|^2 \rho(n) < \infty$. Also we need the following definition. Let H be a Hilbert space and T be a bounded linear operator on H . Then, T is said to be Hilbert Schmidt if there exists an orthonormal basis $\{e_n\}$ in H such that $\sum \|Te_n\|^2 < \infty$.

Throughout in the present paper we denote by e_n , $n = 0, 1, 2, \dots$, the function $e_n(e^{it}) = e^{int}$. We note that $\{e_n\}$ forms an orthonormal basis for H^2 .

2. A NEW CLASS OF COMPOSITION OPERATORS.

Let $\phi: D \rightarrow D$ be analytic and let $C_\phi: H^p(D) \rightarrow H^p(D)$ be defined by $(C_\phi f)(z) = f(\phi(z))$, z in D . The operator C_ϕ is known as a composition operator on $H^p(D)$ and is extensively studied in the literature [4]. In the present paper we introduce and study a new class of composition operators P_ϕ on $H^2(T)$ where $\phi: T \rightarrow \bar{D}$ may be 'non-analytic' also. That is ϕ nonvanishing negative Fourier coefficients.

DEFINITION. Let $\phi: T \rightarrow \bar{D}$ satisfy the following properties:

- (a) for every set $E \subset T$, of linear measure zero, $\phi^{-1}(E) = \{z \in T: \phi(z) = w, w \in E\}$ is also a set of linear measure zero and
- (b) for every f in $H^2(T)$, $f \circ \phi$ is in $L^2(T)$.

Then, define $P_\phi: H^2(T) \rightarrow H^2(T)$ by $P_\phi f = P(f \circ \phi)$ where P is the projection of $L^2(T)$ into $H^2(T)$.

Here some explanations are in order. We observe that a function f in $H^2(T)$ can be extended analytically into D as described in Section 1. So with the condition (a), $f \circ \phi$ is defined almost everywhere on T . Further, let f be represented by the Fourier series $\sum_{n=0}^{\infty} a_n e^{in\theta}$. Then by the Weierstrass theorem, $\sum_{n=0}^{\infty} a_n (\phi(e^{i\theta}))^n$ converges pointwise to $f(\phi(e^{i\theta}))$ for all θ such that $\phi(e^{i\theta}) \in D$ and by a result of Carleson [5] $\sum_{n=0}^{\infty} a_n (\phi(e^{i\theta}))^n$ converges pointwise to $f(\phi(e^{i\theta}))$ for almost all θ such that $\phi(e^{i\theta}) \in T$. Hence $\sum_{n=0}^{\infty} a_n (\phi(e^{i\theta}))^n$ converges pointwise almost everywhere on T to $f(\phi(e^{i\theta}))$. Thus throughout in this paper we write $\sum_{n=0}^{\infty} a_n (\phi(e^{i\theta}))^n$ in place of $f(\phi(e^{i\theta}))$.

We note that if ϕ satisfies the conditions of the definition, then by the Closed Graph Theorem, P_ϕ is a bounded operator. So a natural question is: under what conditions on ϕ , $f \circ \phi$ is in $L^2(T)$ for all f in $H^2(T)$. The present paper primarily deals with this question.

In the following sections we first obtain bounds for the norm of P_ϕ under suitable conditions on ϕ . Then we consider ϕ defined by $\phi(e^{it}) = ae^{it} + be^{-it}$ and study

conditions on a and b such that $f \circ \phi \in L^2(T)$ for all f in $H^2(T)$. In the last section we have discussed the compactness of P_ϕ with the help of some examples.

3. NORM OF P_ϕ .

We have the following results.

THEOREM 1. Let $\phi: T \rightarrow \bar{D}$ be such that

$$\int_0^{2\pi} \frac{dt}{1 - |\phi(e^{it})|^2} = M(\phi) < \infty \tag{3.1}$$

Then, P_ϕ is Hilbert Schmidt and $\|P_\phi\| \leq (M(\phi)/2\pi)^{1/2}$.

PROOF. Let f in $H^2(T)$ be given by $f(z) = \sum a_n z^n$, z in T . Then,

$$\begin{aligned} |f(\phi(e^{i\theta}))|^2 &= \left| \sum a_n (\phi(e^{i\theta}))^n \right|^2 \leq (\sum |a_n|^2) (\sum |\phi(e^{i\theta})|^{2n}) \\ &= \|f\|^2 \frac{1}{1 - |\phi(e^{i\theta})|^2} \quad \text{a.e.} \end{aligned}$$

So,

$$\|P_\phi f\|^2 \leq \|f \circ \phi\|_2^2 \leq \|f\|_2^2 \frac{M(\phi)}{2\pi}$$

and we get $\|P_\phi\| \leq (M(\phi)/2\pi)^{1/2}$.

Next, with the orthonormal basis e_n , $n = 0, 1, 2, \dots$, of $H^2(T)$, we have

$$\sum_{r=0}^{\infty} \|P_\phi(e_r)\|^2 \leq \sum_{n=0}^{\infty} \|e_n \circ \phi\|^2 = \sum_{n=0}^{\infty} \frac{1}{2\pi} \int_0^{2\pi} |\phi(e^{it})|^{2n} dt = (M(\phi)/2\pi) < \infty.$$

Thus, P_ϕ is Hilbert Schmidt.

COROLLARY 2. If $\phi: T \rightarrow D$ is continuous then P_ϕ is Hilbert Schmidt.

PROOF. The condition (3.1) is trivially satisfied if ϕ is continuous.

By an example in the next section we will show that (3.1) is only a sufficient condition for P_ϕ to be Hilbert Schmidt. We need the following lemma due to Gabriel [6] for the proof of our next theorem.

LEMMA. Let Γ be a rectifiable convex curve in the closed unit disc. Then, for every f in H^2

$$\int_{\Gamma} |f(w)|^2 |dw| \leq 4\pi \|f\|_2^2$$

THEOREM 2. Let $\phi: T \rightarrow \bar{D}$ be such that

(i) ϕ describes a closed rectifiable convex curve in \bar{D} and

(ii) $m = \inf |\phi'(e^{it})| > 0$, $0 \leq t \leq 2\pi$,

Then, $\|P_\phi\| \leq (2/m)^{1/2}$.

PROOF. By lemma and the condition (ii) we have

$$4\pi \|f\|_2^2 \geq \int_0^{2\pi} |f(\phi(e^{it}))|^2 |\phi'(t)| dt \geq m \int_0^{2\pi} |(f \circ \phi)(e^{it})|^2 dt$$

so that

$$\|P_\phi f\|^2 \leq \frac{1}{2\pi} \int_0^{2\pi} |(f \circ \phi)(e^{it})|^2 dt \leq \frac{2}{m} \|f\|_2^2$$

The conditions (i) and (ii) in the above theorem are not necessary for P_ϕ to be bounded. As an example consider

$$\phi(t) = (e^{it}) = \begin{cases} e^{it} & 0 < t < \pi \\ 0 & \pi \leq t \leq 2\pi \end{cases}$$

so that ϕ does not satisfy any of the conditions (i) or (ii) of the theorem.

Now,

$$\frac{1}{2\pi} \int_0^{2\pi} |f(\phi)(e^{it})|^2 dt = \frac{1}{2\pi} \int_0^\pi |f(e^{it})|^2 dt + \frac{1}{2\pi} \int_\pi^{2\pi} |f(0)|^2 dt \leq \|f\|_2^2.$$

This shows that P_ϕ is bounded with $\|P_\phi\| \leq \sqrt{2}$.

4. A FAMILY OF COMPOSITION OPERATORS.

In this section we study the properties of P_ϕ for the particular family of functions $\phi : T \rightarrow \bar{D}$ given by

$$\phi(z) = az + b\bar{z}, \quad z \in T \tag{4.1}$$

where $|a| + |b| \leq 1$. We note that if $|a| \neq |b|$ then the curve traced by ϕ is an ellipse containing the origin in its interior. Also $m = \inf |ae^{-it} - b| \geq ||a| - |b|| > 0$. Hence by Theorem 2, P_ϕ is bounded. It turns out that P_ϕ has many interesting properties for different values of the parameters a and b . We need the following technical lemma.

LEMMA 1. For all n, k in Z_+

$$\binom{n+2k}{k} < 2^{n+2k} \tag{4.2}$$

PROOF. We shall prove (4.2) by method of induction on n . Let $n = 0$ so that we have to show

$$\binom{2k}{k} < 2^{2k} \quad \text{for } k = 1, 2, 3, \dots \tag{4.3}$$

We establish (4.3), also by the process of induction on k . For $k = 1$, $\binom{2}{1} = 2 < 2^2$ is trivially true. Next, assume that

$$\binom{2k}{k} < 2^{2k}, \quad \text{i.e.} \quad \frac{(2k)!}{(k!) (k!)} < 2^{2k}$$

To complete induction on k we consider

$$\binom{2(k+1)}{k+1} = \frac{(2k+2)!}{(k+1)! (k+1)!} < 2^{2k} \cdot 2 \frac{(2k+1)}{(k+1)} < 2^{2(k+1)}.$$

Thus, (4.3) is true for all $k = 1, 2, 3, \dots$. Next, let $n = 1$. Then,

$$\binom{2k+1}{k} = \binom{2k}{k} \frac{(2k+1)}{(k+1)} < 2^k \cdot 2 = 2^{k+1}.$$

Now, assume that $\binom{n+2k}{k} < 2^{n+2k}$. To complete the induction we consider

$$\binom{n+1+2k}{k} = \frac{(n+2k)!}{(k!) (n+k)!} \frac{(n+1+2k)}{(n+1+k)} < 2^{n+2k} \cdot 2 < 2^{n+1+2k}.$$

Thus (4.3) is true for all $k = 1, 2, \dots$ and $n = 0, 1, 2, \dots$.

THEOREM 3. Let $\phi: T \rightarrow \bar{D}$ be given by $\phi(z) = az + b\bar{z}$, z in T .

(i) If $|a| + |b| \leq 1$, $|a| \neq |b|$, $b \neq 0$ and $|a| < \frac{1}{2}$ then P_ϕ is Hilbert Schmidt.

(ii) If $|a| + |b| = \frac{1}{2}$ then $f \circ \phi$ need not be in L^2 for all f in H^2 so that P_ϕ is not defined on the whole of H^2 .

(iii) The inequality $|a| < \frac{1}{2}$ in (i) is best possible.

PROOF. Consider the orthonormal basis e_n , $n = 0, 1, 2, \dots$, for H^2 . With respect to this basis P_ϕ has a matrix representation

$$t_{mn} = \begin{cases} 0 & \text{if } n < m \\ \binom{n+2k}{k} a^n (ab)^k & \text{if } n - m = 2k \\ 0 & \text{if } n - m = 2k + 1 \end{cases}$$

where m, n , and $k \in \mathbb{Z}_+$. Now

$$\begin{aligned} \sum |t_{m,n}|^2 &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \binom{n+2k}{k}^2 |a|^{2n} |ab|^{2k} \\ &= \sum_{n=0}^{\infty} (|a_n|^{2n} + \sum_{k=1}^{\infty} \binom{n+2k}{k}^2 |a|^{2n} |ab|^{2k}) \\ &= \frac{1}{1-|a|^2} + \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \binom{n+2k}{k}^2 |a|^{2n} |ab|^{2k} \end{aligned}$$

We use Lemma 1 to show that the second sum in the right hand side is convergent.

$$\begin{aligned} \sum_{k=1}^{\infty} \sum_{n=0}^{\infty} \binom{n+2k}{k}^2 |a|^{2n} |ab|^{2k} &\leq \sum \sum 2^{2n+4k} |a|^{2n} |ab|^{2n} = \sum \sum |2a|^{2n} |4ab|^{2k} \\ &= \frac{1}{1-|2a|^2} \cdot \frac{1}{1-|4ab|^2} < \infty, \end{aligned}$$

since $|a| < \frac{1}{2}$, which also implies $|ab| < \frac{1}{4}$. This proves that P_ϕ is Hilbert

Schmidt. (ii) For the proof of (ii) consider the H^2 function $f(z) = (1-z)^{-\alpha}$, $0 < \alpha < \frac{1}{2}$.

For $a = b = \frac{1}{2}$, $\phi(e^{i\theta}) = \cos \theta$. Thus,

$$f(\phi(e^{i\theta})) = f(\cos \theta) = \frac{1}{(1-\cos \theta)^\alpha} = \frac{1}{(2^\alpha \sin^2 \frac{\theta}{2})^\alpha} \quad \text{a.e.}$$

and

$$\int_0^{2\pi} |(f(\phi(e^{i\theta})))|^2 d\theta = \int_0^{\pi/2} \frac{2^{2-2\alpha}}{\sin^{4\alpha} \theta} d\theta \geq \int_0^{\pi/2} \frac{2^{2-2\alpha}}{\theta^{4\alpha}} d\theta = \infty$$

if $\alpha > \frac{1}{4}$. In the above we have made use of the well known inequality $\frac{2\theta}{\pi} < \sin \theta < \theta$ for $0 < \theta < \frac{\pi}{2}$.

(iii) In view of (ii), for the proof of (iii), it is sufficient to show that P_ϕ is not Hilbert Schmidt if $a + b = 1$ and $a > b$. In fact, we show that under the above

condition $\sum_m |t_{m,m}|^2 = \infty$.

Observe that

$$e_n(\phi(e^{i\theta})) = (ae^{i\theta} + be^{-i\theta})^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k e^{i(n-k)\theta} e^{-ik\theta} .$$

Since $(a + b)^n = 1$, we have $\sum_{k=0}^n \binom{n}{k} a^{n-k} b^k = 1$. Hence considering this sum as the inner product of two vectors

$$[\binom{n}{0} a^n b^0 , \binom{n}{1} a^{n-1} b^1 , \dots , \binom{n}{n} a^0 b^n \text{ | and | } 1, 1 \dots\dots\dots, 1]$$

We see that, since $||[1, 1, \dots, 1]||^2 = (n+1)$, by Cauchy Schwarz inequality

$$\sum_{k=0}^n | \binom{n}{k} a^{n-k} b^k |^2 \geq \frac{1}{(n+1)} .$$

Further, we observe that if $a > b$ then $\binom{n}{r} a^{n-r} b^r > \binom{n}{n-r} a^r b^{n-r}$ so that over half of the above sum is from terms where $n-k \geq k$ and so $||P_\phi e_n||^2 \geq 1/2(n+1)$, leading us to $\sum_n |t_{n,n}|^2 = \infty$. This completes the proof of the theorem.

Also, with the help of the same function $\phi(z) = az + b\bar{z}$, we show that the condition (3.1) of Theorem 1 is not a necessary condition for P_ϕ to be a Hilbert Schmidt.

For this take a, b in \mathbb{R} , $ab > 0$ and $|a| + |b| = 1$. Then,

$$\int_0^{2\pi} \frac{dt}{1 - |\phi(e^{it})|^2} = \int_0^{2\pi} \frac{dt}{(1 - (a-b)^2) \sin^2 t} = \infty .$$

However, in view of Theorem 3, it follows that P_ϕ is Hilbert Schmidt.

In the following theorem we present a sufficient condition on f in H^2 to ensure that $fo\phi$ is in $L^2(T)$.

THEOREM 4. Let $\phi: T \rightarrow \bar{D}$ be given by $\phi(z) = az + b\bar{z}$. $|a| = |b| = \frac{1}{2}$ and f in H^2 be given by $f(z) = \sum_{n=0}^\infty a_n z^n$. Further if $a_n = 0$ ($\frac{1}{n^\alpha}$) with $\alpha > \frac{3}{4}$, then $fo\phi \in L^2$.

PROOF. First let $a = b = \frac{1}{2}$, so that $\phi(e^{i\theta}) = \cos \theta$. Now,

$$|f(\phi(e^{i\theta}))|^2 \leq | \sum_{n=0}^\infty \frac{\cos^n \theta}{n^\alpha} |^2 \tag{4.4}$$

We know that

$$\frac{1}{(1-z)^{\beta+1}} = \sum_{n=0}^\infty \binom{n+\beta}{n} z^n \tag{4.5}$$

and [7]

$$\binom{n+\beta}{n} \sim \frac{n^\beta}{\Gamma(\beta+1)} . \tag{4.6}$$

Taking $\beta = -\alpha$ in (4.5) and (4.6), we get

$$\sum_{n=0} \frac{\cos^n \theta}{n^\alpha} \sim \frac{\Gamma(-\alpha+1)}{(1-\cos\theta)^{(1-\alpha)}} .$$

Thus, to complete the proof, it is sufficient to show that

$$\int_0^{2\pi} (1-\cos\theta)^{(2\alpha-2)} d\theta = \int_0^{2\pi} \sin^{(4\alpha-4)} \frac{\theta}{2} d\theta < \infty .$$

However, this is true because of the condition $\alpha > \frac{3}{4}$. To dispose of the general case

we observe that if $2a = e^{i\alpha}$ and $2b = e^{i\gamma}$ then $\phi(e^{i\theta}) = e^{i(\alpha+\gamma)/2} \cos(\frac{\alpha-\gamma+2\theta}{2})$ and this leads to similar calculations as above.

Taking cue from the above theorem we next show that $f \circ \phi \in L^2$ for $f \in H^2(\rho(n))$ for a suitable choice of the sequence $\rho(n)$.

THEOREM 5. Let $\phi: T \rightarrow \bar{D}$ be given by $\phi(e^{i\theta}) = ae^{i\theta} + be^{-i\theta}$, $|a| = |b| = \frac{1}{2}$ and $\rho(n) = n^\beta$. Then,

(i) $f \circ \phi$ is in L^2 for all f in $H^2(\rho(n))$ if $\beta > \frac{1}{2}$,

(ii) for each $\beta < \frac{1}{2}$ there is a function f_β in $H^2(\rho(n))$ such that $f_\beta \circ \phi$ is not in L^2

PROOF. As in the previous theorem we assume $a = b = \frac{1}{2}$ so that $f(\phi(e^{i\theta})) = f(\cos \theta)$.

Let f in $H^2(\rho)$ be given by $f(z) = \sum_{n=0}^{\infty} a_n z^n$. We have

$$|f(\phi(e^{i\theta}))|^2 = \left| \sum_{n=0}^{\infty} a_n \cos^n \theta \right|^2 \leq \left(\sum_{n=0}^{\infty} |a_n|^2 \rho(n) \right) \left(\sum_{n=0}^{\infty} \frac{\cos^{2n} \theta}{\rho(n)} \right).$$

Now using (4.5) and (4.6) as in the previous theorem it can be shown that

$$\sin^{2\beta-2} \theta \sim \sum_{n=0}^{\infty} \frac{n^{-\beta}}{\Gamma(-\beta+1)} \cos^{2n} \theta.$$

Thus $f \circ \phi$ is in L^2 if $\beta > \frac{1}{2}$.

For the proof of (ii) consider the function

$$f(z) = \frac{1}{(1-z)^{\alpha+1}} = \sum A_n^\alpha z^n.$$

By (4.6),

$$\sum |A_n^\alpha|^2 n^\beta \sim \sum n^{2\alpha+\beta}.$$

The sum on the right hand converges if $2\alpha + \beta < -1$ i.e. $\alpha < -(\beta+1)/2$. Thus, f is in $H^2(\rho)$, $\rho(n) = n^\beta$ for $\alpha < -(\beta+1)/2$. However,

$$\int_0^{2\pi} |f(\phi(e^{i\theta}))|^2 d\theta = \frac{1}{2^{2\alpha+1}} \int_0^\pi \frac{1}{\sin^{4\alpha+4} \theta} d\theta = \infty$$

if $\alpha \geq -\frac{3}{4}$. Thus, for given $\beta < \frac{1}{2}$, if we chose $\alpha = -(3+2\beta+2)/8$, f is in $H^2(\rho)$ but $f \circ \phi$ is not in L^2 .

REMARK. The case $\rho(n) = n^{\frac{1}{2}}$, remains open in the above theorem. However, in the next theorem we prove the same result for a sequence $\rho(n)$ having faster rate of growth than $n^{1/2}$ but with slower rate than $n^{1/2+\epsilon}$ for any $\epsilon > 0$.

THEOREM 6. Let $\phi: T \rightarrow \bar{D}$ be as in the previous theorem and $\rho(n) = n^{1/2}(\log n)^\beta$. Then,

(i) $f \circ \phi$ is in L^2 for all f in $H^2(\rho(n))$ if $\beta > 1$,

(ii) for each $\beta < 0$, there is a function f_β in $H^2(\rho(n))$ such that $f_\beta \circ \phi$ is not in L^2 .

PROOF (i) Let f , given by $f(z) = \sum_{n=0}^{\infty} a_n z^n$, be in $H^2(\rho(n))$ so that

$$\|f\|_{\rho} = \sum n^{1/2} (\log n)^{\beta} |a_n|^2 < \infty .$$

By Cauchy Schwarz inequality,

$$|f(\phi(e^{i\theta}))|^2 \leq \sum_{n=0}^{\infty} n^{1/2} (\log n)^{\beta} |a_n|^2 \left(\sum_{n=0}^{\infty} \frac{\cos^{2n} \theta}{n^{1/2} (\log n)^{\beta}} \right) . \tag{4.7}$$

It is known [7, p. 192] that if

$$\frac{1}{(1-z)^{\alpha+1}} \left(\log \frac{a}{1-z} \right)^{\beta} = \sum_{n=0}^{\infty} A_n^{(\alpha, \beta)} z^n \tag{4.8}$$

then for $a > 2$, $\alpha \neq -1, -2, -3, \dots$, $\alpha, \beta \in \mathbb{R}$

$$A_n^{(\alpha, \beta)} \sim \frac{n^{\alpha}}{\Gamma(\alpha+1)} (\log n)^{\beta}$$

so that

$$A_n^{(-1/2, -\beta)} \sim \frac{1}{\sqrt{\pi} n^{1/2} (\log n)^{\beta}}$$

i.e.

$$\frac{1}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(\cos^2 \theta)^n}{n^{1/2} (\log n)^{\beta}} \sim \frac{1}{(1-\cos^2 \theta)^{1/2}} \left(\log \frac{a}{1-\cos^2 \theta} \right)^{-\beta} = \frac{2^{-\beta}}{\sin \theta} \left(\log \frac{b}{\sin \theta} \right)^{-\beta} ,$$

$$b = \sqrt{a}$$

$$\tag{4.9}$$

In view of (4.7) and (4.9), in order to show that $f \circ \phi$ is in L^2 , it is sufficient to show that

$$\int_0^{2\pi} \frac{1}{\sin \theta} \left(\log \frac{1}{\sin \theta} \right)^{-\beta} d\theta < \infty .$$

Further, because of the inequality $(2\theta/\pi) < \sin \theta < \theta$, it is sufficient to show the integrability, in an interval $(0, \delta)$, of the function

$$h(\theta) = \frac{1}{\theta} \left(\log \frac{1}{\theta} \right)^{-\beta}$$

Making the substitution $\log \left(\frac{1}{\theta} \right) = u$, we get

$$\lim_{\epsilon \rightarrow 0} \int_{\epsilon}^{\delta} \frac{1}{\theta} \left(\log \frac{1}{\theta} \right)^{-\beta} d\theta = \lim_{\epsilon \rightarrow 0} \frac{(\log \frac{1}{\epsilon})^{1-\beta} - (\log \frac{1}{\delta})^{1-\beta}}{1-\beta}$$

Thus, the above integral converges if $\beta > 1$. This completes the proof of (i).

(ii) Let $\rho(n) = n^{1/2} (\log n)^{-\beta}$, $\beta > 0$. Now consider the function

$$f(z) = \frac{1}{(1-z)^{1/4}} \left(\log \frac{a}{1-z} \right)^{\gamma} = \sum_{n=0}^{\infty} A_n^{\gamma} z^n$$

where $-1/2 > \gamma < (\beta-1)/2$. We first observe that f is in $H^2(\rho(n))$. In fact, a comparison of f with (4.8) shows that

$$A_n^{\gamma} \sim \frac{n^{-3/4}}{\Gamma(1/4)} (\log n)^{\gamma} .$$

Thus,

$$\sum |A_n^{\gamma}|^2 \rho(n) \sim \sum n^{-1} (\log n)^{2\gamma-\beta}$$

and the right hand side series converges because $\gamma < (\beta-1)/2$. Now,

$$\int_0^{2\pi} |f(\phi(e^{i\theta}))|^2 d\theta = 2^{\gamma-1/2} \int_0^{\pi/2} \frac{1}{\sin \theta} (\log \frac{b}{\sin \theta})^{2\gamma} d\theta$$

where $b = \frac{a}{2}$. The above integral diverges with the integral

$$\int_0^{\pi/2} \frac{1}{\theta} (\log \frac{1}{\theta})^{2\gamma} d\theta$$

because of the condition $\gamma > \frac{1}{2}$. Thus, we prove that although $f \in H^2(\rho(n))$, $f \circ \phi$ is not in L^2 .

REMARK. The case $\rho(n) = n^{\frac{1}{2}}(\log n)^\beta$, $0 \leq \beta < 1$ remains unsettled.

We conclude this section by showing that P_ϕ is an unbounded operator on H^2 for $\phi(e^{it}) = (e^{it} + e^{-it})/2 = \cos t$. This we do by exhibiting a sequence of function f_n in H^2 for which $\lim_{n \rightarrow \infty} \|P_\phi f_n\| = \infty$

Let $f_n(z) = \sum_{k=1}^n \frac{z^k}{k}$ and $f(z) = \sum_{k=1}^{\infty} \frac{z^k}{k} = \log \frac{1}{1-z}$ so that

$$g_n(t) = f_n(\phi(e^{it})) = \sum_{k=1}^n \frac{\cos^k t}{k} \quad \text{and} \quad g(t) = f(\phi(e^{it})) = \sum_{k=1}^{\infty} \frac{\cos^k t}{k}.$$

Observe that g is in L^1 but is not in L^2 . Let a_k and $a_n^{(n)}$ respectively be the k^{th} Fourier coefficients of g and g_n . Since

$$\lim_{n \rightarrow \infty} \int_0^{2\pi} |g_n(t) - g(t)| dt = 0$$

we get

$$\lim_{n \rightarrow \infty} a_n^{(n)} = a_k. \quad \text{Now,}$$

$$\lim_{n \rightarrow \infty} \|P_\phi\|_2^2 = \lim_{n \rightarrow \infty} \sum_{k=-n}^n |a_k^{(n)}|^2 = \sum_{k=-\infty}^{\infty} |a_k|^2 = \frac{1}{2\pi} \int |g(e^{it})|^2 dt = \infty.$$

5. COMPACTNESS OF P_ϕ

In this section we discuss some examples illustrating cases when P_ϕ is compact and when it is not. Let $\phi_1, \phi_2, \phi_3 : T \rightarrow \bar{D}$, be defined by

(i) $\phi_1(e^{it}) = ae^{-it}, \quad |a| = 1 \quad 0 \leq t \leq 2\pi$

(ii) $\phi_2(e^{it}) = \begin{cases} e^{it} & , & 0 \leq t < \pi \\ 0 & , & \pi \leq t \leq 2\pi \end{cases}$

(iii) $\phi_3(e^{it}) = \begin{cases} e^{it} & , & 0 \leq t \leq \pi \\ ae^{it} + be^{-it} & , & \pi \leq t \leq 2\pi \end{cases} \quad a+b = 1 \quad a \neq b, \quad a, b \geq 0$

(i) P_{ϕ_1} is a finite rank, hence a compact, operator. For, if f in H^2 is given by

$$f(z) = \sum_{n=0}^{\infty} a_n z^n \quad \text{then} \quad (P_{\phi_1} f)(e^{it}) = P \left(\sum_{n=0}^{\infty} a_n a^n e^{-int} \right) = a_0.$$

For composition operators with analytic ϕ Schwartz [8] has shown that if $C_\phi: H^p(D) \rightarrow H^p(D)$ is compact then $|\phi(e^{it})| < 1$ a.e. where $\phi(e^{it})$ is the radial limit of $\phi(z)$. We observe in this example that P_{ϕ_1} deviates in behaviour from C_ϕ . (ii) We have shown at the end of Section 3 that $\|P_{\phi_2}\| < \sqrt{2}$. We show here that P_{ϕ_2} is not compact.

By Riemann-Lebesgue Lemma the sequence e_n , $n = 0, 1, 2, \dots$ converges to zero weakly in H^2 . However, $P_{\phi_2}(e_n) = P(e_n \circ \phi_2)$, does not converge strongly to zero. For, if the Fourier series of $e_n \circ \phi_2$ is given by $(e_n \circ \phi_2)(e^{it}) = \sum_{m=-\infty}^{\infty} a_m e^{imt}$, then by direct computation it can be seen that

$$a_m = \begin{cases} \frac{1}{\pi(n-m)} & \text{if } n-m \text{ is odd} \\ 0 & \text{if } n-m \text{ is even} \\ \frac{1}{2} & \text{if } n=m \end{cases}$$

and $\|P_{\phi_2}(e_n)\|_2^2 = \sum_{m=0}^{\infty} |a_m|^2 > \frac{1}{4}$. Thus, P_{ϕ_2} is not compact.

By a similar argument as in (ii) it can be shown that P_{ϕ_3} is bounded but not a compact operator.

ACKNOWLEDGEMENT. The authors thank the referee for his useful suggestions regarding the definition of P_ϕ .

REFERENCES

1. DUREN, P.L., H^p Spaces, Academic Press, New York, 1970.
2. HOFFMAN, K., Banach Spaces of Analytic Functions, Prentice-Hall, Englewood Cliffs, New Jersey, 1962.
3. HALMOS, P.R., A Hilbert Space Problem Book, Springer-Verlag, New York, Inc., Graduate texts in Mathematics, Vol. 19.
4. NORDGREN, E.A., Composition Operators in Hilbert Spaces, Lecture notes in Mathematics, 639, Springer-Verlag, Berlin, 1978.
5. CARLESON, L., On Convergence Growth of Partial Sums of Fourier Series. Acta Math, 116 (1966), 135-157.
6. GABRIEL, R.M., Some Results Concerning the Moduli of Regular Functions Along Curves of Certain Types, Proc. London Math. Soc. 28(1928), 121-127.
7. ZYGMOND, A., Trigonometric Series, 2nd Ed., Cambridge University Press, 1959.
8. SCHWARTZ, H., Composition Operators on H^p , Dissertation, University of Toledo, 1969.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

