

Research Article

Physiological and Biochemical Responses of *Ulva prolifera* and *Ulva linza* to Cadmium Stress

He-ping Jiang,^{1,2} Bing-bing Gao,¹ Wen-hui Li,¹ Ming Zhu,^{1,3} Chun-fang Zheng,²
Qing-song Zheng,^{1,2} and Chang-hai Wang¹

¹ College of Natural Resources and Environmental Science, Key Laboratory of Marine Biology, Nanjing Agricultural University, Nanjing, Jiangsu 210095, China

² Zhejiang Mariculture Research Institute, Zhejiang Key Laboratory of Exploitation and Preservation of Coastal Bio-Resource, Wenzhou, Zhejiang 325005, China

³ College of Oceanography, Huaihai Institute of Technology, Lianyungang, Jiangsu 222005, China

Correspondence should be addressed to Qing-song Zheng; qszheng@njau.edu.cn

Received 2 January 2013; Accepted 3 February 2013

Academic Editors: J. Huang and Z. Wang

Copyright © 2013 He-ping Jiang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Responses of *Ulva prolifera* and *Ulva linza* to Cd²⁺ stress were studied. We found that the relative growth rate (RGR), Fv/Fm, and actual photochemical efficiency of PSII (Yield) of two *Ulva* species were decreased under Cd²⁺ treatments, and these reductions were greater in *U. prolifera* than in *U. linza*. *U. prolifera* accumulated more cadmium than *U. linza* under Cd²⁺ stress. While *U. linza* showed positive osmotic adjustment ability (OAA) at a wider Cd²⁺ range than *U. prolifera*. *U. linza* had greater contents of N, P, Na⁺, K⁺, and amino acids than *U. prolifera*. A range of parameters (concentrations of cadmium, Ca²⁺, N, P, K⁺, Cl⁻, free amino acids (FAAs), proline, organic acids and soluble protein, Fv/Fm, Yield, OAA, and K⁺/Na⁺) could be used to evaluate cadmium resistance in *Ulva* by correlation analysis. In accordance with the order of the absolute values of correlation coefficient, contents of Cd²⁺ and K⁺, Yield, proline content, Fv/Fm, FAA content, and OAA value of *Ulva* were more highly related to their adaptation to Cd²⁺ than the other eight indices. Thus, *U. linza* has a better adaptation to Cd²⁺ than *U. prolifera*, which was due mainly to higher nutrient content and stronger OAA and photosynthesis in *U. linza*.

1. Introduction

Heavy metal contamination is an environmental problem in the margin sea [1]. As the economy in Asian countries continues to grow, the release of heavy metals and other contaminants has increased noticeably [2, 3]. Due to their acute toxicity, cadmium (Cd), lead, and mercury are among the most hazardous metals to the environment and living things [4].

Cd, an oxophilic and sulfophilic element, forms complexes with various organic particles and thereby triggers a wide range of reactions that collectively put the aquatic ecosystems at risk. Cadmium also poses a serious threat to human health due to its accumulation in the food chain [5, 6]. It has been classified as group (I) a human carcinogen by the International Agency for Research on Cancer (IARC) [7]. Cadmium toxicity may be characterized by a variety of

syndromes and effects, including renal dysfunction, hypertension, hepatic injury, lung damage, and teratogenic effects [8]. To remove Cd pollutants, various treatment technologies, such as precipitation, ion exchange, adsorption, and biosorption, have been employed [9]. Biosorption is one of the promising techniques for removal of heavy metals. Biosorption utilizes the ability of biological materials to accumulate heavy metals from waste streams by either metabolically mediated or purely physicochemical pathways of uptake [10]. Among the biological materials investigated for heavy metal removal, marine macroalgae have high uptake capacities for a number of heavy metal ions [11, 12].

Green algae species of Ulvaceae, especially the members of the green algal genus *Ulva*, have been considered as monitors of heavy metals in estuaries [13–15]. Numerous studies have shown that green macroalgae such as *Ulva lactuca* are able to absorb Cd. These studies mainly focused

on metabolism-independent Cd accumulation [6], synthetic surfactants exerting impact on uptake of Cd [12], effect of pH, contact time, biomass dosage and temperature on the Cd uptake kinetics [2], and induced oxidative stress by Cd [7]. However, little information is available regarding physiological responses of different *Ulva* species to increased Cd^{2+} concentrations.

In this study, *Ulva prolifera* and *Ulva linza* were studied for their responses to different Cd^{2+} concentrations. Their growth, chlorophyll fluorescence parameters, osmotic adjustment ability, and accumulation of inorganic ions and organic solutes were investigated in indoor seawater culture systems. The specific objective of this study was to determine if there was species variation in Cd^{2+} adaptation, and what were the major physiological parameters involved in the adaptation.

2. Materials and Methods

2.1. The Seaweed Collection, Cultivation, and Cd^{2+} Treatment. Green algae were collected from the sea in Dafeng (*Ulva prolifera*) and Lianyungang (*Ulva linza*), Jiangsu province, China. Upon arrival in the laboratory, the seaweeds were washed with distilled water and then cultured in 250 mL flasks containing 200 mL of sterilized artificial seawater (33.33 psu, pH 8.0) enriched with VSE medium [16] for 5 d. The composition of artificial seawater was (g L^{-1}) HCO_3^- 0.25, SO_4^{2-} 3.84, Cl^- 17.45, Ca^{2+} 0.76, Mg^{2+} 1.00, K^+ 0.57, and Na^+ 9.46. The composition of VSE nutrient solution was (mg L^{-1}) NaNO_3 42.50, $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ 10.75, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 0.28, $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ 0.02, $\text{Na}_2\text{EDTA} \cdot 2\text{H}_2\text{O}$ 3.72, vitamin B_1 0.20, Biotin 0.001, and vitamin B_{12} 0.001. After 5 d acclimation, healthy samples (0.5 g fresh weight) were cultured in 250 mL flasks with 200 mL medium as described earlier. CdCl_2 was added to each flask at the following concentrations: 0, 5, 10, 20, 40, 80, or 120 $\mu\text{mol L}^{-1}$. After 7 d treatment, *U. prolifera* and *U. linza* were harvested and analyzed for selected parameters as described later. All experiments were performed in three replicates. During the preculture and the treatment, seaweeds were grown in a GXZ intelligent light incubator at temperature of $20 \pm 1^\circ\text{C}$, light intensity of 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$, and photoperiod of 12/12 h. The culture medium was altered every other day.

2.2. Measurement of Relative Growth Rate (RGR). Fresh weight was determined by weighing the algae after blotting by absorbent paper. RGR was calculated according to the formula $\text{RGR} (\% \text{d}^{-1}) = [\ln(M_t/M_0)]/t \times 100\%$, where M_0 and M_t are the fresh weights (g) at days 0 and 7, respectively [17].

2.3. Measurement of Osmotic Adjustment Ability (OAA). Saturated osmotic potential was measured by the freezing-point depression principle. Seaweeds were placed in double-distilled water for 8 h and then rinsed 5 times with double-distilled water. After blotting dry with absorbent paper, seaweeds were dipped into liquid nitrogen for 20 min. The frozen seaweeds were thawed in a syringe for 50 min, and the

seaweed sap was then collected by pressing the seaweed in the syringe [18]. The π_{100} was measured by using a fully automatic freezing-point osmometer (8P, Shanghai, China). OAA was calculated by the following equation:

$$\Delta\pi_{100} = \pi_{100}^{\mu} - \pi_{100}^{\text{s}}, \quad (1)$$

whereby π_{100}^{μ} was the π_{100} of control seaweeds, and π_{100}^{s} was the π_{100} of Cd^{2+} -stressed seaweeds.

2.4. Measurements of Chlorophyll (Chl) and Carotenoid (Car) Contents. Determination of Chl and Car was carried out by the method of Häder et al. [19]. Weighed 0.1 g fresh seaweeds were cut with scissors and extracted with 95% (v/v) ethanol (10 mL) in the dark for 24 h. The absorbance of pigment extract was measured at wavelengths of 470, 649, and 665 nm with a spectrophotometer. From the measured absorbance, concentrations of Chl a, Chl b, and Car were calculated on a weight basis.

2.5. Determination of Chlorophyll Fluorescence Parameters. A PHYTO-PAM Phytoplankton Analyzer (PAM 2003, Walz, Effeltrich, Germany) was used to determine *in vivo* chlorophyll fluorescence from chlorophyll in photosystem II (PSII) using different experimental protocols [19]. Before determination, samples were adapted for 15 min in the total darkness to complete reoxidation of PSII electron acceptor molecules. The maximal photochemical efficiency of PSII (Fv/Fm) and the actual photochemical efficiency of PSII in the light (Yield) were then determined.

2.6. Measurement of Nitrogen (N) and Phosphorous (P) Concentrations. Dried samples were ground in a mortar and pestle. Total N in seaweed tissue was analyzed by an N gas analyzer using an induction furnace and thermal conductivity. Total P in seaweed tissue was quantitatively determined by Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-AES, Optima 2100 DV, PerkinElmer, USA) following nitric acid/hydrogen peroxide microwave digestion. The total amounts of N and P in the seaweed tissue were calculated by multiplying N and P contents in tissue as a proportion of dry weight by the total dry weight of the sample [20].

2.7. Measurement of Inorganic Elements. After 7 d, seaweeds were harvested, washed, and oven-dried at 65°C for 3 d. A 50 mg sample was ashed in a muffle furnace. The ash was dissolved in 8 mL of $\text{HNO}_3 : \text{HClO}_4$ (3:1, v:v) and diluted to 50 mL with distilled water. The contents of Cd, Na, K, Ca, and Mg were determined by Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-AES, Optima 2100 DV, PerkinElmer, USA) [21]. To determine Cl content, the ash was dissolved in 100 mL distilled water and analyzed by potentiometric titration with silver nitrate (AgNO_3) [18]. Total nitrate was measured as described previously [22] with nitrate extracted from the tissue by boiling fresh seaweeds (20 mg) in distilled water (400 μL) for 20 min. The nitrate concentrations in the samples were measured spectrophotometrically at 540 nm.

2.8. Measurement of Organic Solutes. Soluble sugars (SS) determination was carried out by the anthrone method [23]. Water extract of fresh seaweeds was added to 0.5 mL of 0.1 mol L⁻¹ anthrone-ethyl acetate and 5 mL H₂SO₄. The mixture was heated at 100°C for 1 min, and its absorbance at 620 nm was read after cooling to room temperature. A calibration curve with sucrose was used as a standard. Soluble proteins (SPs) were measured by Coomassie Brilliant Blue G-250 staining [24]. Fresh seaweeds (0.5 g) were homogenized in 1 mL phosphate buffer (pH 7.0). The crude homogenate was centrifuged at 5,000 g for 10 min. An aliquot of 0.5 mL of freshly prepared trichloroacetic acid (TCA) was added and mixture centrifuged at 8,000 g for 15 min. The pellets were dissolved in 1 mL of 0.1 mol L⁻¹ NaOH, and 5 mL of Bradford reagent was added. Absorbance was recorded at 595 nm using bovine serum albumin as a standard. Free amino acids (FAAs) were extracted and determined following the method of Zhou and Yu [23]. A total of 0.5 g fresh tissue was homogenized in 5 mL 10% (w/v) acetic acid, extracts were supplemented with 1 mL distilled water and 3 mL ninhydrin reagent, then boiled for 15 min and fast cooled, and the volume was made up to 5 mL with 60% (v/v) ethanol. Absorbance was read at 570 nm. The content of total free amino acids was calculated from a standard curve prepared using leucine. Proline (PRO) concentration was determined spectrophotometrically by adopting the ninhydrin method of Irigoyen et al. [25]. We first homogenized 300 mg fresh leaf samples in sulphosalicylic acid. To the extract, 2 mL each of ninhydrin and glacial acetic acid were added. The samples were heated at 100°C. The mixture was extracted with toluene, and the free toluene was quantified spectrophotometrically at 528 nm using L-proline as a standard. Organic acids (OAs) were extracted with boiling distilled water. The concentration of total OA was determined by 0.01 mmol L⁻¹ NaOH titration method, with phenolphthalein as indicator [26].

2.9. Statistical Analyses. All experiments were performed in three replicates. The data are presented as the mean ± SD. Data were analyzed using SPSS statistical software. Significant differences between means were determined by Duncan's multiple range test. Unless otherwise stated, differences were considered statistically significant when $P \leq 0.05$. Statistical analysis on two-way variance analysis (ANOVA), and correlation coefficient was performed using Microsoft Excel.

3. Results

3.1. Effect of Cadmium Stress on RGR and OAA of *U. prolifera* and *U. linza*. Compared to the control, treatments with 5 μmol L⁻¹ Cd²⁺ for 7 d did not change RGR of *U. linza*, but significantly decreased RGR of *U. prolifera*. The RGR of both *Ulva* species was significantly decreased as Cd²⁺ concentration increased. After 7 d exposure to 10, 20, 40, 80; or 120 μmol L⁻¹ Cd²⁺, RGR of *U. linza* decreased by 53, 75, 116, 177, and 277%, respectively; *U. prolifera* decreased by 93, 139, 271, and 357%, respectively. *U. prolifera* died at 120 μmol L⁻¹ Cd²⁺ on day 7 (Figure 1).




FIGURE 1: Effects of different concentrations of Cd²⁺ (0, 5, 10, 20, 40, 80, and 120 μmol L⁻¹) on relative growth rate (RGR) in *U. prolifera* and *U. linza*.

The OAA of both species was enhanced by low Cd²⁺ concentration treatments. The enhancement occurred at 5 and 10 μmol L⁻¹ for *U. prolifera* and 5, 10 and 20 μmol L⁻¹ for *U. linza* (Figure 2). However, OAA was negative when *U. prolifera* was treated by 20, 40, and 80 μmol L⁻¹ Cd²⁺, and *U. linza* treated by 40 and 80 μmol L⁻¹ Cd²⁺ (Figure 2).

3.2. Effect of Cadmium Stress on Cadmium Content in *U. prolifera* and *U. linza*. Cadmium contents in *U. prolifera* and *U. linza* increased as Cd²⁺ concentrations increased (Figure 3). At 5, 10, 20, 40, and 80 μmol L⁻¹ Cd²⁺, Cd contents in *U. prolifera* was 32, 78, 114, 140, and 165 times of the Cd²⁺ = 0 treatment, respectively, and 10, 26, 44, 65, and 79 times of its control treatment in *U. linza*, respectively.

3.3. Effect of Cadmium Stress on Chl and Car Contents in *U. prolifera* and *U. linza*. Both Chl and Car contents decreased with the increased Cd²⁺ concentration. There was no significant change in Chl and Car when both species were treated by 5 and 10 μmol L⁻¹ Cd²⁺ for 7 d. However, significant declines in Chl and Car contents were observed when they were exposed to 20, 40, or 80 μmol L⁻¹ Cd²⁺. Compared to the control treatment, Chl contents decreased by 18, 25, and 45% at 20, 40, and 80 μmol L⁻¹ Cd²⁺ in *U. prolifera*, respectively; and the decreases were 16, 20, and 39% in *U. linza*, respectively (Figure 4(a)). The Car content declined by 16, 29 and 54% at 20, 40 and 80 μmol L⁻¹ Cd²⁺ in *U. prolifera*, respectively; and by 13, 16, and 44% in *U. linza*, respectively (Figure 4(b)).




FIGURE 2: Effects of different concentrations of Cd²⁺ (5, 10, 20, 40, and 80 μmol L⁻¹) on osmotic adjustment ability (OAA) of *U. prolifera* and *U. linza*.




FIGURE 3: Effects of different concentrations of Cd²⁺ (0, 5, 10, 20, 40, and 80 μmol L⁻¹) on cadmium concentration of *U. prolifera* and *U. linza*.

3.4. Effect of Cadmium Stress on Chlorophyll Fluorescence Parameters of *U. prolifera* and *U. linza*. Compared to the control treatment, Fv/Fm of *U. prolifera* and *U. linza* were not significantly affected by the treatments of 5 or 10 μmol L⁻¹ Cd²⁺. However, Fv/Fm of both *Ulva* species fell significantly

when Cd²⁺ concentrations reached 20 μmol L⁻¹. In comparison with the control, Fv/Fm of *U. prolifera* decreased 17, 22, and 31% at 20, 40, and 80 μmol L⁻¹ Cd²⁺; whereas Fv/Fm of *U. linza* decreased 9, 10, and 15% after exposure to 20, 40, or 80 μmol L⁻¹ Cd²⁺, respectively (Figure 5(a)). For actual photochemical efficiency of PSII (Yield) of *U. prolifera*, there was an obvious decrease when Cd²⁺ concentrations rose from 20 to 80 μmol L⁻¹; whereas Yield of *U. linza* showed no significant decline until Cd²⁺ concentration was 80 μmol L⁻¹ (Figure 5(b)).

3.5. Effect of Cadmium Stress on Contents of N and P in *U. prolifera* and *U. linza*. Contents of N and P in both *Ulva* species showed a declining trend after an initial increase. The highest N content was recorded at 10 μmol L⁻¹ Cd²⁺ in *U. prolifera* and at 20 μmol L⁻¹ Cd²⁺ in *U. linza*. N contents in *U. linza* in all Cd²⁺ treatments were higher than those of control; however, in *U. prolifera*, N contents at 20, 40, or 80 μmol L⁻¹ Cd²⁺ were significantly decreased compared to the control (Figure 6(a)).

U. prolifera had the highest P concentration at 5 μmol L⁻¹ Cd²⁺; but the highest P concentration was observed when *U. linza* was treated by 10 μmol L⁻¹ Cd²⁺. The P contents decreased 31, 40, and 54% at 20, 40, and 80 μmol L⁻¹ Cd²⁺ in *U. prolifera*, respectively. Compared to the control, the P concentration of *U. linza* at 20 μmol L⁻¹ Cd²⁺ increased significantly, and then decreased by 11 and 27% under 40, and 80 μmol L⁻¹ Cd²⁺, respectively (Figure 6(b)).

3.6. Effect of Cadmium Stress on Inorganic Elements of *U. prolifera* and *U. linza*. The Na⁺ content of *U. prolifera* grown at 5 or 10 μmol L⁻¹ Cd²⁺ was not significantly different from the control, and it increased by 42, 67, and 83% at 20, 40, and 80 μmol L⁻¹ Cd²⁺, respectively. However, in *U. linza*, 5, 10, 20, and 40 μmol L⁻¹ Cd²⁺ had no significant influence on Na⁺ content, and 80 μmol L⁻¹ Cd²⁺ increased Na⁺ content by 36% (Table 1). The K⁺ content of *U. prolifera* grown at 5 or 10 μmol L⁻¹ Cd²⁺ remained unaffected compared to the control; it decreased significantly by 41, 45, and 62% at 20, 40, and 80 μmol L⁻¹ Cd²⁺, respectively. In *U. linza*, 5, 10, and 20 μmol L⁻¹ Cd²⁺ had no significant influence on K⁺ content, whereas 40 and 80 μmol L⁻¹ Cd²⁺ decreased K⁺ content by 34 and 50%, respectively (Table 1). The Ca²⁺ content of *U. prolifera* grown at 5, 10, 20, or 40 μmol L⁻¹ Cd²⁺ remained unaffected, but increased significantly (24%) at 80 μmol L⁻¹ Cd²⁺. However, in *U. linza*, 5 and 10 μmol L⁻¹ Cd²⁺ had no significant influence on Ca²⁺ contents, whereas 20, 40, and 80 μmol L⁻¹ Cd²⁺ increased Ca²⁺ content by 22, 39, and 50%, respectively (Table 1). The Mg²⁺ content of *U. prolifera* grown at 5, 10, 20, 40 or 80 μmol L⁻¹ Cd²⁺ remained unaffected. With increasing Cd²⁺ concentrations, Mg²⁺ contents of *U. linza* showed an increasing trend after an initial decline (Table 1). The Cl⁻ contents appeared to have a declining trend with increasing Cd²⁺ concentration similarly to Mg concentrations. However, no obvious difference in Cl⁻




FIGURE 4: Effects of different concentrations of Cd²⁺ (0, 5, 10, 20, 40, and 80 µmol L⁻¹) on chlorophyll content (a) and carotenoid content (b) in *U. prolifera* and *U. linza*.

contents among all Cd²⁺ treatments was noted in the two *Ulva* species (Table 1). Nitrate content in *U. prolifera* showed an uptrend with increasing Cd²⁺ concentration; however, with increasing Cd²⁺ concentrations, nitrate content of *U. linza* showed a decline trend after an initial increase. We also found that nitrate contents of *U. linza* were much more than those of *U. prolifera* under all treatments except for 80 µmol L⁻¹ Cd²⁺ treatment (Table 1).

The K⁺/Na⁺ and Ca²⁺/Na⁺ ratios in *U. prolifera* were not influenced by 5 and 10 µmol L⁻¹ Cd²⁺, but they showed declining trends at 20, 40, and 80 µmol L⁻¹ Cd²⁺ (Table 1). In *U. linza*, 5 and 10 µmol L⁻¹ Cd²⁺ had no significant influence on the K⁺/Na⁺ ratio, whereas 20, 40, and 80 µmol L⁻¹ Cd²⁺ decreased that ratio by 6, 45, and 64%, respectively. However, in *U. prolifera*, 20, 40, and 80 µmol L⁻¹ Cd²⁺ decreased the K⁺/Na⁺ ratio by 55, 65, and 78%. No Cd²⁺ treatment significantly changed the Ca²⁺/Na⁺ ratio in *U. linza*.

3.7. Effect of Cadmium Stress on Organic Solutes in *U. prolifera* and *U. linza*. With increasing Cd²⁺ concentration, soluble sugar (SS) content appeared to have an ascending trend after an initial decline in both *Ulva* species. In *U. prolifera*, 40 µmol L⁻¹ Cd²⁺ did not change the SS content, and 80 µmol L⁻¹ Cd²⁺ increased SS concentration by 27% compared to the control. However, in *U. linza*, 40 and 80 µmol L⁻¹ Cd²⁺ increased SS content by 40 and 90%, respectively

(Table 2). In *U. prolifera* and *U. linza*, 5 µmol L⁻¹ Cd²⁺ significantly increased free amino acid (FAA) content by 25 and 16%, respectively. However, 10 µmol L⁻¹ Cd²⁺ had no obvious change on FAA contents of the two *Ulva* species. Treatments with 20, 40, and 80 µmol L⁻¹ Cd²⁺ significantly decreased FAA content by 52, 79, and 87% in *U. prolifera* and by 2, 25, and 43% in *U. linza* (Table 2). Proline (PRO) content was greatly enhanced by Cd²⁺ treatments in both *Ulva* species. At 5, 10, 20, 40, and 80 µmol L⁻¹ Cd²⁺, PRO content was increased 154, 431, 715, 1031, and 1069%, respectively, in *U. prolifera*; and increased 147, 420, 726, 1040, and 1147%, respectively, in *U. linza* (Table 2). Organic acid (OA) content in *U. prolifera* was not affected at 5, 10 and 20 µmol L⁻¹ Cd²⁺, and OA concentration in *U. linza* was not affected at 5, 10, 20, and 40 µmol L⁻¹ Cd²⁺. Treatments with 40 and 80 µmol L⁻¹ Cd²⁺ decreased OA content by 29 and 47%, respectively, in *U. prolifera*, whereas in *U. linza* only 80 µmol L⁻¹ Cd²⁺ decreased OA content by 27% (Table 2). The soluble protein (SP) content in the two *Ulva* species was not affected at 5, 10 and 20 µmol L⁻¹ Cd²⁺ and was decreased at 40 and 80 µmol L⁻¹ Cd²⁺. Treatments with 40 and 80 µmol L⁻¹ Cd²⁺ significantly decreased SP content by, respectively, 16 and 42% in *U. prolifera* and by 8 and 25% in *U. linza* (Table 2).

3.8. Correlation Analysis between RGR and Other Physiological and Biochemical Indexes under Cadmium Stress. Correlation




FIGURE 5: Effects of different concentrations of Cd^{2+} (0, 5, 10, 30, 40, and $80 \mu\text{mol L}^{-1}$) on Fv/Fm (a) and Yield (actual photochemical efficiency of PSII) (b) of *U. prolifera* and *U. linza*.

analysis indicated that RGR of both *Ulva* species was insignificantly related to contents of Chl, Car, Na^+ and Mg^{2+} , and the $\text{Ca}^{2+}/\text{Na}^+$ ratio. In contrast, RGR was highly negative correlated with the contents of Cd^{2+} , Ca^{2+} , SS, and PRO, and highly positive correlated with the contents of N, P, K, Cl, FAA, OA and SP, K^+/Na^+ ratio, OAA, Fv/Fm, and Yield (Table 3).

4. Discussion

Plant growth can be suppressed by Cd [7, 17]. It was reported that *Ulva lactuca* was sensitive to cadmium, as obviously shown by growth reduction and lethal effects at $40 \mu\text{mol L}^{-1}$ Cd^{2+} within 6 days [27]. In the study presented here, *U. prolifera* and *U. linza*, the dominant free-floating *Ulva* species of green tide bloom in the Yellow Sea of China [28], showed sensitivity to Cd^{2+} (reduction in RGR, Fv/Fm, and Yield). Furthermore, this reduction was found to be more pronounced in *U. prolifera* than *U. linza*. After 7 d, *U. prolifera* died at $120 \mu\text{mol L}^{-1}$ Cd^{2+} , whereas *U. linza* was still alive (Figures 1 and 4). This result indicated that *U. linza* had better adaptation to Cd^{2+} toxicity than *U. prolifera*.

It is known that marine macroalgae can concentrate heavy metals to a large extent [2, 29]. In this study, Cd accumulation in *U. prolifera* and *U. linza* increased significantly in response to increased Cd^{2+} concentrations. However, *U. prolifera* accumulated more Cd than *U. linza* (Figure 3). In general, plant accumulation of a given metal

is a function of uptake capacity and intracellular binding sites [30]. The cell walls of plant cells contain proteins and different carbohydrates that can bind metal ions. After the binding sites in the cell wall become saturated, intracellular Cd accumulation mediated by metabolic processes may lead to cell toxicity [31].

Ulva species are widely distributed in the coastal intertidal zones where had full change on salinity level. Thus, many *Ulva* species have strong OAA to cope with variable and heterogeneous environments. Similarly to a number of other stresses, heavy metal toxicity can decrease cell water content and lower the cell water potential (ψ_w) through increased net concentrations of solutes (osmotic adjustment), which is a common response to water stress and an important mechanism for maintaining cell water content and, thus, turgor [18, 32]. In our experiments, OAA of *U. linza* had positive values in the treatments with 5, 10, or $20 \mu\text{mol L}^{-1}$ Cd^{2+} , whereas *U. prolifera* had positive OAA only at 5 and $10 \mu\text{mol L}^{-1}$ Cd^{2+} (Figure 2). When OAA values in *Ulva* were positive, that is, OAA contributed to maintaining turgor, *Ulva* could continue growing, and RGR was positive. However, when OAA in *Ulva* was negative resulting in turgor loss, the growth was stopped, and RGR was negative. Correlation analysis also showed that RGR was positively related to OAA, suggesting that OAA played an important role in maintaining algal growth. Also, good osmotic adjustment enabled plants to maintain high photosynthetic activity (Figure 5).

Cadmium is a nonessential element for plant growth, and it inhibits uptake and transport of many macro- and




FIGURE 6: Effects of different concentrations of Cd²⁺ (0, 5, 10, 20, 40, 80 μmol L⁻¹) on contents of N (a) and P (b) of *U. prolifera* and *U. linza*.

m micronutrients, inducing nutrient deficiency [7, 17]. Contradictory data can be found in the literature on the effects exerted by Cd²⁺ on terrestrial plant. Cadmium was reported to reduce uptake of N, P, K, Ca, Mg, Fe, Zn, Cu, Mn, Ni, and Na in many crop plants [33], whereas other authors found reduced K uptake but unchanged P uptake or even an increase in K content of several crop varieties under Cd²⁺ stress [34, 35]. Obata and Umebayashi [36] reported that Cd²⁺ treatment increased Cu content in the roots of pea, rice, and maize, but unchanged Cu content in cucumber and pumpkin plants. With Cd²⁺ stress, Maksimović et al. [37] observed a reduction in the maize root influx and root-shoot transport of Cu, Zn, and Mn, a reduction in the root-shoot transport of Fe, but an increase in Fe influx and Ca and Mg transport. In this study, the response of total N and P concentrations in tissues of the two *Ulva* species to Cd²⁺ treatments was positively related to their Cd resistance. We found that the treatment with low concentration of Cd²⁺ enhanced N and P contents, but high concentrations of Cd²⁺ ($\geq 20 \mu\text{mol L}^{-1}$) decreased N and P contents in both *Ulva* species. The maintenance of total N and total P was more pronounced in less Cd-sensitive *U. linza* than Cd-sensitive *U. prolifera* (Figure 6). This suggests that the maintenance of a normal level of total N content upon challenge with Cd is likely to be a feature in relative Cd-resistant marine macroalga, similarly to terrestrial plants [38]. In *Ulva*, we found that the contents of K⁺, Ca²⁺, and Cl⁻ were related to RGR, especially K⁺ reduction caused *Ulva* growth reduction significantly (Table 1). Thus, the K⁺/Na⁺ ratio in both *Ulva* species decreased significantly with increasing Cd²⁺ treatment concentrations, and

Cd²⁺-sensitive *U. prolifera* showed a greater K⁺/Na⁺ decline than Cd²⁺-sensitive *U. linza* (Table 1).

We measured a decline in soluble sugar (SS) concentration at low Cd²⁺ treatment concentrations and an increase at high Cd²⁺ concentrations in both *Ulva* species. Moreover, the SS increase of *U. linza* is more marked than that of *U. prolifera*. In other studies, the decline in SS concentration corresponded with the photosynthetic inhibition or stimulation of respiration rate, affecting carbon metabolism and leading to production of other osmotica [39]. The accumulating soluble sugars in plants growing in presence of Cd²⁺ could provide an adaptive mechanism via maintaining a favorable osmotic potential under adverse conditions of Cd²⁺ toxicity [40].

Soluble protein (SP) content in organisms is an important indicator of metabolic changes and responds to a wide variety of stresses [41]. In this work, SP contents in *U. prolifera* and *U. linza* declined with increasing Cd²⁺ treatment concentrations. Free amino acid (FAA) contents in both *Ulva* species first increased and then declined, with such a decline more pronounced in *U. prolifera* than in *U. linza*. The decreased protein content together with the increased free amino acid content suggest that the protein synthesizing machinery was impaired due to the Cd²⁺ effect [42].

PRO accumulation in plant tissues in response to a number of stresses, including drought, salinity, extreme temperatures, ultraviolet radiation, or heavy metals, is well documented [43]. In this study, even though PRO content was increased in Cd²⁺-treated *Ulva*, its absolute amount was relatively low. Under assumed localization of inorganic ions

TABLE 1: Effects of different concentrations of Cd²⁺ (0, 5, 10, 30, 40, and 80 μmol L⁻¹) on inorganic ion content (mmol g⁻¹ DW), K⁺/Na⁺ and Ca²⁺/Na⁺ of *U. prolifera* and *U. linza*.

Cd ²⁺ treatment μmol L ⁻¹	Na ⁺ mmol g ⁻¹ DW	K ⁺ mmol g ⁻¹ DW	Ca ²⁺ mmol g ⁻¹ DW	Mg ²⁺ mmol g ⁻¹ DW	Cl ⁻ mmol g ⁻¹ DW	NO ₃ ⁻ mmol g ⁻¹ DW	K ⁺ /Na ⁺	Ca ²⁺ /Na ⁺
<i>U. prolifera</i>								
0	0.12 ± 0.01 c	0.64 ± 0.04 a	0.20 ± 0.02 b	0.82 ± 0.04 a	0.15 ± 0.01 a	0.34 × 10 ⁻³ ± 0.03 × 10 ⁻³ c	5.01 ± 0.12 a	1.70 ± 0.08 a
5	0.13 ± 0.02 c	0.62 ± 0.05 a	0.23 ± 0.01 ab	0.78 ± 0.05 a	0.11 ± 0.01 b	0.49 × 10 ⁻³ ± 0.06 × 10 ⁻³ c	4.86 ± 0.21 a	1.66 ± 0.07 a
10	0.12 ± 0.01 c	0.63 ± 0.04 a	0.23 ± 0.02 ab	0.76 ± 0.05 a	0.10 ± 0.01 b	0.78 × 10 ⁻³ ± 0.06 × 10 ⁻³ b	5.10 ± 0.14 a	1.84 ± 0.12 a
20	0.17 ± 0.02 b	0.38 ± 0.03 b	0.23 ± 0.02 ab	0.75 ± 0.04 a	0.09 ± 0.01 b	1.41 × 10 ⁻³ ± 0.08 × 10 ⁻³ a	2.26 ± 0.15 b	1.39 ± 0.10 b
40	0.20 ± 0.02 ab	0.35 ± 0.03 b	0.25 ± 0.02 ab	0.79 ± 0.04 a	0.10 ± 0.01 b	1.40 × 10 ⁻³ ± 0.11 × 10 ⁻³ a	1.74 ± 0.11 c	1.24 ± 0.08 bc
80	0.22 ± 0.01 a	0.24 ± 0.02 c	0.26 ± 0.02 a	0.73 ± 0.05 a	0.10 ± 0.01 b	1.43 × 10 ⁻³ ± 0.04 × 10 ⁻³ a	1.12 ± 0.08 d	1.14 ± 0.07 c
<i>U. linza</i>								
0	0.25 ± 0.02 b	0.74 ± 0.04 a	0.18 ± 0.02 c	0.78 ± 0.04 a	0.16 ± 0.01 a	0.86 × 10 ⁻³ ± 0.08 × 10 ⁻³ d	3.02 ± 0.15 a	0.72 ± 0.09 a
5	0.24 ± 0.03 b	0.74 ± 0.04 a	0.17 ± 0.02 c	0.75 ± 0.03 ab	0.12 ± 0.01 b	1.21 × 10 ⁻³ ± 0.10 × 10 ⁻³ c	3.10 ± 0.23 a	0.73 ± 0.08 a
10	0.24 ± 0.02 b	0.73 ± 0.04 a	0.18 ± 0.01 c	0.72 ± 0.04 ab	0.10 ± 0.02 b	1.89 × 10 ⁻³ ± 0.07 × 10 ⁻³ a	3.04 ± 0.12 a	0.75 ± 0.06 a
20	0.24 ± 0.01 b	0.68 ± 0.03 a	0.22 ± 0.02 b	0.64 ± 0.03 b	0.10 ± 0.01 b	2.07 × 10 ⁻³ ± 0.12 × 10 ⁻³ a	2.85 ± 0.11 b	0.83 ± 0.07 a
40	0.26 ± 0.02 b	0.49 ± 0.03 c	0.25 ± 0.02 ab	0.68 ± 0.03 b	0.11 ± 0.01 b	1.65 × 10 ⁻³ ± 0.05 × 10 ⁻³ b	1.67 ± 0.07 c	0.88 ± 0.08 a
80	0.34 ± 0.02 a	0.37 ± 0.02 d	0.27 ± 0.02 a	0.72 ± 0.04 ab	0.12 ± 0.02 b	1.12 × 10 ⁻³ ± 0.11 × 10 ⁻³ c	1.10 ± 0.05 d	0.75 ± 0.06 a

The data in the same column are statistically different if labeled with different letters according to Duncan's multiple range test ($P \leq 0.05$).

TABLE 2: Effects of different concentration of Cd²⁺ (0, 5, 10, 30, 40, and 80 μmol L⁻¹) on organic solute content of *U. prolifera* and *U. linza*.

	Cd ²⁺ treatment μmol L ⁻¹	SS mmol g ⁻¹ DW	FAA mmol g ⁻¹ DW	PRO mmol g ⁻¹ DW	OA mmol g ⁻¹ DW	SP mg g ⁻¹ DW
<i>U. prolifera</i>	0	0.15 ± 0.02 b	1.03 ± 0.05 b	0.13 × 10 ⁻³ ± 0.02 × 10 ⁻³ e	0.17 ± 0.01 a	42.15 ± 2.33 a
	5	0.15 ± 0.02 b	1.29 ± 0.12 a	0.33 × 10 ⁻³ ± 0.02 × 10 ⁻³ d	0.17 ± 0.01 a	41.38 ± 2.76 a
	10	0.12 ± 0.01 bc	1.10 ± 0.04 ab	0.69 × 10 ⁻³ ± 0.04 × 10 ⁻³ c	0.19 ± 0.02 a	40.45 ± 1.86 a
	20	0.10 ± 0.01 c	0.49 ± 0.11 c	1.06 × 10 ⁻³ ± 0.07 × 10 ⁻³ b	0.16 ± 0.02 a	38.39 ± 2.75 ab
	40	0.14 ± 0.01 b	0.22 ± 0.05 d	1.47 × 10 ⁻³ ± 0.09 × 10 ⁻³ a	0.12 ± 0.01 b	35.53 ± 2.63 b
	80	0.19 ± 0.01 a	0.13 ± 0.06 d	1.52 × 10 ⁻³ ± 0.12 × 10 ⁻³ a	0.09 ± 0.01 c	24.35 ± 1.88 c
<i>U. linza</i>	0	0.10 ± 0.01 cd	1.23 ± 0.03 b	0.15 × 10 ⁻³ ± 0.05 × 10 ⁻³ f	0.11 ± 0.02 a	39.27 ± 1.22 a
	5	0.10 ± 0.01 cd	1.43 ± 0.09 a	0.37 × 10 ⁻³ ± 0.02 × 10 ⁻³ e	0.12 ± 0.01 a	38.89 ± 2.37 ab
	10	0.07 ± 0.01 d	1.21 ± 0.10 b	0.78 × 10 ⁻³ ± 0.03 × 10 ⁻³ d	0.13 ± 0.01 a	38.52 ± 2.67 ab
	20	0.10 ± 0.01 c	1.20 ± 0.06 b	1.24 × 10 ⁻³ ± 0.08 × 10 ⁻³ c	0.14 ± 0.02 a	37.13 ± 1.89 ab
	40	0.14 ± 0.02 b	0.97 ± 0.06 c	1.71 × 10 ⁻³ ± 0.07 × 10 ⁻³ b	0.12 ± 0.01 a	35.95 ± 2.41 b
	80	0.19 ± 0.01 a	0.76 ± 0.08 d	1.87 × 10 ⁻³ ± 0.15 × 10 ⁻³ a	0.08 ± 0.01 b	29.34 ± 1.87 c

Different letters in the same column indicate statistical difference according to Duncan's multiple range test ($P \leq 0.05$). "SS, FAA, PRO, OA, and SP" in the table indicate the content of soluble sugar, free amino acid, proline, organic acid, and soluble protein, respectively.

TABLE 3: Correlation coefficients between RGR and other indices for *U. prolifera* and *U. linza*.

Index	Correlation coefficient
Chl content	0.072
Car content	0.198
Fv/Fm	0.830**
Yield	0.858**
Cd ²⁺ content	-0.899**
N content	0.561**
P content	0.687**
OAA	0.766**
Na ⁺ content	-0.138
K ⁺ content	0.881**
Ca ²⁺ content	-0.677**
Mg ²⁺ content	0.060
Cl ⁻ content	0.444**
K ⁺ /Na ⁺	0.627**
Ca ²⁺ /Na ⁺	-0.079
SS content	-0.617**
FAA content	0.828**
PRO content	-0.841**
OA content	0.731**
SP content	0.752**

*Significant at 5% level, ** significant at 1% level (two-tailed, $n = 18$).

in the vacuole and organic solutes in the cytoplasm, FAA and PRO may be mainly in the cytoplasm, accounting for about 5%–10% volume in mature cells [44]. A small amount of FAA and PRO accumulating in the cytoplasm can increase concentration significantly and play an important role in balancing vacuolar osmotic potential [44]. It has often been suggested that PRO accumulation may contribute to osmotic adjustment at the cellular level [39]. In addition, PRO as a

compatible solute may protect enzymes from dehydration and inactivation [18].

In conclusion, exposing *U. prolifera* and *U. linza* to different concentrations of Cd²⁺ resulted in the changes in growth, pigment content, chlorophyll fluorescence parameters, Cd accumulation, OAA, and concentration of N, P, main inorganic ions, and organic solutes. These changes make *U. linza* better adapted to withstanding Cd²⁺ stress in comparison with *U. prolifera*. Our results highlight the role of osmotic adjustment in *Ulva* during Cd²⁺ stress as an important mechanism enabling *Ulva* to maintain photosynthetic activity and, thus, growth under Cd²⁺ stress.

Authors' Contribution

H. Jiang and B. Gao both contributed equally to this paper.

Acknowledgments

The authors thank professor Zed Rengel, University of Western Australia, for his helpful comments on this study. Also, they thank professor Jianjun Chen, University of Florida, for his good revision on this paper. This work was supported by Open Foundation of Key Laboratory of Exploitation and Preservation of Coastal Bio-resources of Zhejiang Province (2010F30003), the National Key Project of Scientific and Technical Supporting Programs funded by Ministry of Science and Technology of China (no. 2009BADA3 B04-8). The authors also acknowledge members of our laboratory for assistance in this work.

References

- [1] L. Zhu, J. Xu, F. Wang, and B. Lee, "An assessment of selected heavy metal contamination in the surface sediments from the South China Sea before 1998," *Journal of Geochemical Exploration*, vol. 108, no. 1, pp. 1–14, 2011.

- [2] A. Sari and M. Tuzen, "Biosorption of Pb(II) and Cd(II) from aqueous solution using green alga (*Ulva lactuca*) biomass," *Journal of Hazardous Materials*, vol. 152, no. 1, pp. 302–308, 2008.
- [3] B. Xu, X. Yang, Z. Gu, Y. Zhang, Y. Chen, and Y. Lv, "The trend and extent of heavy metal accumulation over last one hundred years in the Liaodong Bay, China," *Chemosphere*, vol. 75, no. 4, pp. 442–446, 2009.
- [4] P. Lodeiro, B. Cordero, J. L. Barriada, R. Herrero, and M. E. Sastre De Vicente, "Biosorption of cadmium by biomass of brown marine macroalgae," *Bioresource Technology*, vol. 96, no. 16, pp. 1796–1803, 2005.
- [5] B. Volesky and Z. R. Holan, "Biosorption of heavy metals," *Biotechnology Progress*, vol. 11, no. 3, pp. 235–250, 1995.
- [6] E. A. Webster, A. J. Murphy, J. A. Chudek, and G. M. Gadd, "Metabolism-independent binding of toxic metals by *Ulva lactuca*: cadmium binds to oxygen-containing groups, as determined by NMR," *BioMetals*, vol. 10, no. 2, pp. 105–117, 1997.
- [7] M. Kumar, P. Kumari, V. Gupta, P. A. Anisha, C. R. K. Reddy, and B. Jha, "Differential responses to cadmium induced oxidative stress in marine macroalga *Ulva lactuca* (Ulvales, Chlorophyta)," *BioMetals*, vol. 23, no. 2, pp. 315–325, 2010.
- [8] S. Hajjaligol, M. A. Taher, and A. Malekpour, "A new method for the selective removal of cadmium and zinc ions from aqueous solution by modified clinoptilolite," *Adsorption Science and Technology*, vol. 24, no. 6, pp. 487–496, 2006.
- [9] P. X. Sheng, Y. P. Ting, J. P. Chen, and L. Hong, "Sorption of lead, copper, cadmium, zinc, and nickel by marine algal biomass: characterization of biosorptive capacity and investigation of mechanisms," *Journal of Colloid and Interface Science*, vol. 275, no. 1, pp. 131–141, 2004.
- [10] A. El-Sikaily, A. E. Nemr, A. Khaled, and O. Abdelwehab, "Removal of toxic chromium from wastewater using green alga *Ulva lactuca* and its activated carbon," *Journal of Hazardous Materials*, vol. 148, no. 1-2, pp. 216–228, 2007.
- [11] S. Karthikeyan, R. Balasubramanian, and C. S. P. Iyer, "Evaluation of the marine algae *Ulva fasciata* and *Sargassum* sp. for the biosorption of Cu(II) from aqueous solutions," *Bioresource Technology*, vol. 98, no. 2, pp. 452–455, 2007.
- [12] K. Masakorala, A. Turner, and M. T. Brown, "Influence of synthetic surfactants on the uptake of Pd, Cd and Pb by the marine macroalga, *Ulva lactuca*," *Environmental Pollution*, vol. 156, no. 3, pp. 897–904, 2008.
- [13] P. J. Say, I. G. Burrows, and B. A. Whitton, "Enteromorpha as a monitor of heavy metals in estuaries," *Hydrobiologia*, vol. 195, pp. 119–126, 1990.
- [14] P. Malea, J. W. Rijstenbil, and S. Haritonidis, "Effects of cadmium, zinc and nitrogen status on non-protein thiols in the macroalgae *Enteromorpha* spp. from the Scheldt Estuary (SW Netherlands, Belgium) and Thermaikos Gulf (N Aegean Sea, Greece)," *Marine Environmental Research*, vol. 62, no. 1, pp. 45–60, 2006.
- [15] A. Lin, S. Shen, G. Wang et al., "Comparison of chlorophyll and photosynthesis parameters of floating and attached *Ulva prolifera*," *Journal of Integrative Plant Biology*, vol. 53, no. 1, pp. 25–34, 2011.
- [16] C. Yarish, T. Chopin, R. Wilkes, A. C. Mathieson, X. G. Fei, and S. Lu, "Domestication of Nori for northeast America: the Asian experience," *Bulletin of Aquaculture Association of Canada*, vol. 1, pp. 11–17, 1999.
- [17] J. R. Xia, Y. J. Li, J. Lu, and B. Chen, "Effects of copper and cadmium on growth, photosynthesis, and pigment content in *Gracilaria lemaneiformis*," *Bulletin of Environmental Contamination and Toxicology*, vol. 73, no. 6, pp. 979–986, 2004.
- [18] Q. Zheng, Z. Liu, G. Chen, Y. Gao, Q. Li, and J. Wang, "Comparison of osmotic regulation in dehydration and salinity-stressed sunflower seedlings," *Journal of Plant Nutrition*, vol. 33, no. 7, pp. 966–981, 2010.
- [19] D. P. Häder, M. Lebert, and E. W. Helbling, "Effects of solar radiation on the Patagonian macroalga *Enteromorpha linza* (L.) J. Agardh—Chlorophyceae," *Journal of Photochemistry and Photobiology B*, vol. 62, no. 1-2, pp. 43–54, 2001.
- [20] K. Kamer and P. Fong, "Nitrogen enrichment ameliorates the negative effects of reduced salinity on the green macroalga *Enteromorpha intestinalis*," *Marine Ecology Progress Series*, vol. 218, pp. 87–93, 2001.
- [21] F. B. Wu, J. Dong, Q. Q. Qiong, and G. P. Zhang, "Subcellular distribution and chemical form of Cd and Cd-Zn interaction in different barley genotypes," *Chemosphere*, vol. 60, no. 10, pp. 1437–1446, 2005.
- [22] M. B. Gilliam, M. P. Sherman, J. M. Griscavage, and L. J. Ignarro, "A spectrophotometric assay for nitrate using NADPH oxidation by *Aspergillus* nitrate reductase," *Analytical Biochemistry*, vol. 212, no. 2, pp. 359–365, 1993.
- [23] Q. Zhou and B. Yu, "Changes in content of free, conjugated and bound polyamines and osmotic adjustment in adaptation of vetiver grass to water deficit," *Plant Physiology and Biochemistry*, vol. 48, no. 6, pp. 417–425, 2010.
- [24] M. M. Bradford, "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding," *Analytical Biochemistry*, vol. 72, no. 1-2, pp. 248–254, 1976.
- [25] J. J. Irigoyen, D. W. Emerich, and M. Sanchezdiaz, "Water-stress induced changes in concentrations of proline and total soluble sugars in nodulated alfalfa (*Medicago sativa*) plants," *Plant Physiology*, vol. 84, no. 1, pp. 55–60, 1992.
- [26] J. Song, G. Feng, C. Y. Tian, and F. S. Zhang, "Osmotic adjustment traits of *Suaeda physophora*, *Haloxylon ammodendron* and *Haloxylon persicum* in field or controlled conditions," *Plant Science*, vol. 170, no. 1, pp. 113–119, 2006.
- [27] J. W. Markham, B. P. Kremer, and K. R. Sperling, "Cadmium effects on growth and physiology of *Ulva lactuca*," *Helgoländer Meeresuntersuchungen*, vol. 33, no. 1-4, pp. 103–110, 1980.
- [28] D. Liu, J. K. Keesing, Z. Dong et al., "Recurrence of the world's largest green-tide in 2009 in Yellow Sea, China: *Porphyra yezoensis* aquaculture rafts confirmed as nursery for macroalgal blooms," *Marine Pollution Bulletin*, vol. 60, no. 9, pp. 1423–1432, 2010.
- [29] H. A. Baumann, L. Morrison, and D. B. Stengel, "Metal accumulation and toxicity measured by PAM—Chlorophyll fluorescence in seven species of marine macroalgae," *Ecotoxicology and Environmental Safety*, vol. 72, no. 4, pp. 1063–1075, 2009.
- [30] M. P. Benavides, S. M. Gallego, and M. L. Tomaro, "Cadmium toxicity in plants," *Brazilian Journal of Plant Physiology*, vol. 17, no. 1, pp. 21–34, 2005.
- [31] S. Hu, C. H. Tang, and M. Wu, "Cadmium accumulation by several seaweeds," *Science of the Total Environment*, vol. 187, no. 2, pp. 65–71, 1996.
- [32] M. T. Milone, C. Sgherri, H. Clijsters, and F. Navari-Izzo, "Antioxidative responses of wheat treated with realistic concentration of cadmium," *Environmental and Experimental Botany*, vol. 50, no. 3, pp. 265–276, 2003.

- [33] L. Sanità Di Toppi and R. Gabbrielli, "Response to cadmium in higher plants," *Environmental and Experimental Botany*, vol. 41, no. 2, pp. 105–130, 1999.
- [34] F. F. Nocito, L. Pirovano, M. Cocucci, and G. A. Sacchi, "Cadmium-induced sulfate uptake in maize roots," *Plant Physiology*, vol. 129, no. 4, pp. 1872–1879, 2002.
- [35] Z. Cieccko, S. Kalembasa, M. Wyszowski, and E. Rolka, "The effect of elevated cadmium content in soil on the uptake of nitrogen by plants," *Plant, Soil and Environment*, vol. 50, no. 7, pp. 283–294, 2004.
- [36] H. Obata and M. Umebayashi, "Effects of cadmium on mineral nutrient concentrations in plants differing in tolerance for cadmium," *Journal of Plant Nutrition*, vol. 20, no. 1, pp. 97–105, 1997.
- [37] I. Maksimović, R. Kastori, L. Krstić, and J. Luković, "Steady presence of cadmium and nickel affects root anatomy, accumulation and distribution of essential ions in maize seedlings," *Biologia Plantarum*, vol. 51, no. 3, pp. 589–592, 2007.
- [38] T. Ghnaya, I. Slama, D. Messedi, C. Grignon, M. H. Ghorbel, and C. Abdelly, "Effects of Cd^{2+} on K^+ , Ca^{2+} and N uptake in two halophytes *Sesuvium portulacastrum* and *Mesembryanthemum crystallinum*: consequences on growth," *Chemosphere*, vol. 67, no. 1, pp. 72–79, 2007.
- [39] R. John, P. Ahmad, K. Gadgil, and S. Sharma, "Effect of cadmium and lead on growth, biochemical parameters and uptake in *Lemna polyrrhiza* L.," *Plant, Soil and Environment*, vol. 54, no. 6, pp. 262–270, 2008.
- [40] S. Verma and R. S. Dubey, "Effect of cadmium on soluble sugars and enzymes of their metabolism in rice," *Biologia Plantarum*, vol. 44, no. 1, pp. 117–123, 2001.
- [41] P. K. Singh and R. K. Tewari, "Cadmium toxicity induced changes in plant water relations and oxidative metabolism of *Brassicajuncea* L. plants," *Journal of Environmental Biology*, vol. 24, no. 1, pp. 107–112, 2003.
- [42] S. Jana and M. A. Choudhuri, "Synergistic effects of heavy metal pollutants on senescence in submerged aquatic plants," *Water, Air, and Soil Pollution*, vol. 21, no. 1–4, pp. 351–357, 1984.
- [43] D. Delmail, P. Labrousse, P. Hourdin, L. Larcher, C. Moesch, and M. Botineau, "Differential responses of *Myriophyllum alterniflorum* DC (Haloragaceae) organs to copper: physiological and developmental approaches," *Hydrobiologia*, vol. 664, no. 1, pp. 95–105, 2011.
- [44] R. Munns and M. Tester, "Mechanisms of salinity tolerance," *Annual Review of Plant Biology*, vol. 59, pp. 651–681, 2008.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

