
IJMMS 31:12 (2002) 731–743
 PII. S0161171202103012
<http://ijmms.hindawi.com>
 © Hindawi Publishing Corp.

SUBGROUPS OF QUASI-HNN GROUPS

R. M. S. MAHMOOD and M. I. KHANFAR

Received 20 March 2001 and in revised form 20 February 2002

We extend the structure theorem for the subgroups of the class of HNN groups to a new class of groups called quasi-HNN groups. The main technique used is the subgroup theorem for groups acting on trees with inversions.

2000 Mathematics Subject Classification: 20F65, 20E07, 20E08.

1. Introduction. Quasi-HNN groups have appeared in [3]. In this paper, we use the results of [8] to obtain the structure theorem for the subgroups of quasi-HNN groups and give some applications. We will use the terminology and notation of [3].

Let G be a group, and let I, J be two indexed sets. Let $\{A_i : i \in I\}$, $\{B_i : i \in I\}$, and $\{C_j : j \in J\}$ be families of subgroups of G .

For each $i \in I$, let $\phi_i : A_i \rightarrow B_i$ be an onto isomorphism and for each $j \in J$, let $\alpha_j : C_j \rightarrow C_j$ be an automorphism such that α_j^2 is an inner automorphism determined by $c_j \in C_j$ and c_j is fixed by α_j . That is, $\alpha_j(c_j) = c_j$ and $\alpha_j^2(c) = c_j c c_j^{-1}$ for all $c \in C_j$.

The group G^* with the presentation

$$G^* = \langle G, t_i, t_j \mid \text{rel } G, t_i A_i t_i^{-1} = B_i, t_j C_j t_j^{-1} = C_j, t_j^2 = c_j, i \in I, j \in J \rangle \quad (1.1)$$

is called a *quasi-HNN group* with base G and associated pairs (A_i, B_i) and (C_j, C_j) , $i \in I$, $j \in J$ of subgroups of G , where $\langle G \mid \text{rel } G \rangle$ stands for any presentation of G , $t_i A_i t_i^{-1} = B_i$ stands for the set of relations $t_i w(a) t_i^{-1} = w(\phi_i(a))$, and $t_j C_j t_j^{-1} = C_j$ stands for the set of relations $t_j w(c) t_j^{-1} = w(\alpha_j(c))$, where $w(a)$, $w(\phi_i(a))$, $w(c)$, and $w(\alpha_j(c))$ are words in the generating symbols of the presentation of G of values a , $\phi_i(a)$, c , and $\alpha_j(c)$, respectively, where a runs over the generators of A_i and c over the generators of C_j .

It is proved in [3] that G is embedded in G^* (the imbedding theorem of quasi-HNN groups), and every element g of G^* , $g \neq 1$ can be written as a reduced word of G^* . That is, $g = g_0 t_{k_1}^{e_1} g_1 t_{k_2}^{e_2} g_2 \cdots t_{k_n}^{e_n} g_n$, where $g_s \in G$, $e_s = \pm 1$, $k_s \in I \cup J$, for $s = 1, \dots, n$, such that g contains no subword of the following forms:

- (1) $t_{k_s} a t_{k_s}^{-1}$, $a \in A_i$, or
- (2) $t_{k_s}^{-1} b t_{k_s}$, $b \in B_i$, or
- (3) $t_{k_s}^e c t_{k_s}^\delta$, $c \in C_j$, $e, \delta = \pm 1$, or
- (4) $t_{k_s}^{\pm 2}$, for some $k_s \in J$.

We note that if $J = \emptyset$, then $G^* = \langle G, t_i \mid \text{rel } G, t_i A_i t_i^{-1} = B_i, i \in I \rangle$ is an HNN group with base G and associated pairs (A_i, B_i) , $i \in I$ of subgroups of G .

If $I = \emptyset$, then $G^* = \langle G, t_j \mid \text{rel } G, t_j C_j t_j^{-1} = C_j, t_j^2 = c_j, j \in J \rangle$. In this case we call G^* a *pure quasi-HNN group* with base G and associated pairs of subgroups (C_j, C_j) , $j \in J$ of G . The notation of a tree product of groups as defined in [2] will be needed.

In this paper, we form the subgroup theorem for the quasi-HNN group G^* , defined above, by applying the subgroup theorem for groups acting on trees with inversions obtained by [8]. In particular, if H is a subgroup of the quasi-HNN group G^* defined above, then H is itself a quasi-HNN group (with possibly trivial free part); its base is a tree product π with vertices of the form $xGx^{-1} \cap H$ and amalgamated subgroups either trivial or conjugate of the A_i and C_j intersected with H ; moreover, every conjugate of G intersected with H is either trivial or is conjugate in H to a vertex of π .

This paper is divided into seven sections. In Section 2, we introduce the concepts of graphs and the actions of groups on graphs. In Section 3, we have a summary of the structure of groups acting on trees with inversions. In Section 4, we have a summary of the structure of the subgroup theorem of the groups acting on trees with inversions. In Section 5, we associate a tree on which a quasi-HNN group acts and then we form the subgroup theorem for quasi-HNN groups. In Section 6, we apply the results of Section 5 to find the structures of the subgroups of HNN group. In Section 7, we apply the results of Section 5 to find the structures of the subgroups of pure quasi-HNN group.

2. Basic concepts. We begin by giving preliminary definitions. We denote by a graph X a pair of disjoint sets $V(X)$ and $E(X)$ with $V(X)$ nonempty, together with a mapping $E(X) \rightarrow V(X) \times V(X)$, $\gamma \rightarrow (o(\gamma), t(\gamma))$, and a mapping $E(X) \rightarrow E(X)$, $\gamma \rightarrow \bar{\gamma}$, satisfying the conditions that $\overline{\bar{\gamma}} = \gamma$, and $o(\bar{\gamma}) = t(\gamma)$, for all $\gamma \in E(X)$. The case $\bar{\gamma} = \gamma$ is possible for some $\gamma \in E(X)$. For $\gamma \in E(X)$, $o(\gamma)$, and $t(\gamma)$ are called the ends of γ , and $\bar{\gamma}$ is called the inverse of γ . If A is a set of edges of X , we define \bar{A} to be the set of inverses of the edges of A . That is, $\bar{A} = \{\bar{\gamma} : \gamma \in A\}$. For definitions of subgraphs, trees, morphisms of graphs, and $\text{Aut}(X)$, the set of all automorphisms of the graph X which is a group under the composition of morphisms of graphs, see [5] or [9]. We say that a group G acts on a graph X , if there is a group homomorphism $\phi : G \rightarrow \text{Aut}(X)$. If $x \in X$ (vertex or edge) and $g \in G$, we write $g(x)$ for $(\phi(g))(x)$. If $\gamma \in E(X)$ and $g \in G$, then $g(o(\gamma)) = o(g(\gamma))$, $g(t(\gamma)) = t(g(\gamma))$, and $g(\bar{\gamma}) = \overline{g(\gamma)}$. The case $g(\gamma) = \bar{\gamma}$ for some $g \in G$ and some $\gamma \in E(X)$ may occur. That is, G acts with inversions on X .

We have the following notations related to the action of the group G on the graph X :

- (1) if $x \in X$ (vertex or edge), we define $G(x) = \{g(x) : g \in G\}$, and this set is called the orbit of x containing x ;
- (2) if $x, y \in X$, we define $G(x, y) = \{g \in G : g(x) = y\}$ and $G(x, x) = G_x$, the stabilizer of x . Thus $G(x, y) \neq \emptyset$ if and only if x and y are in the same G orbit. It is clear that if $v \in V(X)$, $\gamma \in E(X)$, and $u \in \{o(\gamma), t(\gamma)\}$, then $G(v, \gamma) = \emptyset$, $G_{\bar{\gamma}} = G_\gamma$, and $G_\gamma \leq G_u$. If $H \leq G$ and $x \in X$, then it is clear that $H_x = H \cap G_x$;
- (3) let X^G be the set of elements of X fixed by G . That is, $X^G = \{x \in X : G = G_x\}$.

3. The structure of groups acting on trees with inversions. In this section, we summarize the presentation for groups of groups acting on trees with inversions obtained by [5].

DEFINITION 3.1. Let G be a group acting on a graph X , and T and Y be two subtrees of X such that $T \subseteq Y$ and $X^G = \emptyset$. Then T is called a tree of representatives for the action of G on X if T contains exactly one vertex from each G vertex orbit, and Y is called a fundamental domain for the action G on X , if each edge of Y has at least one end in T , and Y contains exactly one edge (say) y from each G edge orbit such that $G(\bar{y}, y) = \emptyset$, and exactly one pair x and \bar{x} from each G -edge orbit such that $G(\bar{x}, x) \neq \emptyset$. It is clear that the properties of T and Y imply that if u and v are two vertices of T such that $G(u, v) \neq \emptyset$, and if x and y are two edges of Y such that $G(x, y) \neq \emptyset$, then $u = v$ and $x = y$ or $x = \bar{y}$.

Let Y be as above. Define the following subsets Y_0, Y_1, Y_2 of edges of Y as follows:

- (1) $Y_0 = E(T)$, the set of edges of T ;
- (2) $Y_1 = \{y \in E(Y) : o(y) \in V(T), t(y) \notin V(T), G(\bar{y}, y) = \emptyset\}$;
- (3) $Y_2 = \{x \in E(Y) : o(x) \in V(T), t(x) \notin V(T), G(\bar{x}, x) \neq \emptyset\}$.

It is clear that $\bar{Y}_0 = Y_0, E(Y) = Y_0 \cup Y_1 \cup Y_2 \cup \bar{Y}_1 \cup \bar{Y}_2$, and G acts with inversions on X if and only if $Y_2 \neq \emptyset$.

For the rest of this section G will be a group acting on a graph X , T be a tree of representatives for the action of the group G on X , and Y be a fundamental domain for the action of G on X such that $T \subseteq Y$ and $X^G = \emptyset$.

We have the following definitions.

DEFINITION 3.2. For each vertex v of X , define v^* to be the unique vertex of T such that $G(v, v^*) \neq \emptyset$. That is, v and v^* are in the same G vertex orbit. It is clear that if v is a vertex of T , then $v^* = v$ and in general, for any two vertices u and v of X such that $G(u, v) \neq \emptyset$, we have $u^* = v^*$.

DEFINITION 3.3. For each edge y of $Y_0 \cup Y_1 \cup Y_2$ define $[y]$ to be an element of $G(t(y), (t(y))^*)$. That is, $[y]$ satisfies the condition $[y]((t(y))^*) = t(y)$, to be chosen as follows:

$$[y] = 1 \quad \text{if } y \in Y_0, \quad y = \bar{y} \quad \text{if } y \in Y_2. \tag{3.1}$$

Define $[\bar{y}]$ to be the element

$$[\bar{y}] = \begin{cases} [y] & \text{if } y \in Y_0 \cup Y_2, \\ [y]^{-1} & \text{if } y \in Y_1, \end{cases} \tag{3.2}$$

$[y]$ is called *the value of the edge y* .

DEFINITION 3.4. For each edge y of Y define $-y$ to be the edge $-y = [y]^{-1}(y)$ if $y \in Y_1$, otherwise $-y = y$. It is clear that $t(-y) = (t(y))^*$ and $G_{-y} \leq G_{(t(y))^*}$.

DEFINITION 3.5. By a reduced word w of G we mean an expression of the form $w = g_0 \cdot y_1 \cdot g_1 \cdot y_2 \cdot g_2 \cdots y_n \cdot g_n, n \geq 0, y_i \in E(Y), \text{ for } i = 1, 2, \dots, n$ such that

- (1) $g_0 \in G_{(o(y_1))^*}$;
- (2) $g_i \in G_{(t(y_i))^*}$, for $i = 1, 2, \dots, n$;
- (3) $(t(y_i))^* = (o(y_{i+1}))^*$, for $i = 1, 2, \dots, n-1$;
- (4) w contains no expression of the form $y_i \cdot g_i y_i^{-1}$ if $g_i \in G_{-y_i}$ and $G(y_i, \bar{y}_i) = \emptyset$, or $y_i \cdot g_i \cdot y_i$ if $g_i \in G_{y_i}$ and $G(y_i, \bar{y}_i) \neq \emptyset$.

If $w = g_0$, then w is called a trivial word of G . If $(o(y_1))^* = (t(y_n))^*$ then w is called a closed word of G .

The value of w denoted $[w]$ is defined to be the element

$$[w] = g_0[y_1]g_1[y_2]g_2 \cdots [y_n]g_n \text{ of } G. \tag{3.3}$$

Before we state the main result of this section, we have the following notations: let $v \in V(T)$, $m \in Y_0$, $y \in Y_1$, and $x \in Y_2$, then

- (1) $\langle G_v \mid \text{rel } G_v \rangle$ stands for any presentation of G_v ;
- (2) $G_m = G_m$ stands for the set of relations $w(g) = w'(g)$, where $w(g)$ and $w'(g)$ are words in the generating symbols of $G_{o(m)}$ and $G_{t(m)}$, respectively of value g , and g is in the set of the generators of G_m ;
- (3) $y \cdot [y]^{-1}G_y[y] \cdot y^{-1} = G_y$ stands for the set of relations: $yw([y]^{-1}g[y])y^{-1} = w(g)$, where $w([y]^{-1}g[y])$ and $w(g)$ are words in the generating symbols of $G_{(t(y))^*}$ and $G_{(o(y))^*}$ of values $[y]^{-1}g[y]$ and g , respectively, where g is in the set of the generators of G_y ;
- (4) $x \cdot G_x \cdot x^{-1} = G_x$ stands for the set of relations: $xw(g)x^{-1} = w'(g)$, where $w(g)$ and $w'(g)$ are words in the generating symbols of $G_{o(x)}$ of values g and $[x]g[x]^{-1}$, respectively, where g is in the set of the generators of G_x ;
- (5) $x^2 = [x]^2$ stands for the of relation $x^2 = w([x]^2)$ where $w([x]^2)$ is a word in the set of the generators of G_x .

THEOREM 3.6. *Let G, X, T, Y, Y_0, Y_1 , and Y_2 be as above. Then the following are equivalent:*

- (i) X is tree;
- (ii) G is generated by the generators of G_v and by the elements $[y]$ and $[x]$ and G has the presentation

$$G = \langle G_v, y, x \mid \text{rel } G_v, G_m = G_m, y \cdot [y]^{-1}G_y[y] \cdot y^{-1} = G_y, x \cdot G_x \cdot x^{-1} = G_x, x^2 = [x]^2 \rangle \tag{3.4}$$

via the map $G_v \rightarrow G_v, y \rightarrow [y]$, and $x \rightarrow [x]$ where $v \in V(T)$, $m \in Y_0$, $y \in Y_1$, and $x \in Y_2$;

- (iii) every element of G is the value of a closed and reduced word of G . Moreover, if w is a nontrivial closed and reduced word of G , then $[w]$ is not the identity element of G .

PROOF. (i) implies (ii) follows from [5, Theorem 5.1].

(ii) implies (iii) follows from [6, Corollary 1].

(iii) implies (i) follows from [7, Corollary 3.6]. □

4. Structure theorem for subgroups of groups acting on trees with inversions.

In this section, we summarize the methods for obtaining generators and presentation for subgroups of groups acting on trees with inversions obtained by [8, Theorem 3, page 28]. In [4, Theorem 2.5, page 91] generators and presentation for a subgroup B of a given group A with given generators and presentation is obtained by a method called the Reidemeister-Schreier methods. In view of such methods we have the following

remarks related to subgroups of groups acting on trees with inversions: let G be a group acting on a tree X , v be a vertex of X , z be an edge of X and, H be a subgroup of G , then

- (1) the generators and presentation for the vertex stabilizer G_v of v under G are arbitrary;
- (2) the generators for the edge stabilizer G_z of z under G are arbitrary;
- (3) the generators and presentation for G are those of [Theorem 3.6\(ii\)](#);
- (4) the generators and presentation for the vertex stabilizer $H_v = H \cap G_v$ of v under H are those obtained by the Reidemeister-Schreier methods from the generators and presentation for G_v ;
- (5) the generators for the edge stabilizer $H_z = H \cap G_z$ of z under H are those obtained by the Reidemeister-Schreier methods from the generators of G_z ;
- (6) the generators and presentation for H are given in [Theorem 4.4](#).

Now we proceed to obtain the generators and presentation for subgroups of groups acting on trees with inversions as follows.

Let G, X, T, Y, Y_0, Y_1 , and Y_2 be as above such that X is a tree, and H be a subgroup of G . We have the following definitions.

DEFINITION 4.1. (1) For each $g \in G$ and $y \in Y_0 \cup Y_1 \cup Y_2$, define D_y^g to be any double coset representative system for $G_{o(y)} \text{ mod } (G_{o(y)} \cap g^{-1}Hg, G_y)$ containing 1 but otherwise arbitrary.

(2) For each $v \in V(T)$, let D_v be a double coset representative system for $G \text{ mod } (H, G_v)$ satisfying the condition that if $g \in D_v$ and

$$g = g_0[y_1]g_1[y_2]g_2 \cdots [y_n]g_n, \tag{4.1}$$

where $g_0 \cdot y_1 \cdot g_1 \cdot y_2 \cdot g_2 \cdots y_n \cdot g_n$ is a closed and reduced word of G , then $g_0[y_1]g_1[y_2]g_2 \cdots [y_i] \in D_{(o(y_{i+1}))^*}$, and $g_i \in D_{y_i}^{f_i}$, where $f_i = g_0[y_1]g_1[y_2]g_2 \cdots [y_i]$. For more details see [\[8, page 23\]](#).

(3) For each $y \in Y_0 \cup Y_1 \cup Y_2$, define D^y to be the set $D^y = \{de : d \in D_{o(y)}, e \in D_y^d, de[y] \notin D_{(t(y))^*}\}$, and define D_y to be the set $D_y = \{de : d \in D_{o(y)}, e \in D_y^d, de[y] \in D_{(t(y))^*}\}$.

DEFINITION 4.2. The collection of double coset representative $\{D_v\}$, and the associated collections $\{D^y\}$, and $\{D_y\}$ defined above will be called a *cross* for $G \text{ mod } H$. For the existence of a cross for $G \text{ mod } H$ we refer the readers to [\[8, page 23\]](#).

PROPOSITION 4.3. *For each $y \in Y_0 \cup Y_1 \cup Y_2$ and $g \in D^y$ there exist unique elements $\overline{g[y]} \in D_{(t(y))^*}$, $\overline{g[y]} \in D_y^{g[y]}$, and $a_y \in [y]^{-1}G_y[y]$ such that $g[y](\overline{g[y]}\overline{g[y]}a_y)^{-1} \in H$.*

PROOF. See [\[8, page 28\]](#). □

Note that $H \cap \overline{g[y]}\overline{g[y]}G_y\overline{g[y]}^{-1}\overline{g[y]}^{-1}$ is a subgroup of $H \cap gG_{o(y)}g^{-1}$.

Before we proceed to state the main result of this section, we introduce the following convention on notation on the generators and defining relations of H :

(A) For any vertex v of T , $\langle H \cap dG_vd^{-1} \mid \text{rel}(H \cap dG_vd^{-1}) \rangle$ is the presentation of the subgroup $H \cap dG_vd^{-1}$ of G_v obtained by the Reidemeister-Schreier methods.

(B) For any $y \in Y_0 \cup Y_1 \cup Y_2$, the relations

$$g(y) \cdot (H \cap \overline{g[y]g[y]G_y \overline{g[y]}^{-1} \overline{g[y]}^{-1}}) \cdot (g(y))^{-1} = H \cap gG_y g^{-1}, \tag{4.2}$$

where $g \in D^y$ are similar to the relations of G as in [Theorem 3.6\(ii\)](#).

The main result of this section is the following theorem. For the proof see [\[8, Theorem 3\]](#).

THEOREM 4.4. *Let G be a group acting on a tree X , and let T be a tree of representatives for the action of the group G on X and Y be a fundamental domain for the action of G on X such that $T \subseteq Y$ and $X^G = \emptyset$. Let H be a subgroup of G , and $\{D_u\}$, $\{D^z\}$, and $\{D_z\}$ be a cress for $G \bmod H$, for all $u \in V(T)$, and all $z \in Y_0 \cup Y_1 \cup Y_2$. Then we have the following:*

(I) H is generated by the following:

- (1) the generators of the subgroups $H \cap dG_v d^{-1}$ of G_v , where $v \in V(T)$ and $d \in D_v$;
- (2) the elements $g(\overline{g} \overline{g} a_m)^{-1}$, where $m \in Y_0$ and $g \in D^m$;
- (3) the elements $g[y](\overline{g[y]g[y]a_y})^{-1}$, where $y \in Y_1$ and $g \in D^y$;
- (4) the elements $g[x](\overline{g[x]g[x]a_x})^{-1}$, where $x \in Y_2$ and $g \in D^x$ such that $H \cap [x]G_x g^{-1} = \emptyset$;
- (5) the elements $g[x]g^{-1}$, where $x \in Y_2$ and $g \in D^x$ such that $H \cap g[x]G_x g^{-1} \neq \emptyset$.

(II) H has the presentation $\langle P \mid R \rangle$, where P is the set of generating symbols of the following forms:

- (1) the generating symbols of the subgroups $H \cap dG_v d^{-1}$ of G_v , where $d \in D_v$;
- (2) the edges $g(m)$, where $m \in Y_0$ and $g \in D^m$;
- (3) the edges $g(y)$, where $y \in Y_1$ and $g \in D^y$;
- (4) the edges $g(x)$, where $x \in Y_2$ and $g \in D^x$ such that $H \cap g[x]G_x g^{-1} \neq \emptyset$.

(III) R is the set of relations of the following forms:

- (1) the relations, $\text{rel}(H \cap dG_v d^{-1})$ of the subgroups $H \cap dG_v d^{-1}$ of G_v , where $d \in D_v$;
- (2) the relations $H \cap gG_m g^{-1} = H \cap gG_m g^{-1}$, where $m \in Y_0$ and $g \in D^m$;
- (3) the relations $H \cap gG_y g^{-1} = H \cap gG_y g^{-1}$, where $y \in Y_0$ and $g \in D_y$;
- (4) the relations $H \cap gG_x g^{-1} = H \cap gG_x g^{-1}$, where $x \in Y_2$ and $g \in D_x$ such that $H \cap g[x]G_x g^{-1} \neq \emptyset$;
- (5) the relations $g(m) \cdot (H \cap \overline{g} \overline{g} G_m \overline{g}^{-1} \overline{g}^{-1}) \cdot (g(m))^{-1} = H \cap gG_m g^{-1}$, where $m \in Y_0$ and $g \in D^m$;
- (6) the relations $g(y) \cdot (H \cap \overline{g[y]g[y]G_y \overline{g[y]}^{-1} \overline{g[y]}^{-1}}) \cdot (de(y))^{-1} = H \cap gG_y g^{-1}$, where $y \in Y_1$ and $g \in D^y$;
- (7) the relations $g(x) \cdot (H \cap \overline{g[x]g[x]G_x \overline{g[x]}^{-1} \overline{g[x]}^{-1}}) \cdot (g(x))^{-1} = H \cap gG_x g^{-1}$, where $x \in Y_2$ and $g \in D^x$ such that $H \cap g[x]G_x g^{-1} \neq \emptyset$;
- (8) the relations $g(x) \cdot (H \cap gG_x g^{-1}) \cdot (g(x))^{-1} = H \cap gG_x g^{-1}$, where $x \in Y_2$ and $g \in D^x$ such that $H \cap g[x]G_x g^{-1} \neq \emptyset$;
- (9) the relations $(g(x))^2 = g[x]^2 g^{-1}$, where $x \in Y_2$ and $g \in D^x$ such that $H \cap g[x]G_x g^{-1} \neq \emptyset$.

DEFINITION 4.5. A group is called *quasi free group* if it is a free product of a free group and a number of cyclic groups of order 2.

We have the following corollaries of [Theorem 4.4](#).

COROLLARY 4.6. *If $H \cap G_v = \{1\}$ for all $v \in V(X)$, then H is a quasi free group.*

COROLLARY 4.7. *If $H \cap G_y = \{1\}$ for all $y \in E(X)$, then H is a free product of a quasi-HNN group and the intersections of H with G_v for all $v \in V(T)$.*

COROLLARY 4.8. *If H is a nontrivial free product, then H is either infinite cyclic group, a finite cyclic group of order 2, or H is contained in G_v for some $v \in V(X)$.*

COROLLARY 4.9. *If H is a normal subgroup of G such that $H \cap G_v$ is quasi free group for all $v \in V(X)$, and $H \cap G_y = \{1\}$ for all $y \in E(X)$, then H is a quasi free group.*

COROLLARY 4.10. *If H has the property that $D^z = \emptyset$, for all $z \in Y_0 \cup Y_1 \cup Y_2$, then H is a tree product of the subgroups $H \cap dG_v d^{-1}$ of G_v , where $v \in V(T)$ and $d \in D_v$ with amalgamation subgroups $H \cap gG_m g^{-1}$, $m \in Y_0$, $g \in D_m$, $H \cap gG_y g^{-1}$, $y \in Y_1$, $g \in D_y$, and $H \cap gG_x g^{-1}$, $x \in Y_2$, $g \in D_x$ such that $H \cap g[x]G_x g^{-1} \neq \emptyset$.*

5. Subgroups of quasi-HNN groups. In this section, we construct a tree on which a quasi-HNN group acts with inversions, and then we formulate its subgroups.

The following lemma is essential for the proof of the main theorem of the paper.

LEMMA 5.1. *A group is a quasi-HNN group if and only if there is a tree on which the group acts with inversion and is transitive on the set of vertices. Moreover, the stabilizer of any vertex is conjugate to the base and the stabilizer of any edge is conjugate to an associate subgroup of the base.*

PROOF. Let G be a group acting with inversions on a tree X such that G is transitive on $V(X)$. Let T be a tree of representatives, and Y be a fundamental domain for the action of G on X such that $T \subseteq Y$. Since G is transitive on $V(X)$, therefore T consists of exactly one vertex v (say) without edges. So $Y_0 = \emptyset$. Since G acts with inversions on X , therefore $Y_2 \neq \emptyset$. Then by [Theorem 3.6](#), G has the presentation

$$G = \langle G_v, \gamma, x \mid \text{rel } G_v, \gamma \cdot [\gamma]^{-1} G_\gamma [\gamma] \cdot \gamma^{-1} = G_\gamma, x \cdot G_x \cdot x^{-1} = G_x, x^2 = [x]^2 \rangle \quad (5.1)$$

via the map $G_v \rightarrow G_v$, $\gamma \rightarrow [\gamma]$, and $x \rightarrow [x]$ where $v \in V(T)$, $\gamma \in Y_1$, and $x \in Y_2$. Then G is a quasi-HNN group of base G_v , and associated isomorphic pairs $([\gamma]^{-1} G_\gamma [\gamma], G_\gamma)$ of subgroups of G_v via the isomorphism $\phi_\gamma : [\gamma]^{-1} G_\gamma [\gamma] \rightarrow G_\gamma$ defined by $\phi_\gamma([\gamma]^{-1} g [\gamma]) = g$, for all $g \in G_\gamma$ and all $\gamma \in Y_1$ such that $o(\gamma) = v$, and associated isomorphic pairs (G_x, G_x) of subgroups of G_v via the mapping $\alpha_x : G_x \rightarrow G_x$ defined by $\alpha_x(g) = [x]g[x]^{-1}$, for all $g \in G_x$ and all $x \in Y_2$ such that $o(x) = v$. Let $g_x = [x]^2$. It is clear that α_x is an automorphism of G_x , and α_x^2 is inner automorphism of G_x determined by g_x , and $\alpha_x([x]^2) = [x]^2$.

Conversely, let G^* be the quasi-HNN group

$$G^* = \langle G, t_i, t_j \mid \text{rel } G, t_i A_i t_i^{-1} = B_i, t_j C_j t_j^2 = C_j, t_j^2 = c_j, i \in I, j \in J \rangle \quad (5.2)$$

of base G , and associated pairs (A_i, B_i) , and (C_j, C_j) , $i \in I, j \in J$ of subgroups of G .

Now, we construct a tree X on which G^* acts on X with inversions such that G^* is transitive on $V(X)$, and call it the standard tree associated with the quasi-HNN group. In this construction, define

$$V(X) = \{gG : g \in G^*\}, \tag{5.3}$$

and define

$$E(X) = \{(gB_i, t_i) : g \in G^*, i \in I\} \cup \{(gA_i, t_i^{-1}) : g \in G^*, i \in I\} \cup \{(gC_j, t_j) : g \in G^*, j \in J\}. \tag{5.4}$$

The terminals of the edges are defined as follows:

$$o(gB_i, t_i) = o(gA_i, t_i^{-1}) = o(gC_j, t_j) = gG, \tag{5.5}$$

$$t(gB_i, t_i) = gt_iG, \quad t(gA_i, t_i^{-1}) = gt_i^{-1}G, \quad t(gC_j, t_j) = gt_jG.$$

The inverses of the edges are defined as follows:

$$\overline{(gB_i, t_i)} = (gt_iA_i, t_i^{-1}), \quad \overline{(gA_i, t_i^{-1})} = (gt_i^{-1}B_i, t_i), \quad \overline{(gC_j, t_j)} = (gt_jC_j, t_j). \tag{5.6}$$

From above X is a graph. G^* acts on X as follows: let $g' \in G^*$. Then for any vertex gG of X we have $g'(gG) = g'gG$, and for any edge (gB_i, t_i) , or (gA_i, t_i^{-1}) , or (gC_j, t_j) of X , we have

$$g'(gB_i, t_i) = (g'gB_i, t_i), \quad g'(gA_i, t_i^{-1}) = (g'gA_i, t_i^{-1}), \quad g'(gC_j, t_j) = (g'gC_j, t_j). \tag{5.7}$$

The action of G^* on the vertices of X is transitive because for any two vertices aG and bG of X we have $ba^{-1}(aG) = bG$. That is, the element ba^{-1} of G^* maps the vertex aG to the vertex bG . Then there is exactly one G vertex orbit. We take $T = \{G\}$ to be the tree of representatives for the action of G^* on X . G^* acts on X with inversions because the element $t_j \in G^*$ maps the edge (C_j, t_j) to its inverse $\overline{(C_j, t_j)}$. That is,

$$t_j(C_j, t_j) = (t_jC_j, t_j) = \overline{(C_j, t_j)}. \tag{5.8}$$

Let λ_i, β_i , and γ_j stand for the edges (B_i, t_i) , (A_i, t_i^{-1}) , and (C_j, t_j) , respectively. Then

$$o(\lambda_i) = o(\beta_i) = o(\gamma_j) = G, \quad t(\lambda_i) = t_iG, \quad t(\beta_i) = t_i^{-1}G, \quad t(\gamma_j) = t_jG, \tag{5.9}$$

$$\bar{\lambda}_i = (t_iA_i, t_i^{-1}), \quad \bar{\beta}_i = (t_i^{-1}B_i, t_i), \quad \bar{\gamma}_j = (t_jC_j, t_j).$$

From above, we see that λ_i and $\bar{\lambda}_i$, and β_i and $\bar{\beta}_i$ are in different G^* edge orbits, while γ_j and $\bar{\gamma}_j$ are in the same G^* edge orbit. Moreover, any edge of X is of the form $g(\lambda_i)$, $g(\bar{\lambda}_i)$, or $g(\gamma_j)$ for some $g \in G^*$.

Let Y be the subgraph of X consisting of the edges λ_i, β_i , and γ_j , and their terminals and inverses for all $i \in I$ and all $j \in J$. Thus, $V(Y) = \{G, t_i G, t_i^{-1} G, t_j G : i \in I, j \in J\}$ and $E(Y) = \{\lambda_i, \beta_i, \gamma_j, \bar{\lambda}_i, \bar{\beta}_i, \bar{\gamma}_j : i \in I, j \in J\}$. It is clear that T is a tree of representatives for the action G^* on X , and Y is a fundamental domain for the action of G^* on X and $T \subseteq Y$. Moreover, $Y_0 = \emptyset$, $Y_1 = \{\lambda_i : i \in I\}$, and $Y_2 = \{\gamma_j : j \in J\}$.

From above, it is easy to verify that the stabilizer of the vertex v , where $v = G$ is $G_v^* = G$ and the stabilizers of the edges λ_i, β_i , and γ_j are $G_{\lambda_i}^* = B_i, G_{\beta_i}^* = A_i$, and $G_{\gamma_j}^* = C_j$, respectively. This implies that the stabilizer of any vertex is conjugate to the base G and the stabilizer of any edge is conjugate to an associate subgroup A_i, B_i , or C_j of the base G . The values of the edges λ_i, β_i , and γ_j are $[\lambda_i] = t_i, [\beta_i] = t_i^{-1}$, and $[\gamma_j] = t_j$, respectively. By [Theorem 3.6](#), the presentation of G^* and the action of G^* on X implies that X is a tree.

This completes the proof. □

REMARK 5.2. The tree X , constructed above, will be called the standard tree of the quasi-HNN group G^* .

In view of [Lemma 5.1](#) and [Definition 4.2](#) the following concepts is clear.

Let H be a subgroup of the quasi-HNN group

$$G^* = \langle G, t_i, t_j \mid \text{rel } G, t_i A_i t_i^{-1} = B_i, t_j C_j t_j^{-1} = C_j, t_j^2 = c_j, i \in I, j \in J \rangle \tag{5.10}$$

of base G and associated pairs (A_i, B_i) , and (C_j, C_j) , $i \in I, j \in J$ of subgroups of G .

We have the following concepts:

- (1) for each $i \in I, j \in J$, and $d \in G$, let D_i^d and D_j^d be double coset representative system for $G \text{ mod } (G \cap d^{-1} H d, B_i)$, and $G \text{ mod } (G \cap d^{-1} H d, C_j)$, respectively containing 1, but otherwise arbitrary;
- (2) let D be a double coset representative system for $G^* \text{ mod } (H, D)$ satisfying the condition that if $g = g_0 t_{k_1}^{e_1} g_1 t_{k_2}^{e_2} g_2 \cdots t_{k_n}^{e_n} g_n \in D$, then $w_s \in D$ and $g_s \in D_{k_s}^{w_s}$, where

$$w_s = g_0 t_{k_1}^{e_1} g_1 t_{k_2}^{e_2} g_2 \cdots t_{k_s}^{e_s} \quad \text{for } s = 1, \dots, n; \tag{5.11}$$

- (3) for each $s \in I \cup J$, let $D^s = \{de : d \in D, e \in D_s^d, \det_s \notin D\}$, and $D_s = \{de : d \in D, e \in D_s^d, \det_s \in D\}$;
- (4) for each $s \in I \cup J$, and $g \in D^s$, let $\overline{gt_s} \in D$ be the representative of gt_s in D , and $\overline{gt_s} \in D_s^{\overline{gt_s}}$ be the representative of gt_s in $D_s^{\overline{gt_s}}$;
- (5) for each $i \in I$, and $g \in D^i$, let k_i be an element of A_i such that $gt_i(\overline{gt_i} \overline{gt_i} k_i)^{-1} \in H$;
- (6) for each $j \in J$, and $g \in D^j$, let k_j be an element of C_j such that $gt_j(\overline{gt_j} \overline{gt_j} k_j)^{-1} \in H$.

DEFINITION 5.3. The collection of double coset representative $\{D\}$, and the associated collections $\{D_s^d\}$, $\{D^s\}$, and $\{D_s\}$ defined above will be called a cress for $G^* \text{ mod } H$.

The main result of this section is the following theorem.

THEOREM 5.4. *Let H be a subgroup of the quasi-HNN group*

$$G^* = \langle G, t_i, t_j \mid \text{rel } G, t_i A_i t_i^{-1} = B_i, t_j C_j t_j^{-1} = C_j, t_j^2 = c_j, i \in I, j \in J \rangle \quad (5.12)$$

with base G and associated pairs (A_i, B_i) , and (C_j, C_j) , $i \in I, j \in J$ of subgroups of G .

Let $\{D\}, \{D_s^d\}, \{D^s\}$, and $\{D_s\}$, $s \in I \cup J$, be a cress for $G^* \text{ mod } H$.

Then, we have the following:

(I) H is generated by the following:

- (1) the generators of the subgroup $H \cap dGd^{-1}$ of G , for all $d \in D$;
- (2) the elements $gt_i(\overline{gt_i \overline{gt_i k_i}})^{-1}$, where $i \in I$ and $g \in D^i$;
- (3) the elements $gt_j(\overline{gt_j \overline{gt_j k_j}})^{-1}$, where $j \in J, g \in D^j$, and $H \cap gt_j C_j g^{-1} = \emptyset$;
- (4) the elements $gt_j g^{-1}$, where $j \in J, g \in D^j$, and $H \cap gt_j C_j g^{-1} \neq \emptyset$.

(II) H has the presentation $\langle P \mid R \rangle$, where P is the set of generating symbols of the following forms:

- (1) the generating symbols of the subgroup $H \cap dGd^{-1}$ of G , for all $d \in D$;
- (2) the symbols $g(\lambda_i)$, where $i \in I$ and $g \in D^i$;
- (3) the symbols $g(\lambda_j)$, where $j \in J$ and $g \in D^j$.

(III) R is the set of relations of the following forms:

- (1) the relations $\text{rel}(H \cap dGd^{-1})$ of the subgroup $H \cap dGd^{-1}$ of G , for all $d \in D$;
- (2) the relations $H \cap gA_i g^{-1} = H \cap gB_i g^{-1}$, where $i \in I$ and $g \in D^i$;
- (3) the relations $H \cap gC_j g^{-1} = H \cap gC_j g^{-1}$, where $j \in J$ and $g \in D^j$;
- (4) the relations $g(\lambda_i) \cdot (H \cap \overline{gt_i \overline{gt_i A_i \overline{gt_i}^{-1} \overline{gt_i}^{-1}}}) \cdot (g(\lambda_i))^{-1} = H \cap gB_i g^{-1}$, where $i \in I$ and $g \in D^i$;
- (5) the relations $g(\lambda_j) \cdot (H \cap \overline{gt_j \overline{gt_j C_j \overline{gt_j}^{-1} \overline{gt_j}^{-1}}}) \cdot (g(\lambda_j))^{-1} = H \cap g(C_j)g^{-1}$, where $j \in J$ and $g \in D^j$ such that $H \cap gt_j C_j g^{-1} = \emptyset$;
- (6) the relations $g(\lambda_j) \cdot (H \cap gC_j g^{-1}) \cdot (g(\lambda_j))^{-1} = H \cap g(C_j)g^{-1}$, where $j \in J$ and $g \in D^j$ such that $H \cap gt_j C_j g^{-1} \neq \emptyset$;
- (7) the relations $(g(\lambda_j))^2 = gc_j g^{-1}$, where $j \in J$ and $g \in D^j$ such that $H \cap gt_j C_j g^{-1} \neq \emptyset$.

PROOF. Let X be the standard tree constructed in Lemma 5.1 on which G^* acts. Then $T = \{G\}, Y_0 = \emptyset, Y_1 = \{\lambda_i : i \in I\}$, and $Y_2 = \{\lambda_j : j \in J\}$. Let $v = G$. Then $D = D_v, D^i = D^{\lambda_i}, D^j = D^{\lambda_j}, D_i = D_{\lambda_i}$, and $D_j = D_{\lambda_j}$. Since $Y_0 = \emptyset$, therefore X does not contain the edges m of Theorem 4.4. Therefore H is generated by the generators of the forms (I)(1), (3), (4), and (5) of Theorem 4.4. Similarly H has the presentation of generating symbols (II)(1), (3), and (4), and relations (II)(1), (6), (7), (8), and (9) of Theorem 4.4.

This completes the proof. □

We have the following corollaries of Theorem 5.4.

COROLLARY 5.5. *Any subgroup H of G^* having trivial intersection with each conjugate of the base G is a quasi free.*

COROLLARY 5.6. *Any subgroup H of G^* having trivial intersection with the conjugates of B_i and C_j is the free product of a quasi free group and the intersection of H with certain conjugates of G .*

COROLLARY 5.7. *If H is a normal subgroup of G^* , $H \cap gGg^{-1}$ is a quasi free group, $H \cap gA_i g^{-1} = \{1\}$, and $H \cap gC_j g^{-1} = \{1\}$ for all $i \in I$, $j \in J$, and $g \in G^*$, then H is quasi free.*

COROLLARY 5.8. *If H has the property that $D^s = \emptyset$, for all $s \in I \cup J$, then H is a tree product of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$ with amalgamation subgroups $H \cap gA_i g^{-1}$, $H \cap gB_i g^{-1}$, where $i \in I$, $g \in D_i$, and $H \cap gC_j g^{-1}$, where $j \in J$ and $g \in D_j$.*

6. Subgroups of HNN groups. This section is an application of [Theorem 5.4](#). First we start by finding the structures of subgroups of HNN groups. For different methods of finding the structures of subgroups of HNN groups we refer the readers to [1].

By taking $J = \emptyset$ of [Theorem 5.4](#), we have the following theorem and corollaries.

THEOREM 6.1. *Let H be a subgroup of the HNN group,*

$$G^* = \langle G, t_i \mid \text{rel } G, t_i A_i t_i^{-1} = B_i, i \in I \rangle, \tag{6.1}$$

with base G and associated pairs (A_i, B_i) , $i \in I$ subgroups of G .

Let $\{D\}$, $\{D_i^d\}$, $\{D^i\}$, and $\{D_i\}$, $i \in I$, be a cress for $G^* \text{ mod } H$.

Then, we have the following:

(I) H is generated by the following:

- (1) the generators of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$;
- (2) the elements $gt_i(\overline{gt_i \overline{gt_i} k_i})^{-1}$, where $i \in I$ and $g \in D^i$.

(II) H has the presentation $\langle P \mid R \rangle$, where P is the set of generating symbols of the following forms:

- (1) the generating symbols of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$;
- (2) the symbols $g(\lambda_i)$, where $i \in I$ and $g \in D^i$.

(III) R is the set of relations of the following forms:

- (1) the relations $\text{rel}(H \cap dGd^{-1})$ of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$;
- (2) the relations $H \cap gA_i g^{-1} = H \cap gB_i g^{-1}$, where $i \in I$ and $g \in D_i$;
- (3) the relations $g(\lambda_i) \cdot (H \cap \overline{gt_i \overline{gt_i} A_i \overline{gt_i}^{-1} gt_i^{-1}}) \cdot (g(\lambda_i))^{-1} = H \cap gB_i g^{-1}$, where $i \in I$ and $g \in D^i$.

COROLLARY 6.2. *Any subgroup of G^* having trivial intersection with each conjugate of the base is free.*

COROLLARY 6.3. *Any subgroup H of G^* having trivial intersection with the conjugates of B_i is the free product of a free group and the intersection of H with certain conjugates of G .*

COROLLARY 6.4. *If H is a normal subgroup of G^* , $H \cap gGg^{-1}$ is a free group, and $H \cap gA_i g^{-1} = \{1\}$, for all $i \in I$ and $g \in G^*$, then H is free.*

COROLLARY 6.5. *If H has the property that $D^i = \emptyset$, for all $i \in I$, then H is a tree product of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$ with amalgamation subgroups $H \cap gA_i g^{-1}$ and $H \cap gB_i g^{-1}$, where $i \in I$ and $g \in D_i$.*

7. Subgroups of pure quasi-HNN groups. This section is an application of [Theorem 5.4](#). First, we start by finding the structures of subgroups of the pure quasi-HNN groups.

By taking $I = \emptyset$ in [Theorem 5.4](#), we have the following theorem and corollaries.

THEOREM 7.1. *Let H be a subgroup of the pure quasi-HNN group,*

$$G^* = \langle G, t_j \mid \text{rel } G, t_j C_j t_j^{-1} = C_j, t_j^2 = c_j, j \in J \rangle, \quad (7.1)$$

with base G and associated pairs (C_j, C_j) , $j \in J$ of subgroups of G .

Let $\{D\}$, $\{D_j^d\}$, $\{D^j\}$, and $\{D_j\}$, $j \in J$, be a cross for $G^* \text{ mod } H$.

Then, we have the following:

- (I) H is generated by the following:
- (1) the generators of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$;
 - (2) the elements $gt_j(\overline{gt_j} \overline{gt_j} k_j)^{-1}$ where $j \in J$, $g \in D^j$, and $H \cap gt_j C_j g^{-1} = \emptyset$;
 - (3) the elements $gt_j g^{-1}$, where $j \in J$, $g \in D^j$, and $H \cap gt_j C_j g^{-1} \neq \emptyset$.
- (II) H has the presentation $\langle P \mid R \rangle$, where P is the set of generating symbols of the following forms:
- (1) the generating symbols of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$;
 - (2) the symbols $g(\lambda_j)$, where $j \in J$ and $g \in D^j$.
- (III) R is the set of relations of the following forms:
- (1) the relations $\text{rel}(H \cap dGd^{-1})$ of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$;
 - (2) the relations $H \cap gC_j g^{-1} = H \cap gC_j g^{-1}$, where $j \in J$ and $g \in D_j$;
 - (3) the relations $g(\gamma_j) \cdot (H \cap \overline{gt_j} \overline{gt_j} C_j \overline{gt_j}^{-1} \overline{gt_j}^{-1}) \cdot (g(\gamma_j))^{-1} = H \cap gC_j g^{-1}$, where $j \in J$ and $g \in D^j$ such that $H \cap gt_j C_j g^{-1} = \emptyset$;
 - (4) the relations $g(\gamma_j) \cdot (H \cap gC_j g^{-1}) \cdot (g(\gamma_j))^{-1} = H \cap gC_j g^{-1}$, where $j \in J$ and $g \in D^j$ such that $H \cap gt_j C_j g^{-1} \neq \emptyset$;
 - (5) the relations $(g(\gamma_j))^2 = gc_j g^{-1}$, where $j \in J$ and $g \in D^j$ such that $H \cap gt_j C_j g^{-1} \neq \emptyset$.

We have the following corollaries of [Theorem 7.1](#).

COROLLARY 7.2. *Any subgroup of G^* having trivial intersection with each conjugate of the base G is quasi free.*

COROLLARY 7.3. *Any subgroup H of G^* having trivial intersection with the conjugates of C_j is the free product of a quasi free group and the intersection of H with certain conjugates of G .*

COROLLARY 7.4. *If H is a normal subgroup of G^* , $H \cap gGg^{-1}$ is a quasi free group, and $H \cap gC_j g^{-1} = \{1\}$ for all $j \in J$ and all $g \in G^*$, then H is quasi free.*

COROLLARY 7.5. *If H has the property that $D^j = \emptyset$, for all $j \in J$, then H is a tree product of the subgroups $H \cap dGd^{-1}$ of G , for all $d \in D$ with amalgamation subgroups $H \cap gC_j g^{-1}$ where $j \in J$ and $g \in D_j$.*

ACKNOWLEDGMENTS. The authors would like to thank Dr. A. H. M. Hoare for his help in this work. Also, the authors are grateful to the referee for his sincere evaluation and constructive comments which improved the paper considerably.

REFERENCES

- [1] D. E. Cohen, *Subgroups of HNN groups*, J. Austral. Math. Soc. **17** (1974), 394–405.
- [2] J. Fischer, *The subgroups of a tree product of groups*, Trans. Amer. Math. Soc. **210** (1975), 27–50.
- [3] M. I. Khanfar and R. M. S. Mahmud, *On quasi HNN groups*, to appear in J. Univ. Kuwait Sci.
- [4] W. Magnus, A. Karrass, and D. Solitar, *Combinatorial Group Theory*, 2nd revised ed., Dover Publications, New York, 1976.
- [5] R. M. S. Mahmud, *Presentation of groups acting on trees with inversions*, Proc. Roy. Soc. Edinburgh Sect. A **113** (1989), no. 3-4, 235–241.
- [6] ———, *The normal form theorem of groups acting on trees with inversions*, J. Univ. Kuwait Sci. **18** (1991), 7–16.
- [7] ———, *On a condition for a graph to be a tree*, JKAU: Sci. **7** (1995), 97–109.
- [8] ———, *The subgroup theorem for groups acting on trees*, Kuwait J. Sci. Engrg. **25** (1998), no. 1, 17–33.
- [9] J.-P. Serre, *Arbres, Amalgames, SL_2* , Astérisque, no. 46, Société Mathématique de France, Paris, 1977 (French), English translation, *Trees*, Springer-Verlag, Berlin, 1980.

R. M. S. MAHMOOD: AJMAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, ABU DHABI, UNITED ARAB EMIRATES

E-mail address: rasheedmsm@yahoo.com

M. I. KHANFAR: DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN

E-mail address: mohammadkhanfar2002@hotmail.com



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

