

RESEARCH

Open Access

Hybrid projection methods for a bifunction and relatively asymptotically nonexpansive mappings

Wenxin Wang^{1*} and Jianmin Song²

*Correspondence:

hdwangwx@sohu.com¹Department of Applied Mathematics and Physics, North China Electric Power University, Baoding, 071003, China
Full list of author information is available at the end of the article**Abstract**

The purpose of this paper is to investigate a bifunction equilibrium problem and a fixed point problem of relatively asymptotically nonexpansive mappings based on a generalized projection method. A weak convergence theorem for common solutions is established in a uniformly smooth and uniformly convex Banach space.

Keywords: bifunction; equilibrium problem; fixed point; generalized projection; relatively asymptotically nonexpansive mapping

1 Introduction and preliminaries

Let E be a real Banach space, E^* be the dual space of E , and C be a nonempty subset of E . Let F be a bifunction from $C \times C$ to \mathbb{R} , where \mathbb{R} denotes the set of real numbers. Recall the following equilibrium problem: Find $\bar{x} \in C$ such that

$$F(\bar{x}, y) \geq 0, \quad \forall y \in C. \quad (1.1)$$

From now on, we use $EP(F)$ to denote the solution set of equilibrium problem (1.1) and assume that F satisfies the following conditions:

(A1) $F(x, x) = 0, \forall x \in C$;

(A2) F is monotone, i.e., $F(x, y) + F(y, x) \leq 0, \forall x, y \in C$;

(A3)

$$\limsup_{t \downarrow 0} F(tz + (1-t)x, y) \leq F(x, y), \quad \forall x, y, z \in C;$$

(A4) for each $x \in C$, $y \mapsto F(x, y)$ is convex and weakly lower semi-continuous.

Let $U_E = \{x \in E : \|x\| = 1\}$ be the unit sphere of E . Then the Banach space E is said to be smooth iff

$$\lim_{t \rightarrow 0} \frac{\|x + ty\| - \|x\|}{t}$$

exists for each $x, y \in U_E$. It is also said to be uniformly smooth iff the above limit is attained uniformly for $x, y \in U_E$. It is well known that if E is uniformly smooth, then J is uniformly norm-to-norm continuous on each bounded subset of E . Recall that E is said to be uniformly convex iff $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$ for any two sequences $\{x_n\}$ and $\{y_n\}$ in E such that

$\|x_n\| = \|y_n\| = 1$ and $\lim_{n \rightarrow \infty} \|\frac{x_n + y_n}{2}\| = 1$. It is well known that E is uniformly smooth if and only if E^* is uniformly convex.

Recall that a Banach space E enjoys the Kadec-Klee property if for any sequence $\{x_n\} \subset E$, and $x \in E$ with $x_n \rightharpoonup x$, and $\|x_n\| \rightarrow \|x\|$, then $\|x_n - x\| \rightarrow 0$ as $n \rightarrow \infty$. For more details on the Kadec-Klee property, the readers can refer to [1] and the references therein. It is well known that if E is a uniformly convex Banach space, then E enjoys the Kadec-Klee property.

Let $T : C \rightarrow C$ be a mapping. From now on, we use $F(T)$ to denote the fixed point set of T . Recall that T is said to be closed if for any sequence $\{x_n\} \subset C$ such that $\lim_{n \rightarrow \infty} x_n = x_0$ and $\lim_{n \rightarrow \infty} Tx_n = y_0$, then $Tx_0 = y_0$. In this paper, we use \rightarrow and \rightharpoonup to denote the strong convergence and the weak convergence, respectively.

Recall that the normalized duality mapping J from E to 2^{E^*} is defined by

$$Jx = \{f^* \in E^* : \langle x, f^* \rangle = \|x\|^2 = \|f^*\|^2\},$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. Next, we assume that E is a smooth Banach space. Consider the functional defined by

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2, \quad \forall x, y \in E.$$

Observe that in a Hilbert space H the equality is reduced to $\phi(x, y) = \|x - y\|^2$, $x, y \in H$. As we all know, if C is a nonempty closed convex subset of a Hilbert space H and $P_C : H \rightarrow C$ is the metric projection of H onto C , then P_C is nonexpansive. This fact actually characterizes Hilbert spaces and, consequently, it is not available in more general Banach spaces. In this connection, Alber [2] recently introduced a generalized projection operator Π_C in a Banach space E which is an analogue of the metric projection P_C in Hilbert spaces. Recall that the generalized projection $\Pi_C : E \rightarrow C$ is a map that assigns to an arbitrary point $x \in E$ the minimum point of the functional $\phi(x, y)$, that is, $\Pi_C x = \bar{x}$, where \bar{x} is the solution to the minimization problem

$$\phi(\bar{x}, x) = \min_{y \in C} \phi(y, x).$$

Existence and uniqueness of the operator Π_C follow from the properties of the functional $\phi(x, y)$ and strict monotonicity of the mapping J . In Hilbert spaces, $\Pi_C = P_C$. It is obvious from the definition of a function ϕ that

$$(\|x\| - \|y\|)^2 \leq \phi(x, y) \leq (\|y\| + \|x\|)^2, \quad \forall x, y \in E. \tag{1.2}$$

Remark 1.1 If E is a reflexive, strictly convex, and smooth Banach space, then $\phi(x, y) = 0$ if and only if $x = y$; for more details, see [2] and the references therein.

Recall that a point p in C is said to be an asymptotic fixed point of a mapping T iff C contains a sequence $\{x_n\}$ which converges weakly to p so that $\lim_{n \rightarrow \infty} \|x_n - T^n x_n\| = 0$. The set of asymptotic fixed points of T will be denoted by $\tilde{F}(T)$.

Recall that a mapping T is said to be relatively nonexpansive iff

$$\tilde{F}(T) = F(T) \neq \emptyset, \quad \phi(p, Tx) \leq \phi(p, x), \quad \forall x \in C, \forall p \in F(T).$$

Recall that a mapping T is said to be relatively asymptotically nonexpansive iff

$$\tilde{F}(T) = F(T) \neq \emptyset, \quad \phi(p, T^n x) \leq (1 + \mu_n)\phi(p, x), \quad \forall x \in C, \forall p \in F(T), \forall n \geq 1,$$

where $\{\mu_n\} \subset [0, \infty)$ is a sequence such that $\mu_n \rightarrow 0$ as $n \rightarrow \infty$.

Remark 1.2 The class of relatively nonexpansive mappings was first considered in Butnariu *et al.* [3]. The class of relatively asymptotically nonexpansive mappings was first considered in Agarwal *et al.* [4] and the references therein.

Recently, many authors investigated fixed point problems of a (relatively) nonexpansive mapping based on hybrid projection methods; for more details, see [5–37] and the references therein. However, most of the results are on strong convergence. In this article, we investigate a bifunction equilibrium problem and a fixed point problem of relatively asymptotically nonexpansive mappings based on a generalized projection method. A weak convergence theorem for common solutions is established in a uniformly smooth and uniformly convex Banach space.

The following lemmas play an important role in this paper.

Lemma 1.3 [37, 38] *Let C be a closed convex subset of a uniformly smooth and uniformly convex Banach space E . Let F be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)-(A4). Let $r > 0$ and $x \in E$. Then there exists $z \in C$ such that $F(z, y) + \frac{1}{r}\langle y - z, Jz - Jx \rangle \geq 0, \forall y \in C$. Define a mapping $S_r : E \rightarrow C$ by $S_r x = \{z \in C : F(z, y) + \frac{1}{r}\langle y - z, Jz - Jx \rangle, \forall y \in C\}$. Then the following conclusions hold:*

- (a) S_r is single-valued;
- (b) S_r is a firmly nonexpansive-type mapping, i.e., for all $x, y \in E$,

$$\langle S_r x - S_r y, JS_r x - JS_r y \rangle \leq \langle S_r x - S_r y, Jx - Jy \rangle;$$

- (c) $F(S_r) = EP(F)$ is closed and convex;
- (d) S_r is relatively nonexpansive;
- (e) $\phi(q, S_r x) + \phi(S_r x, x) \leq \phi(q, x), \forall q \in F(S_r)$.

Lemma 1.4 [4] *Let E be a uniformly smooth and uniformly convex Banach space. Let C be a nonempty closed and convex subset of E . Let $T : C \rightarrow C$ be a relatively asymptotically nonexpansive mapping. Then $F(T)$ is a closed convex subset of C .*

Lemma 1.5 [2] *Let E be a reflexive, strictly convex, and smooth Banach space, let C be a nonempty, closed, and convex subset of E , and let $x \in E$. Then*

$$\phi(y, \Pi_C x) + \phi(\Pi_C x, x) \leq \phi(y, x), \quad \forall y \in C.$$

Lemma 1.6 [2] *Let C be a nonempty, closed, and convex subset of a smooth Banach space E , and let $x \in E$. Then $x_0 = \Pi_C x$ if and only if*

$$\langle x_0 - y, Jx - Jx_0 \rangle \geq 0, \quad \forall y \in C.$$

Lemma 1.7 [39] *Let E be a smooth and uniformly convex Banach space, and let $r > 0$. Then there exists a strictly increasing, continuous, and convex function $g : [0, 2r] \rightarrow \mathbb{R}$ such that $g(0) = 0$ and*

$$\|tx + (1-t)y\|^2 \leq t\|x\|^2 + (1-t)\|y\|^2 - t(1-t)g(\|x-y\|)$$

for all $x, y \in B_r = \{x \in E : \|x\| \leq r\}$ and $t \in [0, 1]$.

Lemma 1.8 [40] *Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be three nonnegative sequences satisfying the following condition:*

$$a_{n+1} \leq (1 + b_n)a_n + c_n, \quad \forall n \geq n_0,$$

where n_0 is some nonnegative integer. If $\sum_{n=1}^{\infty} b_n < \infty$ and $\sum_{n=1}^{\infty} c_n < \infty$, then the limit of the sequence $\{a_n\}$ exists. If, in addition, there exists a subsequence $\{\alpha_{n_i}\} \subset \{a_n\}$ such that $\alpha_{n_i} \rightarrow 0$, then $a_n \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 1.9 [41] *Let E be a smooth and uniformly convex Banach space, and let $r > 0$. Then there exists a strictly increasing, continuous, and convex function $g : [0, 2r] \rightarrow \mathbb{R}$ such that $g(0) = 0$ and $g(\|x-y\|) \leq \phi(x, y)$ for all $x, y \in B_r$.*

2 Main results

Theorem 2.1 *Let E be a uniformly smooth and uniformly convex Banach space, and let C be a nonempty closed and convex subset of E . Let F be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)-(A4). Let $T : C \rightarrow C$ be a relatively asymptotically nonexpansive mapping with the sequence $\{\mu_{n,1}\}$, and let $S : C \rightarrow C$ be a relatively asymptotically nonexpansive mapping with the sequence $\{\mu_{n,2}\}$. Assume that $\Phi := F(T) \cap F(S) \cap EP(F)$ is nonempty. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{cases} y_0 \in E \text{ chosen arbitrarily,} \\ x_n \in C \text{ such that } F(x_n, x) + \frac{1}{r_n} \langle x - x_n, Jx_n - Jy_n \rangle \geq 0, \quad \forall x \in C, \\ y_{n+1} = J^{-1}(\alpha_n Jx_n + \beta_n J T^n x_n + \gamma_n J S^n x_n), \quad \forall n \geq 0, \end{cases}$$

where $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$ are real sequences in $[0, 1]$ and $\{r_n\}$ is a real number sequence in $[r, \infty)$, where $r > 0$ is some real number. Assume that J is weakly sequentially continuous and the following restrictions hold:

- (i) $\alpha_n + \beta_n + \gamma_n = 1$;
- (ii) $\sum_{n=1}^{\infty} \mu_n < \infty$;
- (iii) $\liminf_{n \rightarrow \infty} \alpha_n \beta_n > 0$, $\liminf_{n \rightarrow \infty} \alpha_n \gamma_n > 0$.

Then the sequence $\{x_n\}$ converges weakly to $\bar{x} \in \Phi$, where $\bar{x} = \lim_{n \rightarrow \infty} \Pi_{\Phi} x_n$.

Proof Set $\mu_n = \max\{\mu_{n,1}, \mu_{n,2}\}$. Fixing $p \in \Phi$, we find that

$$\begin{aligned} \phi(p, x_{n+1}) &= \phi(p, S_{r_{n+1}} y_{n+1}) \\ &\leq \phi(p, y_{n+1}) \end{aligned}$$

$$\begin{aligned}
 &= \|p\|^2 - 2\langle p, \alpha_n Jx_n + \beta_n JT^n x_n + \gamma_n JS^n x_n \rangle \\
 &\quad + \|\alpha_n Jx_n + \beta_n JT^n x_n + \gamma_n JS^n x_n\|^2
 \end{aligned} \tag{2.1}$$

$$\begin{aligned}
 &\leq \|p\|^2 - 2\alpha_n \langle p, Jx_n \rangle - 2\beta_n \langle p, JT^n x_n \rangle - 2\gamma_n \langle p, JS^n x_n \rangle \\
 &\quad + \alpha_n \|x_n\|^2 + \beta_n \|T^n x_n\|^2 + \gamma_n \|S^n x_n\|^2 \\
 &= \alpha_n \phi(p, x_n) + \beta_n \phi(p, T^n x_n) + \gamma_n \phi(p, S^n x_n) \\
 &\leq \phi(p, x_n) + \beta_n \mu_n \phi(p, x_n) + \gamma_n \mu_n \phi(p, x_n) \\
 &\leq (1 + \mu_n) \phi(p, x_n).
 \end{aligned} \tag{2.2}$$

In view of Lemma 1.8, we obtain that $\lim_{n \rightarrow \infty} \phi(p, x_n)$ exists. This implies that the sequence $\{x_n\}$ is bounded. In the light of Lemma 1.7, we find that

$$\begin{aligned}
 \phi(p, x_{n+1}) &= \phi(p, S_{r_{n+1}} y_{n+1}) \\
 &\leq \|p\|^2 - 2\langle p, \alpha_n Jx_n + \beta_n JT^n x_n + \gamma_n JS^n x_n \rangle \\
 &\quad + \|\alpha_n Jx_n + \beta_n JT^n x_n + \gamma_n JS^n x_n\|^2 \\
 &\leq \|p\|^2 - 2\alpha_n \langle p, Jx_n \rangle - 2\beta_n \langle p, JT^n x_n \rangle - 2\gamma_n \langle p, JS^n x_n \rangle \\
 &\quad + \alpha_n \|x_n\|^2 + \beta_n \|T^n x_n\|^2 + \gamma_n \|S^n x_n\|^2 - \alpha_n \beta_n g(\|JT^n x_n - Jx_n\|) \\
 &\leq \phi(p, x_n) + \beta_n \mu_n \phi(p, x_n) + \gamma_n \mu_n \phi(p, x_n) - \alpha_n \beta_n g(\|JT^n x_n - Jx_n\|) \\
 &\leq (1 + \mu_n) \phi(p, x_n) - \alpha_n \beta_n g(\|JT^n x_n - Jx_n\|).
 \end{aligned}$$

It follows that

$$\alpha_n \beta_n g(\|JT^n x_n - Jx_n\|) \leq (1 + \mu_n) \phi(p, x_n) - \phi(p, x_{n+1}).$$

This finds from the restrictions (ii) and (iii) that

$$\lim_{n \rightarrow \infty} g(\|JT^n x_n - Jx_n\|) = 0.$$

This implies that

$$\lim_{n \rightarrow \infty} \|JT^n x_n - Jx_n\| = 0.$$

Since J^{-1} is uniformly norm-to-norm continuous on bounded sets, we find that

$$\lim_{n \rightarrow \infty} \|T^n x_n - x_n\| = 0.$$

In the same way, we find that

$$\lim_{n \rightarrow \infty} \|S^n x_n - x_n\| = 0.$$

Since $\{x_n\}$ is bounded, we see that there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $\{x_{n_i}\}$ converges weakly to $p \in C$. It follows that $p \in F(T) \cap F(S)$. Next, we prove that $p \in EP(F)$.

Let $r = \sup_{n \geq 1} \{\|x_n\|, \|y_n\|\}$. In view of Lemma 1.9, we find that there exists a continuous, strictly increasing and convex function h with $h(0) = 0$ such that

$$h(x, y) \leq \phi(x, y), \quad \forall x, y \in B_r.$$

It follows from (2.1) that

$$\begin{aligned} h(\|x_n - y_n\|) &\leq \phi(x_n, y_n) \\ &\leq \phi(p, y_n) - \phi(p, x_n) \\ &\leq \phi(p, x_{n-1}) - \phi(p, x_n) + \mu_{n-1}\phi(p, x_{n-1}). \end{aligned}$$

This implies that

$$\lim_{n \rightarrow \infty} h(\|x_n - y_n\|) = 0.$$

It follows from the property of h that

$$\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0.$$

Since J is uniformly norm-to-norm continuous on bounded sets, one has

$$\lim_{n \rightarrow \infty} \|Jx_n - Jy_n\| = 0.$$

Since $\{r_n\}$ is a real number sequence in $[r, \infty)$, where $r > 0$ is some real number, one finds that

$$\lim_{n \rightarrow \infty} \frac{\|Jx_n - Jy_n\|}{r_n} = 0.$$

Notice that $x_n = S_{r_n}y_n$, one sees that

$$F(x_n, x) + \frac{1}{r_n} \langle x - x_n, Jx_n - Jy_n \rangle \geq 0, \quad \forall x \in C.$$

By replacing n by n_i , one finds from (A2) that

$$\begin{aligned} \|x - x_{n_i}\| \frac{\|Jx_{n_i} - Jy_{n_i}\|}{r_{n_i}} &\geq \frac{1}{r_{n_i}} \langle x - x_{n_i}, Jx_{n_i} - Jy_{n_i} \rangle \\ &\geq F(x, x_{n_i}). \end{aligned}$$

Letting $i \rightarrow \infty$ in the above inequality, one obtains from (A4) that

$$F(x, p) \leq 0, \quad \forall x \in C.$$

For $0 < t < 1$ and $y \in C$, define $x_t = tx + (1-t)p$. It follows that $x_t \in C$, which yields that $F(x_t, p) \leq 0$. It follows from (A1) and (A4) that

$$0 = F(x_t, x_t) \leq tF(x_t, x) + (1-t)F(x_t, p) \leq tF(x_t, x).$$

That is,

$$F(x_t, x) \geq 0.$$

Letting $t \downarrow 0$, we obtain from (A3) that $F(p, x) \geq 0, \forall x \in C$. This implies that $p \in EP(F)$. This completes the proof that $p \in F(T) \cap F(S) \cap EP(F)$. Define $z_n = \Pi_{F(T) \cap F(S) \cap EP(F)} x_n$. It follows from (2.1) that

$$\phi(z_n, x_{n+1}) \leq (1 + \mu_n)\phi(z_n, x_n). \tag{2.3}$$

This in turn implies from Lemma 1.5 that

$$\begin{aligned} \phi(z_{n+1}, x_{n+1}) &= \phi(\Pi_{F(T) \cap F(S) \cap EP(F)} x_{n+1}, x_{n+1}) \\ &\leq \phi(z_n, x_{n+1}) - \phi(z_n, \Pi_{F(T) \cap F(S) \cap EP(F)} x_{n+1}) \\ &\leq \phi(z_n, x_{n+1}) - \phi(z_n, z_{n+1}) \\ &\leq \phi(z_n, x_{n+1}). \end{aligned}$$

It follows from (2.3) that

$$\phi(z_{n+1}, x_{n+1}) \leq (1 + \mu_n)\phi(z_n, x_n).$$

This finds from Lemma 1.8 that the sequence $\{\phi(z_n, x_n)\}$ is a convergence sequence. It follows from (2.1) that

$$\phi(p, x_{n+m}) \leq \phi(p, x_n) + L \left(\sum_{i=1}^m \mu_{n+m-i} \right), \tag{2.4}$$

where $L = \sup_{n \geq 1} \phi(p, x_n)$. Since $z_n \in F(T) \cap F(S) \cap EP(F)$, we find that

$$\phi(z_n, x_{n+m}) \leq \phi(z_n, x_n) + M \left(\sum_{i=1}^m \mu_{n+m-i} \right),$$

where $M = \sup_{n \geq 1} \phi(z_n, x_n)$. Since $z_{n+m} = \Pi_{F(T) \cap F(S) \cap EP(F)} x_{n+m}$, we find from Lemma 1.5 that

$$\phi(z_n, z_{n+m}) + \phi(z_{n+m}, x_{n+m}) \leq \phi(z_n, x_{n+m}) \leq \phi(z_n, x_n) + M \left(\sum_{i=1}^m \mu_{n+m-i} \right).$$

It follows that

$$\phi(z_n, z_{n+m}) \leq \phi(z_n, x_n) - \phi(z_{n+m}, x_{n+m}) + M \left(\sum_{i=1}^m \mu_{n+m-i} \right).$$

In view of Lemma 1.9, we find that there exists a continuous, strictly increasing, and convex function g with

$$g(\|z_n - z_m\|) \leq \phi(z_n, z_m) \leq \phi(z_n, x_n) - \phi(z_{n+m}, x_{n+m}) + M \left(\sum_{i=1}^m \mu_{n+m-i} \right).$$

This shows that $\{z_n\}$ is a Cauchy sequence. Since $F(T) \cap F(S) \cap EP(F)$ is closed, one sees that $\{z_n\}$ converges strongly to $z \in F(T) \cap F(S) \cap EP(F)$. Since $p \in F(T) \cap F(S) \cap EP(F)$, we find from Lemma 1.6 that

$$\langle z_{n_k} - p, Jx_{n_k} - Jz_{n_k} \rangle \geq 0.$$

Notice that J is weakly sequentially continuous. Letting $k \rightarrow \infty$, we find that $\langle z - p, Jp - Jz \rangle \geq 0$. It follows from the monotonicity of J that $\langle z - p, Jp - Jz \rangle \leq 0$. Since the space is uniformly convex, we find that $z = p$. This completes the proof. \square

Remark 2.2 Theorem 2.1 improves Theorem 2.5 in Qin *et al.* [36] on the mappings from the class of relatively nonexpansive mappings to the class of relatively asymptotically nonexpansive mappings.

If $T = S$, then Theorem 2.1 is reduced to the following.

Corollary 2.3 *Let E be a uniformly smooth and uniformly convex Banach space, and let C be a nonempty closed and convex subset of E . Let F be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)-(A4). Let $T : C \rightarrow C$ be a relatively asymptotically nonexpansive mapping with the sequence $\{\mu_n\}$. Assume that $\Phi := F(T) \cap EP(F)$ is nonempty. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{cases} y_0 \in E \text{ chosen arbitrarily,} \\ x_n \in C \text{ such that } F(x_n, x) + \frac{1}{r_n} \langle x - x_n, Jx_n - Jy_n \rangle \geq 0, \quad \forall x \in C, \\ y_{n+1} = J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JT^n x_n), \quad \forall n \geq 0, \end{cases}$$

where $\{\alpha_n\}$ is a real sequence in $[0, 1]$ and $\{r_n\}$ is a real number sequence in $[r, \infty)$, where $r > 0$ is some real number. Assume that J is weakly sequentially continuous and the following restrictions hold:

- (i) $\sum_{n=1}^{\infty} \mu_n < \infty$;
- (ii) $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$.

Then the sequence $\{x_n\}$ converges weakly to $\bar{x} \in \Phi$, where $\bar{x} = \lim_{n \rightarrow \infty} \Pi_{\Phi} x_n$.

Remark 2.4 Corollary 2.3 is an improvement of Theorem 4.1 in Zembayashi and Takahashi [37] on the nonlinear mapping.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Both authors contributed equally and read and approved the final manuscript.

Author details

¹Department of Applied Mathematics and Physics, North China Electric Power University, Baoding, 071003, China.

²Department of Mathematics and Sciences, Shijiazhuang University of Economics, Shijiazhuang, Hebei 050031, China.

Acknowledgements

This work is supported by the Fundamental Research Funds for the Central Universities (Grant No.: 9161013002). The authors are grateful to the editor and the anonymous reviewers' suggestions which improved the contents of the article.

Received: 8 July 2013 Accepted: 4 September 2013 Published: 09 Nov 2013

References

1. Cioranescu, I: *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*. Kluwer Academic, Dordrecht (1990)
2. Alber, YI: Metric and generalized projection operators in Banach spaces: properties and applications. In: Kartsatos, AG (ed.) *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type*. Dekker, New York (1996)
3. Butnariu, D, Reich, S, Zaslavski, AJ: Weak convergence of orbits of nonlinear operators in reflexive Banach spaces. *Numer. Funct. Anal. Optim.* **24**, 489-508 (2003)
4. Agarwal, RP, Cho, YJ, Qin, X: Generalized projection algorithms for nonlinear operators. *Numer. Funct. Anal. Optim.* **28**, 1197-1215 (2007)
5. Qin, X, Agarwal, RP: Shrinking projection methods for a pair of asymptotically quasi- ϕ -nonexpansive mappings. *Numer. Funct. Anal. Optim.* **31**, 1072-1089 (2010)
6. Ye, J, Huang, J: Strong convergence theorems for fixed point problems and generalized equilibrium problems of three relatively quasi-nonexpansive mappings in Banach spaces. *J. Math. Comput. Sci.* **1**, 1-18 (2011)
7. Kang, SM, Cho, SY, Liu, Z: Convergence of iterative sequences for generalized equilibrium problems involving inverse-strongly monotone mappings. *J. Inequal. Appl.* **2010**, Article ID 827082 (2010)
8. Shen, J, Pang, LP: An approximate bundle method for solving variational inequalities. *Commun. Optim. Theory* **1**, 1-18 (2012)
9. Zegeye, H, Shahzad, N: Strong convergence theorem for a common point of solution of variational inequality and fixed point problem. *Adv. Fixed Point Theory* **2**, 374-397 (2012)
10. Qin, X, Chang, SS, Cho, YJ: Iterative methods for generalized equilibrium problems and fixed point problems with applications. *Nonlinear Anal.* **11**, 2963-2972 (2010)
11. Noor, MA, Noor, KI, Waseem, M: Decomposition method for solving system of linear equations. *Eng. Math. Lett.* **2**, 34-41 (2012)
12. Luo, H, Wang, Y: Iterative approximation for the common solutions of a infinite variational inequality system for inverse-strongly accretive mappings. *J. Math. Comput. Sci.* **2**, 1660-1670 (2012)
13. Qin, X, Cho, SY, Kang, SM: On hybrid projection methods for asymptotically quasi- ϕ -nonexpansive mappings. *Appl. Math. Comput.* **215**, 3874-3883 (2010)
14. Chang, SS, Wang, L, Tang, YK, Zao, YH, Wang, B: Strong convergence theorems of quasi- ϕ -asymptotically nonexpansive semi-groups in Banach spaces. *Fixed Point Theory Appl.* **2012**, Article ID 15 (2012)
15. Chang, SS, Kim, JK, Wang, L: Total quasi- ϕ -asymptotically nonexpansive semigroups and strong convergence theorems in Banach spaces. *Fixed Point Theory Appl.* **2012**, Article ID 153 (2012)
16. Cho, SY, Li, W, Kang, SM: Convergence analysis of an iterative algorithm for monotone operators. *J. Inequal. Appl.* **2013**, Article ID 199 (2013)
17. Zuo, P, Chang, SS, Liu, M: On a hybrid algorithm for a family of total quasi- ϕ -asymptotically nonexpansive mappings in Banach spaces. *Fixed Point Theory Appl.* **2012**, Article ID 70 (2012)
18. Kim, JK: Strong convergence theorems by hybrid projection methods for equilibrium problems and fixed point problems of the asymptotically quasi- ϕ -nonexpansive mappings. *Fixed Point Theory Appl.* **2011**, Article ID 10 (2011)
19. Kim, JK: Hybrid projection algorithms for generalized equilibrium problems and strictly pseudocontractive mappings. *J. Inequal. Appl.* **2010**, Article ID 312602 (2010)
20. Cho, SY, Kang, SM: Approximation of fixed points of pseudocontraction semigroups based on a viscosity iterative process. *Appl. Math. Lett.* **24**, 224-228 (2011)
21. Ali, B, Minjibir, M: Convergence of a hybrid iterative method for finite families of generalized quasi- ϕ -asymptotically nonexpansive mappings. *Fixed Point Theory Appl.* **2012**, Article ID 121 (2012)
22. Qin, X, Cho, SY, Kang, SM: Strong convergence of shrinking projection methods for quasi- ϕ -nonexpansive mappings and equilibrium problems. *J. Comput. Appl. Math.* **234**, 750-760 (2010)
23. Li, Y, Liu, H: Strong convergence theorems for modifying Halpern-Mann iterations for a quasi- ϕ -asymptotically nonexpansive multi-valued mapping in Banach spaces. *Fixed Point Theory Appl.* **2013**, Article ID 132 (2013)
24. Zhang, J, Su, Y, Chen, Q: Simple projection algorithm for a countable family of weak relatively nonexpansive mappings and applications. *Fixed Point Theory Appl.* **2012**, Article ID 205 (2012)
25. Qin, X, Agarwal, RP, Cho, SY, Kang, SM: Convergence of algorithms for fixed points of generalized asymptotically quasi- ϕ -nonexpansive mappings with applications. *Fixed Point Theory Appl.* **2012**, Article ID 58 (2012)
26. Zhou, H, Gao, X: A strong convergence theorem for a family of quasi- ϕ -nonexpansive mappings in a Banach space. *Fixed Point Theory Appl.* **2009**, Article ID 351265 (2009)
27. Qin, X, Cho, SY, Kang, SM: An extragradient-type method for generalized equilibrium problems involving strictly pseudocontractive mappings. *J. Glob. Optim.* **49**, 679-693 (2011)
28. Kohsaka, F, Takahashi, W: Block iterative methods for a finite family of relatively nonexpansive mappings in Banach spaces. *Fixed Point Theory Appl.* **2007**, Article ID 021972 (2007)
29. Hao, Y: Some results on a modified Mann iterative scheme in a reflexive Banach space. *Fixed Point Theory Appl.* **2013**, Article ID 227 (2013)
30. Qin, X, Cho, YJ, Kang, SM, Zhou, H: Convergence of a modified Halpern-type iteration algorithm for quasi- ϕ -nonexpansive mappings. *Appl. Math. Lett.* **22**, 1051-1055 (2009)
31. Wang, X, Hu, C, Guan, J: Strong convergence theorems for fixed point problems of infinite family of asymptotically quasi- ϕ -nonexpansive mappings and a system of equilibrium problems. *Fixed Point Theory Appl.* **2013**, Article ID 221 (2013)
32. Kim, JK: Some results on generalized equilibrium problems involving strictly pseudocontractive mappings. *Acta Math. Sci.* **31**, 2041-2057 (2011)
33. Su, Y, Qin, X: Monotone CQ iteration processes for nonexpansive semigroups and maximal monotone operators. *Nonlinear Anal.* **68**, 3657-3664 (2008)
34. Cho, SY, Kang, SM: Approximation of common solutions of variational inequalities via strict pseudocontractions. *Acta Math. Sci.* **32**, 1607-1618 (2012)
35. Wangkeeree, R: A new hybrid approximation algorithm based on the shrinking projection method for two asymptotically quasi- ϕ -nonexpansive mappings. *J. Comput. Anal. Appl.* **14**, 298-313 (2012)

36. Qin, X, Cho, YJ, Kang, SM: Convergence theorems of common elements for equilibrium problems and fixed point problems in Banach spaces. *J. Comput. Appl. Math.* **225**, 20-30 (2009)
37. Zembayashi, WK, Takahashi, W: Strong and weak convergence theorems for equilibrium problems and relatively nonexpansive mappings in Banach spaces. *Nonlinear Anal.* **70**, 45-57 (2009)
38. Blum, E, Oettli, W: From optimization and variational inequalities to equilibrium problems. *Math. Stud.* **63**, 123-145 (1994)
39. Zălinescu, C: On uniformly convex functions. *J. Math. Anal. Appl.* **95**, 344-374 (1983)
40. Chang, SS, Cho, YJ, Zhou, H: *Iterative Methods for Nonlinear Operator Equations in Banach Spaces*. Nova Science Publishers, Huntington (2002)
41. Kamimura, S, Takahashi, W: Strong convergence of a proximal-type algorithm in a Banach space. *SIAM J. Optim.* **13**, 938-945 (2002)

10.1186/1687-1812-2013-294

Cite this article as: Wang and Song: Hybrid projection methods for a bifunction and relatively asymptotically nonexpansive mappings. *Fixed Point Theory and Applications* 2013, **2013**:294

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
