

## RESEARCH

## Open Access

# A new application of quasi power increasing sequences. I

Hüseyin Bor\*

\*Correspondence:  
hbor33@gmail.com  
P.O. Box 121, Bahçelievler, Ankara,  
TR-06502, Turkey**Abstract**

In (Rocky Mt. J. Math. 38:801-807, 2008), we proved a theorem dealing with an application of quasi- $\sigma$ -power increasing sequences. In the present paper, we prove that theorem under less and more weaker conditions. This theorem also includes some new and known results.

**MSC:** 40D15; 40F05; 40G99; 46A45

**Keywords:** Hölder's inequality; sequence spaces; absolute summability; increasing sequences

## 1 Introduction

A positive sequence  $(b_n)$  is said to be almost increasing if there exists a positive increasing sequence  $c_n$  and two positive constants  $A$  and  $B$  such that  $Ac_n \leq b_n \leq Bc_n$  (see [1]). A sequence  $(\lambda_n)$  is said to be of bounded variation, denoted by  $(\lambda_n) \in \mathcal{BV}$ , if  $\sum_{n=1}^{\infty} |\Delta \lambda_n| = \sum_{n=1}^{\infty} |\lambda_n - \lambda_{n+1}| < \infty$ . A positive sequence  $X = (X_n)$  is said to be a quasi- $\sigma$ -power increasing sequence if there exists a constant  $K = K(\sigma, X) \geq 1$  such that  $Kn^\sigma X_n \geq m^\sigma X_m$  holds for all  $n \geq m \geq 1$ . It should be noted that every almost increasing sequence is a quasi- $\sigma$ -power increasing sequence for any nonnegative  $\sigma$ , but the converse may not be true as can be seen by taking an example, say  $X_n = n^{-\sigma}$  for  $\sigma > 0$  (see [2]). Let  $(\varphi_n)$  be a sequence of complex numbers and let  $\sum a_n$  be a given infinite series with partial sums  $(s_n)$ . We denote by  $z_n^\alpha$  and  $t_n^\alpha$  the  $n$ th Cesàro means of order  $\alpha$ , with  $\alpha > -1$ , of the sequences  $(s_n)$  and  $(na_n)$ , respectively, that is,

$$z_n^\alpha = \frac{1}{A_n^\alpha} \sum_{v=0}^n A_{n-v}^{\alpha-1} s_v, \quad (1)$$

$$t_n^\alpha = \frac{1}{A_n^\alpha} \sum_{v=1}^n A_{n-v}^{\alpha-1} v a_v, \quad (2)$$

where

$$A_n^\alpha = \binom{n+\alpha}{n} = \frac{(\alpha+1)(\alpha+2)\cdots(\alpha+n)}{n!} = O(n^\alpha), \quad A_{-n}^\alpha = 0 \quad \text{for } n > 0. \quad (3)$$

The series  $\sum a_n$  is said to be summable  $\varphi - |C, \alpha|_k$ ,  $k \geq 1$  and  $\alpha > -1$ , if (see [3])

$$\sum_{n=1}^{\infty} |\varphi_n (z_n^\alpha - z_{n-1}^\alpha)|^k = \sum_{n=1}^{\infty} n^{-k} |\varphi_n t_n^\alpha|^k < \infty. \quad (4)$$

In the special case, if we take  $\varphi_n = n^{1-\frac{1}{k}}$ , then  $\varphi - |C, \alpha|_k$  summability is the same as  $|C, \alpha|_k$  summability (see [4]). Also, if we take  $\varphi_n = n^{\delta+1-\frac{1}{k}}$ , then  $\varphi - |C, \alpha|_k$  summability reduces to  $|C, \alpha; \delta|_k$  summability (see [5]).

### 2 Known result

In [6], we have proved the following theorem.

**Theorem A** *Let  $(\lambda_n) \in \mathcal{BV}$  and let  $(X_n)$  be a quasi- $\sigma$ -power increasing sequence for some  $\sigma$  ( $0 < \sigma < 1$ ). Suppose also that there exist sequences  $(\beta_n)$  and  $(\lambda_n)$  such that*

$$|\Delta\lambda_n| \leq \beta_n, \tag{5}$$

$$\beta_n \rightarrow 0 \text{ as } n \rightarrow \infty, \tag{6}$$

$$\sum_{n=1}^{\infty} n|\Delta\beta_n|X_n < \infty, \tag{7}$$

$$|\lambda_n|X_n = O(1) \text{ as } n \rightarrow \infty. \tag{8}$$

*If there exists an  $\epsilon > 0$  such that the sequence  $(n^{\epsilon-k}|\varphi_n|^k)$  is non-increasing and if the sequence  $(w_n^\alpha)$  defined by (see [7])*

$$w_n^\alpha = \begin{cases} |t_n^\alpha|, & \alpha = 1, \\ \max_{1 \leq v \leq n} |t_v^\alpha|, & 0 < \alpha < 1, \end{cases} \tag{9}$$

*satisfies the condition*

$$\sum_{n=1}^m \frac{(|\varphi_n|w_n^\alpha)^k}{n^k} = O(X_m) \text{ as } m \rightarrow \infty, \tag{10}$$

*then the series  $\sum a_n\lambda_n$  is summable  $\varphi - |C, \alpha|_k, k \geq 1, 0 < \alpha \leq 1$  and  $k\alpha + \epsilon > 1$ .*

It should be remarked that we have added the condition ‘ $(\lambda_n) \in \mathcal{BV}$ ’ in the statement of Theorem A because it is necessary.

### 3 The main result

The aim of this paper is to prove Theorem A under less and weaker conditions. Now, we will prove the following theorem.

**Theorem** *Let  $(X_n)$  be a quasi- $\sigma$ -power increasing sequence for some  $\sigma$  ( $0 < \sigma < 1$ ). If there exists an  $\epsilon > 0$  such that the sequence  $(n^{\epsilon-k}|\varphi_n|^k)$  is non-increasing and if the conditions from (5) to (8) are satisfied and if the condition*

$$\sum_{n=1}^m \frac{(|\varphi_n|w_n^\alpha)^k}{n^k X_n^{k-1}} = O(X_m) \text{ as } m \rightarrow \infty \tag{11}$$

*is satisfied, then the series  $\sum a_n\lambda_n$  is summable  $\varphi - |C, \alpha|_k, k \geq 1, 0 < \alpha \leq 1$ , and  $k(\alpha - 1) + \epsilon > 1$ .*

**Remark** It should be noted that condition (11) is the same as condition (10) when  $k = 1$ . When  $k > 1$ , condition (11) is weaker than condition (10), but the converse is not true. As in [8] we can show that if (10) is satisfied, then we get that

$$\sum_{n=1}^m \frac{(|\varphi_n|w_n^\alpha)^k}{n^k X_n^{k-1}} = O\left(\frac{1}{X_1^{k-1}}\right) \sum_{n=1}^m \frac{(|\varphi_n|w_n^\alpha)^k}{n^k} = O(X_m).$$

If (11) is satisfied, then for  $k > 1$  we obtain that

$$\sum_{n=1}^m \frac{(|\varphi_n|w_n^\alpha)^k}{n^k} = \sum_{n=1}^m X_n^{k-1} \frac{(|\varphi_n|w_n^\alpha)^k}{n^k X_n^{k-1}} = O(X_m^{k-1}) \sum_{n=1}^m \frac{(|\varphi_n|w_n^\alpha)^k}{n^k X_n^{k-1}} = O(X_m^k) \neq O(X_m).$$

Also, it should be noted that the condition ‘ $(\lambda_n) \in \mathcal{BV}$ ’ has been removed.

We need the following lemmas for the proof of our theorem.

**Lemma 1** [9] *If  $0 < \alpha \leq 1$  and  $1 \leq v \leq n$ , then*

$$\left| \sum_{p=0}^v A_{n-p}^{\alpha-1} a_p \right| \leq \max_{1 \leq m \leq v} \left| \sum_{p=0}^m A_{m-p}^{\alpha-1} a_p \right|. \tag{12}$$

**Lemma 2** [2] *Under the conditions on  $(X_n)$ ,  $(\beta_n)$  and  $(\lambda_n)$ , as expressed in the statement of the theorem, we have the following:*

$$n\beta_n X_n = O(1) \quad \text{as } n \rightarrow \infty, \tag{13}$$

$$\sum_{n=1}^{\infty} \beta_n X_n < \infty. \tag{14}$$

#### 4 Proof of the theorem

Let  $(T_n^\alpha)$  be the  $n$ th  $(C, \alpha)$ , with  $0 < \alpha \leq 1$ , mean of the sequence  $(na_n \lambda_n)$ . Then, by (2), we have

$$T_n^\alpha = \frac{1}{A_n^\alpha} \sum_{v=1}^n A_{n-v}^{\alpha-1} v a_v \lambda_v. \tag{15}$$

First, applying Abel’s transformation and then using Lemma 1, we get that

$$\begin{aligned} T_n^\alpha &= \frac{1}{A_n^\alpha} \sum_{v=1}^{n-1} \Delta \lambda_v \sum_{p=1}^v A_{n-p}^{\alpha-1} p a_p + \frac{\lambda_n}{A_n^\alpha} \sum_{v=1}^n A_{n-v}^{\alpha-1} v a_v, \\ |T_n^\alpha| &\leq \frac{1}{A_n^\alpha} \sum_{v=1}^{n-1} |\Delta \lambda_v| \left| \sum_{p=1}^v A_{n-p}^{\alpha-1} p a_p \right| + \frac{|\lambda_n|}{A_n^\alpha} \left| \sum_{v=1}^n A_{n-v}^{\alpha-1} v a_v \right| \\ &\leq \frac{1}{A_n^\alpha} \sum_{v=1}^{n-1} A_v^\alpha w_v^\alpha |\Delta \lambda_v| + |\lambda_n| w_n^\alpha \\ &= T_{n,1}^\alpha + T_{n,2}^\alpha. \end{aligned}$$

To complete the proof of the theorem, by Minkowski's inequality, it is sufficient to show that

$$\sum_{n=1}^{\infty} n^{-k} |\varphi_n T_{n,r}^\alpha|^k < \infty \quad \text{for } r = 1, 2.$$

Now, when  $k > 1$ , applying Hölder's inequality with indices  $k$  and  $k'$ , where  $\frac{1}{k} + \frac{1}{k'} = 1$ , we get that

$$\begin{aligned} \sum_{n=2}^{m+1} n^{-k} |\varphi_n T_{n,1}^\alpha|^k &\leq \sum_{n=2}^{m+1} n^{-k} (A_n^\alpha)^{-k} |\varphi_n|^k \left\{ \sum_{v=1}^{n-1} A_v^\alpha w_v^\alpha |\Delta \lambda_v| \right\}^k \\ &\leq \sum_{n=2}^{m+1} n^{-k} n^{-\alpha k} |\varphi_n|^k \sum_{v=1}^{n-1} v^{\alpha k} (w_v^\alpha)^k |\Delta \lambda_v|^k \times \left\{ \sum_{v=1}^{n-1} 1 \right\}^{k-1} \\ &= O(1) \sum_{v=1}^m v^{\alpha k} (w_v^\alpha)^k (\beta_v)^k \sum_{n=v+1}^{m+1} \frac{n^{\epsilon-k} |\varphi_n|^k}{n^{k(\alpha-1)+\epsilon+1}} \\ &= O(1) \sum_{v=1}^m v^{\alpha k} (w_v^\alpha)^k \beta_v (\beta_v)^{k-1} v^{\epsilon-k} |\varphi_v|^k \sum_{n=v+1}^{m+1} \frac{1}{n^{k(\alpha-1)+\epsilon+1}} \\ &= O(1) \sum_{v=1}^m v^{\alpha k} (w_v^\alpha)^k \beta_v (\beta_v)^{k-1} v^{\epsilon-k} |\varphi_v|^k \int_v^\infty \frac{dx}{x^{k(\alpha-1)+\epsilon+1}} \\ &= O(1) \sum_{v=1}^m \beta_v (\beta_v)^{k-1} (w_v^\alpha |\varphi_v|)^k \\ &= O(1) \sum_{v=1}^m \beta_v \left( \frac{1}{v X_v} \right)^{k-1} (w_v^\alpha |\varphi_v|)^k \\ &= O(1) \sum_{v=1}^{m-1} \Delta(v \beta_v) \sum_{r=1}^v \frac{(|\varphi_r| w_r^\alpha)^k}{r^k X_r^{k-1}} + O(1) m \beta_m \sum_{v=1}^m \frac{(|\varphi_v| w_v^\alpha)^k}{v^k X_v^{k-1}} \\ &= O(1) \sum_{v=1}^{m-1} |\Delta(v \beta_v)| X_v + O(1) m \beta_m X_m \\ &= O(1) \sum_{v=1}^{m-1} |(v+1) \Delta \beta_v - \beta_v| X_v + O(1) m \beta_m X_m \\ &= O(1) \sum_{v=1}^{m-1} v |\Delta \beta_v| X_v + O(1) \sum_{v=1}^{m-1} \beta_v X_v + O(1) m \beta_m X_m \\ &= O(1) \quad \text{as } m \rightarrow \infty \end{aligned}$$

by virtue of the hypotheses of the theorem and Lemma 2. Finally, we have that

$$\begin{aligned} \sum_{n=1}^m n^{-k} |\varphi_n T_{n,2}^\alpha|^k &= \sum_{n=1}^m |\lambda_n| |\lambda_n|^{k-1} n^{-k} (w_n^\alpha |\varphi_n|)^k \\ &= O(1) \sum_{n=1}^m |\lambda_n| \left( \frac{1}{X_n} \right)^{k-1} n^{-k} (w_n^\alpha |\varphi_n|)^k \end{aligned}$$

$$\begin{aligned} &= O(1) \sum_{n=1}^{m-1} \Delta |\lambda_n| \sum_{n=1}^m \frac{(|\varphi_n| w_n^\alpha)^k}{n^k X_n^{k-1}} + O(1) |\lambda_m| \sum_{n=1}^m \frac{(|\varphi_n| w_n^\alpha)^k}{n^k X_n^{k-1}} \\ &= O(1) \sum_{n=1}^{m-1} |\Delta \lambda_n| X_n + O(1) |\lambda_m| X_m \\ &= O(1) \sum_{n=1}^{m-1} \beta_n X_n + O(1) |\lambda_m| X_m = O(1) \quad \text{as } m \rightarrow \infty, \end{aligned}$$

by virtue of the hypotheses of the theorem and Lemma 2. This completes the proof of the theorem. If we take  $\epsilon = 1$  and  $\varphi_n = n^{1-\frac{1}{k}}$  (resp.  $\epsilon = 1$ ,  $\alpha = 1$  and  $\varphi_n = n^{1-\frac{1}{k}}$ ), then we get a new result dealing with  $|C, \alpha|_k$  (resp.  $|C, 1|_k$ ) summability factors. Also, if we set  $\epsilon = 1$  and  $\varphi_n = n^{\delta+1-\frac{1}{k}}$ , then we get another new result concerning the  $|C, \alpha; \delta|_k$  summability factors. Finally, if we take  $(X_n)$  as an almost increasing sequence, then we get the result of Bor and Seyhan under weaker conditions (see [10]).

#### Competing interests

The author declares that he has no competing interests.

#### Acknowledgements

Dedicated to Professor Hari M Srivastava.

Received: 22 December 2012 Accepted: 9 February 2013 Published: 26 February 2013

#### References

1. Bari, NK, Stečkin, SB: Best approximation and differential properties of two conjugate functions. *Tr. Mosk. Mat. Obs.* **5**, 483-522 (1956) (in Russian)
2. Leindler, L: A new application of quasi power increasing sequences. *Publ. Math. (Debr.)* **58**, 791-796 (2001)
3. Balci, M: Absolute  $\varphi$ -summability factors. *Commun. Fac. Sci. Univ. Ankara, Ser. A1* **29**, 63-80 (1980)
4. Flett, TM: On an extension of absolute summability and some theorems of Littlewood and Paley. *Proc. Lond. Math. Soc.* **7**, 113-141 (1957)
5. Flett, TM: Some more theorems concerning the absolute summability of Fourier series. *Proc. Lond. Math. Soc.* **8**, 357-387 (1958)
6. Bor, H, Özarslan, HS: A study on quasi power increasing sequences. *Rocky Mt. J. Math.* **38**, 801-807 (2008)
7. Pati, T: The summability factors of infinite series. *Duke Math. J.* **21**, 271-284 (1954)
8. Sulaiman, WT: On a new application of almost increasing sequences. *Bull. Math. Anal. Appl.* **4**(3), 29-33 (2012)
9. Bosanquet, LS: A mean value theorem. *J. Lond. Math. Soc.* **16**, 146-148 (1941)
10. Bor, H, Seyhan, H: A note on almost increasing sequences. *Comment. Math. Prace Mat.* **39**, 37-42 (1999)

doi:10.1186/1029-242X-2013-69

**Cite this article as:** Bor: A new application of quasi power increasing sequences. I. *Journal of Inequalities and Applications* 2013 **2013**:69.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)