



The diagnostic accuracy of acute phase proteins and proinflammatory cytokines in sheep with pneumonic pasteurellosis

Wael M. El-Deeb^{1,2} and Ahmed M. Elmoslemany^{1,3}

¹Department of Clinical Studies, College of Veterinary Medicine and Animal Resources, King Faisal University, Al-Ahsa, Saudi Arabia

²Department of Internal Medicine, Infectious Diseases and Fish Diseases, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt

³Faculty of Veterinary Medicine, Hygiene and Preventive Medicine, Kafrelsheikh University, Kafrelsheikh, Egypt

ABSTRACT

The goal of this study was to assess the diagnostic accuracy of acute phase proteins and proinflammatory cytokines in sheep with pneumonic pasteurellosis. Blood samples were collected from 56 sheep (36 naturally infected with *Pasteurella multocida* and 20 healthy controls) belonging to one farm in Eastern region, Saudi Arabia. Serum samples were evaluated for acute phase proteins (Haptoglobin (Hp), serum amyloid A (SAA) and fibrinogen (Fb)), and the proinflammatory cytokines (interleukins (IL-1 α , IL-1 β , and IL-6), tumor necrosis factor-alpha (TNF- α), and interferon-gamma (IFN- γ)). Additionally, nasopharyngeal swabs and bronchoalveolar lavages were collected from all animals for bacteriological examinations. Receiver operating characteristic curve was used to assess the diagnostic performance of each parameter. All parameters showed moderate to high degree of positive correlation with case-control status. There was no significant difference in the area under the curve (AUC) among acute phase proteins; however, both Hp and SAA showed better sensitivity and specificity than Fb. The proinflammatory cytokines (IL1- α , IL1- β , and IL6) showed similar and highly accurate diagnostic performance (AUC > 0.9), whereas IFN- γ was moderately accurate (AUC = 0.79). In conclusion, this study confirms the value of acute phase proteins and cytokines as diagnostic biomarkers of naturally occurring pneumonic pasteurellosis in sheep.

Submitted 2 March 2016

Accepted 1 June 2016

Published 19 July 2016

Corresponding author

Wael M. El-Deeb,
drwaeledeeb@yahoo.com,
Weldeeb@KFU.edu.sa

Academic editor

María Ángeles Esteban

Additional Information and
Declarations can be found on
page 8

DOI 10.7717/peerj.2161

© Copyright
2016 El-Deeb and Elmoslemany

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Agricultural Science, Microbiology, Veterinary Medicine

Keywords Pneumonic pasteurellosis, Acute phase proteins, Cytokines, Sheep, Diagnostic accuracy, Haptoglobin

INTRODUCTION

Pneumonia is an inflammatory response of the alveoli in the lungs to infective agents, resulting in lung consolidation. It is a common disease of sheep in all major sheep-producing countries (Yener *et al.*, 2009). The disease is multifactorial, resulting from dynamic interactions between host, infectious agent and environmental factors. When the bacterial population reach a certain threshold, host susceptibility, non-specific defense mechanisms and lung defense mechanisms become compromised, allowing diseases to occur (Bruere, West & Ridler, 2002).

Pneumonic pasteurellosis and systemic pasteurellosis are the two main diseases caused by *Pasteurella* species in ovine ([Donachie, 2000](#); [Rad et al., 2011](#)). Although *Pasteurella* species are opportunistic pathogens which commonly inhabit the upper respiratory tract of ruminants ([Dowling et al., 2002](#); [Ackermann & Brogden, 2000](#); [Yener et al., 2009](#)), they become pathogenic when the host is exposed to either stressful environment or infection with primary respiratory pathogens such as viral agents or *Mycoplasma* spp. Infection with pneumonic pasteurellosis can cause serious economic losses in sheep population resulting from fatalities in acute outbreaks and reduced productivity of chronically infected animals.

Acute phase proteins (APPs) are plasma proteins which increase or decrease in concentration in response to infection, inflammation and internal or external challenges. Monitoring changes in APPs levels have been shown to provide valuable diagnostic and prognostic information during infection and inflammation. However, there are substantial variations in acute phase response between different species ([Eckersall, 2000](#)). In small ruminants, some APPs levels change similarly in both sheep and goat, whereas other APPs show different magnitude of response between the two species ([Gonzalez et al., 2008](#)).

In bovine and camels, several studies have been conducted to elucidate the role of APPs in different disorders including parasitic and bacterial infection ([El-Deeb & Iacob, 2012](#); [El-Deeb, Fouda & El-Bahr, 2014](#); [El-Deeb, 2015](#); [El-Deeb & Elmoslemany, 2016](#); [El-Deeb & Buczinski, 2015](#)). Conversely, only a limited number of studies have been done regarding specific bacterial, viral or parasitic infections in ovine species ([El-Deeb, 2013](#); [El-Deeb & Tharwat, 2015](#)). Furthermore, these studies focused mainly on changes in the level of APPs but provided little information on their diagnostic accuracy. Therefore, the goal of this study was to assess the diagnostic accuracy of APPs and inflammatory cytokines in sheep infected with pneumonic pasteurellosis.

MATERIALS AND METHODS

Study animals

This study was conducted initially on 73 Naimi sheep, three to four years old (53 cases and 20 healthy control) belonging to a flock of free grazing sheep ($n = 543$) in Al-Ahsa region, Saudi Arabia. The project was approved by the animal care committee at King Faisal University (number 130031). The flock was examined following reports of respiratory disease problems. All sheep were clinically examined with direct observation and recording of clinical signs. Additionally, nasopharyngeal swabs and bronchoalveolar lavages were collected from sheep exhibiting signs of clinical disease for bacteriologic examination. Sheep were classified into cases and controls based on clinical signs, culture results and autopsy for dead animals. Cases were defined as sheep with clinical signs of pneumonia and positive culture for *Pasteurella multocida* (*P. multocida*). Healthy controls were sheep with no clinical signs and negative culture results. Seven animals (with positive nasal swabs, negative bronchoalveolar lavages, and with no clinical signs of pneumonia) and ten animals (with other bacterial infections) were excluded from the study to avoid misclassification of control sheep as being cases. Finally, only 36 sheep that satisfied case definition and 20 healthy controls (randomly selected from healthy sheep) were considered for further analysis in this study.

Sampling

Serum samples were collected from all 56 sheep and stored at -20°C until biochemical analyses were performed. Moreover, nasopharyngeal swabs and bronchoalveolar lavage were collected from all animals under investigation ($N = 73$). Finally, heart-blood and lung samples (from necropsied sheep ($n = 6$)) which died after initial serum sampling were also collected for bacteriological examinations.

Determination of acute phase proteins and inflammatory cytokines

Hp levels were detected in serum samples via preservation of the peroxidase activity of haemoglobin, which is directly proportional to the amount of Hp (Tridelta Development Plc.). SAA was measured in serum samples by a solid phase sandwich ELISA (Tridelta Development Plc.).

The levels of inflammatory cytokines IL-6, TNF- α , IL-1 α , IL-1 β , and IFN- γ were measured from serum samples using commercially available sheep ELISA Kits (CUSABIO).

Isolation and identification of *P. multocida*

All samples were cultured for isolation and identification of *P. multocida* based on cultural, morphological and biochemical characteristics using standard bacteriological techniques (Carter & Cole Jr, 1990).

Statistical analysis

Descriptive statistics were calculated as mean, median, and 25th and 75th percentiles for each parameter by disease status (pneumonic cases vs. controls). The mean ranks for each parameter were compared between cases and control groups using non-parametric Wilcoxon rank-sum (Mann–Whitney) test due to significant deviation of the data from normality and failure to normalize data with transformation. Differences between cases and control were considered statistically significant at $P < 0.05$. Correlation among different parameters was assessed using Spearman's correlation coefficient.

Selection of cut-off points that optimize sensitivity and specificity for each parameter were determined using non-parametric receiver operating characteristic curve (ROC). The ROC curves were constructed by plotting 1-specificity (x -axis) versus sensitivity (y -axis) for all possible threshold values of the parameter to be evaluated. The area under the curve (AUC) indicates the overall accuracy of the tested parameter. A biomarker with no predictive value would have an AUC of 0.5 (represented by the diagonal line in the ROC plot Fig. 1), while a biomarker with perfect ability to predict disease would have an AUC of one. Values of AUC between 0.5 and one are interpreted as low ($0.5 > \text{AUC} \leq 0.7$), moderate ($0.7 > \text{AUC} \leq 0.9$), or high ($0.9 > \text{AUC} < 1$), accuracy (Swets, 1988). The two-graph ROC (TGROC) plot was also used to graph variation of sensitivity and specificity of biomarker across a range of cut-offs. All analyses were done using Stata version 13 (Stata Corp).

RESULTS

Table 1 shows significantly higher levels of Hp, SAA, and Fb in cases compared to controls, though the increase was more pronounced (several folds) in both Hp and SAA levels.




Figure 1 Receiver operating characteristic plot: comparison of the area under the curve (AUC) for haptoglobin (Hp), serum amyloid A (SAA), and fibrinogen (Fb).

Table 1 Summary statistics of the level of acute phase proteins (APP) and proinflammatory cytokines in control and pneumonic sheep.

| Parameter | Control (N = 20) | | | | Cases (N = 36) | | | | P ^a value |
|---------------|------------------|--------|-------|-------|----------------|--------|-------|-------|----------------------|
| | Mean | Median | 25% | 75% | Mean | Median | 25% | 75% | |
| Hp (g/L) | 0.048 | 0.054 | 0.047 | 0.062 | 1.65 | 1.87 | 1.65 | 2.14 | <0.0001 |
| SAA (μg/mL) | 4.32 | 4.56 | 3.95 | 4.87 | 26.89 | 29.36 | 28.26 | 30.75 | <0.0001 |
| Fb (g/L) | 2.34 | 2.36 | 2.27 | 2.41 | 3.47 | 3.65 | 2.51 | 4.12 | <0.0001 |
| IL1-α (pg/ml) | 13.48 | 13.75 | 12.36 | 14.74 | 24.75 | 26.35 | 24.96 | 27.81 | <0.0001 |
| IL1-β (pg/ml) | 18.49 | 18.52 | 17.26 | 19.45 | 29.30 | 30.35 | 30.12 | 31.80 | <0.0001 |
| IL6 (pg/ml) | 10.63 | 10.40 | 9.51 | 11.40 | 16.27 | 17.25 | 15.47 | 18.24 | <0.0001 |
| TNF-α (pg/ml) | 8.63 | 8.57 | 8.05 | 9.25 | 17.57 | 19.25 | 18.31 | 19.84 | <0.0001 |
| IFN-γ (pg/ml) | 10.02 | 10.31 | 9.07 | 10.79 | 14.53 | 16.30 | 10.36 | 17.25 | <0.0001 |

Notes.

^aP value resulting from non-parametric Wilcoxon Mann-Whitney test.

Additionally, a significant increase ($P < 0.05$) in the level of proinflammatory cytokines was also observed in the cases.

Table 2 shows correlation coefficients (r) among study parameters. All measured biomarkers had a significant ($P < 0.05$) positive correlation with pneumonic cases and most of the studied biomarkers were moderately correlated. SAA showed the highest correlation with pneumonic cases ($r = 0.74$). A high correlation ($r \geq 0.75$) was observed between Hp & IL1-α, and also among different interleukins. A low correlation ($r \leq 0.34$) was observed between IFN-γ and most of the other parameters.

The results of the ROC analysis for evaluation of the overall diagnostic accuracy of APPs and cytokines are shown in Table 3. Comparison of the AUC indicated no significant difference ($P = 0.49$) in the AUC among Hp (0.90), SAA (0.96), and Fb (0.93) (Fig. 1).

Table 2 Correlation matrix among acute phase proteins (APPs) and proinflammatory cytokines in 56 sheep (20 control and 36 with pneumonic pasteurellosis).

| Parameter | Case control | Hp | SAA | Fb | IL1- α | IL1- β | IL6 | TNF- α |
|---------------|--------------|------|------|------|---------------|--------------|------|---------------|
| Hp | 0.66 | | | | | | | |
| SAA | 0.74 | 0.51 | 1.00 | | | | | |
| Fb | 0.71 | 0.53 | 0.61 | 1.00 | | | | |
| IL1- α | 0.68 | 0.75 | 0.38 | 0.48 | 1.00 | | | |
| IL1- β | 0.70 | 0.71 | 0.47 | 0.50 | 0.71 | 1.00 | | |
| IL6 | 0.68 | 0.73 | 0.42 | 0.51 | 0.78 | 0.77 | 1.00 | |
| TNF- α | 0.72 | 0.50 | 0.54 | 0.56 | 0.58 | 0.49 | 0.51 | 1.00 |
| IFN- γ | 0.48 | 0.33 | 0.33 | 0.48 | 0.25 | 0.31 | 0.25 | 0.34 |

Table 3 Test characteristics of acute phase proteins (APPs) and proinflammatory cytokines for diagnosis of pneumonic pasteurellosis in sheep.

| Parameter | Threshold | Sensitivity (%) | Specificity (%) | % correctly classified | AUC (95% CI) ^a |
|--------------------------|--------------|-----------------|-----------------|------------------------|---------------------------|
| Hp (g/L) | ≥ 0.065 | 89 | 100 | 93 | 0.90 (0.78–0.96) |
| SAA ($\mu\text{g/mL}$) | ≥ 5.26 | 89 | 95 | 91 | 0.96 (0.88–0.99) |
| Fb (g/L) | ≥ 2.45 | 86 | 85 | 86 | 0.93 (0.83–0.98) |
| IL1- α (pg/ml) | ≥ 16.25 | 86 | 95 | 89 | 0.91 (0.80–0.97) |
| IL1- β (pg/ml) | ≥ 29 | 86 | 100 | 91 | 0.92 (0.85–0.99) |
| IL6 (pg/ml) | ≥ 15 | 86 | 100 | 91 | 0.91 (0.83–0.99) |
| TNF- α (pg/ml) | ≥ 17 | 83 | 100 | 89 | 0.93 (0.87–1.00) |
| IFN- γ (pg/ml) | ≥ 15 | 69 | 100 | 80 | 0.79 (0.67–0.91) |

Notes.^a95% CI, 95% confidence interval.

However, both Hp and SAA showed better sensitivity and specificity than Fb at selected cut-offs (Table 3). Using ROC analysis and TGROC, the optimum cut-offs for discrimination between pneumonic and control sheep for Hp, SAA and Fb were 0.065, 5.26 and 2.45 g/L, respectively (Table 3). Figure 2 illustrates the optimal cut-off and variation in sensitivity and specificity of Fb at different cut-off points.

The proinflammatory cytokines IL1- α , IL1- β , and IL6 showed similar and highly accurate diagnostic performance (AUC > 0.9), whereas IFN- γ was moderately accurate in differentiating between pneumonic and control sheep (AUC = 0.79) (Table 3).

DISCUSSION

To the best of our knowledge, our data is the first to explore the diagnostic accuracy of APPs and inflammatory cytokines in sheep with pneumonic pasteurellosis under natural field condition.

Overall, there is limited information on the relationship between APPs and naturally occurring pneumonic pasteurellosis in sheep. Hp acts as a major APP in ruminants, it could be detected in subclinically diseased animals and its serum level is used as a




Figure 2 Two-graph receiver operating characteristic (TG-ROC) plot showing optimal cut-off and variation in sensitivity and specificity over various cut-off points of fibrinogen (Fb).

biological marker of disease severity (Godson *et al.*, 1996). In our study, Hp concentration increased 34 times in pneumonic sheep indicating a major response to the infection. The increased plasma level of Hp could be induced by tissue damage resulting from infection or inflammation (Blackmore, 1988). Hp has bacteriostatic effects by binding free haemoglobin, thus depriving bacteria from iron required for their growth (Eaton *et al.*, 1982). Our results are consistent with previous studies on calves with respiratory diseases due to natural (Humblet *et al.*, 2004; Nikunen *et al.*, 2007) or experimental (Dowling *et al.*, 2002) infection with *P. multocida*. Similar results were also observed in sheep with experimentally induced pneumonia (Pfeffer & Rogers, 1989) and buffalo calves with bacterial bronchopneumonia (El-Deeb, 2011; El-Bahr & El-Deeb, 2013). In contrast, some studies demonstrated limited association between Hp and respiratory diseases in feedlot cattle (Wittum *et al.*, 1996; Young *et al.*, 1996). This study also showed that SAA level increased approximately 7 folds in pneumonic sheep compared to healthy one indicating a moderate response, this could be attributed to the physiological role of SAA in host defense during inflammation (Murata, Shimada & Yoshioka, 2004; Orro *et al.*, 2011; Urieli-Shoval, Linke & Matzner, 2000). SAA has a role in alteration of cholesterol metabolism under inflammatory conditions (Pannen & Robotham, 1995). Moreover, SAA binds Gram-negative bacteria (Hari-Dass *et al.*, 2005), possibly to facilitate the uptake by macrophages and neutrophils (Larson *et al.*, 2005). Similarly, marked elevation of SAA was reported in pneumonic calves (Horadagoda *et al.*, 1994; Nikunen *et al.*, 2007; Orro *et al.*, 2011). Although, some studies suggested that SAA is a

more sensitive biomarker for viral infections (Heegaard *et al.*, 2000) and acute inflammation (Horadagoda *et al.*, 1994). Others indicated that Hp might be preferred to SAA in detecting respiratory disease in calves under field conditions (Angen *et al.*, 2009). In the former study, SAA showed more rapid response, whereas Hp was more correlated with severity of clinical signs and degree of lung consolidation (Heegaard *et al.*, 2000). Fb is classified as a minor APP, which is characterized by a minor response with a maximum increase of about twice the normal concentrations during infection or inflammation (Hirvonen, Hietakorpi & Saloniemä, 1997). Fb is also considered a consistent marker of bacterial infection and inflammation in domestic ruminants (Gonzalez *et al.*, 2008; Nikunen *et al.*, 2007; Pfeiffer *et al.*, 1993; Youssef, El-khodery & Abdo, 2015). In our study, the concentration of serum Fb in pneumonic sheep was approximately twice that of the control group. This elevation may be attributed to the involvement of Fb in modulating hemostasis, inflammatory response, and the tissue repairing process (Feldman *et al.*, 2000).

The different types of inflammatory cytokines are the principal stimulators of APPs gene expression, and each kind of cytokines recruits a different type of APPs (Baumann & Gauldie, 1994). Thus, the elevated levels of APPs seen in this study reveal the secretion of different amounts or types of inflammatory cytokines. Comparable results regarding such high levels of IL1- β , TNF- α and IFN- γ were detected in pigs with viral respiratory disease (Van Reeth & Nauwynck, 2000) and cattle with bacterial infection (Pace *et al.*, 1993; Horadagoda *et al.*, 1994; Morse *et al.*, 1999; Kasimanickam *et al.*, 2013; El-Deeb & Elmoslemany, 2016). In addition, the expression of IL1- β and TNF- α were elevated in the respiratory airways and lung lesions of diseased calves with pneumonic pasteurellosis (Malazdrewich *et al.*, 2001).

The ability of APPs to discriminate between pneumonic and healthy sheep was evaluated using ROC analysis. Under the conditions of this study, all APPs showed a high degree of discrimination between pneumonic and control sheep ($AUC \geq 0.9$) according to guidelines reported by Swets (1988). However, both Hp and SAA showed better sensitivity and specificity than Fb resulting in better overall correct classification of pneumonic and control sheep. Selection of cut-off point for each parameters was based on optimizing sensitivity and specificity which lead to the best overall correct classification. The ability of Hp and SAA to discriminate between control and clinical cases may be related to the large difference in their concentration between control and clinical cases. The proinflammatory cytokines IL1- α , IL1- β , and IL6 showed similar and highly accurate diagnostic performance ($AUC > 0.9$), whereas IFN- γ was moderately accurate in differentiation between control and pneumonic sheep ($AUC = 0.79$). Our results also indicate a high diagnostic accuracy of the measured cytokines. However, APPs may serve as better biomarkers of inflammation considering the very short half-life of cytokines.

CONCLUSIONS

The results of this study indicate that pneumonic pasteurellosis caused by *P. multocida* in sheep was associated with a significant increase in APPs and cytokines, with the greatest increase being observed in Hp. There was a moderate to high correlation between disease

status and APPs and cytokines concentrations. Finally, this study highlights the value of acute phase proteins and cytokines as diagnostic biomarkers of naturally occurring pneumonic pasteurellosis in sheep.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The research is funded by the Deanship of Scientific Research, King Faisal University (Project No. 130031). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Deanship of Scientific Research, King Faisal University: 130031.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Wael M. El-Deeb conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Ahmed M. Elmoslemany analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The research is approved by the Deanship of Scientific Research, King Faisal University (Approval number: 130031).

Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as [Data S1](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2161#supplemental-information>.

REFERENCES

- Ackermann MR, Brogden KA. 2000. Response of the ruminant respiratory tract to *Mannheimia (Pasteurella) haemolytica*. *Microbes and Infection* 2:1079–1088
[DOI 10.1016/S1286-4579\(00\)01262-4](https://doi.org/10.1016/S1286-4579(00)01262-4).

- Angen Ø, Thomsen J, Larsen LE, Larsen J, Kokotovic B, Heegaard PM, Enemark J. 2009.** Respiratory disease in calves: microbiological investigations on trans-tracheally aspirated bronchoalveolar fluid and acute phase protein response. *Veterinary Microbiology* **137**:165–171 DOI [10.1016/j.vetmic.2008.12.024](https://doi.org/10.1016/j.vetmic.2008.12.024).
- Baumann H, Gauldie J. 1994.** The acute phase response. *Immunology Today* **15**:74–80 DOI [10.1016/0167-5699\(94\)90137-6](https://doi.org/10.1016/0167-5699(94)90137-6).
- Blackmore DJ. 1988.** *Animal clinical biochemistry-the future*. Cambridge: Cambridge University Press.
- Bruere A, West D, Ridler A. 2002.** *Enzootic pneumonia. The sheep: health, disease & production: written for veterinarians and farmers*. Palmerston North: Veterinary Continuing Education Massey University, 100–108.
- Carter GR, Cole Jr JR. 1990.** *Diagnostic procedure in veterinary bacteriology and mycology*. Fifth edition. San Diego: Academic Press.
- Donachie E. 2000.** Bacteriology of bovine respiratory disease. *Cattle Practice* **8**:5–7.
- Dowling A, Hodgson J, Schock A, Donachie W, Eckersall P, McKendrick I. 2002.** Experimental induction of pneumonic pasteurellosis in calves by intratracheal infection with *Pasteurella multocida* biotype A: 3. *Research in Veterinary Science* **73**:37–44 DOI [10.1016/S0034-5288\(02\)00037-1](https://doi.org/10.1016/S0034-5288(02)00037-1).
- Eaton JW, Brandt P, Mahoney JR, Lee JT. 1982.** Haptoglobin—a natural bacteriostat. *Science* **215**:691–693 DOI [10.1126/science.7036344](https://doi.org/10.1126/science.7036344).
- Eckersall P. 2000.** Recent advances and future prospects for the use of acute phase proteins as markers of disease in animals. *Revue de Médecine Vétérinaire* **151**:577–584.
- El-Bahr SM, El-Deeb WM. 2013.** Acute phase proteins, lipid profile and proinflammatory cytokines in healthy and bronchopneumonic water buffalo calves. *American Journal of Biochemistry and Biotechnology* **9**(1):34–40 DOI [10.3844/ajbbsp.2013.34.40](https://doi.org/10.3844/ajbbsp.2013.34.40).
- El-Deeb WM. 2011.** Clinico-biochemical investigation of bacterial bronchopneumonia in water buffalo calves: acute phase response and lipoprotein profiles. *Lucrări Stiintifice USAMV Iasi, Medicină Veterinară* **54**(4):412–417.
- El-Deeb WM. 2013.** Clinicobiochemical investigations of gangrenous mastitis in Does: immunological responses and oxidative stress biomarkers. *Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology)* **14**(1):33–40 DOI [10.1631/jzus.B1200123](https://doi.org/10.1631/jzus.B1200123).
- El-Deeb WM. 2015.** Acute phase response and oxidative stress biomarkers in pneumonic camel-calves (*Camelus dromedarius*). *Bulgarian Veterinary Journal* **18**(3):258–269 DOI [10.15547/bjvm.853](https://doi.org/10.15547/bjvm.853).
- El-Deeb WM, Buczinski S. 2015.** The diagnostic and prognostic importance of oxidative stress biomarkers and acute phase proteins in urinary tract infection (UTI) in camels. *PeerJ* **3**:e1363 DOI [10.7717/peerj.1363](https://doi.org/10.7717/peerj.1363).
- El-Deeb WM, Elmoslemany AM. 2016.** Acute phase proteins as biomarkers of Urinary Tract Infection in dairy cows: diagnostic and prognostic accuracy. *Japanese Journal of Veterinary Research* **64**(1):57–66 DOI [10.14943/jjvr.64.1.57](https://doi.org/10.14943/jjvr.64.1.57).

- El-Deeb WM, Fouda TA, El-Bahr SM. 2014. Clinicobiochemical investigations of Partuberculosis of camels in Saudi Arabia: proinflammatory cytokines and oxidative stress markers. *Pakistan Veterinary Journal* **34**(4):484–488.
- El-Deeb WM, Iacob O. 2012. Serum Acute phase proteins in control and *Theileria annulata* infected water buffaloes (*Bubalus bubalis*). *Veterinary Parasitology* **23**; **190**(1–2):12–18 DOI [10.1016/j.vetpar](https://doi.org/10.1016/j.vetpar.2012.05.001).
- El-Deeb W, Tharwat M. 2015. Lipoproteins profile, acute phase proteins, proinflammatory cytokines and oxidative stress biomarkers in sheep with pneumonic pasteurellosis. *Comparative Clinical Pathology* **24**:581–588 DOI [10.1007/s00580-014-1949-z](https://doi.org/10.1007/s00580-014-1949-z).
- Feldman B, Zinkl J, Jain N, Schalm S. 2000. *Veterinary hematology*. Philadelphia: Lippincott Williams & Wilkins.
- Godson DL, Campos M, Attah-Poku SK, Redmond MJ, Cordeiro DM, Sethi MS, Harland RJ, Babiuk LA. 1996. Serum haptoglobin as an indicator of the acute phase response in bovine respiratory disease. *Veterinary Immunology and Immunopathology* **51**:277–292 DOI [10.1016/0165-2427\(95\)05520-7](https://doi.org/10.1016/0165-2427(95)05520-7).
- Gonzalez FH, Tecles F, Martinez-Subiela S, Tvarijonaviciute A, Soler L, Ceron JJ. 2008. Acute phase protein response in goats. *Journal of Veterinary Diagnostic Investigation* **20**:580–584 DOI [10.1177/104063870802000507](https://doi.org/10.1177/104063870802000507).
- Hari-Dass R, Shah C, Meyer DJ, Raynes JG. 2005. Serum amyloid A protein binds to outer membrane protein A of gram-negative bacteria. *The Journal of Biological Chemistry* **280**:18562–18567 DOI [10.1074/jbc.M500490200](https://doi.org/10.1074/jbc.M500490200).
- Heegaard PM, Godson DL, Toussaint MJ, Tjørnehøj K, Larsen LE, Viuff B, Rønsholt L. 2000. The acute phase response of haptoglobin and serum amyloid A (SAA) in cattle undergoing experimental infection with bovine respiratory syncytial virus. *Veterinary Immunology and Immunopathology* **77**:151–159 DOI [10.1016/S0165-2427\(00\)00226-9](https://doi.org/10.1016/S0165-2427(00)00226-9).
- Hirvonen J, Hietakorpi S, Saloniemi H. 1997. Acute phase response in emergency slaughtered dairy cows. *Meat Science* **46**:249–257.
- Horadagoda A, Eckersall P, Hodgson J, Gibbs H, Moon G. 1994. Immediate responses in serum TNF- α and acute phase protein concentrations to infection with *Pasteurella haemolytica* A1 in calves. *Research in Veterinary Science* **57**:129–132 DOI [10.1016/0034-5288\(94\)90094-9](https://doi.org/10.1016/0034-5288(94)90094-9).
- Humblet M, Coghe J, Lekeux P, Godeau J. 2004. Acute phase proteins assessment for an early selection of treatments in growing calves suffering from bronchopneumonia under field conditions. *Research in Veterinary Science* **77**:41–47 DOI [10.1016/j.rvsc.2004.02.009](https://doi.org/10.1016/j.rvsc.2004.02.009).
- Kasimanickam RK, Kasimanickam VR, Olsen JR, Jeffress EJ, Moore DA, Kastelic JP. 2013. Associations among serum pro- and anti-inflammatory cytokines, metabolic mediators, body condition, and uterine disease in postpartum dairy cows. *Reproductive Biology and Endocrinology* **11**:103 DOI [10.1186/1477-7827-11-103](https://doi.org/10.1186/1477-7827-11-103).
- Larson MA, Weber A, Weber AT, McDonald TL. 2005. Differential expression and secretion of bovine serum amyloid A (SAA3) by mammary epithelial cells stimulated

- with prolactin or lipopolysaccharide. *Veterinary Immunology Immunopathology* **107**:255–264 DOI [10.1016/j.vetimm.2005.05.006](https://doi.org/10.1016/j.vetimm.2005.05.006).
- Malazdrewich C, Ames TR, Abrahamsen MS, Maheswaran SK. 2001.** Pulmonary expression of tumor necrosis factor alpha, interleukin-1 beta, and interleukin-8 in the acute phase of bovine pneumonic pasteurellosis. *Veterinary Pathology* **38**:297–310 DOI [10.1354/vp.38-3-297](https://doi.org/10.1354/vp.38-3-297).
- Morsey M, Van-Kessel A, Mori Y, Popowych Y, Godson D, Campos M, Babiuk L. 1999.** Cytokine profiles following interaction between bovine alveolar macrophages and *Pasteurella haemolytica*. *Microbial Pathogenesis* **26**:325–331 DOI [10.1006/mpat.1999.0274](https://doi.org/10.1006/mpat.1999.0274).
- Murata H, Shimada N, Yoshioka M. 2004.** Current research on acute phase proteins in veterinary diagnosis: an overview. *The Veterinary Journal* **168**:28–40 DOI [10.1016/S1090-0233\(03\)00119-9](https://doi.org/10.1016/S1090-0233(03)00119-9).
- Nikunen S, Härtel H, Orro T, Neuvonen E, Tanskanen R, Kivelä S, Sankari S, Aho P, Pyörälä S, Saloniemi H. 2007.** Association of bovine respiratory disease with clinical status and acute phase proteins in calves. *Comparative Immunology, Microbiology and Infectious Diseases* **30**:143–151 DOI [10.1016/j.cimid.2006.11.004](https://doi.org/10.1016/j.cimid.2006.11.004).
- Orro T, Pohjanvirta T, Rikula U, Huovilainen A, Alasuutari S, Sihvonen L, Pelkonen S, Soveri T. 2011.** Acute phase protein changes in calves during an outbreak of respiratory disease caused by bovine respiratory syncytial virus. *Comparative Immunology, Microbiology and Infectious Diseases* **34**:23–29 DOI [10.1016/j.cimid.2009.10.005](https://doi.org/10.1016/j.cimid.2009.10.005).
- Pace L, Kreeger J, Bailey K, Turnquist S, Fales W. 1993.** Serum levels of tumor necrosis factor- α in calves experimentally infected with *Pasteurella haemolytica* A1. *Veterinary Immunology and Immunopathology* **35**:353–364 DOI [10.1016/0165-2427\(93\)90044-5](https://doi.org/10.1016/0165-2427(93)90044-5).
- Pannen BHJ, Robotham JL. 1995.** The acute phase response. *New Horiz* **3**:183–197.
- Pfeffer A, Rogers KM. 1989.** Acute phase response of sheep: changes in the concentrations of ceruloplasmin, fibrinogen, haptoglobin and the major blood cell types associated with pulmonary damage. *Research in Veterinary Science* **46**:118–124.
- Pfeffer A, Rogers K, O'keeffe L, Osborn P. 1993.** Acute phase protein response, food intake, liveweight change and lesions following intrathoracic injection of yeast in sheep. *Research in Veterinary Science* **55**:360–366 DOI [10.1016/0034-5288\(93\)90108-R](https://doi.org/10.1016/0034-5288(93)90108-R).
- Rad M, Movassaghi AR, Sharifi K, Naseri Z, Seifi H. 2011.** Two outbreaks of *Pasteurella multocida* septicemia in neonatal lambs. *Comparative Clinical Pathology* **20**:57–59 DOI [10.1007/s00580-009-0936-2](https://doi.org/10.1007/s00580-009-0936-2).
- Swets JA. 1988.** Measuring the accuracy of diagnostic systems. *Science* **240**:1285–1293 DOI [10.1126/science.3287615](https://doi.org/10.1126/science.3287615).
- Urieli-Shoval S, Linke RP, Matzner Y. 2000.** Expression and function of serum amyloid A, a major acute-phase protein, in normal and disease states. *Current Opinion in Hematology* **7**:64–69 DOI [10.1097/00062752-200001000-00012](https://doi.org/10.1097/00062752-200001000-00012).
- Van Reeth K, Nauwynck H. 2000.** Proinflammatory cytokines and viral respiratory disease in pigs. *Veterinary Research* **31**:187–213 DOI [10.1051/vetres:2000113](https://doi.org/10.1051/vetres:2000113).

- Wittum TE, Young CR, Stanker LH, Griffin DD, Perino LJ, Littledike ET. 1996.** Haptoglobin response to clinical respiratory tract disease in feedlot cattle. *American Journal of Veterinary Research* 57:646–649.
- Yener Z, Ilhan F, Ilhan Z, Saglam Y. 2009.** Immunohistochemical detection of *Mannheimia (Pasteurella) haemolytica* antigens in goats with natural pneumonia. *Veterinary Research Communications* 33:305–313 DOI [10.1007/s11259-008-9178-z](https://doi.org/10.1007/s11259-008-9178-z).
- Young CR, Wittum TE, Stanker LH, Perino LJ, Griffin DD, Littledike ET. 1996.** Serum haptoglobin concentrations in a population of feedlot cattle. *American Journal of Veterinary Research* 57:138–141.
- Youssef MA, El-khodery SA, Abdo M. 2015.** A comparative study on selected acute phase proteins (APPs) and immunoglobulins in buffalo and bovine calves with respiratory disease. *Comparative Clinical Pathology* 24:515–520 DOI [10.1007/s00580-014-1933-7](https://doi.org/10.1007/s00580-014-1933-7).