



# Sediment microbial taxonomic and functional diversity in a natural salinity gradient challenge Remane's "species minimum" concept

Christina Pavlouidi<sup>1,2,3</sup>, Jon B. Kristoffersen<sup>1</sup>, Anastasis Oulas<sup>1,4</sup>,  
Marleen De Troch<sup>2</sup> and Christos Arvanitidis<sup>1</sup>

<sup>1</sup>Institute of Marine Biology, Biotechnology and Aquaculture (IMBBC), Hellenic Centre for Marine Research (HCMR), Heraklion, Crete, Greece

<sup>2</sup>Marine Biology Research Group, Department of Biology, Faculty of Sciences, Ghent University, Ghent, Belgium

<sup>3</sup>Microbial Ecophysiology Group, Faculty of Biology/Chemistry and MARUM, University of Bremen, Bremen, Germany

<sup>4</sup>Bioinformatics Group, The Cyprus Institute of Neurology and Genetics, Nicosia, Cyprus

## ABSTRACT

Several models have been developed for the description of diversity in estuaries and other brackish habitats, with the most recognized being Remane's Artenminimum ("species minimum") concept. It was developed for the Baltic Sea, one of the world's largest semi-enclosed brackish water body with a unique permanent salinity gradient, and it argues that taxonomic diversity of macrobenthic organisms is lowest within the horohalinicum (5 to 8 psu). The aim of the present study was to investigate the relationship between salinity and sediment microbial diversity at a freshwater-marine transect in Amvrakikos Gulf (Ionian Sea, Western Greece) and assess whether species composition and community function follow a generalized concept such as Remane's. DNA was extracted from sediment samples from six stations along the aforementioned transect and sequenced for the 16S rRNA gene using high-throughput sequencing. The metabolic functions of the OTUs were predicted and the most abundant metabolic pathways were extracted. Key abiotic variables, i.e., salinity, temperature, chlorophyll-a and oxygen concentration etc., were measured and their relation with diversity and functional patterns was explored. Microbial communities were found to differ in the three habitats examined (river, lagoon and sea) with certain taxonomic groups being more abundant in the freshwater and less in the marine environment, and *vice versa*. Salinity was the environmental factor with the highest correlation to the microbial community pattern, while oxygen concentration was highly correlated to the metabolic functional pattern. The total number of OTUs showed a negative relationship with increasing salinity, thus the sediment microbial OTUs in this study area do not follow Remane's concept.

Submitted 4 May 2017  
Accepted 24 July 2017  
Published 13 October 2017

Corresponding author  
Christina Pavlouidi,  
[cpavloud@hcmr.gr](mailto:cpavloud@hcmr.gr)

Academic editor  
Hilary Morrison

Additional Information and  
Declarations can be found on  
page 18

DOI 10.7717/peerj.3687

 Copyright  
2017 Pavlouidi et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

**Subjects** Biodiversity, Marine Biology, Microbiology

**Keywords** Amvrakikos Gulf, Salinity gradient, Illumina sequencing, Prokaryotes, Sediment, 16S rRNA

## INTRODUCTION

Salinity is considered as one of the most influential environmental factors, not only for the distribution of benthic and pelagic organisms (*Remane, 1934; Rolston & Dittmann, 2009; Palmer et al., 2011; Darr, Gogina & Zettler, 2014*), but also for microbial community composition (e.g., *Barcina, Lebaron & Vives-Rego, 2006; Wu et al., 2006; Lozupone & Knight, 2007; Logares et al., 2009*). Bacterial abundance, activity and growth can be affected by salinity (*Ben-Dov, Brenner & Kushmaro, 2007; Caporaso et al., 2011*) and, in certain cases salinity can induce mortality of bacteria, thus regulating bacterial abundance in some estuaries (*Painchaud, Therriault & Legendre, 1995*). Specifically, salinity fluctuations, and their subsequent effect on aquatic biota, are more noticeable in estuaries and other brackish water bodies, as these habitats are characterized by a more or less pronounced salinity gradient (*Telesh, Schubert & Skarlato, 2013*).

It has been suggested that in brackish water ecosystems, taxonomic diversity of macrobenthic organisms is lowest within the horohalinicum, which occurs at salinity 5 to 8 psu, because the number of brackish specialist species does not compensate for the decline of the marine and freshwater species richness (*Remane, 1934*). This concept, referred to as the Remane's Artenminimum ('species minimum') concept originated from the Baltic Sea, one of the world's largest semi-enclosed, brackish water body with a unique permanent salinity gradient (*Telesh, Schubert & Skarlato, 2011*). Despite being developed for the Baltic Sea, Remane's concept became the recognized model for the description of diversity in estuaries and other brackish habitats (*McLusky & Elliott, 2004*). However, alternative models challenging Remane's concept have also been developed. In certain cases a reverse curve has been observed, with the peak of species occurring in the horohalinicum (*Telesh, Schubert & Skarlato, 2011*) while in others a linear decrease (*Attrill, 2002*) or even no change (*Herlemann et al., 2011*) in the number of species across the salinity gradient were observed.

Remane's concept can be projected in other aquatic bodies with similar evolution to the Baltic Sea, such as the Amvrakikos Gulf (Ionian Sea, Western Greece) (*Ferentinos et al., 2010*). The Gulf was formed in the Middle Quaternary period (*Anastasakis, Piper & Tziavos, 2007*); the marine transgression took place at approximately 11 ka BP and the Gulf attained its present shape at approximately 4 ka BP (*Kapsimalis et al., 2005*). In addition, the low tidal range (on average 5 cm) and the low energy wave regime prevailing in Amvrakikos Gulf (*Ferentinos et al., 2010*), render the latter as a Baltic Sea analogue in the Mediterranean Sea.

In the light of projected climate changes and the subsequent sea level rise and saltwater intrusion that will occur, microbial populations in freshwater wetlands near the coast will be subjected to elevated salinities (*Chambers, Reddy & Osborne, 2011*). Due to the long-term effect of sea level rise, saltwater intrusion can affect ecosystems on timescales of decades (*Neubauer, Franklin & Berrier, 2013*). It is therefore crucial to explore the current status of microbial communities in wetlands in order to comprehend the impact of such acute changes in their diversity patterns, since they are involved in biogeochemical processes that are crucial for maintaining the planet in a habitable state (e.g., *Falkowski, Fenchel & Delong, 2008*).

Currently, state-of-the-art studies of microbial communities through high throughput sequencing of the 16S rRNA gene, i.e., marker gene metagenomics ([Oulas et al., 2015](#)), allow documentation of the high diversity of microbial communities in a variety of habitats, e.g., wetlands ([Wang et al., 2012](#); [Jiang et al., 2013](#)), estuaries ([Bobrova et al., 2016](#)), lakes ([Zhang et al., 2015](#)), rivers ([Staley et al., 2013](#)) and coastal lagoons ([Highton et al., 2016](#)). The substantial data derived from such techniques can be used to test ecological hypotheses, on which sound conclusions can be drawn; for example if microorganisms have a biogeography ([Zinger et al., 2011](#)) or if their biodiversity is driven by changes in elevation ([Fierer et al., 2011](#)).

The aim of the present study was to investigate the sediment prokaryotic diversity along a transect river-lagoon-open sea, i.e., from freshwater to marine, in Amvrakikos Gulf in order to test whether it follows a generalized concept, such as Remane's, both in terms of Operational Taxonomic Unit (OTU) composition as well as on distribution of metabolic functions. If the applicability of the concept is confirmed in Amvrakikos Gulf, it would enhance its transferability to other brackish water bodies than the Baltic Sea. In addition, confirmation of Remane's concept for prokaryotes would mean that small-bodied, fast-developing and rapidly evolving microbes respond in a similar way as benthic macroorganisms in a salinity gradient.


## MATERIALS & METHODS

### Field sampling

The transect river-lagoon-open sea, i.e., from freshwater to marine, chosen for the present study is naturally occurring at Amvrakikos Gulf (Ionian Sea, Western Greece) ([Fig. 1](#)). The Gulf is a semi-enclosed embayment connected with the Ionian Sea via a narrow channel, the Preveza (Aktio) Strait ([Kapsimalis et al., 2005](#)) and is characterized by a fjord-like oceanographic regime ([Ferentinos et al., 2010](#)). The wetlands of Amvrakikos Gulf are listed in both the Ramsar international convention and the Natura 2000 Network.

The northwest part of the Amvrakikos Gulf is formed by the rivers Arachthos and Louros ([Poulos, Collins & Ke, 1993](#); [Poulos, Collins & Lykousis, 1995](#); [Vasileiadou et al., 2012](#)). Arachthos river is the main source of freshwater from November to April ([Poulos, Collins & Ke, 1993](#)) and its flow is controlled by two hydroelectric dams: the first being an earthfill dam, of 107 m height, mainly built for flood control and the second being a concrete dam, of 42 m height, primarily having a regulatory role, that is to ensure the constant flow of water throughout the year.

Two stations were chosen at the Arachthos river, thus representing the freshwater conditions: one being close to its mouth (39.029690N, 21.025880E) and one in the upper limit of saltwater intrusion when the dams are closed, close to the village of Neochori (39.070770N, 21.025060E). One more station was chosen at the Arachthos delta (39.039199N, 21.094200E), as an intermediate between the freshwater and the marine realm. In addition, two stations were chosen in the Logarou lagoon due to its vicinity to Arachthos, representing the brackish stations: one at the inner part of the lagoon (39.045528N, 20.902283E), closer to the terrestrial end, and one near the



**Figure 1** Map showing the location of the sampling stations. Map showing the location of the six sampling stations and the location of the river dams (Map data: ©2017 Google).

channel connecting the lagoon to the Gulf (39.037458N, 20.878626E). Finally, Kalamitsi station (38.966250N, 20.690783E) was chosen from the marine realm, i.e., outside of the Amvrakikos Gulf.

Sampling was carried out in winter of 2014 (November–December), as described in *Pavlouidi et al. (2016)*. For the Logarou lagoon and Arachthos river stations, salinity, water temperature and dissolved oxygen concentration were measured in the water overlaying the sediments by means of a portable multi-parameter (WTW Multi 3420 SET G). For the Arachthos delta and Kalamitsi station, water abiotic variables (fluorescence, salinity, water temperature and dissolved oxygen concentration) were recorded with a Sea-Bird Electronics 25 CTD probe. Fluorescence was regarded as a proxy for chlorophyll-a concentration.

Sediment samples from the Logarou lagoon and the Arachthos river were collected by means of a modified manually-operated box-corer, with a sampling surface of 156.25 square centimeters and sediment penetration depth of 25 centimeters, deployed from fishing boats specifically used at the lagoons. Samples from the Arachthos delta and Kalamitsi stations were collected with a Smith McIntyre Grab operated from the R/V *Philia*. The permission to conduct the field study was provided by the Amvrakikos Wetlands Management Body.

Cylindrical sampling corers (internal sampling surface 15.90 square centimeters) were placed inside the box-corer and the Smith McIntyre Grab in order to collect sub-samples of the sediment's upper layer (0–2 cm). Three replicate units, each retrieved from a different box-corer to avoid pseudoreplication, were taken for each analysis from each sampling station, in order to determine variability within and between stations. Samples for molecular analysis (about 15 cm<sup>3</sup> each), i.e., DNA extractions, were placed in 50 ml

falcon tubes (Sarstedt, Nümbrecht, Germany) and were stored at  $-20^{\circ}\text{C}$ , until further processing in the laboratory.

Samples were also collected from cylindrical corers for the measurement of the Particulate Organic Carbon (POC), chloroplast pigments concentration (chlorophyll-a and phaeopigments) and sediment granulometry (for the latter, the sampling depth was four centimeters to allow comparison with previously published data on the study area). To quantify chloroplast pigments concentration in the water column, three replicate water samples were collected from the Logarou lagoon and the Arachthos river by means of Niskin bottles (5 lt). The aforementioned samples were processed at the Chemistry Lab of the IMBBC (HCMR), based on standard techniques (chloroplast pigments: [Yentsch & Menzel, 1963](#); POC: [Hedges & Stern, 1984](#); granulometry: [Gray & Elliott, 2009](#)).

### DNA extraction, PCR amplification and 16S rRNA sequencing

DNA was extracted using the PowerSoil DNA Isolation Kit (MO BIO Laboratories, Inc., Carlsbad, CA, USA), as recommended by the manufacturer. About  $0.5 (\pm 0.2)$  grams of wet sediment were used from each sample and the quality of the extracted DNA was evaluated by gel electrophoresis.

PCR amplification was performed targeting the V3–V4 region of the 16S rRNA gene using the bacterial primer pair S-DBact-0341-b-S-17 (or 341F) and S-D-Bact-0785-a-A-21-B (or 805RB), which has been referred to as the most promising primer pair ([Herlemann et al., 2011](#); [Klindworth et al., 2013](#)), with a revision for detection of SAR11 bacterioplankton ([Apprill et al., 2015](#)).

The Two-Step PCR Approach was used for this study. The first-step PCR was performed with the aforementioned primers containing a universal 5' tail as specified in the Nextera library protocol from Illumina. The amplification reaction mix of the first PCR contained  $6\ \mu\text{l}$  5x KAPA HiFi Fidelity buffer,  $0.9\ \mu\text{l}$  BSA ( $2\ \mu\text{g}/\mu\text{l}$ ),  $0.75\ \mu\text{l}$  KAPA dNTP Mix ( $10\ \text{mM}$ ),  $1.5\ \mu\text{l}$  from each primer ( $10\ \mu\text{M}$ ),  $0.75\ \mu\text{l}$  KAPA HiFi HotStart DNA polymerase ( $1\ \text{U}/\mu\text{l}$ ) in a final volume of  $30\ \mu\text{l}$  per reaction. DNA template concentration was about  $10\ \text{ng}/\mu\text{l}$ . The first PCR protocol used was the following:  $95^{\circ}\text{C}$  for 5 min; 25 cycles at  $98^{\circ}\text{C}$  for 20 s,  $57^{\circ}\text{C}$  for 2 min,  $72^{\circ}\text{C}$  for 1 min;  $72^{\circ}\text{C}$  for 7 min.

The resulting PCR amplicons ( $\sim 531\ \text{bp}$ ) were purified using Agencourt AMPure XP magnetic beads (Beckman Coulter, Brea, CA, USA), quantified using Qubit fluorometric quantitation (Thermo Scientific Fisher, Waltham, MA, USA) and were used as templates for the second-step PCR in order to include the indexes (barcodes), as well as the Illumina adaptors. The amplification reaction mix of the second PCR contained  $6\ \mu\text{l}$  5x KAPA HiFi Fidelity buffer,  $0.75\ \mu\text{l}$  KAPA dNTP Mix ( $10\ \text{mM}$ ),  $3\ \mu\text{l}$  from each primer ( $10\ \mu\text{M}$ ),  $0.75\ \mu\text{l}$  KAPA HiFi HotStart DNA polymerase ( $1\ \text{U}/\mu\text{l}$ ) in a final volume of  $30\ \mu\text{l}$  per reaction. DNA template concentration was about  $20\ \text{ng}/\mu\text{l}$ . The second PCR protocol used was the following:  $95^{\circ}\text{C}$  for 3 min; 8 cycles at  $98^{\circ}\text{C}$  for 20 s,  $55^{\circ}\text{C}$  for 30 s,  $72^{\circ}\text{C}$  for 30 s;  $72^{\circ}\text{C}$  for 5 min. Amplifications were carried out using T100 Thermal Cycler (BIORAD, Hercules, CA, USA). Again, the resulting PCR amplicons ( $\sim 600\ \text{bp}$ ) were purified and quantified as mentioned previously, mixed in equimolar amounts and sequenced using a MiSeq Reagent Kit v3 ( $2 \times 300$ -cycles) at the IMBBC (HCMR).

All the raw sequence files of this study were submitted to the European Nucleotide Archive (ENA) (Leinonen et al., 2011) with the study accession number PRJEB20211 (available at <http://www.ebi.ac.uk/ena/data/view/PRJEB20211>).

## Data analyses

The raw sequence reads retrieved from all the sediment samples were quality trimmed using *sickle* (Joshi & Fass, 2011), to where the average quality score dropped below 20 ( $-q\ 20$ ) as well as where read length was below 10 bp ( $-l\ 10$ ). *SPAdes assembler* (Bankevich et al., 2012), which incorporates *BayesHammer* (Nikolenko, Korobeynikov & Alekseyev, 2013), was used for the creation of error-corrected paired-end reads, since this strategy along with overlapping paired-end reads reduces errors for MiSeq (Schirmer et al., 2015).

Afterwards, *pandaseq* (Masella et al., 2012) was used to overlap the paired-end reads using a minimum overlap of 20 ( $-o\ 20$ ). The overlapped sequences were combined and dereplicated. Then, using *USEARCH* (Edgar, 2010), reads were sorted based on abundance, singletons were discarded and OTU clustering and *de novo* chimera removal were performed. Following the relevant recommendation, a reference-based chimera filtering step was performed using *UCHIME* (Edgar et al., 2011) using the “Gold” database as a reference.

Reads, including singletons, were then mapped back to OTUs, using the 97% similarity threshold level. Afterwards, they were aligned using *MAFFT* (Kato et al., 2005) and a phylogenetic tree was created using *FastTree* (Price, Dehal & Arkin, 2010). Finally, taxonomic profiles of the OTUs were generated using *RDP classifier* (Wang et al., 2007).

The metabolic function of the OTUs was predicted using the *Tax4Fun* package (Aßhauer et al., 2015), which transforms the SILVA-based OTUs into a taxonomic profile of KEGG organisms ( $fctProfiling = T$ ), normalized by the 16S rRNA copy number ( $normCopyNo = T$ ). The method for pre-computing the functional reference profiles was the ultrafast protein classification tool (Meinicke, 2015) ( $refProfile = UProC$ ) and the functional reference profiles were computed based 400 bp reads ( $shortReadMode = F$ ). The 100 most abundant metabolic pathways were extracted.

The number of sequences assigned to an OTU represented its relative abundance at a given replicate sample of each sampling station. A matrix containing the microbial OTUs as variables and sampling stations as samples was constructed. A second matrix was constructed, with the predicted metabolic functions of the microbial OTUs as variables and sampling stations as samples. Both matrices were subsequently used to calculate triangular similarity matrices using the Bray–Curtis similarity coefficient (e.g., Clarke & Warwick, 1994). In order to test whether the microbial community pattern and the metabolic function pattern could be differentiated based on the sampled habitat, we performed non-metric multidimensional scaling (nMDS) (Clarke, 1993) and permutational multivariate analysis of variance (PERMANOVA) (Anderson, 2001). The design considered two factors: “habitat” and “location” (999 permutations), with the latter being nested in the former.

A third matrix was also constructed, with the respective abiotic parameters as variables and the sampling stations as samples, which was normalized and used as an input for the BIO-ENV analysis (Clarke & Ainsworth, 1993) by employing the Spearman’s rank

coefficient. The analysis was performed to identify the subsets of abiotic parameters that were associated with the community (i.e., OTU) and the metabolic function matrices. In order to account for factor confounding, and estimate the amount of variation each factor might explain in the community as well the metabolic function similarity matrices, variation partitioning analysis was performed. The amount of variation in both matrices that was due to salinity uniquely, while taking the other environmental factors into consideration, was estimated and its significance was tested using distance-based redundancy analysis (db-RDA) and Analysis of Variance (ANOVA).

A suite of diversity indices (Margalef's species richness, Pielou's evenness, Shannon-Wiener (Pielou, 1969), Chao-1, Abundance Coverage Estimator (ACE)) was calculated for each sampling station. The indices were subsequently tested for significant differences between the different locations and habitats using the Kruskal–Wallis test. The Mann–Whitney *U* test (Mann & Whitney, 1947) was used for the *post-hoc* pairwise comparisons; a Bonferroni-correction was applied and the level of significance for the results was lowered to 0.008, in the case of the locations, and to 0.017, in the case of the habitats.

Linear regression was used to examine the significance of relationships between OTU diversity, i.e., the average number of OTUs as well as the average values of diversity indices, and salinity. Shapiro–Wilk test was used to assess normality in the residuals (Shapiro & Wilk, 1965), while homoscedastic residual variances were confirmed by examining plots of the standardized residuals (Draper & Smith, 1981).


The non-parametric Kruskal–Wallis test (Kruskal & Wallis, 1952) was used to determine which of the predicted metabolic pathways were statistically significant between the different habitats, i.e., “lagoon”, “river” and “sea”. The *p* value threshold was adjusted from the initial 0.05, using the Benjamini–Hochberg procedure (Benjamini & Hochberg, 1995). In addition, the relative abundance values of the main microbial taxa were compared among the locations and habitats of the sampling stations, using the Kruskal–Wallis test.

The *vegan* package (Oksanen et al., 2008) was used for the nMDS (*metaMDS* function), PERMANOVA (*adonis* function), BIO-ENV (*bioenv* function), variation partitioning analysis (*varpart* function) and db-RDA (*capscale* function), for the calculation of diversity indices and for the generation of rarefaction curves. Groups in the nMDS plots were displayed using the *ordiellipse* and *veganCovEllipse* functions. Linear regressions, Shapiro–Wilk, Mann–Whitney, Kruskal–Wallis and ANOVA tests were conducted using *stats* package (R Core Team, 2015). Graphs were constructed using the *ggplot2* package (Wickham, 2009). The aforementioned analyses were performed using R version 3.2.1 (R Core Team, 2015).

## RESULTS

### Microbial community composition

The results of the processing of the sequences are shown in Table S1. The 1,893,500 overlapped reads were dereplicated (1,578,523 remained) and singletons were removed (111,223 remained). After *de novo* chimera removal (7,088 OTUs) and chimera removal using the “Gold” database as a reference, the final number of OTUs was 7,050. The corresponding rarefaction curve is shown in Fig. S1.




**Figure 2** nMDS of the microbial OTUs. nMDS of the similarity matrix of the sampling stations based on the microbial OTUs abundances. Ellipses according to habitat, signs according to location. AR, Arachthos; ARO, Arachthos Neochori; ARDelta, Arachthos Delta; LOin, Logarou station inside the lagoon; LOout, Logarou station in the channel connecting the lagoon to the gulf; Kal, Kalamitsi.

The nMDS of the microbial OTUs (Fig. 2) showed that their spatial pattern differs both by habitat and location, which was also confirmed by the PERMANOVA results (habitat: F.Model = 19.416,  $p < 0.01$ ; location: F.Model = 10.647,  $p < 0.01$ ).

The relative abundance percentages of the microbial taxa, at the phylum level, did not show a significant differentiation between the different locations or habitats (Kruskal–Wallis:  $p > 0.05$  for all cases). However, as shown in Fig. 3 where the relative abundance percentages of each replicate sample have been averaged per sampling station, there are certain differences that can be observed. For example, the Bacteroidetes showed an increasing abundance when moving from the inner station of the river (~9%) to the more brackish stations (~28%), and decrease again in the marine station (~11%). A similar trend was observed for the Proteobacteria, but in this case the higher abundance was found in the marine environment (~49%). The abundance of Archaea was quite low in all the





**Figure 3** Bar chart showing the abundances of the main microbial taxa, at the phylum level, at the sampling stations. AR, Arachthos; ARO, Arachthos Neochori; ARDelta, Arachthos Delta; LOin, Logarou station inside the lagoon; LOout, Logarou station in the channel connecting the lagoon to the gulf; Kal, Kalamitsi.

sampling stations; higher values were observed at the inner station ( $\sim 17\%$ ) of the river and decrease towards the marine station. Cyanobacteria/Chloroplasts were more abundant in the marine station ( $\sim 10\%$ ) and were also found in the Arachthos delta and inner station of Logarou lagoon at lower abundance ( $\sim 5\%$ ).

When the most abundant phyla are examined in more detail, again using the relative abundance percentages of each replicate sample averaged per sampling station, there are certain differences observed at the sampling stations, although non-significant (Kruskal–Wallis:  $p > 0.05$  for all the cases). In the case of Bacteroidetes (Fig. S2), Flavobacteria were less abundant at the inner station of the river ( $\sim 17\%$ ); their abundance was higher in the other riverine stations ( $\sim 37\%$ ) and in the lagoonal stations ( $\sim 54\%$ ) while remaining stable at the marine station ( $\sim 49\%$ ). Spingobacteria exhibited the lower abundance in the Arachthos delta ( $\sim 7\%$ ) and the highest in Kalamitsi ( $\sim 28\%$ ).

Regarding the Proteobacteria phylum (Fig. S3), Alphaproteobacteria exhibited higher abundances at the mouth of the Arachthos river ( $\sim 49\%$ ) and at the marine station ( $\sim 56\%$ ). Betaproteobacteria were almost exclusively present in the inner station and mouth of the Arachthos river ( $\sim 30\%$  and  $\sim 11\%$  respectively). Gammaproteobacteria were mostly abundant in the lagoonal stations ( $\sim 54\%$ ) and Deltaproteobacteria in the Arachthos delta ( $\sim 49\%$ ), which was the station with the lowest oxygen concentration in the water overlaying the sediment (0.24 mg/l; Table S6).

Diversity measured as Shannon-Wiener, Pielou's evenness and Margalef's species richness indices (Table 1) was significantly different between the locations and habitats

**Table 1** Diversity indices of the samples.

	OTUs	N	H'(ln)	J'	d	Chao-1	ACE
R_AR_A	2,414	51,617	6.64	0.85	222.36	2,771.31	2,736.48
R_AR_B	3,070	74,057	7	0.87	273.71	3,450.95	3,353.58
R_AR_C	2,810	87,640	6.77	0.85	246.82	3,145.4	3,109.94
R_ARO_A	2,271	49,793	6.66	0.86	209.88	2,643.01	2,570.08
R_ARO_B	2,419	77,520	6.66	0.86	214.78	2,625.88	2,582.98
R_ARO_C	2,480	74,292	6.59	0.84	221.03	2,773.28	2,717.73
R_ARDelta_A	2,098	42,365	6.66	0.87	196.83	2,388.94	2,319.51
R_ARDelta_B	2,479	49,442	6.61	0.85	229.26	2,821.49	2,737.81
R_ARDelta_C	2,247	38,209	6.57	0.85	212.87	2,614.22	2,565.73
L_LOin_A	2,510	70,514	6.39	0.82	224.75	2,849.83	2,770.58
L_LOin_B	2,082	64,400	5.9	0.77	187.94	2,445.6	2,399.07
L_LOin_C	2,237	57,953	6.16	0.8	203.88	2,567.4	2,532.17
L_LOout_A	1,850	60,250	5.66	0.75	168	2,271.05	2,200.13
L_LOout_B	2,487	80,942	6.38	0.82	219.97	2,841.38	2,766.75
L_LOout_C	2,088	83,935	5.95	0.78	184.07	2,365.67	2,323.44
S_Kal_A	1,732	52,975	5.9	0.79	159.13	1,906.93	1,903.36
S_Kal_B	2,004	48,280	6.40	0.84	185.72	2,278.12	2,237.63
S_Kal_C	1,974	51,129	6.34	0.84	181.98	2,211.46	2,182.35

**Notes.**


OTUs, total number of OTUs; N, total microbial relative abundance values; H', Shannon-Wiener; J', Pielou's evenness; d, Margalef's species richness; ACE, Abundance Coverage Estimator; R, River; L, Lagoon; S, Sea; AR, Arachthos; ARO, Arachthos Neochori; ARDelta, Arachthos Delta; LOin, Logarou station inside the lagoon; LOout, Logarou station in the channel connecting the lagoon to the gulf; Kal, Kalamitsi; A, B, C, replicate samples.

(Kruskal–Wallis test, [Table S2](#)). In addition, the number of OTUs, Chao-1 and ACE were also significantly different between the different habitats (Kruskal–Wallis test, [Table S2](#)). The results of the values of the Mann–Whitney *U* tests for the *post-hoc* comparisons ([Table S3](#)) show that the main driver for the difference between the habitats is the difference of the riverine and the marine environment and secondly the difference of the former with the lagoonal environment.


### Functional community composition

The retrieved KEGG metabolic profiles, and their abundance in each sample, are provided in [Table S4](#). For certain microbial OTUs a KEGG profile could not be retrieved, thus these OTUs constitute the fraction of unexplained taxonomic units (FTU) ([Aßhauer et al., 2015](#)), i.e., the amount of sequences assigned to a taxonomic unit and not transferable to KEGG reference organisms. As shown in [Fig. S4](#) and [Table S5](#), this fraction was highest in the lagoonal samples (67.51%) and lowest in the marine samples (38.91%). From the riverine samples (56.85%), the highest FTU was observed at the Arachthos Delta (71.32%). Due to the aforementioned FTU values, the interpretation of the results should be done cautiously as there are many OTUs for which retrieval of a metabolic profile was not possible.

However, the metabolic profiles retrieved from the OTUs were significantly different between the three habitats (Kruskal–Wallis,  $p < 0.05$ ). Specifically, as shown in [Fig. 4](#), there were 13 metabolic pathways that were responsible for the between-habitat dissimilarity.



**Figure 4** The metabolic pathways that were significantly different between the three habitats.



**Figure 5** nMDS of the KEGG metabolic profiles. nMDS of the similarity matrix of the sampling stations based on the abundances of KEGG metabolic profiles. Ellipses according to habitat, signs according to location. AR, Arachthos; ARO, Arachthos Neochori; ARDelta, Arachthos Delta; LOin, Logarou station inside the lagoon; LOout, Logarou station in the channel connecting the lagoon to the gulf; Kal, Kalamitsi.

In the nMDS of the KEGG metabolic profiles (Fig. 5), it is depicted that the different habitats were functionally distinctive. This was supported by the PERMANOVA results (habitat: F.Model = 10.743,  $p < 0.01$ ; location: F.Model = 11.206,  $p < 0.01$ ).

### Correlation with abiotic parameters

The physicochemical variables of the sampling stations are given in Table S6. According to the results shown on Table 2, the abiotic variable that was best correlated with the microbial community pattern is salinity ( $\rho = 0.88$ ). When the metabolic function pattern is considered, oxygen concentration was the variable showing the highest correlation with the former ( $\rho = 0.73$ ), although the combination of salinity and oxygen concentration was also highly correlated with the metabolic pattern ( $\rho = 0.63$ ). In addition, as shown by the variation partitioning analysis for all the combinations physicochemical variables of that resulted in models with adjusted  $R^2$  of residuals less than 0.60 (Table 3), the combination of salinity, POC and temperature explained 56% of the total variation in the

**Table 2** The environmental variables best correlated with the community and the metabolic function pattern.

	Water				Sediment				$\rho$	
	Salinity (psu)	O <sub>2</sub> (mg/l)	Chl-a	Temperature (°C)	POC (ug/g)	Sand (%)	Silt & Clay (%)	Phaeopigments (ug/g)		CPE (ug/g)
Community pattern	+									0.88
	+				+					0.85
	+	+			+					0.79
	+	+			+		+			0.72
	+	+			+		+		+	0.66
	+	+		+	+		+	+		0.64
	+	+		+	+	+	+	+		0.58
	+	+	+	+	+	+	+	+		0.54
Metabolic function pattern		+								0.73
	+	+								0.63
		+		+			+			0.57
	+	+				+	+			0.54

**Notes.**

POC, Particulate Organic Carbon; CPE, Chloroplasic pigment equivalents; Chl-a, Chlorophyll-a concentration (ug/l) and Fluorescence;  $\rho$ , Spearman rank correlation coefficient.

community similarity matrix. When the explanatory variables were regarded separately, salinity accounted for 27% of the variation, followed by POC (21%) and temperature (14%). In metabolic function pattern, 50% of the variation was explained by salinity and oxygen concentration; salinity alone accounted for 12% of the variation while oxygen concentration for 33%.

Regarding the relationship between the number of OTUs and the salinity values, as shown in Fig. 6, a linear decrease of the former was observed from the freshwater to the marine stations; salinity explained over 65% of the variation in the number of OTUs. Robust models were also evident when the data were divided into taxonomic groups (Table S7), with salinity explaining from 70% up to 98% of the variation of the OTUs diversity, in the case of Firmicutes and TM7 respectively. The residuals of all significant models presented in Table S7 showed no evidence of heteroscedasticity and were found to be normally distributed.

## DISCUSSION

### Microbial community composition

Based on the results of the present study, it is suggested that the microbial community diversity pattern differs by habitat and location, thus indicating that, at least to some extent, each habitat hosts a different microbial community from the others, regarding both the number of OTUs and their composition, as has been also shown from similar studies in the Baltic Sea (Herlemann et al., 2011). Furthermore, the abiotic variable that is best correlated with the microbial community pattern is salinity, as it has been shown from studies on bacterioplankton (e.g., Kirchman et al., 2005; Nemergut et al., 2011; Fortunato et al., 2012),

**Table 3** The percentage of variation explained (adjusted  $R^2$ ) of each explanatory physicochemical variable, as well as their combinations.

	Community pattern	Metabolic function pattern
Salinity	27% <sup>***</sup>	12% <sup>+</sup>
O <sub>2</sub>		33% <sup>**</sup>
Temperature	14% <sup>**</sup>	
POC	21% <sup>***</sup>	
Salinity + O <sub>2</sub>		50% <sup>***</sup> (residuals: 50%)
Salinity + Temperature	47% <sup>***</sup> (residuals: 53%)	
Salinity + POC	45% <sup>***</sup> (residuals: 55%)	
POC + Temperature	36% <sup>***</sup>	
Salinity + POC + Temperature	56% <sup>***</sup> (residuals: 44%)	
Salinity with Temperature as condition variable	32% <sup>***</sup>	
Salinity with POC as condition variable	24% <sup>***</sup>	
Salinity with O <sub>2</sub> as condition variable		16% <sup>**</sup>
Salinity with Temperature and POC as condition variables	20% <sup>***</sup>	
Temperature with Salinity as condition variable	20% <sup>***</sup>	
Temperature with POC as condition variable	15% <sup>***</sup>	
Temperature with Salinity and POC as condition variables	11% <sup>**</sup>	
POC with Salinity as condition variable	18% <sup>***</sup>	
POC with Temperature as condition variable	22% <sup>***</sup>	
POC with Salinity and Temperature as condition variables	10% <sup>**</sup>	
O <sub>2</sub> with Salinity as condition variable		37% <sup>***</sup>

**Notes.**


POC, Particulate Organic Carbon.

\* $p < 0.05$ .\*\* $p < 0.01$ .\*\*\* $p < 0.001$ .

as well as sediment bacterial communities (Bolhuis & Stal, 2011; Severin, Confurius-Guns & Stal, 2012; Bolhuis, Fillinger & Stal, 2013; Pavlouidi et al., 2016).

The majority of the observed OTUs was classified as Bacteroidetes and Proteobacteria. Bacteroidetes have been found to be omnipresent along estuarine gradients (e.g., Bouvier & Del Giorgio, 2002), although in certain cases they dominated higher salinity communities (Campbell & Kirchman, 2013; Dupont et al., 2014) which has been attributed to their ability for degradation of complex organic matter (Blümel, Süling & Imhoff, 2007). In the present study, their abundance indeed increased in the brackish stations but decreased in the marine station, which could be due to the lower availability of organic matter in the latter, as reflected in the concentration of particulate organic carbon.

Alphaproteobacteria dominate marine communities (Edmonds et al., 2009; Campbell & Kirchman, 2013; Herlemann et al., 2016) but they are also ubiquitous in freshwater



**Figure 6** Linear regression between the number of OTUs (averaged per sampling station) and the salinity of the sampling stations. Color according to habitat. AR, Arachthos; ARO, Arachthos Neochori; ARDelta, Arachthos Delta; LOin, Logarou station inside the lagoon; LOout, Logarou station in the channel connecting the lagoon to the gulf; Kal, Kalamitsi.

habitats (Zhang *et al.*, 2014), which justifies their high abundance in both the mouth of the Arachthos river and the Kalamitsi stations. Betaproteobacteria were almost exclusively present in the inner station and mouth of the Arachthos river; this complies with the general trend of Betaproteobacteria dominating freshwater habitats (Campbell & Kirchman, 2013; Zhang *et al.*, 2014) and declining with increasing salinity (Wu *et al.*, 2006). Their absence from the other sampling stations can be explained by the fact that they are typical freshwater bacteria (e.g., De Bie *et al.*, 2001), adapted to live in low salt concentrations and low osmotic pressure (Zhang *et al.*, 2014).

Gammaproteobacteria are generally more abundant in higher salinities (Wu *et al.*, 2006; Zhang *et al.*, 2014; Herlemann *et al.*, 2016), which has been attributed to their opportunistic life strategies (Pinhassi & Berman, 2003) and the low salinity conditions being unfavorable for their growth (Zhang *et al.*, 2014). However, Gammaproteobacteria can dominate brackish habitats (Edmonds *et al.*, 2009) as has also been shown from previous studies in the same sampling stations (Pavlouidi *et al.*, 2016) and from the results of the present study.

Deltaproteobacteria were mostly abundant in the Arachthos delta; this can be attributed to the hypoxic conditions prevailing in this sampling station (Crump *et al.*, 2007) and to their generally documented abundance in brackish habitats (Edmonds *et al.*, 2009; Pavlouidi *et al.*, 2016).

The low abundance of Archaea could be attributed to the primers used for the present study, which were not specific for amplification of the Archaeal communities. In addition, the low abundances found can be traced to their inability to grow under estuarine environmental conditions since they are primarily of allochthonous origin (Bouvier & Del Giorgio, 2002). However, the abundances of Archaea were decreasing with increasing salinity; in the marine environment, Archaea are generally limited to shallow or deep-sea

anaerobic sediments and extreme environments (DeLong, 1992), which could explain their absence from the marine station in the present study.

It is evident that sea level rise, and the subsequent saltwater intrusion that it will cause, will impose an osmotic stress in microbial populations in the riverine stations (Chambers, Reddy & Osborne, 2011). This may eventually shift the community in a more brackish state, although this will depend on the time-scale and intensity of saltwater intrusion. Microbial communities in fluctuating environments have been shown to be rather resilient, which causes variations in their expected functional response to change (Hawkes & Keitt, 2015). However, saltwater intrusion may induce an indirect effect in microbial communities, for example by causing changes in vegetation (Nelson et al., 2015) or by increasing sulfate concentrations, thus by stimulating its reduction (Chambers, Reddy & Osborne, 2011).

As far as the relationship between the number of OTUs and salinity values is concerned, it has been shown that it is negative, i.e., diversity decreases with the increase of salinity. Thus, the sediment microbial communities in the Amvrakikos Gulf salinity gradient do not appear to follow Remane's concept but rather the linear model proposed by Attrill (2002) with the species minimum at the point of maximum salinity range. This is in contrast with the constant relationship observed by Herlemann et al. (2011) for the Baltic Sea bacterioplankton along the salinity gradient. The divergence from the species-minimum concept could be attributed to the different life strategies of micro- and macroorganisms, as has been suggested by Telesh, Schubert & Skarlato (2013). In addition, microorganisms can be transported with water movement, apart from experiencing adaptation only at the molecular and cellular level (Telesh, Schubert & Skarlato, 2015). Thus, they experience salinity stress differently from benthic animals with reduced mobility (Skarlato & Telesh, 2017), which causes their deviation from the recognized models for macrobenthic organisms.

Although a decreasing trend of microbial diversity along gradients of increasing salinity has been observed (Rodriguez-Valera et al., 1985; Benlloch et al., 2002), the results of our study showed that this was only evident for the total number of OTUs; the other diversity indices did not show a statistically significant negative relationship with salinity (data not shown).

### Functional community composition

The higher number of FTU observed in the lagoonal samples is indicative of the high microbial community diversity of this habitat. It also reflects the need for more extensive studies in order for an enhanced taxonomic resolution to be achieved for sequences found in lagoons. Similarly, the lower FTU was found in the marine samples, thus reflecting the number of studies that have been conducted in the marine environment so far which allowed for more OTUs to be transferred to KEGG reference organisms. Analogous studies in Chesapeake Bay, which is also a large water body with a pronounced salinity gradient, have also suggested that protein identification in environmental samples is rather challenging. Sequence databases have derived from cultured organisms and thus, it is unlikely that significant matches can be found between the bacterioplankton metaproteome and the queried databases (Kan et al., 2005). Hence, more studies on isolation and cultivation of



microorganisms from these habitats are needed in order to enrich the available information on KEGG and similar databases.

Although retrieval of a metabolic profile was not possible for the majority of the OTUs in certain cases, it is shown that different habitats were functionally distinctive and that salinity and oxygen concentration were highly correlated with the retrieved metabolic pattern. This is in accordance with studies suggesting that the actual patterns of composition and metabolism transition are strongly linked to hydrological conditions (*Bouvier & Del Giorgio, 2002*). Furthermore, salinity has been found to be the main factor explaining almost all differences in the key metabolic capabilities of the Baltic Sea bacterial communities (*Dupont et al., 2014*), so a similar relationship could be expected for the Mediterranean analogue of the Baltic Sea. In addition, salinity has been shown to influence proteome profiles from bacterioplankton communities of the Chesapeake Bay, since samples with higher salinity, i.e., closer to marine origin, are more similar and, at the same time distinct from, inner Bay samples (*Kan et al., 2005*). Similar results have been found for the San Francisco Bay, where salinity is one of the key variables influencing the abundance and activity of ammonia-oxidizing prokaryotes and denitrifiers (*Mosier & Francis, 2008; Mosier & Francis, 2010*).

## CONCLUSIONS

From the results of the present study, it can be concluded that the sediment microbial OTUs of the Amvrakikos Gulf salinity gradient do not follow the Remane's concept, i.e., there is no decrease in the intermediate salinities, but rather a negative trend. In addition, different taxonomic groups were more abundant in the freshwater stations while others were more abundant in the marine environment. Salinity was also found to influence the metabolic function patterns that were retrieved for the sampling stations. However, future studies are needed to decipher the metabolic capabilities of all the OTUs found at the habitats under study and investigate in depth the impact of salinity at their functional potential. Furthermore, experimental studies are needed in order to directly examine the effect of salinity on microbial community composition and investigate how the latter will respond when subjected to salinities varying from freshwater to marine in the light of climate change and sea level rise.

## ACKNOWLEDGEMENTS

The authors would like to thank the Amvrakikos Wetlands Management Body (<http://www.amvrakikos.eu/>), and especially Mr Spyridon Konstas and Mr Demetrios Barellos, for providing access to the studied lagoons and for their valuable support during our sampling campaign. In addition, the authors would like to thank the captain and crew of the R/V *Philia* and Dr. Manolis Tsapakis for providing the samples and CTD data from the Arachthos delta and the Kalamitsi station. Also, the authors would like to thank the members of the Biodiversity lab of HCMR, and specifically Dr. Georgios Chatzigeorgiou, Dr. Eva Chatzinikolaou, Ms. Kleoniki Keklikoglou and Dr. Katerina Vasileiadou for their valuable assistance during the sampling campaign.

Furthermore, the authors would like to thank Dr. Umer Zeeshan Ijaz for providing access to the Orion IT cluster, where the analyses were performed, and for his bioinformatics tutorial entitled “Illumina Amplicons OTU Construction with Noise Removal” (available at <http://www.tinyurl.com/JCBioinformatics>). This work is part of a MARES Doctoral Research (MARES\_12\_13). MARES is a Joint Doctorate programme selected under Erasmus Mundus coordinated by Ghent University (FPA 2011-0016) (<http://www.mares-eu.org/>).

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the LifeWatchGreece (European Strategy Forum on Research Infrastructures) project (<http://www.lifewatchgreece.eu/>) [384676-94/GSRT/NSRF(C&E)] and the EU BON (Building the European Biodiversity Observation Network) project (<http://www.eubon.eu/>), funded by the European Union under the 7th Framework programme, Contract No. 308454. There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

LifeWatchGreece (European Strategy Forum on Research Infrastructures) project: 384676-94/GSRT/NSRF(C&E).

EU BON (Building the European Biodiversity Observation Network) project.  
European Union under the 7th Framework programme: 308454.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Christina Pavloudi conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Jon B. Kristoffersen and Anastasis Oulas contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Marleen De Troch reviewed drafts of the paper.
- Christos Arvanitidis conceived and designed the experiments, reviewed drafts of the paper.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

The permission to conduct the field study was provided by the Amvrakikos Wetlands Management Body (<http://www.amvrakikos.eu/>).

## DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

All the raw sequence files of this study were submitted to the European Nucleotide Archive (ENA) with the study accession number PRJEB20211 (Available at <http://www.ebi.ac.uk/ena/data/view/PRJEB20211>).

## Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied in the [Tables S1–S7](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3687#supplemental-information>.

## REFERENCES

- Anastasakis G, Piper DJW, Tziavos C. 2007.** Sedimentological response to neotectonics and sea-level change in a delta-fed, complex graben: gulf of Amvrakikos, western Greece. *Marine Geology* **236**:27–44 DOI [10.1016/j.margeo.2006.09.014](https://doi.org/10.1016/j.margeo.2006.09.014).
- Anderson MJ. 2001.** A new method for non-parametric multivariate analysis of variance. *Austral Ecology* **26**:32–46 DOI [10.1111/j.1442-9993.2001.01070.pp.x](https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x).
- Apprill A, McNally S, Parsons R, Weber L. 2015.** Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology* **75**:129–137 DOI [10.3354/ame01753](https://doi.org/10.3354/ame01753).
- Aßhauer KP, Wemheuer B, Daniel R, Meinicke P. 2015.** Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data. *Bioinformatics* **31**:2882–2884 DOI [10.1093/bioinformatics/btv287](https://doi.org/10.1093/bioinformatics/btv287).
- Attrill MJ. 2002.** A testable linear model for diversity trends in estuaries. *Journal of Animal Ecology* **71**:262–269 DOI [10.1046/j.1365-2656.2002.00593.x](https://doi.org/10.1046/j.1365-2656.2002.00593.x).
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012.** SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* **19**:455–477 DOI [10.1089/cmb.2012.0021](https://doi.org/10.1089/cmb.2012.0021).
- Barcina I, Lebaron P, Vives-Rego J. 2006.** Survival of allochthonous bacteria in aquatic systems: a biological approach. *FEMS Microbiology Ecology* **23**:1–9 DOI [10.1111/j.1574-6941.1997.tb00385.x](https://doi.org/10.1111/j.1574-6941.1997.tb00385.x).
- Ben-Dov E, Brenner A, Kushmaro A. 2007.** Quantification of sulfate-reducing bacteria in industrial wastewater, by real-time polymerase chain reaction (PCR) using *dsrA* and *apsA* genes. *Microbial Ecology* **54**:439–451 DOI [10.1007/s00248-007-9233-2](https://doi.org/10.1007/s00248-007-9233-2).
- Benjamini Y, Hochberg Y. 1995.** Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**:289–300.

- Benlloch S, Lopez-Lopez A, Casamayor EO, Ovreas L, Goddard V, Daae FL, Smerdon G, Massana R, Joint I, Thingstad F, Pedros-Alio C, Rodriguez-Valera F. 2002.** Prokaryotic genetic diversity throughout the salinity gradient of a coastal solar saltern. *Environmental Microbiology* **4**:349–360  
[DOI 10.1046/j.1462-2920.2002.00306.x](https://doi.org/10.1046/j.1462-2920.2002.00306.x).
- Blümel M, Süling J, Imhoff JF. 2007.** Depth-specific distribution of Bacteroidetes in the oligotrophic Eastern Mediterranean Sea. *Aquatic Microbial Ecology* **46**:209–224  
[DOI 10.3354/ame046209](https://doi.org/10.3354/ame046209).
- Bobrova O, Kristoffersen JB, Oulas A, Ivanytsia V. 2016.** Metagenomic 16s rRNA investigation of microbial communities in the Black Sea estuaries in South-West of Ukraine. *Acta Biochimica Polonica* **63**:315–319 [DOI 10.18388/abp.2015\\_1145](https://doi.org/10.18388/abp.2015_1145).
- Bolhuis H, Fillinger L, Stal LJ. 2013.** Coastal microbial mat diversity along a natural salinity gradient. *PLOS ONE* **8**:e63166 [DOI 10.1371/journal.pone.0063166](https://doi.org/10.1371/journal.pone.0063166).
- Bolhuis H, Stal LJ. 2011.** Analysis of bacterial and archaeal diversity in coastal microbial mats using massive parallel 16S rRNA gene tag sequencing. *The ISME Journal* **5**:1701–1712 [DOI 10.1038/ismej.2011.52](https://doi.org/10.1038/ismej.2011.52).
- Bouvier TC, Del Giorgio PA. 2002.** Compositional changes in free-living bacterial communities along a salinity gradient in two temperate estuaries. *Limnology and Oceanography* **47**:453–470 [DOI 10.4319/lo.2002.47.2.0453](https://doi.org/10.4319/lo.2002.47.2.0453).
- Campbell BJ, Kirchman DL. 2013.** Bacterial diversity, community structure and potential growth rates along an estuarine salinity gradient. *The ISME Journal* **7**:210–220  
[DOI 10.1038/ismej.2012.93](https://doi.org/10.1038/ismej.2012.93).
- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N, Knight R. 2011.** Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences of the United States of America* **108**(Suppl. 1):4516–4522 [DOI 10.1073/pnas.1000080107](https://doi.org/10.1073/pnas.1000080107).
- Chambers LG, Reddy KR, Osborne TZ. 2011.** Short-term response of carbon cycling to salinity pulses in a freshwater wetland. *Soil Science Society of America Journal* **75**:2000–2007 [DOI 10.2136/sssaj2011.0026](https://doi.org/10.2136/sssaj2011.0026).
- Clarke KR. 1993.** Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* **18**:117–143  
[DOI 10.1111/j.1442-9993.1993.tb00438.x](https://doi.org/10.1111/j.1442-9993.1993.tb00438.x).
- Clarke KR, Ainsworth M. 1993.** A method of linking multivariate community structure to environmental variables. *Marine Ecology-Progress Series* **92**:205–219  
[DOI 10.3354/meps092205](https://doi.org/10.3354/meps092205).
- Clarke KR, Warwick R. 1994.** Similarity-based testing for community pattern: the two-way layout with no replication. *Marine Biology* **118**:167–176  
[DOI 10.1007/BF00699231](https://doi.org/10.1007/BF00699231).
- Crump BC, Peranteau C, Beckingham B, Cornwell JC. 2007.** Respiratory succession and community succession of bacterioplankton in seasonally anoxic estuarine waters. *Applied and Environmental Microbiology* **73**:6802–6810 [DOI 10.1128/AEM.00648-07](https://doi.org/10.1128/AEM.00648-07).
- Darr A, Gogina M, Zettler ML. 2014.** Functional changes in benthic communities along a salinity gradient—a western Baltic case study. *Journal of Sea Research* **85**:315–324  
[DOI 10.1016/j.seares.2013.06.003](https://doi.org/10.1016/j.seares.2013.06.003).

- De Bie MJM, Speksnijder AGCL, Kowalchuk GA, Schuurman T, Zwart G, Stephen JR, Diekmann OE, Laanbroek HJ. 2001.** Shifts in the dominant populations of ammonia-oxidizing  $\beta$  subclass Proteobacteria along the eutrophic Schelde estuary. *Aquatic Microbial Ecology* **23**:225–236 DOI [10.3354/ame023225](https://doi.org/10.3354/ame023225).
- DeLong EF. 1992.** Archaea in coastal marine environments. *Proceedings of the National Academy of Sciences of the United States of America* **89**:5685–5689 DOI [10.1073/PNAS.89.12.5685](https://doi.org/10.1073/PNAS.89.12.5685).
- Draper NR, Smith H. 1981.** *Applied regression analysis*. New York: John Wiley & Sons.
- Dupont CL, Larsson J, Yooseph S, Ininbergs K, Goll J, Asplund-Samuelsson J, McCrow JP, Celepli N, Allen LZ, Ekman M, Lucas AJ, Hagström Å, Thiagarajan M, Brindefalk B, Richter AR, Andersson AF, Tenney A, Lundin D, Tovchigrechko A, Nylander JAA, Bami D, Badger JH, Allen AE, Rusch DB, Hoffman J, Norrby E, Friedman R, Pinhassi J, Venter JC, Bergman B. 2014.** Functional tradeoffs underpin salinity-driven divergence in microbial community composition. *PLOS ONE* **9**:e89549 DOI [10.1371/journal.pone.0089549](https://doi.org/10.1371/journal.pone.0089549).
- Edgar RC. 2010.** Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**:2460–2461 DOI [10.1093/bioinformatics/btq461](https://doi.org/10.1093/bioinformatics/btq461).
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011.** UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27**:2194–2200 DOI [10.1093/bioinformatics/btr381](https://doi.org/10.1093/bioinformatics/btr381).
- Edmonds JW, Weston NB, Joye SB, Mou X, Moran MA. 2009.** Microbial community response to seawater amendment in low-salinity tidal sediments. *Microbial Ecology* **58**:558–568 DOI [10.1007/s00248-009-9556-2](https://doi.org/10.1007/s00248-009-9556-2).
- Falkowski PG, Fenchel T, DeLong EF. 2008.** The microbial engines that drive earth's biogeochemical cycles. *Science* **320**:1034–1039 DOI [10.1126/science.1153213](https://doi.org/10.1126/science.1153213).
- Ferentinos G, Papatheodorou G, Geraga M, Iatrou M, Fakiris E, Christodoulou D, Dimitriou E, Koutsikopoulos C. 2010.** Fjord water circulation patterns and dysoxic/anoxic conditions in a Mediterranean semi-enclosed embayment in the Amvrakikos Gulf, Greece. *Estuarine, Coastal and Shelf Science* **88**:473–481 DOI [10.1016/j.ecss.2010.05.006](https://doi.org/10.1016/j.ecss.2010.05.006).
- Fierer N, McCain CMCM, Meir P, Zimmermann M, Rapp JMJM, Silman MRMR, Knight R. 2011.** Microbes do not follow the elevational diversity patterns of plants and animals. *Ecology* **92**:797–804 DOI [10.1890/10-1170.1](https://doi.org/10.1890/10-1170.1).
- Fortunato CS, Herfort L, Zuber P, Baptista AM, Crump BC. 2012.** Spatial variability overwhelms seasonal patterns in bacterioplankton communities across a river to ocean gradient. *The ISME Journal* **6**:554–563 DOI [10.1038/ismej.2011.135](https://doi.org/10.1038/ismej.2011.135).
- Gray JS, Elliott M. 2009.** *Ecology of marine sediments. From science to management*. Oxford: Oxford University Press.
- Hawkes CV, Keitt TH. 2015.** Resilience vs. historical contingency in microbial responses to environmental change. *Ecology Letters* **18**:612–625 DOI [10.1111/ele.12451](https://doi.org/10.1111/ele.12451).
- Hedges J, Stern J. 1984.** Carbon and nitrogen determinations of carbonate-containing solids. *Limnology and Oceanography* **29**:657–663 DOI [10.4319/lo.1984.29.3.0657](https://doi.org/10.4319/lo.1984.29.3.0657).

- Herlemann DPR, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF. 2011.** Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *The ISME Journal* 5:1571–1579 DOI 10.1038/ismej.2011.41.
- Herlemann DPR, Lundin D, Andersson AF, Labrenz M, Jürgens K. 2016.** Phylogenetic signals of salinity and season in bacterial community composition across the salinity gradient of the baltic sea. *Frontiers in Microbiology* 7:1–13 DOI 10.3389/fmicb.2016.01883.
- Highton MP, Roosa S, Crawshaw J, Schallenberg M, Morales SE. 2016.** Physical factors correlate to microbial community structure and nitrogen cycling gene abundance in a nitrate fed eutrophic lagoon. *Frontiers in Microbiology* 7:Article 1691 DOI 10.3389/fmicb.2016.01691.
- Jiang X-T, Peng X, Deng G-H, Sheng H-F, Wang Y, Zhou H-W, Tam NF-Y. 2013.** Illumina sequencing of 16s rRNA tag revealed spatial variations of bacterial communities in a mangrove wetland. *Microbial Ecology* 66:96–104 DOI 10.1007/s00248-013-0238-8.
- Joshi N, Fass J. 2011.** Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files. Version 1.33. Available at <https://github.com/najosh>.
- Kan J, Hanson TE, Ginter JM, Wang K, Chen F. 2005.** Metaproteomic analysis of Chesapeake Bay microbial communities. *Saline Systems* 1:Article 7 DOI 10.1186/1746-1448-1-7.
- Kapsimalis V, Pavlakis P, Poulos SE, Alexandri S, Tziavos C, Sioulas A, Filippas D, Lykousis V. 2005.** Internal structure and evolution of the Late Quaternary sequence in a shallow embayment: the Amvrakikos Gulf, NW Greece. *Marine Geology* 222–223:399–418 DOI 10.1016/j.margeo.2005.06.008.
- Katoh K, Kuma K, Toh H, Miyata T. 2005.** MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* 33:511–518 DOI 10.1093/nar/gki198.
- Kirchman DL, Dittel AI, Malmstrom RR, Cottrell MT. 2005.** Biogeography of major bacterial groups in the Delaware estuary. *Limnology and Oceanography* 50:1697–1706 DOI 10.4319/lo.2005.50.5.1697.
- Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO. 2013.** Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research* 41:e1 DOI 10.1093/nar/gks808.
- Kruskal WH, Wallis WA. 1952.** Use of ranks in one-criterion variance analysis. *Journal of the American Statistical Association* 47:583–621 DOI 10.1080/01621459.1952.10483441.
- Leinonen R, Akhtar R, Birney E, Bower L, Cerdeno-Tárraga A, Cheng Y, Clel I, Faruque N, Goodgame N, Gibson R, Hoad G, Jang M, Pakseresht N, Plaister S, Radhakrishnan R, Reddy K, Sobhany S, Ten Hoopen P, Vaughan R, Zalunin V, Cochrane G. 2011.** The european nucleotide archive. *Nucleic Acids Research* 39:D28–D31 DOI 10.1093/nar/gkq967.

- Logares R, Bråte J, Bertilsson S, Clasen JL, Shalchian-Tabrizi K, Rengefors K. 2009.** Infrequent marine–freshwater transitions in the microbial world. *Trends in Microbiology* **17**:414–422 DOI [10.1016/j.tim.2009.05.010](https://doi.org/10.1016/j.tim.2009.05.010).
- Lozupone CA, Knight R. 2007.** Global patterns in bacterial diversity. *Proceedings of the National Academy of Sciences of the United States of America* **104**:11436–11440 DOI [10.1073/pnas.0611525104](https://doi.org/10.1073/pnas.0611525104).
- Mann HB, Whitney DR. 1947.** On a test of whether one of two random variables is stochastically larger than the other. *The Annals of Mathematical Statistics* **18**:50–60 DOI [10.1214/aoms/1177730491](https://doi.org/10.1214/aoms/1177730491).
- Masella AP, Bartram AK, Truszkowski JM, Brown DG, Neufeld JD. 2012.** PAN-DAseq: paired-end assembler for illumina sequences. *BMC Bioinformatics* **13**:31 DOI [10.1186/1471-2105-13-31](https://doi.org/10.1186/1471-2105-13-31).
- McLusky D, Elliott M. 2004.** *The estuarine ecosystem: ecology, threats and management*. Oxford: Oxford University Press.
- Meinicke P. 2015.** UProC: tools for ultra-fast protein domain classification. *Bioinformatics* **31**:1382–1388 DOI [10.1093/bioinformatics/btu843](https://doi.org/10.1093/bioinformatics/btu843).
- Mosier AC, Francis CA. 2008.** Relative abundance and diversity of ammonia-oxidizing archaea and bacteria in the San Francisco Bay estuary. *Environmental Microbiology* **10**:3002–3016 DOI [10.1111/j.1462-2920.2008.01764.x](https://doi.org/10.1111/j.1462-2920.2008.01764.x).
- Mosier AC, Francis CA. 2010.** Denitrifier abundance and activity across the San Francisco Bay estuary. *Environmental Microbiology Reports* **2**:667–676 DOI [10.1111/j.1758-2229.2010.00156.x](https://doi.org/10.1111/j.1758-2229.2010.00156.x).
- Nelson TM, Streten C, Gibb KS, Chariton AA. 2015.** Saltwater intrusion history shapes the response of bacterial communities upon rehydration. *Science of the Total Environment* **502**:143–148 DOI [10.1016/j.scitotenv.2014.08.109](https://doi.org/10.1016/j.scitotenv.2014.08.109).
- Nemergut DR, Costello EK, Hamady M, Lozupone C, Jiang L, Schmidt SK, Fierer N, Townsend AR, Cleveland CC, Stanish L, Knight R. 2011.** Global patterns in the biogeography of bacterial taxa. *Environmental Microbiology* **13**:135–144 DOI [10.1111/j.1462-2920.2010.02315.x](https://doi.org/10.1111/j.1462-2920.2010.02315.x).
- Neubauer SC, Franklin RB, Berrier DJ. 2013.** Saltwater intrusion into tidal freshwater marshes alters the biogeochemical processing of organic carbon. *Biogeosciences* **10**:8171–8183 DOI [10.5194/bg-10-8171-2013](https://doi.org/10.5194/bg-10-8171-2013).
- Nikolenko SI, Korobeynikov AI, Alekseyev MA. 2013.** BayesHammer: bayesian clustering for error correction in single-cell sequencing. *BMC Genomics* **14**:S7 DOI [10.1186/1471-2164-14-S1-S7](https://doi.org/10.1186/1471-2164-14-S1-S7).
- Oksanen J, Kindt R, Legendre P, O’Hara B, Simpson GL, Solymos P, Stevens MHH, Wagner H. 2008.** The vegan package. 10:2008. Available at <https://CRAN.R-project.org/package=vegan>.
- Oulas A, Pavloudi C, Polymenakou P, Pavlopoulos GA, Papanikolaou N, Kotoulas G, Arvanitidis C, Iliopoulos I. 2015.** Metagenomics: tools and insights for analyzing next-generation sequencing data derived from biodiversity studies. *Bioinformatics and Biology Insights* **9**:75–88 DOI [10.4137/BBI.S12462](https://doi.org/10.4137/BBI.S12462).

- Painchaud J, Therriault J, Legendre L. 1995.** Assessment of salinity-related mortality of freshwater bacteria in the Saint Lawrence estuary. *Applied and Environmental Microbiology* **61**:205–208.
- Palmer TA, Montagna PA, Pollack JB, Kalke RD, DeYoe HR. 2011.** The role of freshwater inflow in lagoons, rivers, and bays. *Hydrobiologia* **667**:49–67 DOI [10.1007/s10750-011-0637-0](https://doi.org/10.1007/s10750-011-0637-0).
- Pavlouidi C, Oulas A, Vasileiadou K, Sarropoulou E, Kotoulas G, Arvanitidis C. 2016.** Salinity is the major factor influencing the sediment bacterial communities in a Mediterranean lagoonal complex (Amvrakikos Gulf, Ionian Sea). *Marine Genomics* **28**:71–81 DOI [10.1016/j.margen.2016.01.005](https://doi.org/10.1016/j.margen.2016.01.005).
- Pielou EC. 1969.** *An introduction to mathematical ecology*. New York: Wiley-Interscience.
- Pinhassi J, Berman T. 2003.** Differential growth response of colony-forming alpha- and gamma-proteobacteria in dilution culture and nutrient addition experiments from Lake Kinneret (Israel), the eastern Mediterranean Sea, and the Gulf of Eilat. *Applied and Environmental Microbiology* **69**:199–211 DOI [10.1128/AEM.69.1.199-211.2003](https://doi.org/10.1128/AEM.69.1.199-211.2003).
- Poulos S, Collins MB, Ke X. 1993.** Fluvial/wave interaction controls on delta formation for ephemeral rivers discharging into microtidal waters. *Geo-Marine Letters* **13**:24–31 DOI [10.1007/BF01204389](https://doi.org/10.1007/BF01204389).
- Poulos SE, Collins MB, Lykousis V. 1995.** Late quaternary evolution of amvrakikos gulf, western greece. *Geo-Marine Letters* **15**:9–16 DOI [10.1007/BF01204492](https://doi.org/10.1007/BF01204492).
- Price MN, Dehal PS, Arkin AP. 2010.** FastTree 2—approximately maximum-likelihood trees for large alignments. *PLOS ONE* **5**:e9490 DOI [10.1371/journal.pone.0009490](https://doi.org/10.1371/journal.pone.0009490).
- R Core Team. 2015.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
- Remane A. 1934.** Die brackwasserfauna. In: *Verhandlungen der Deutschen Zoologischen Gesellschaft*. Leipzig: Deutschen Zoologischen Gesellschaft im Frankfurt am Main, 34–74.
- Rodriguez-Valera F, Ventosa A, Juez G, Imhoff JF. 1985.** Variation of environmental features and microbial populations with salt concentrations in a multi-pond saltern. *Microbial Ecology* **11**:107–115 DOI [10.1007/BF02010483](https://doi.org/10.1007/BF02010483).
- Rolston A, Dittmann S. 2009.** *The distribution and abundance of macrobenthic invertebrates in the murray mouth and coorong lagoons 2006 to 2008*. Canberra: CSIRO: Water for a Healthy Country National Research Flagship, 89.
- Schirmer M, Ijaz UZ, D'Amore R, Hall N, Sloan WT, Quince C. 2015.** Insight into biases and sequencing errors for amplicon sequencing with the Illumina MiSeq platform. *Nucleic Acids Research* **43**:e37–e37 DOI [10.1093/nar/gku1341](https://doi.org/10.1093/nar/gku1341).
- Severin I, Confurius-Guns V, Stal LJ. 2012.** Effect of salinity on nitrogenase activity and composition of the active diazotrophic community in intertidal microbial mats. *Archives of Microbiology* **194**:483–491 DOI [10.1007/s00203-011-0787-5](https://doi.org/10.1007/s00203-011-0787-5).
- Shapiro SS, Wilk MB. 1965.** An analysis of variance test for normality (complete samples). *Biometrika* **52**:591–611 DOI [10.1093/biomet/52.3-4.591](https://doi.org/10.1093/biomet/52.3-4.591).
- Skarlato SO, Telesh IV. 2017.** The development of the protistan species-maximum concept for the critical salinity zone. *Russian Journal of Marine Biology* **43**:1–11 DOI [10.1134/S1063074017010114](https://doi.org/10.1134/S1063074017010114).



- Staley C, Unno T, Gould TJ, Jarvis B, Phillips J, Cotner JB, Sadowsky MJ. 2013.** Application of Illumina next-generation sequencing to characterize the bacterial community of the Upper Mississippi River. *Journal of Applied Microbiology* **115**:1147–1158 DOI [10.1111/jam.12323](https://doi.org/10.1111/jam.12323).
- Telesh I, Schubert H, Skarlato S. 2011.** Revisiting Remane’s concept: evidence for high plankton diversity and a protistan species maximum in the horohaliniacum of the Baltic Sea. *Marine Ecology Progress Series* **421**:1–11 DOI [10.3354/meps08928](https://doi.org/10.3354/meps08928).
- Telesh I, Schubert H, Skarlato S. 2013.** Life in the salinity gradient: discovering mechanisms behind a new biodiversity pattern. *Estuarine, Coastal and Shelf Science* **135**:317–327 DOI [10.1016/j.ecss.2013.10.013](https://doi.org/10.1016/j.ecss.2013.10.013).
- Telesh I, Schubert H, Skarlato S. 2015.** Size, seasonality, or salinity: what drives the protistan species maximum in the horohaliniacum? *Estuarine, Coastal and Shelf Science* **161**:102–111 DOI [10.1016/j.ecss.2015.05.003](https://doi.org/10.1016/j.ecss.2015.05.003).
- Vasileiadou K, Sarropoulou E, Tsigenopoulos C, Reizopoulou S, Nikolaidou A, Orfanidis S, Simboura N, Kotoulas G, Arvanitidis C. 2012.** Genetic vs community diversity patterns of macrobenthic species: preliminary results from the lagoonal ecosystem. *Transitional Waters Bulletin* **6**:20–33 DOI [10.1285/i1825229Xv6n2p20](https://doi.org/10.1285/i1825229Xv6n2p20).
- Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007.** Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* **73**:5261–5267 DOI [10.1128/AEM.00062-07](https://doi.org/10.1128/AEM.00062-07).
- Wang Y, Sheng H-F, He Y, Wu J-Y, Jiang Y-X, Tam NF-Y, Zhou H-W. 2012.** Comparison of the levels of bacterial diversity in freshwater, intertidal wetland, and marine sediments by using millions of illumina tags. *Applied and Environmental Microbiology* **78**:8264–8271 DOI [10.1128/AEM.01821-12](https://doi.org/10.1128/AEM.01821-12).
- Wickham H. 2009.** *ggplot2: elegant graphics for data analysis*. New York: Springer.
- Wu QL, Zwart G, Schauer M, Kamst-van Agterveld MP, Hahn MW. 2006.** Bacterioplankton community composition along a salinity gradient of sixteen high-mountain lakes located on the Tibetan Plateau, China. *Applied and Environmental Microbiology* **72**:5478–5485 DOI [10.1128/AEM.00767-06](https://doi.org/10.1128/AEM.00767-06).
- Yentsch CS, Menzel DW. 1963.** A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. *Deep Sea Research and Oceanographic Abstracts* **10**:221–231 DOI [10.1016/0011-7471\(63\)90358-9](https://doi.org/10.1016/0011-7471(63)90358-9).
- Zhang L, Gao G, Tang X, Shao K. 2014.** Can the freshwater bacterial communities shift to the “marine-like” taxa? *Journal of Basic Microbiology* **54**:1264–1272 DOI [10.1002/jobm.201300818](https://doi.org/10.1002/jobm.201300818).
- Zhang J, Yang Y, Zhao L, Li Y, Xie S, Liu Y. 2015.** Distribution of sediment bacterial and archaeal communities in plateau freshwater lakes. *Applied Microbiology and Biotechnology* **99**:3291–3302 DOI [10.1007/s00253-014-6262-x](https://doi.org/10.1007/s00253-014-6262-x).
- Zinger L, Amaral-Zettler LA, Fuhrman JA, Horner-Devine MC, Huse SM, Welch DBM, Martiny JBH, Sogin M, Boetius A, Ramette A. 2011.** Global patterns of bacterial beta-diversity in seafloor and seawater ecosystems. *PLOS ONE* **6**:e24570 DOI [10.1371/journal.pone.0024570](https://doi.org/10.1371/journal.pone.0024570).