

Genetic characterization of Yug Bogdanovac virus

Martin Pfeffer · Meik Dilcher · Robert B. Tesh ·
Frank T. Hufert · Manfred Weidmann

Received: 3 July 2012 / Accepted: 5 September 2012 / Published online: 14 October 2012
© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract We present pyrosequencing data and phylogenetic analysis for the full genome of Yug Bogdanovac virus (YBV), a member of the Vesicular stomatitis virus serogroup of the Rhabdoviridae isolated from a pool of *Phlebotomus perfliewi* sandflies collected in Serbia in 1976. YBV shows very low nucleotide identities to other members of the Vesicular stomatitis virus serogroup and does not contain a reading frame for C'/C proteins.

Keywords Vesiculovirus · Yug Bogdanovac virus · Chandipura virus · Isfahan virus · 454 Pyrosequencing

Yug Bogdanovac virus (YBV) was isolated in 1976 from a pool of 200 unengorged female *Phlebotomus perfliewi* sandflies collected in Serbia. Electron microscopy and serological analysis (complement fixation test, immunofluorescence, plaque reduction neutralization assay) placed it into the Vesicular stomatitis virus (VSV) serogroup of the Rhabdoviridae. Antibodies against YBV were found in humans (6/274 tested) [1] and domestic animals [1–3] but the role of YBV as a human pathogen is unclear. An

antigenic relationship to Chandipura virus (CHDV) isolated in India and West Africa [4–7] and to Isfahan virus (ISFV) isolated in Iran and Turkmenia [8, 9], two members of the VSV serogroup implicated as causes of febrile and neurological diseases (CHDV) in humans [10–12] was described. In order to determine the genome sequence of this European vesiculovirus, YBV was passaged thrice in Vero B4 cells and RNA extraction was performed as described [13]. In order to cover the termini, a self-complementary 3'-FLAC adapter and a 5'-RACE adapter were ligated to the genome prior to pyrosequencing [14]. A MID-barcoded Roche/454 Rapid Library was produced from 300 ng adapter-ligated viral genomic RNA following reverse transcription at 65 °C for 30 min (Transcriptor (Roche)). 82 % of 11,970 reads were YBV genome specific (coverage 294-fold). The 11,202 nucleotide –ssRNA genome shows the typical genomic organization of vesiculoviruses including 3'-leader followed by 5 structural genes, and 5'-trailer sequence (Fig. 1a, GenBank JF911700). The full-length genome of YBV virus shows very low nucleotide identities to the genomes of CHDV (55.7 %), ISFV (55.3 %, [15]), VSIV (52.8 %), and of rabies virus (41.0 %) (Fig. 1b). Nevertheless, the high bootstrap values of the phylogenetic analysis confirm serological grouping of YBV within the genus *Vesiculovirus* of rhabdoviruses (Fig. 1b). In YBV, the C'/C proteins, two small highly basic, non-structural proteins encoded in a second ORF within the P gene of most vesiculoviruses [16, 17] are absent as in Alagoas virus (VSAV) [18]. Both 3'-leader and 5'-positive-sense antigenomic trailer of YBV are highly complementary to each other in the first 32 nucleotides (Fig. 1c, [19]). The genomic data presented here will help to design a YBV-specific RT-PCR which could be used to monitor YBV activity in phlebotomine sandflies in the Balkans.

M. Pfeffer
Institute of Animal Hygiene and Veterinary Public Health,
An den Tierkliniken 1, 04103 Leipzig, Germany

M. Dilcher · F. T. Hufert · M. Weidmann (✉)
Department of Virology, University Medical Center Goettingen,
Kreuzberggring 57, 37075 Goettingen, Germany
e-mail: mweidma@gwdg.de

R. B. Tesh
Department of Pathology, Center for Biodefense and Emerging
Infectious Diseases, University of Texas Medical Branch,
Galveston, TX, USA

