

Meta-analysis of 0 to 8 h post-prandial evolution of ruminal pH

C. Dragomir^{1,4}, D. Sauvant^{2†}, J.-L. Peyraud³, S. Giger-Reverdin² and B. Michalet-Doreau¹

¹Unité de Recherches sur les Herbivores, INRA, 63122, St-Genès Champanelle, France; ²UMR INRA INA P-G, Physiologie de la Nutrition et Alimentation, 16 rue Claude Bernard, 75231 Paris Cedex 05, France; ³UMR INRA/ENSAR Production du Lait, 35590 St Gilles, France; ⁴National R-D Institute for Animal Biology and Nutrition, Calea Bucuresti 1, Balotesti, Ilfov 077015, Romania

(Received 6 December 2006; Accepted 21 April 2008; First published online 8 July 2008)

The objective of this study was to identify relevant descriptors of ruminal pH post-prandial evolution that can replace the mean pH (considered unsatisfactory). These descriptors are to be used in the attempts to predict ruminal pH from dietary characteristics, in order to quantify the potential of a diet to induce subacute ruminal acidosis from its intrinsic characteristics. A total of 219 pH curves, reported as graphics in 48 published articles describing the post-prandial evolution of ruminal pH (first 8 h), were digitized by image analysis then summarized in 15 pH variables. Relationships among pH variables and the principal components (PCs) of pH variability were analyzed in order to identify possible alternatives to mean pH, as the average value of all pH data the curve is composed of. Two groups of pH variables were identified according to their relationship with the most important principal components. A first group, including mean pH, was closely related to PC₁, which accounted for 78% of data variability; hence, correlations between variables of this group were generally high. Of these, threshold-related variables were distinct as their within-study correlations with mean pH were rather moderate (0.69 on average). This suggests they might carry supplementary information that could explain the variation in ruminal pH induced by within-study factors, e.g. diet characteristics. However, caution should be taken in their use because of their truncation at 0 h and their non-normal distribution. Variables from the second group were independent of the PC₁, and thus of the first group of variables, whereas they were mostly related to PC₂ and PC₃. This implies they are complementary to mean pH. Of this second group, the rate of pH decreases or the time period when pH reaches its minimum might be useful to better describe the ruminal status, from the point of view of the risk of subacute ruminal acidosis.

Keywords: meta-analysis, pH-dynamics, rumen, subacute rumen acidosis

Introduction

The quantity of concentrates fed to ruminants is often a compromise between the objective to fully express their productive potential and the concern of avoiding digestive disorders, such as ruminal acidosis. Decrease of pH below a safety threshold value leads to negative effects on rumen fermentation. It may significantly reduce cellulolytic activity in the rumen. A very strong decrease of pH can also trigger the development of ruminal acidosis. According to the severity of this disorder, two forms are distinguished (Owens *et al.*, 1998; Oetzel, 2003): acute and subacute rumen acidosis. The latter one is less obvious (although affecting many animals and causing massive economic losses) and difficult to diagnose; therefore, its occurrence has to be predicted, e.g. from diet characteristics.

Rumen pH is believed to be the most representative index of the ruminal fermentation status. It is commonly

reported in articles on ruminant nutrition, and its daily fluctuations is a crucial element in several published rumen models (Argyle and Baldwin, 1988; Dijkstra *et al.*, 1992; Lescoat and Sauvant, 1995; Pitt *et al.*, 1996) and its influences on the rumen ecosystem (cellulolysis, proteosynthesis, VFA profile, etc.) are well documented.

Although the most frequently reported evidence is the mean of successive during the day values, numerous publications describe pH evolution in graphics, generally within several hours after the morning meal. These graphical data give a better image of the rumen environment, as the ruminal pH is not steady during the day. However, these data cannot be used in quantitative analyses as a curve.

This has raised the problem of finding one or several variables that could accurately describe the various aspects of ruminal pH dynamics. As mean pH seems to be an unsatisfactory descriptor (Sauvant *et al.*, 1999; Kolver and de Veth, 2002), other pH variables describing diurnal variations of pH are candidate to provide useful complementary information, in particular when subacute ruminal acidosis is studied.

† E-mail: sauvant@inapg.inra.fr

Some summarizing variables, as time and area under a threshold value, are sometimes proposed (Mackie and Gilchrist, 1979; Nocek, 1997; Beauchemin *et al.*, 2001). However, to our knowledge, no attempt has been made to analyze and interpret the graphical data of pH evolution, across published data.

The objective of this study was to assess the variability of the available pH curves in order to identify and evaluate other pH variables that describe the post-prandial evolution of rumen pH.

Material and methods

Database description

The database was built from articles published from 1985 to 2002. Inclusion of an article in the database was decided on the basis of the existence of the description of pH evolution during at least 8 h after the morning meal, the major meal of the day. A minimum of four samplings within the first 8 h after the morning meal was imposed and articles where a second meal was fed within these first 8 h were excluded. Also, the various *in vivo* measurements had to be done on the same animals, which had to be already adapted to diets. Articles with unpractical feeding conditions (e.g. ruminal infusions, induced ruminal acidosis or challenge-type trials) were discarded.

All available data concerning diet characteristics (ingredients, chemical composition, digestibility, feeding schedule, ingesta, etc.) or ruminal characteristics (short-chain fatty acids and ammonia concentrations, protozoa density, etc.) were pooled in the database. Only raw data were included (no assumptions or approximations were used).

The database contained 219 treatments, representing diets that were obtained from 48 articles and 56 experiments. Data were mostly obtained on cattle (186 treatments); of these, 79% concerned dairy cows and 21% concerned growing cattle and dry cows. The rest of the data were obtained on sheep and goats. Diets were fed in one (20%), two (71%) or three (9%) meals/day. For 54 treatments, details on feeding were not reported and 12 diets consisted exclusively of concentrates or of forages; of the remaining 153 diets, 60% were fed as total mixed ration (TMR). The level of daily dry mater intake (DMI) was specified for 178 treatments: of these, 83% were fed *ad libitum* (with a maximum of 15% orts). In general, TMR were fed *ad libitum*; in the case of non-mixed rations, concentrates were limited, whereas forages were fed *ad libitum*. When fed separately, concentrates generally preceded forages.

The proportion of concentrates in diets varied from 0% to 100% of DMI. Diets usually contained conserved forages: 59% of them were based on silages, 37% on hays and only 4% on fresh forages. The most frequent silages were corn, alfalfa and grass; as hays, mainly alfalfa and grass were used. Concentrate mixtures generally consisted of barley, corn and protein meals; grains were generally included in ground, rolled, cracked or flaked form.

Calculation of pH variables derived from the original pH graphics

The 219 pH curves corresponding to the 219 treatments had different lengths and measurement intervals; they were reported in various forms and were associated with various feeding schedules. All of them were transformed in digitized scatter graphics, which were used as backgrounds in Visilog 5.4 (Noesis Inc., Crolles, France). A script in C language was used to trace virtual graph scales and the inflexion points of pH curves by following the background (clicking on the important points of the digitized graphics). On this basis, by projecting points from the pH curve to the pH axis, values for pH at sampling times were calculated. Also pH values at every 30 min were calculated using linear interpolation between the measured values. This interval was chosen as a compromise between the needed level of detailing and volume of data to be processed. Also, a prior study of the published pH graphics showed that pH was rarely measured at smaller intervals. The accuracy of pH values acquired from graphics was evaluated by comparing their means to those reported in articles, by taking into account only pH values corresponding to sampling times. Biases greater than 1%, caused by the poor quality of graphics, discarded the corresponding articles from database.

Data of pH curves were then synchronized according to the starting time of the morning meal, as the origin of graphs not always matched the feeding time. Only the values corresponding to the first 8 post-prandial hours were retained (17 values/curve, either measured or interpolated). The choice of the 8 h duration compromised between maximizing the extent of the studied period and maximizing the database size, by respecting the previously mentioned inclusion criteria.

Potential information carried by graphical representation of the pH post-prandial evolution (Figure 1) was summarized by spotting or calculating the following parameters:

- initial pH value, at feeding time (pH_0);
- final pH value, 8 h after the meal (pH_8);
- minimal pH value for the considered period (pH_{min});
- maximal pH value for the considered period (pH_{max});
- mean pH: the arithmetic average of the pH values, using both measured and interpolated values;
- standard deviation (s.d.) of pH, calculated from the same series of values as mean pH ($pH_{s.d.}$);
- area under the curve (a.u.c.), represents the whole area between the pH curve and the time axis, measured in $pH \times h$;
- amplitude of pH perturbation: $\Delta pH_{0-min} = pH_0 - pH_{min}$;
- decrease of pH, 8 h after feeding time: $\Delta pH_{0-8} = pH_0 - pH_8$;
- period when pH is less than a pH threshold, measured in h (time under threshold or $t < pH_t$);
- area between the pH threshold and the pH curve, when the pH value is less than a pH threshold, measured in $pH \times h$ (area under threshold or $a < pH_t$).

Although the proposed pH thresholds vary considerably (from 5.0 to 6.3) in the literature, only the most relevant (having biological significance) and frequently used were selected in this study: 5.5, 5.8, 6.0, 6.2. Thus, *in vitro* trials




Figure 1 Generic curve of pH and potential pH variables.

revealed that pH below certain levels would significantly impair or inhibit some rumen functions such as cellulose or microbial proteosynthesis (Shriver *et al.*, 1986; Slyter and Rumsey, 1991; Russell and Wilson, 1996). *In vivo* trials also confirmed the influence of pH on microorganisms (Mould *et al.*, 1983; Martin and Michalet-Doreau, 1995).

For calculation of time and area under thresholds, it was assumed that pH evolutions within 30 min were linear (Pereira and Armentano, 2000). Limitation of the studied period to 8 post-prandial hours caused supplementary truncation of some of the threshold-related variables, especially when the threshold was high. Thus, in our database, 3% of $t < \text{pH}_{6.0}$ values and 14% of the $t < \text{pH}_{6.2}$ values were truncated at 8 h, as an artifact linked to the limitation of the studied period. The truncated values were discarded from the analyses.

Calculation of pH variables issued from the modelling of pH graphics

As the pH curves generally exhibited a concave asymmetric shape (Figure 1), a third-degree polynomial model was regressed to all series of pH data:

$$\text{pH}_t = at^3 + bt^2 + ct + d, \text{ where } t \text{ was expressed in hours.}$$

The PROC REG procedure of SAS (SAS version 8.1, 1999) was used to fit polynomial equations to each of the pH datasets corresponding to the 219 pH graphics. All four coefficients (a , b , c , d) were taken into account, even if they were statistically not significant in the model. For each pH dataset, the number of measurements, the residual s.d. of the model and the s.d. of the four coefficients were also retained for further statistical analysis.

The four coefficients were used to calculate four other pH variables that cannot be directly derived from the observed pH graphics (Figure 1):

- rate of pH change at 0, 2:30 and 8 h, in pH units/hour (called initslope, 2:30 slope and finslope, respectively);
- time when pH reaches its minimum, measured in hours (t_{\min}).

Rates of pH change (dpH/dt) were obtained by considering $t = 0, 2.5$ or 8, respectively, in the first derivatives of polynomial equations. t_{\min} was calculated by choosing one of the two roots of derivative, depending on the sign of the first derivative. In some cases t_{\min} was outside the studied area (0 to 8 h), it was then considered equal to 8 h.

Statistical analysis

Principal component (PC) analysis was applied to the 17 post-prandial pH values in order to cluster the 219 pH graphics. Global, across- and within-study relationships between pairs of pH variables were studied. Global correlations considered the 219 observations as a whole, irrespective of the study conditions, whereas across-study considered the experiment as an explanatory factor. Pearson correlations were calculated and, when appropriate, within-study regression equations were fitted (SAS version 8.1, 1999; Minitab 13.20, 2000).

Results and discussions

Principal component analysis of pH curves

Principal component analysis allows transformation of a number of possibly correlated variables into a smaller number of uncorrelated variables, called principal components. The first principal component accounts for as much of the variability in the data as possible, and each succeeding component accounts for as much of the remaining variability as possible. This multivariate approach allowed summarizing the 17 pH values composing the pH curves ($\text{pH}_0, \text{pH}_{0:30}, \text{pH}_1, \dots, \text{pH}_8$) into few independent new variables extracting the major variations of data. The principal components with the four highest eigen values were considered for interpretation, on the basis of the main factors that characterize the pH curves: their average, slope, deepness and possible asymmetry (presence of a lag time).

The first principal component (PC_1) had the greatest weight as it accounted for 78.1% of data variability. Correlations between PC_1 and the 17 pH values composing the pH curves were high and exhibited the same sign (Table 1).

This expressed the influence of a 'size factor' and it was consistent with the fact that the 17 pH values were largely auto-correlated (data not shown). This suggests that mean pH is closely related to PC₁, as also illustrated in Figure 2a, with the graphical representation of the pH curves that were the most discriminated at the two extreme parts of this principal component (average of 25 graphics for each part). The types of curves discriminated by mean pH, which

is an expression of PC₁, confirmed that this is an indicator of the level of rumen fluid acidity.

The three other principal components were also investigated in order to establish pH variables that may be complementary to mean pH. The second principal component (PC₂) accounted for 12.5% of data variability. Data presented in Table 1 indicate that the extreme times of the pH graphics presented opposite signs of correlation with PC₂. This suggests that PC₂ mainly reflects the overall slope of pH change. This trend is illustrated in Figure 2b with the evolution of the 50 curves that were the most discriminated at the two extreme parts of the PC₂. It must be stressed that for the decreasing pH curves, pH_{min} was frequently achieved beyond 8 h, whereas, for the increasing type, pH_{min} occurred more early.

The third principal component (PC₃) accounted for 5.5% of the variability. Table 1 indicates that the initial and final phases of the pH graphs have an opposite sign with the middle phase on PC₃. This suggests that pH curves were mainly discriminated by their relative 'deepness'. Figure 2c, built from the 50 pH graphics most differentiated by PC₃, exhibited that fact with flat-shaped curves opposite to deep ones, which suggested a stronger perturbation. It has to be noticed that the flat type of curves rarely reached their minimum within 8 h.

The fourth principal component (PC₄) accounted for only 1.7% of data variability and suggested the existence of an asymmetry of curves. Indeed, graphical expression for the 50 pH curves mostly differentiated by this principal component shows the existence of two opposite groups: pH curves with an initial 'lag-time' v. pH curves that decrease immediately and fairly abruptly, quickly reaching their

Table 1 Global correlations between the principal components of variance and the 17 values composing the pH graphics

	PC ₁	PC ₂	PC ₃	PC ₄
pH ₀	0.658	0.265	0.640	-0.277
pH _{0:30}	0.777	0.424	0.443	0.019
pH ₁	0.779	0.535	0.112	0.246
pH _{1:30}	0.834	0.507	-0.039	0.168
pH ₂	0.863	0.428	-0.178	0.070
pH _{2:30}	0.910	0.316	-0.215	-0.004
pH ₃	0.930	0.204	-0.239	-0.077
pH _{3:30}	0.957	0.116	-0.210	-0.104
pH ₄	0.960	0.024	-0.185	-0.139
pH _{4:30}	0.967	-0.076	-0.136	-0.148
pH ₅	0.966	-0.171	-0.099	-0.099
pH _{5:30}	0.954	-0.260	-0.052	-0.054
pH ₆	0.920	-0.353	-0.009	0.015
pH _{6:30}	0.907	-0.397	0.028	0.057
pH ₇	0.889	-0.431	0.092	0.087
pH _{7:30}	0.863	-0.446	0.155	0.124
pH ₈	0.824	-0.468	0.213	0.149

PC₁, PC₂, ... = principal component 1, 2, ...
 pH₀, pH_{0:30}, ... = pH variables corresponding to pH at 0 h, 30 min, ...




Figure 2 Types of curves discriminated by the first four principal components of variance (PC₁ to PC₄) (one line represents 25 kinetics, corresponding to either pH maximum or minimum of the concerned principal component).

minimum pH (Figure 2d). Moreover, the final slope was higher for the pH curves that began with a lag.

Repartition of the 219 treatments on the four principal components was globally homogenous, suggesting a certain 'continuum' in the data according to the diversity of the pH graphics. To evaluate the weight of the across- and within-study variations on the determination of the components, an analysis of variance was applied on the values of the treatments on the four first principal components. The influence of the publication was highly significant for the four principal components with corresponding *F* ratios of 8.0, 10.2, 9.9 and 7.3. These comparable values obtained for the Fisher ratios suggest that the relative weight of among- and within-study was consistent from one principal component to another. Moreover, the residual variations of the four analyses of variance were normally distributed and, logically, they were not mutually correlated between the four principal components.

Based on their global correlations with principal components (Table 2), the pH variables were clustered in two groups. In Group 1, mean pH, a.u.c., pH_{\min} , pH_8 and the eight threshold-related variables are largely explained by the first principal component of variance ($0.77 < r < 1.00$); in contrast, they are weakly correlated to PC_2 , PC_3 and PC_4 ($0.00 < r < 0.47$). The variable pH_{\max} and, in the same manner, pH_0 were partly discordant with Group 1: their correlations with PC_1 are weaker than those of the above-mentioned pH variables ($r = 0.708$ and 0.658 , respectively). Moreover, they were moderately related to PC_3 ($r = 0.592$ and 0.640 , respectively).

In Group 2, variables expressing pH fluctuations ($\Delta pH_{0-\min}$, ΔpH_{0-8} and $pH_{s.d.}$) and variables issued from modelling the pH graphics (t_{\min} , initslope, 2:30 slope and finslope) were poorly correlated with PC_1 ($0.13 < r < 0.45$); on the contrary, they exhibited much higher correlations with one of the other principal components. Thus, ΔpH_{0-8} , t_{\min} and 2:30 slope were mostly related to PC_2 ($r = 0.756$; 0.752 and 0.704 , respectively), initslope, $\Delta pH_{0-\min}$ and $pH_{s.d.}$ were mostly related to PC_3 ($r = 0.822$; 0.759 and 0.682 , respectively) and the final slope was mostly related to PC_4 ($r = 0.724$).

Group 1: pH variables strongly correlated to PC_1

Mean pH (Table 3) had a relatively low average (6.13 ± 0.018) across the 219 treatments, and a normal distribution (of an Anderson–Darling index of 0.59) with a slight left asymmetry (0.19). Most authors in the database calculated arithmetic and not weighted average; when measurement times were unevenly distributed over the studied period (67% of the cases), this induced biases as also observed by Murphy (1981) and by Pitt and Pell (1997).

Our database revealed an underestimation of mean pH that was small (0.04 pH units on average) but variable (s.d. of 0.05 pH units and biases up to 0.28 pH units). Biases have little influence on within-study comparisons but induce further biases when analyzing data from different trials, e.g. in meta-analyses or rumen modelling. However, irrespective of the accuracy of its calculation, the mean pH alone does not offer information on pH pattern throughout

Table 2 Global correlations between the principal components of variance and the pH variables

Variable	Group of variables*	PC_1	PC_2	PC_3	PC_4
pH_0	1**	-0.658	0.265	0.640	-0.277
pH_8	1	-0.824	-0.468	0.213	0.149
pH_{\min}	1	-0.946	-0.078	-0.166	0.059
pH_{\max}	1**	-0.708	0.255	0.592	-0.220
Mean pH	1	-1.000	-0.013	0.010	0.001
$pH_{s.d.}$	2	0.408	0.417	0.682	-0.114
a.u.c.	1	-0.991	0.003	-0.016	0.011
$\Delta pH_{0-\min}$	2	0.373	0.323	0.759	-0.315
ΔpH_{0-8}	2	0.316	0.756	0.343	-0.415
$t < pH_{5.5}$	1	0.820	0.182	0.315	0.076
$a < pH_{5.5}$	1	0.768	0.179	0.310	0.123
$t < pH_{5.8}$	1	0.882	-0.074	-0.137	-0.034
$a < pH_{5.8}$	1	0.853	0.079	0.141	0.020
$t < pH_{6.0}$	1	0.905	0.085	-0.087	0.191
$a < pH_{6.0}$	1	0.936	0.108	0.052	0.130
$t < pH_{6.2}$	1	0.824	0.133	0.072	0.054
$a < pH_{6.2}$	1	0.979	0.089	0.066	0.104
t_{\min}	2	0.130	0.752	-0.340	-0.031
initslope	2	-0.287	0.357	-0.822	0.227
2:30 slope	2	-0.451	-0.704	-0.481	-0.191
finslope	2	0.177	0.234	0.082	0.724

a.u.c. = area under the curve.

*Group 1 = mostly related to PC_1 ; Group 2 = mostly related to PC_2 .

**Partly discordant with the rest of Group 1.

the day. Moreover, the s.d., which could have brought supplementary information, is rarely reported.

Area under the pH curve (a.u.c.) averaged 48.9 ± 0.15 pH units \times hours and showed a normal distribution, similar to mean pH (Table 3). This variable is sometimes reported as a descriptor of the ruminal pH evolution (Yang *et al.*, 2000 and 2001) but in fact it has no other value than being a slightly more precise way to express mean pH when unevenly spaced measurements are used: by calculating the a.u.c. and by dividing it to time units, a more accurate mean is obtained. As in the present study, mean pH was calculated from all pH values (at 30-min intervals) not only from measured values, the corresponding a.u.c. brought no supplementary information and was discarded from further analysis.

Of the 17 values composing the pH graphics, four were investigated as having possible significance in describing pH evolution: pH_0 , pH_{\max} , pH_{\min} and pH_8 (Table 3). Some information can be drawn when pH_{\max} occurs later than t_0 (14% of the cases in our database) or when these values are much different from pH thresholds. For example, in our database, pH_{\max} was lower than 6.2 in 34 cases, suggesting that the ruminal digestion (e.g. cellulolysis) of concerned animals was impaired. pH_{\max} and pH_{\min} are sometimes proposed besides mean pH as supplementary quantitative descriptors of the ruminal status (Beauchemin and Buchanan-Smith, 1990; Maekawa *et al.*, 2002) but these variables rely on punctual measurements. This makes them subject to bias as a result of the sampling schedule chosen.

Table 3 Variability and distribution of pH variables describing the pH graphics

pH variable	Group of variables	n**	Mean	Overall s.d.	Across-trials s.d.	Within-trial s.d.	Anderson–Darling index
pH ₀	1***	219	6.52	0.29	0.51	0.18	1.142
pH ₈	1	219	6.10	0.33	0.59	0.20	0.448*
pH _{min}	1	219	5.91	0.32	0.55	0.21	0.489*
pH _{max}	1***	219	6.53	0.28	0.49	0.18	0.920
Mean pH	1	219	6.13	0.29	0.50	0.18	0.590*
pH _{s.d.}	2	219	0.19	0.10	0.16	0.07	2.676
a.u.c.	1	219	49.0	2.33	4.13	1.45	0.615*
ΔpH _{0–min}	2	219	0.60	0.33	0.58	0.21	1.034
ΔpH _{0–8}	2	219	0.42	0.31	0.55	0.20	1.403
t < pH _{5.5}	1	24	3.15	1.90	2.22	1.48	0.374*
a < pH _{5.5}	1	24	0.37	0.37	0.42	0.31	0.975
t < pH _{5.8}	1	97	4.05	2.00	2.46	1.71	1.355
a < pH _{5.8}	1	97	0.71	0.77	0.95	0.66	5.828
t < pH _{6.0}	1	152	4.33	2.26	3.32	1.53	2.301
a < pH _{6.0}	1	160	1.11	1.12	1.57	0.85	6.312
t < pH _{6.2}	1	184	5.21	2.02	2.90	1.48	3.639
a < pH _{6.2}	1	219	1.70	1.47	2.39	1.02	6.890
t _{min}	2	219	5.12	1.93	3.51	1.17	4.321
initslope	2	219	−0.243	0.207	0.361	0.136	1.261
2 : 30 slope	2	219	−0.083	0.060	0.104	0.040	1.103
finslope	2	219	0.025	0.124	0.216	0.083	0.555*

a.u.c. = area under the curve.

*Variable with a normal distribution (Anderson–Darling test); s.d. = standard deviation.

**Where n < 219, values were calculated from non-truncated values.

***Partly discordant with the rest of Group 1.

Within Group 1, mean pH, a.u.c., pH_{min} and pH₈ have a distinct behaviour, because of their very low global, across- and within-study s.d.s (Table 3). They might be less sensitive to the influence of dietary factors affecting rumen acidity (concentrate proportions from 0% to 100%, starch content of diets from 0% to 75.7% – DM basis; NDF content from 14.7% to 65.0% – DM basis, total tract digestibility of starch from 66% to almost 100%, etc.).

‘Threshold-related’ pH variables, time under threshold and area under threshold are also highly correlated with PC₁. These variables are claimed to have higher biological significance than mean pH. However, a major limitation in the use of threshold-related variables was their truncation at 0, which occurred in our study when the threshold value was lower than pH_{min}. Throughout the 219 treatments, even for the highest of the studied thresholds (pH = 6.2), many of the corresponding values were equal to 0 (30% for t < pH_{6.2} and 15% for a < pH_{6.2}). Also, truncation at 8 h occurred as a result of the chosen period of 8 post-prandial hours. All truncated values were eliminated from the statistical analysis in this study; therefore, data in Table 3 are based on non-truncated values.

Truncation was in general accompanied by a non-normal distribution of threshold-related variables (Table 3). The cause of this distribution is that, due to the general pattern of pH, area under pH 5.5, 5.8, 6.0 and 6.2 decreased asymptotically to 0 while the mean pH became higher than the pH threshold.

Therefore, caution should be taken when using linear models for statistical analysis that includes threshold-related variables. Different from mean pH, the eight threshold-

related variables showed much higher variability, which increased with the level of threshold (Table 3).

Group 2: pH variables weakly correlated to PC₁

ΔpH_{0–min}, ΔpH_{0–8} and pH_{s.d.} showed large variability (Table 3), which could make them good candidates as descriptors of changes in rumen status when trying to relate them to diet characteristics. However, ΔpH_{0–8} and ΔpH_{0–min} have the disadvantage of being based on punctual measurements of pH; thus they are subjected to errors. In our database, pH_{s.d.} varied strongly, from 0.02 to 0.63 pH units, when only measured points, as in published trials, were used to calculate it. The utility of this variable is impaired because it carries limited information (it expresses the extent of pH variation without detecting its trend). Various authors mention differences between pH values or use the concept of pH fluctuations in commentaries (Malestein *et al.*, 1984; MacLeod *et al.*, 1994; Pitt and Pell, 1997) but these were not translated into explicit quantitative pH variables.

The variables initslope, 2 : 30 slope, finslope and t_{min}, were calculated from coefficients obtained by regression of the pH curves with a third-degree polynomial equation. The equation was chosen on the basis of the general shape of the pH graphics (Figure 1) and of the types of pH curves discriminated by the first four principal components (Figure 2). This choice compromised between the need to fit accuracy and the need to interpret the equations coefficients. Although part of the pH curves tended to be rather linear during the 8 h after the meal, the use of a complete type

Table 4 Global correlations among pH variables

pH variable	pH ₀	pH ₈	pH _{min}	pH _{max}	Mean pH	pH _{s.d.}	a.u.c.	ΔpH _{0-min}	ΔpH ₀₋₈	t < pH _{5.5}	a < pH _{5.5}	t < pH _{5.8}	a < pH _{5.8}	t < pH _{6.0}	a < pH _{6.0}	t < pH _{6.2}	a < pH _{6.2}	t _{min}	initslope	2:30 slope
pH ₈	0.519																			
pH _{min}	0.479	0.798																		
pH _{max}	0.984	0.565	0.528																	
Mean pH	0.661	0.833	0.946	0.710																
pH _{s.d.}	0.316	-0.408	-0.635	0.283	-0.406															
a.u.c.	0.638	0.809	0.939	0.689	0.990	-0.415														
ΔpH _{0-min}	0.424	-0.346	-0.592	0.359	-0.369	0.946	-0.383													
ΔpH ₀₋₈	0.329	-0.636	-0.448	0.265	-0.324	0.736	-0.317	0.765												
t < pH _{5.5}	0.072	-0.400	-0.885	0.007	-0.814	0.276	-0.849	0.294	0.355											
a < pH _{5.5}	0.066	-0.392	-0.924	0.014	-0.763	0.307	-0.794	0.298	0.344	0.945										
t < pH _{5.8}	-0.456	-0.538	-0.713	-0.467	-0.883	-0.070	-0.881	-0.068	0.054	0.696	0.598									
a < pH _{5.8}	-0.196	-0.542	-0.916	-0.212	-0.852	0.294	-0.866	0.274	0.279	0.974	0.943	0.755								
t < pH _{6.0}	-0.540	-0.651	-0.787	-0.578	-0.906	0.084	-0.905	0.049	0.181	0.392	0.284	0.753	0.547							
a < pH _{6.0}	-0.401	-0.673	-0.934	-0.438	-0.934	0.290	-0.942	0.241	0.284	0.940	0.897	0.876	0.966	0.810						
t < pH _{6.2}	-0.417	-0.575	-0.692	-0.445	-0.825	0.294	-0.836	0.263	0.273	0.554	0.435	0.568	0.400	0.756	0.593					
a < pH _{6.2}	-0.484	-0.737	-0.939	-0.523	-0.978	0.378	-0.980	0.318	0.298	0.894	0.842	0.893	0.918	0.892	0.983	0.735				
t _{min}	-0.091	-0.527	-0.166	-0.085	-0.143	0.205	-0.114	0.087	0.500	-0.270	-0.247	-0.152	-0.088	0.053	-0.017	0.146	0.042			
initslope	-0.298	-0.028	0.402	-0.232	0.275	-0.557	0.296	-0.688	-0.238	-0.278	-0.284	-0.057	-0.252	-0.063	-0.206	-0.141	-0.243	0.516		
2:30 slope	-0.154	0.540	0.546	-0.114	0.455	-0.785	0.454	-0.704	-0.735	-0.472	-0.450	-0.076	-0.349	-0.305	-0.398	-0.427	-0.452	-0.429	0.186	
finslope	-0.177	-0.023	-0.138	-0.143	-0.176	0.141	-0.180	-0.020	-0.134	0.405	0.360	0.008	0.081	0.167	0.141	0.245	0.188	0.180	0.284	-0.533

a.u.c. = area under the curve.




Figure 3 Relationships among threshold-related variables describing pH curves. (a) Time under threshold variables (periods while pH is lower than 5.5, 5.8, 6.0 and 6.2), in hours. (b) Area under threshold variables (areas between the line representing pH = 5.5, 5.8, 6.0, 6.2 and the pH curve), in hours \times pH units.

of third-degree polynomial fitting was decided in order to provide a full table of regression coefficients. Of the 219 regressions, 98% were significant; on the other hand, in 40% of the cases coefficients of t^2 and t^3 were not significant. Nevertheless, we intended to use a single approach; hence, all equations and all coefficients were retained for further calculations. Effectively, the cubic equation presented flexibility that allowed to conveniently describe the various shapes of pH curves, including the quasi-linear ones.

The initial and final slopes of pH evolution (initslope, finslope) are expressions of pH rates of decrease and recovery, respectively. However, the initial slope was insufficient to describe pH decrease: pH was not measured at t_0 in 13% of the treatments, and some of the pH curves tended to increase for a while after t_0 . Therefore, pH slope at $t_{2.5}$ (2 : 30 slope), in the middle of the t_0-t_{\min} interval, was considered more appropriate to describe pH decrease, whereas the initial slope was retained as an identifier of pH curves expressing a lag before decrease.

In the literature, the coefficients from fitting the polynomial equation were sometimes directly used for statistical comparisons of the values composing pH graphics (Leiva *et al.*, 2000). On the contrary, the rates of pH change and t_{\min} are seldom used in the literature and, to our knowledge, no attempt was made to use them in analysis of data from the literature.

The concepts of pH decrease and recovery were used, without being quantified, by several authors (Malestein *et al.*, 1984; Pitt and Pell, 1997; Leiva *et al.*, 2000), some of them suggesting possible relationships with ruminal processes. Sometimes, the time when pH reaches its minimum (t_{\min}) is briefly used in text commentaries (Khalili and Huhtanen, 1991; Krause *et al.*, 1998). Only in two recent references (Krause and Combs, 2003; Krause *et al.*, 2003) it was found as an explicit variable used to support the hypothesis on ruminal digestion. Pitt and Pell (1997) reported that t_{\min} could be reasonably predicted within the net carbohydrate and protein system (from DMI, eNDF and the concentration of organic acids in the rumen). However, it has to be mentioned that efficacy of t_{\min} is highly dependent of the frequency of pH measurements.

All variables of Group 2 ($\Delta\text{pH}_{0-\min}$, ΔpH_{0-8} , $\text{pH}_{s,d}$, initslope, 2 : 30 slope, finslope, t_{\min}) expressed large variability (Table 3). Besides, their high correlations with PC₂, PC₃ and PC₄ and low correlations with PC₁ (Table 2) make them good candidates as descriptors of changes in the rumen status.

Correlations among pH variables

As a consequence of their high correlations with PC₁, the global correlations among pH variables from Group 1 were high (Table 4). The across-study and within-study correlations showed the same trend (data not shown). The high




Figure 4 Relationships between time under threshold and area under the same threshold.

correlations between the mean pH and pH_{\min} , pH_{\max} , pH_0 or pH_8 were determined by the auto-correlation of the 17 values composing the pH graphics (as shown by their relationships with PC_1). The auto-correlative process also explains the weaker correlations between pH_{\max} (or pH_0) and the above-mentioned variables: correlations between distant values are weaker, especially at the beginning of the data series.

The global mutual correlations were systematically high among the four areas under threshold variables; they were also significant but lower for the four variables time under threshold (Table 4). This could be determined by the limitation of the studied period to 8 h, which has induced the non-linearity, more stressed for time under threshold than for the area under threshold variables (Figure 3). This is also expressed by the relationships between time and area under the same threshold (Figure 4).

Also, high global correlations linked threshold-related variables and mean pH (Table 4). Beauchemin *et al.* (2003) found much lower correlations between mean pH and time under pH 5.8 or area under pH 5.8, but they used fewer observations, from a single study. It is noteworthy that they used continuous measuring of ruminal pH, whereas our study is mostly based on measurements with time intervals

of more than 30 min. These relationships are detailed in Figure 5. The graphical encoding showed that the global, among- and within- relationships followed approximately the same trends and it also expressed the above-mentioned 'asymptotic' evolutions of the threshold-related pH variables. Generally, global correlations were higher than across-study correlations, which were higher than within-study correlations (0.89, 0.83 and 0.69, respectively, when only significant correlations were considered). This suggests that threshold-related variables may be interesting as substitutes of mean pH, especially for within-study variations (e.g. for studying feed characteristics).

Mean pH and pH_{\min} presented a similar level and pattern of correlations with threshold-related variables. On the contrary, correlations between the threshold-related variables and pH_0 , pH_8 , pH_{\max} , $\text{pH}_{\text{s.d.}}$, $\Delta\text{pH}_{0-\min}$ and ΔpH_{0-8} were moderate and even insignificant as the threshold value decreases.

Strong global, across- and within-study relationships between mean pH and other variables from Group 1 imply that mean pH can be replaced by any of them as descriptor of rumen status. Of these, threshold-related variables offer the advantage of biological significance, as the deleterious effects of the low pH depend on the duration and intensity




Figure 5 Relationships between mean pH and threshold-related parameters (values of threshold-related parameters that are equal to 0 or 8, as a result of the studied period chosen, were excluded from analyses).

of its decrease (Beauchemin, 1991; Krehbiel *et al.*, 1995; Nocek *et al.*, 2002).

Relationships among variables from Group 2 were more heterogeneous as they were related to more than one principal component. High global correlations between $pH_{s.d.}$ and the pH drop, expressed by either ΔpH_{0-min} or ΔpH_{0-8} , confirmed the fact that $pH_{s.d.}$ offers a good image of pH stability in the rumen (Table 4). In fact, the three variables behaved in the same manner regarding their correlations with the principal components (Table 2), their variability (Table 3) and their inter-correlations (Table 4).

Moderate correlations were detected between t_{min} and initial slope (0.516) or slope at 2:30 h (-0.429). A negative moderate correlation (-0.533) also linked the final slope with the slope at 2:30 h (Table 4). The other correlations between variables issued from modelling were weak ($r < 0.28$), which implies they may be complementary in expressing pH evolution.

Global correlations between ΔpH_{0-min} , ΔpH_{0-8} or $pH_{s.d.}$ and the pH slope at 2:30 h were high ($r > 0.70$); their correlations with the initial slope or t_{min} were rather inconsistent and their correlations with the final slope were weak ($r < 0.14$). However, the low correlations of the final slope with other variables may be due to the fact that PC_4 is insufficiently expressed in pH curves described by few measurements. The values of t_{min} were higher for curves presenting high values of ΔpH_{0-8} and low values of pH_8 (Table 4). The drop ΔpH_{0-min} was not related to t_{min} but was closely linked to the slopes at t_0 and $t_{2.5}$, which seems logical. These two slopes were also positively related to pH_{min} and negatively to $pH_{s.d.}$ (Table 4).

Conclusions

The relationships between pH variables and the principal components allowed splitting the pH variables that are

linked to PC₁ (Group 1) and to PC₂, or PC₃ or PC₄ (Group 2). Screening the relevant variables only, this clustered the mean pH or threshold-related variables to pH_{s.d.} and the pH calculated from the regression equations, such as the rates of pH decreasing or t_{\min} .

As the principal components are independent by definition, the two groups of pH variables may be complementary in describing post-prandial ruminal pH evolution. This is consistent with the weak correlations ($r < 0.40$) between any of the variables from Group 1 and those from Group 2, except for the slope at 2 : 30 h.

Of the variables belonging to Group 1, a.u.c., pH_{min} and pH₈ variables have lower relevance as reliable descriptors of rumen pH. On the contrary, threshold-related variables might carry supplementary information about pH dynamics induced by diet characteristics, since they are calculated only from pH values that are relevant to ruminal fermentation and their within-study correlations with mean pH are rather moderate (0.69, on average). However, in our database, which is representative of normal feeding situations, these variables were truncated and showed distributions other than normal, irrespective of the choice of pH threshold. This implies that caution should be taken when they are used as descriptors of ruminal pH evolution, e.g. in meta-analyses or rumen modelling.

The pH slope variables (at 0, 2.5 and 8 h), t_{\min} , $\Delta\text{pH}_{0-\min}$, ΔpH_{0-8} and pH_{s.d.} expressed fairly similar relationships with the principal components of variance (poorly correlated to PC₁, highly correlated to PC₂, PC₃ and PC₄). However, the pH slope variables and t_{\min} are more reliable than $\Delta\text{pH}_{0-\min}$ or ΔpH_{0-8} , the latter being determined from only two measurements. Moreover, simple differences between the critical pH values do not take time into account. Compared to pH_{s.d.}, pH slope variables and t_{\min} offered the advantage of adding information on the nature and mean of pH irregularities.

A third-degree polynomial equation was used to describe the 0 to 8 h post-prandial pH graphics of various forms. We consider that next to pH graphics, the authors should also report the four coefficients estimates derived with this model. Besides a fairly accurate reproduction of the original curves, this would allow the calculation of new pH variables (rate of pH change, t_{\min}) or recalculation of variables, which are based on punctual measurements (e.g. pH_{min}). These variables may be used in quantitative analyses of data from the literature. These new variables are poorly related to the currently used mean pH or threshold-related variables, which means they carry additional information on pH variation; this makes them good candidates as descriptors of short-term ruminal status.

Implications

The evolution of the ruminal pH in the first 8 h after the morning meal can be conveniently modelled by a third-degree polynomial equation whose coefficients allow (either directly or through derived pH variables) the quantitative comparison

of various pH graphics and quantitative reviews across published data. The analysis of variables summarizing post-prandial evolution of ruminal pH revealed that some of them could replace or complement the classical mean pH, as descriptors of short-term ruminal acidity.

References

- Argyle JL and Baldwin RL 1988. Modelling of rumen water kinetics and effects of rumen pH changes. *Journal of Dairy Science* 71, 1178–1188.
- Beauchemin KA 1991. Effects of dietary neutral detergent fiber concentration and alfalfa hay quality on chewing, rumen function and milk production of dairy cows. *Journal of Dairy Science* 74, 3140–3151.
- Beauchemin KA and Buchanan-Smith JG 1990. Effects of fiber source and method of feeding on chewing activities, digestive function and productivity of dairy cows. *Journal of Dairy Science* 73, 749–762.
- Beauchemin KA, Yang WZ and Rode LM 2001. Effects of barley grain processing on the site and extent of digestion of beef feedlot finishing diets. *Journal of Animal Science* 79, 1925–1936.
- Beauchemin KA, Yang WZ and Rode LM 2003. Effects of particle size of alfalfa-based dairy cow diets on chewing activity, ruminal fermentation, and milk production. *Journal of Dairy Science* 86, 630–643.
- Dijkstra J, Neal HDStC, Beaver DE and France J 1992. Simulation of nutrient digestion, absorption and outflow in the rumen: model description. *Journal of Nutrition* 122, 2239–2256.
- Khalili H and Huhtanen P 1991. Sucrose supplements in cattle given grass silage-based diets. 1. Digestion of organic matter and nitrogen. *Animal Feed Science and Technology* 33, 275–287.
- Kolver ES and de Veth MJ 2002. Prediction of ruminal pH from pasture-based diets. *Journal of Dairy Science* 85, 1255–1266.
- Krause KM and Combs DK 2003. Effects of forage particle size, forage source, and grain fermentability on performance and ruminal pH in midlactation cows. *Journal of Dairy Science* 86, 1382–1397.
- Krause M, Beauchemin KA, Rode LM, Farr BI and Nørgaard L 1998. Fibrolytic enzyme treatment of barley grain and source of forage in high-grain diets fed to growing cattle. *Journal of Animal Science* 76, 2912–2920.
- Krause KM, Combs DK and Beauchemin KA 2003. Effects of increasing levels of refined cornstarch in the diet of lactating dairy cows on performance and ruminal pH. *Journal of Dairy Science* 86, 1341–1353.
- Krehbiel CR, Britton RA, Harmon DL, Wester TJ and Stock RA 1995. The effects of ruminal acidosis on volatile fatty acid absorption and plasma activities of pancreatic enzymes in lambs. *Journal of Animal Science* 73, 3111–3121.
- Leiva E, Hall MB and Van Horn HH 2000. Performance of dairy cattle fed citrus pulp or corn products as sources of neutral detergent-soluble carbohydrates. *Journal of Dairy Science* 83, 2866–2875.
- Lescoat P and Sauvant D 1995. Development of a mechanistic model for rumen digestion validated using the duodenal flux of amino acids. *Reproduction, Nutrition, Development* 35, 45–70.
- Mackie RI and Gilchrist FMC 1979. Changes in lactate-producing and lactate-utilizing bacteria in relation to pH in the rumen of sheep during stepwise adaptation to a high-concentrate diet. *Applied Environmental Microbiology* 38, 422–430.
- MacLeod GK, Colucci PE, Moore AD, Grieve DG and Lewis N 1994. The effects of feeding frequency of concentrates and feeding sequence of hay on eating behaviour, ruminal environment and milk production in dairy cows. *Canadian Journal of Animal Science* 74, 103–113.
- Maekawa M, Beauchemin KA and Christensen DA 2002. Effect of concentrate level and feeding management on chewing activities, saliva production, and ruminal pH of lactating dairy cows. *Journal of Dairy Science* 85, 1165–1175.
- Malestein A, Van't Klooster AJh, Pains RA and Connette GM 1984. Concentrate feeding and ruminal fermentation. 3. Influence of concentrate ingredients on pH, DL-lactic acid concentration in rumen fluid of dairy cows and on dry matter intake. *Netherlands Journal of Agricultural Sciences* 32, 9–16.
- Martin C and Michalet-Doreau B 1995. Variations in mass and enzyme activity of rumen microorganisms: effect of barley and buffer supplements. *Journal of the Science of Food and Agriculture* 67, 407–413.

- Minitab 2000. Minitab 13.20 for Windows. Minitab Inc., State College, PA, USA.
- Mould FL, Orskov ER and Mann SO 1983. Associative effects of mixed feeds. 1. Effects of type and level of supplementation and the influence of rumen pH on cellulolysis in vivo and dry matter digestion of various roughages. *Animal Feed Science and Technology* 10, 15–30.
- Murphy MR 1981. Analyzing and presenting pH data. *Journal of Dairy Science* 65, 161–163.
- Nocek JE 1997. Bovine acidosis: implications on laminitis. *Journal of Dairy Science* 80, 1005–1028.
- Nocek JE, Kautz WP, Leedle JAZ and Allman JG 2002. Ruminal supplementation of direct-fed microbials on diurnal pH variation and in situ digestion in dairy cattle. *Journal of Dairy Science* 85, 429–433.
- Oetzel GR 2003. Introduction to ruminal acidosis in dairy cattle. Preconvention Seminar 7: Dairy Herd Problem Investigation Strategies. American Association of bovine practitioners. 36th Annual Conference, 15–17 September 2003, Columbus, Ohio, USA, 11pp.
- Owens FN, Secrist DS, Hill WJ and Gill DR 1998. Acidosis in cattle: a review. *Journal of Animal Science* 76, 275–286.
- Pereira MN and Armentano LE 2000. Partial replacement of forage with nonforage fiber sources in lactating cow diets. II. Digestion and rumen function. *Journal of Dairy Science* 83, 2876–2887.
- Pitt RE and Pell AN 1997. Modeling ruminal pH fluctuations: interactions between meal frequency and digestion rate. *Journal of Dairy Science* 80, 2429–2441.
- Pitt RE, Van Kessel JS, Fox DG, Pell AN, Barry MC and Van Soest PJ 1996. Prediction of ruminal volatile fatty acids and pH within the net carbohydrate and protein system. *Journal of Animal Science* 74, 226–244.
- Russell JB and Wilson DB 1996. Why are ruminal cellulolytic bacteria unable to digest cellulose at low pH? *Journal of Dairy Science* 79, 1503–1509.
- SAS 1999. SAS/STAT guide for personal computers, version 8.1. SAS Institute, Inc., Cary, NC, USA.
- Sauvant D, Meschy F and Mertens D 1999. Les composantes de l'acidose ruminale et les effets acidogènes des rations. *I.N.R.A. Production Animales* 12, 49–60.
- Shriver BJ, Hoover WH, Sargent JP, Crawford RJ and Thayne WV 1986. Fermentation of a high concentrate diet as affected by ruminal pH and digesta flow. *Journal of Dairy Science* 69, 413–419.
- Slyter IL and Rumsey TS 1991. Effect of coliform bacteria, feed deprivation and pH on ruminal D-lactic acid production by steer or continuous-culture microbial populations changed from forage to concentrates. *Journal of Animal Science* 69, 3055–3066.
- Yang WZ, Beauchemin KA and Rode LM 2000. Effects of barley grain processing on extent of digestion and milk production of lactating cows. *Journal of Dairy Science* 83, 554–568.
- Yang WZ, Beauchemin KA and Rode LM 2001. Effects of grain processing, forage to concentrate ratio, and forage particle size on rumen pH and digestion by dairy cows. *Journal of Dairy Science* 84, 2203–2216.