

**A176 MOLECULAR MECHANISMS OF PERIPHERAL CD25
REGULATORY T CELL (TREG) DEVELOPMENT**

I Prots, A Skapenko, P E Lipsky, H Schulze-Koops *Division of Rheumatology, Med. Poliklinik, University of Munich, Munich, Germany and Autoimmunity Branch, NIAMS, Bethesda, MD, USA*

10.1136/ard.2010.129668m

Compromised development and/or function of CD25 regulatory T cells (Tregs) is of central importance for the development of autoimmune diseases. The mechanisms that regulate the generation of peripheral Tregs are, however, largely unknown. To gain insights into Treg differentiation on a molecular level, the authors investigated gene expression profiles of Tregs at different developmental stages.

Human Tregs were generated in vitro from CD4 CD25-naive T cells in the presence of autologous feeder cells and interleukin (IL)4 as described previously by our group. Total RNA was isolated at the start and from purified CD25⁺ and CD25⁻ T cells at days 3, 5, 7 and 10 of culture and used for DNA microarrays. CD25⁻ T cells generated in the absence of IL4 constituted activated effector T cells (Teffs) lacking regulatory capacity and were therefore used to control for Treg-specific gene expression. Differentially expressed transcripts in Tregs and Teffs versus CD25⁺ T cells were determined at each time point by the microarray data analysis using GeneSpring GX10 software.

A decrease in the number of regulated transcripts was observed in Tregs during the development compared with the control cell population. Analysis of the biological processes and pathways differentially regulated between Tregs and the control population revealed an activated and proliferative state of Tregs at early developmental stages, progressing to a transcriptionally more quiescent phenotype at later time points. A lack of significant overlaps of T cell pathways present in Tregs at late development and in Teffs suggests dichotomy of the two populations, in particular of mature cells. The transcription factor *MAFF*, the chemokine receptor *CCR2*, the enzyme prostaglandin D synthase and the neuromediator *npT1* were identified as Treg-specific genes at different developmental stages. Expression of *npT1* transcript was most pronounced in mature Tregs. Moreover, transcription factors of the FoxO family and FoxM1 were identified as molecular pathways specifying Treg development and activation of Teffs. These data offer new insights into Treg development in the periphery and might provide new targets for therapies of disorders caused by dysregulation of Tregs.

10.1136/ard.2010.129668n



Molecular mechanisms of peripheral CD25 regulatory T cell (Treg) development

I Prots, A Skapenko, P E Lipsky, et al.

Ann Rheum Dis 2010 69: A70

doi: 10.1136/ard.2010.129668m

Updated information and services can be found at:

http://ard.bmj.com/content/69/Suppl_2/A70.1.full.html

These include:

**Email alerting
service**

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>